

This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines


Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + Make non-commercial use of the files We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + Refrain from automated querying Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + Maintain attribution The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + Keep it legal Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at http://books.google.com/



Soc. 1996 d. 322

GenPer. 49

Par. 1996 - d. 590

	- ,			
	_			
	•		_	
		•	•	
		•	•	
		•	•	
		•		
•				
		•		
	•			
			•	
			_	
			•	
·				
			•	
-				
•				
•				

-			


		,	
		,	
			•
	•		
	•		
		1	
	-		
		•	
	•		
•			

			•	•
		•		
•				
	•			
		•		

SMITHSONIAN

MISCELLANEOUS COLLECTIONS.

VOL. XXVII.



"TARES. MEWIN ON VATORITE RESIDENCE AND RESIDENCE SOURCE AND RESIDENCE STREET, MEWIN ON VATORITE RESIDENCE AND RESIDENCE SOURCE SOURCE

WASHINGTON:
PUBLISHED BY THE SMITHSONIAN INSTITUTION.
1888.



ADVERTISEMENT.

The present series, entitled "Smithsonian Miscellaneous Collections," is intended to embrace all the publications issued directly by the Smithsonian Institution in octavo form; those in quarto constituting the "Smithsonian Contributions to Knowledge." The quarto series includes memoirs, embracing the records of extended original investigations and researches, resulting in what are believed to be new truths, and constituting positive additions to the sum of human knowledge. The octavo series is designed to contain reports on the present state of our knowledge of particular branches of science; instructions for collecting and digesting facts and materials for research; lists and synopses of species of the organic and inorganic world; museum catalogues; reports of explorations; aids to bibliographical investigations, etc., generally prepared at the express request of the Institution, and at its expense.

The assignment of a work to one or the other of the two series will sometimes depend upon whether the required illustrations can be presented more conveniently in the quarto or the octavo form.

In the Smithsonian Contributions to Knowledge, as well as in the present series, each article is separately paged and indexed, and the actual date of its publication is that given on its special title page, and not that of the volume in which it is placed. In many cases works have been published, and largely distributed, years before their combination into volumes.

While due care is taken on the part of the Smithsonian Institution to insure a proper standard of excellence in its publications, it will be readily understood that it cannot hold itself responsible for the facts and conclusions of the authors, as it is impossible in most cases to verify their statements.


S. F. BAIRD,

Secretary S. I.



TABLE OF CONTENTS.

- ARTICLE I. (358.) THE CONSTANTS OF NATURE. Part IV. ATOMIC WEIGHT DETERMINATIONS: A DIGEST OF THE INVESTIGATIONS PUBLISHED SINCE 1814. By George F. Becker. 1880. Pp. 152.
 - ARTICLE II. (441.) THE CONSTANTS OF NATURE. Part V. A RECAL-CULATION OF THE ATOMIC WEIGHTS. By FRANK WIGGLES-WORTH CLARKE. 1882. Pp. 293.
 - ARTICLE III. (437.) CHECK LIST OF PUBLICATIONS OF THE SMITHSONIAN INSTITUTION, December, 1881. 1881. Pp. 22.
 - ARTICLE IV. (478.) CATALOGUE OF PUBLICATIONS OF THE SMITHSONIAN INSTITUTION (1846–1882) WITH AN ALPHABETICAL INDEX OF ARTICLES IN THE SMITHSONIAN CONTRIBUTIONS TO KNOWLEDGE, MISCELLANEOUS COLLECTIONS, ANNUAL REPORTS, BULLETINS AND PROCEEDINGS OF THE U.S. NATIONAL MUSEUM, AND REPORT OF THE BUREAU OF ETHNOLOGY. BY WILLIAM J. RHEES. 1882. Pp. 342.



SMITHSONIAN MISCELLANEOUS COLLECTIONS.

358

THE

CONSTANTS OF NATURE.

PART IV.

ATOMIC WEIGHT DETERMINATIONS:

A DIGEST

OF THE INVESTIGATIONS PUBLISHED SINCE 1814.

BY

GEORGE F. BECKER.



WASHINGTON: SMITHSONIAN INSTITUTION. AUGUST, 1880.



,

•

ADVERTISEMENT.

The following forms the fourth part of a general work on the "Constants of Nature," of which the first three are as follows:

- Part I and Supplement.—Specific Gravities, Boiling Points and Melting Points, by F. W. Clarke.
- Part II.—A Table of Specific Heats for Solids and Liquids, by F. W. Clarke.
- Part III.—Tables of Expansion by Heat for Solids and Liquids, by F. W. Clarke.

The manuscript of the present work has been presented to the Smithsonian Institution by Mr. G. F. Becker and is published at the expense of its fund.

S. F. BAIRD, Secretary Smithsonian Institution.

Washington, August, 1880.



PREFACE.

Of the fundamental importance of the most accurate attainable knowledge concerning the true atomic weights of the elements there can be no two opinions. If the enormous mass of known facts relating to the properties of matter is ever to be brought under wide generalizations, it is with the simple substances that a beginning must be made, and with the simplest property of these substances, the relative weights of their ultimate particles. Berzelius held this view and the labors of Mendelejeff, Meyer and others leave no question as to the fact of a relation between the atomic weights and the properties of simple and compound matter. Accurate information on the subject, however, is not easily attainable; different writers on chemistry follow different authorities, and some even take a mean . between the results arrived at by experimenters of different degrees of skill and accuracy, or assume some convenient number without experimental foundation. Nowhere, to my knowledge, is there even an approximately complete list of the determinations that have been made.

Forced back, myself, upon the original memoirs for information, I believed that I should do other chemists a service in presenting to them a short but systematic digest of each investigation on the subject, including the following points, so far as they could be ascertained: The nature of the material experimented upon, and the method of its preparation; the experimental method adopted to effect the determination, and the number of experiments; the mean result reached by the experiments, and the extreme difference between the results; such a record of the constants employed in the calculation as will enable any one to recalculate the results for different constants; and the place in literature where the original paper is to be found.

The following pages are the result. From the information

cases, I think, be able to decide which determination offers the best guarantees for accuracy, or at least between which determinations his choice must lie, forming his judgment to a great extent independently of the comparative reputation of the observers—not always a safe guide where one is, in a general way, the unquestionable superior of the other—and no guide at all when the names carry on the whole an equal weight. As a record of the direction investigations have taken and of analytical methods of the most exact character also, I hope that this digest may not be without value.

As this compilation would serve rather to mislead than to assist investigators, unless it be accurate and practically exhaustive, it seems proper to explain the manner in which it has been prepared. Believing it best to work independently of any previous compilations, I selected as my base the three great German journals—Poggendorff's Annalen, Liebig's Annalen, and Erdmann's Journal für Praktische Chemie. My choice was determined not only by the position these journals take in chemico-physical science, but by the fact that their indices are admirable, and their tone cosmopolitan; all of them, until lately, having furnished their readers with the scientific news of the time, and with abstracts from and translations of the important papers published elsewhere and in whatever language, as well as with original investigations. The indices of these journals I read through from beginning to end, making an extract of every entry which bore on the subject of atomic weights, or which I suspected might do so. In studying the articles thus reached, every reference to other atomic weight determinations was preserved, and the originals, so far as possible, sought out; a task in which the Royal Society's Catalogue of Scientific Papers was of the greatest assistance. Having exhausted the supply of information in these journals, I turned to Berzelius' Jahresbericht, and to its continuation edited by Kopp, Liebig et al., and made a study of their contents by the same method. Later, I made a similar systematic study of the Annales de Chimie et de Physique, the Bericht der Deutschen Chemischen Gesellschaft, the Chemical PREFACE. 3

News, Fresenius' Zeitschrift für Analytische Chemie, the Journal of the Chemical Society, the Proceedings of the Royal Society, and the Philosophical Transactions, and of Silliman's American Journal of Science. I have also made some use of the Philosophical Magazine, and a great deal of use of the Paris Comptes Rendus. These publications are not so indexed as to make their contents readily available; but what appears in the Comptes Rendus is pretty sure to be noticed elsewhere, and I scarcely think that any determinations there published have escaped me. I have also made use of the Bibliothèque Universelle, Archives des Sciences of Geneva, (an incomplete set, unfortunately,) the Zeitschrift für Berg-Hütten-und Salinen-Wesen im Preussischen Staate, Thomson's Annals of Philosophy, Gilbert's Annalen der Physik und der Physikalischen Chemie, the British Association Reports, the Transactions of the Royal Society of Edinburgh, the Transactions of the Academies of Brussels and of St. Petersburg, and have consulted numerous works on chemistry, particularly Berzelius' Lehrbuch der Chemie and Gmelin-Kraut's Handbuch der Chemie.

I have not thought it necessary, or even desirable, to extend my search for atomic weight determinations further back than Wollaston's famous "Table of Equivalents," published in the Philosophical Transactions for 1814. It is true that numerous determinations had been made before that time, but, with the exception of those mentioned by Wollaston, few which can be of either interest or value to the chemist of the present day, except from a purely historical point of view. From Wollaston's table onwards, I have not felt that the purposes of this paper permitted of any selection between atomic weight determinations, however valueless many of them might appear to my own judgment. Indeed, it has cost me more labor to put many ill-made and ill-reported investigations into proper form for this digest than was required for a majority of those determinations upon which I set the highest value. In the attempt to make a complete collection of the determinations since the time indicated, a few may have escaped my search; but, if so, they must have fallen singularly dead upon the chemical world, and would be unlikely to repay further labor in seeking them. On the other hand, I have rigidly excluded atomic weights calculated from analyses never designed so to be used. Any chemist, upon whose experiments we could rely, would proceed in a very different manner in making an atomic weight determination, from that which he would select for an ordinary analysis, and to put his credit at stake by calculating atomic weights from analyses not designed for this use is alike unfair to him and to the scientific public, which is asked to receive as an atomic weight determination what really is not such.

The purpose of this paper is distinctly not critical, and the remarks I have added to, or inserted in, the digest are simply explanatory. I have, however, frequently mentioned criticisms which have appeared in literature when they seemed pertinent.

As for the accuracy with which the digests have been made, I may state that the preponderating importance of this point has been constantly before my mind. In the effort to crowd the maximum amount of information into the fewest words, I have had occasion to refer to most of the papers digested a number of times, and at long intervals. I have always taken advantage of such occasions, as well as those on which I have met with a reprint, translation or abstract of a determination, to verify the rough draughts of my digests. Only in a couple of instances have I thus discovered a trifling error. On the other hand, I have been able to detect and point out numerous misprints and miscalculations in the original sources. While, therefore, I cannot hope entirely to have escaped error in the thousands of values I have copied, and the almost equal number of calculations I have made, I have strong hopes that the accuracy of this digest will be found at least on a par with that of the original papers.

When, as is the case with provoking frequency, chemists have given their analytical data, but have omitted to state the atomic weights, or other constants, assumed in calculating their results, I have recalculated their data with accepted constants, which I have in each case stated. I have also, in many instances, recalculated determinations of importance,

PREFACE. 5

in which constants varying considerably from those now received were assumed. I have further reduced the determinations originally given in terms of O = 100, or of O = 15.96, to O = 16. No confusion, however, will be found between the numbers for which the original investigators are responsible and my own. All values which I have calculated are in italics, or, with my explanations, enclosed in square brackets. The only arithmetical operation I have permitted myself to perform without these indications is a multiplication or division by two; and even in such cases it will usually appear from the digest itself that this operation has been performed.

The abbreviations of the literary references are essentially those adopted in the Royal Society's Catalogue of Scientific Papers. The first reference in each case is to the source upon which I have depended. When two references are necessary, they are connected by the word and. When my authority is not the original source, that to which it is accredited in my authority is also mentioned.

In conclusion, I shall be grateful to any one who, by drawing my attention to omissions or mistakes, will assist me in perfecting a labor which has occupied all my available time for twenty months.

BERKELEY, CAL., April, 1878.

Postscript.

In preparing the following paper, I designed making it preliminary to a discussion of the various determinations and of the value to be assigned to each, and in this work I had already made some progress. After presenting this paper to the Institution, however, I learned that Prof. F. W. Clarke had been for some time engaged on a similar undertaking, and to him I gladly resigned the discussion of the data here compiled. The two papers will appear in the same form, and may be regarded as complementary.

G. F. B.



ATOMIC WEIGHT DETERMINATIONS.

ALUMINIUM.

The specific heat of aluminium, as determined by Regnault and by Kopp, and the vapor density of volatile compounds, as determined by Deville and Troost and by Odling, indicate that the atomic weight of this element is about 27.5. (Gmelin-Kraut, Handbuch der Chemie, 1, 39; and L. Meyer, Moderne Theorien der Chemie, 50.)

J. J. Berzelius: 27.267 (O = 16).

100 parts of anhydrous aluminic sulphate decomposed by heat, gave 29.934 parts of oxide. Preparation not described. Number of experiments, probably 1. In Berzelius' Lehrbuch these data are calculated for S=200.75, and give Al=170.9 (O=100,) or 27.344 (O=16.) [If S=32, the data give Al=27.267.] (Poggend. Ann., 8, 1826, 187.*)

T. Thomson: $30 \text{ (O} = 16).\dagger$

.Thomson found, probably from analysis of the sulphate, (see appendix,) that 125 Al = 100 O. Thomson supposed aluminic oxide to be a protoxide. [If it is a sesqui-oxide, the data give Al at 30.] (Thomson's System of Chemistry, 7th ed., 1, 1831, 454.)

W. W. MATHER: 20.55 (O = 16).

According to this chemist 0.646 grammes of chloride, prepared according to Woehler, gave 2.055 grammes argen-

^{*}This article by Berzelius, which contains the particulars of a large part of his earlier atomic weight determinations, will be referred to frequently in the course of this paper. It is unfortunately full of misprints, all of which are, by no means, corrected in the table of errata at the end of the volume. The correctly printed values of the atomic weights discussed in it are to be found in *Poggend. Ann.*, 10, 1827, 389.

[†] It must be remarked, in justice to Dr. Thomson, that his atomic weight determinations are, properly speaking, of a different nature from those of other chemists. So thoroughly persuaded was he of the truth of Prout's hypothesis, (that the atomic weights of the elements are all exact multiples of that of hydrogen,) that his experiments were directed merely towards accertaining which multiple, in any case, was to be adopted.

tic chloride, and 0.2975 aluminic oxide. (Silliman's Amer. Journ., 27 1835, 138, 241.) Berzelius points out the inconsistency of these data. (Berzelius' Jahresbericht, 15, 1835, 138.)

C. Tissier: 27.12 (O = 16).

Determined by dissolving aluminium in chlorhydric acid, evaporating to dryness with excess of nitric acid and decomposing the nitrate by heat. The aluminium employed contained 0.135 per cent. sodium. 1.935 of this metal gave 3.645 oxide. [If Na = 23, these data give Al = 27.12.] The metal was prepared by heating aluminic fluoride with purified sodium in a graphite crucible. (Paris Comptes Rend., 46, 1858, 1105.)

J. Dumas: 27.446 (O = 16).

Determined by six experiments on the titration of aluminic chloride with argentic nitrate. The mean result was $Al = 13.723 \ (0=8)$; extreme difference 0.09. The aluminic chloride, which had been prepared on a large scale, was purified by sublimation over iron-filings and over aluminium filings, and by a third sublimation in a current of hydrogen over aluminium filings, after which it was melted. Experiments on the oxidation of aluminium were found unsatisfactory on account of the difficulty of obtaining the metal pure. They gave Al at from 13.74 to 13.89. Dumas takes Ag = 108; Cl = 35.5. (Ann. de Chim. et de Phys., (3,) 55, 1859, 151.)

W. Odling: 27.5 (O = 16).

Determined from the vapor density of aluminium methide and ethide at 220° and upwards. (*Phil. Mag.*, (4,) 29, 1865, 316.)

—. Isnard:
$$27 (O = 16)$$
.

Pure aluminium dissolved in chlorhydric acid, evaporated and heated to redness, gives \$\forall \text{ of its weight in oxide. (Paris Comptes Rend., 66, 1868, 508.)

Pelouze and Fremy give 27.357 (O = 16); 170.98 (O = 100,) for the atomic weight of aluminium, and assert that this value is derived from the composition of potash-alum, but they give no authority for the value. The experiments were made by precipitation with barium chloride. (Traité de Chimie, 3d ed., 1, 50.)

ANTIMONY.

From the specific heat of antimony, as determined by Bunsen, Regnault, and others, and from the vapor density of volatile compounds, as determined by Mitscherlich, Loewig and Schweizer and others, it is certain that the atomic weight must be about 120. (Gmelin-Kraut, l. c.; and L. Meyer, l. c.)

J. J. Berzelius: 129.03 (O = 16); 806.452 (O = 100).

100 parts of pure antimony, oxidized with nitric acid, evaporated to dryness, and heated to redness, gave 124.8 antimonic antimoniate. The number of experiments and the preparation of the metal are not given. (*Poggend. Ann.*, 8, 1826, 23.)

R. Schneider: 120.3 (O = 16); 751.9 (O = 100).

Determined by experiments on the reduction of native antimonic ter-sulphide in a current of hydrogen. The only foreign substance to be found in the mineral was silicic acid, which was determined in each case. The temperature was kept as low as possible, and the amount of sulphide volatilized, and of that undecomposed by the process, was determined. The mean composition, as ascertained by eight experiments, was 71.48 antimony—extreme difference, 0.078; and 28.52 per cent. sulphur. The atomic weight was calculated from the mean for S = 200. (Poggend. Ann., 98, 1856, 293.) Schneider published a preliminary note in Poggend. Ann., 97, 1856, 483, in which, from a portion of the above-mentioned experiments, he deduced the value 120.25.

H. Rose and Weber: 120.626 (O = 16).

Rose published this determination expressly as a confirmation of Schneider's value. Antimony ter-chloride was dissolved in water containing tartaric acid, and decomposed by hydrogen sulphide. Sulphur was removed from the filtrate by ferric sulphate, and the chlorine determined with argentic nitrate. 2.162 antimony chloride were found equivalent to 4.097 argentic chloride. [If Ag = 107.93 and Cl = 35.457, these data give Sb = 120.626; or, for O = 100, Sb = 753.92.] Rose, adopting some other values gets 1508.67 [twice 754.34.] He also recalculates some earlier

analyses of the ter-chloride, and the penta-chloride (*Poggend. Ann.*, 3, 1825, 443) made by himself by the same method, which give respectively 1512.91 and 1508.6. (*Poggend. Ann.*, 98, 1856, 455.)

W. P. Dexter: 122.336 (O = 16); 764.6 (O = 100).

Attempts were made to determine the atomic weight of antimony from its reducing action on the chloride of gold, but no constant result was obtained. Berzelius' method (vide supra) was, therefore, adopted. From the mean of ten irreproachable experiments Dexter deduces the value 1529.2; extreme difference, 3. The metal was prepared as follows: From antimony tartrate, sodium metantimonate was prepared, and antimonic acid separated out with nitric acid. The antimonic acid was reduced with carbon, and melted with another portion of antimonic acid to remove traces of sodium, etc. It was also heated in a current of hydrogen to remove traces of oxide. The investigation was carried out in Bunsen's laboratory, and with his assistance. (Poggend. Ann., 100, 1857, 563.)

J. Dumas: 122 (O = 16).

Neither the reduction of cervantite nor of the sulphide, nor the oxidation of metallic antimony gave accordant results. Dumas, therefore, resorted to the analysis of the ter-chloride with argentic nitrate. The chloride was prepared by three different methods, and was dissolved in water acidulated with tartaric acid. Seven experiments gave an average of 121.975; extreme difference, 0.69. Ag = 108; Cl = 35.5. (Ann. de Chim. et de Phys., (3,) 1859, 175.)

F. Kessler: 122.24 (O = 16).

In four experiments crystals of antimony ter-oxide were employed. This oxide had been sublimed in a current of pure, dry carbonic acid. A known weight of the compound was nearly oxidized in a chlorhydric acid solution by a known, slightly insufficient, weight of potassic chlorate. The remainder was titrated with a standard solution of potassic bi-chromate, and countertitrated with ferrous chloride. The mean result was Sb = 122.16. In three experiments metallic antimony was employed. It was prepared by reducing the precipitate formed when ammonic hydrate is added to stibium-ammonium tartrate. The metal was oxidized in chlorhydric acid solution by potassic chlorate, (not weighed,) and reduced to antimony ter-chloride by

stannous chloride. The excess of this reagent was chloridized by mercuric chloride, calomel being separated by filtration. The experiment was continued exactly as in the cases where the oxide was taken to start with. The mean of the experiments on metallic antimony was 122.34. The mean of the seven experiments above described is 122.24; extreme difference, 0.94. K = 39.12; Cl = 107.97. Kessler also made experiments by Rose's method, but got discordant results. (Poggend. Ann., 113, 1861, 145.)

B. Unger: 119.76 (O = 16).

Determined by analysis of sodium sulph-antimonate, (Schlippe's Salt.) (Kopp's Jarresbericht, 1871, 325; Arch. der Pharm., (2,) 147, 193; 148, 1.) A single determination by a method from which great accuracy could not be expected. S = 32; Na = 23. (J. P. Cooke, Jr., in Proc. Amer. Acad., 13, 6.)

J. P. COOKE Jr.: 120 (O = 16).

Cooke objects to the determinations of Dexter and Dumas, on the ground that there is no sufficient evidence of the absence of higher or lower compounds of the same elements in the salts employed.

In two experiments antimony was dissolved and precipitated as sulphide, which was heated to 240° before weighing. The formation of free S was prevented, occluded tartaric acid was determined, but occluded oxy-chloride was neglected. The experiments gave each Sb = 120.6 for S= 32. In thirteen experiments Sb was dissolved in a minimum of nitric acid, and the solution boiled over bullets of Sb to complete saturation. The sulphide was then precipitated in an atmosphere of carbon di-oxide. The precipitate contained no free S. The oxy-chloride was driven off at 180° and determined. The tartaric acid was decomposed at 210° and determined. The errors are opposed and min-The mean of the weighings of sulphide, dried at 180°, gave Sb = 119.994 for S = 32; extreme difference, 1.01. The mean of weighings of sulphide heated to 210° gave Sb = 120.295; extreme difference, 1.07. General mean Sb = 120.145. Fifteen analyses of antimonious bromide gave the Br contents at 66.6665 per cent. for Ag = 108, Br = 80, with an extreme difference of 0.195. This composition gives Sb In seven experiments the iodide was analyzed. I = 127 and Ag = 108, it gave a mean of 76.051 per cent. Sb, or Sb = 120. It was also shown that the chloride cannot be prepared free from oxy-chloride, and that its Sb and Cl contents correspond to Sb = 120. Metallic Sb was prepared by reduction of sodic antimoniate, or of oxide, with potassic cyanide, or by Liebig's method. In all cases it was fused for several hours under its own oxide. The haloid salts were purified by fractional recrystallization and distillation, in part in a current of carbon di-oxide. (*Proc. Am. Acad.*, 13, 1877, 1.)

ARSENIC.

The specific heat of metallic arsenic, as determined by Regnault, and the vapor density of a number of volatile compounds, as determined by Dumas, Mitscherlich, Bunsen, and others, prove that the atomic weight of this element must be in the neighborhood of 75. (Gmelin-Kraut, l. c.; and L. Meyer, l. c.)

J. J. Berzelius: 75.1 (O = 16); 469.4 (O = 100).

2.203 grammes of arsenious acid, heated with sulphur in a distilling apparatus in such a manner that sulphurous acid, but no sulphur, could escape, set free 1.069 grammes sulphurous acid. If S = 200.75, the value follows. (Poggend. Ann., 8, 1826, 22; and Lehrbuch, 5 ed., 3, 1205.)

J. Dumas: 75 (O = 16).

Dumas found the vapor density of arsine 2.695. [This value multiplied by 28.94278 gives As = (sensibly) 75.] (Ann. de Chim. et de Phys., 33, 1826, 337.)

J. Pelouze: 75 (O = 16); 468.75 (O = 100).

A known weight of arsenic ter-chloride was introduced into a nitric acid solution of a known weight of perfectly pure silver, the chloride being in slight excess. The excess of chloride was then titrated with decimal silver solution.* As the mean of three experiments Pelouze found As = 937.50; extreme difference, 0.8. Ag = 1349.01; Cl = 443.2. The ter-chloride was repeatedly distilled to free it from excess of chlorine. It was colorless, dissolved com-

^{*}This method, which has been frequently employed in the determination of atomic weights, will be referred to as "Pelouze's method."

13

pletely in chlorine, and boiled between 184° and 135°. (Paris Comptes Rend., 20, 1845, 1047.)

J. Dumas: 74.94 (0 = 16).

Determined by four experiments on the titration of arsenic ter-chloride with argentic nitrate, the ter-chloride being prepared in several lots. The number is the mean of the experiments; the extreme difference being 0.15. Dumas takes Ag = 108; Cl = 35.5. (Annal. de Chimie et de Physique, (3,) 55, 1859, 174.)

F. Kessler: 75.2 (O = 16).

In six experiments arsenious acid was titrated with potassic bichromate and counter-titrated with ferrous chloride. The number so obtained was 75.15. In twelve experiments a known weight of arsenious acid was oxidized in caustic potash solution by potassic chlorate, the arsenious acid being slightly in excess, acidified with chlorhydric acid and the excess of arsenious acid titrated with potassic bichromate and counter-titrated with ferrous chloride. The oxidizing action of the potassic bichromate was experimentally The number obtained from these experiments determined. was 75.24. Five experiments were made with acid instead of alkaline solutions of arsenious acid; they gave 75.15. The arsenious acid was colorless, transparent, volatilized without any residue, and was thoroughly dessicated. Kessler assumed K = 39.12; Cl = 107.97. (Poggend. Annal., 95, 1855, 210; 113, 1861, 140.)

BARIUM.

The specific heat of barium compounds, especially of the chloride, as determined by Regnault and by Kopp, shows that the atomic weight of this element lies in the neighborhood of 137. (Gmelin-Kraut, l. c.)

Wollaston and Klaproth. 139.2 (O = 16); 870 (O = 100).

Klaproth found that 100 parts of carbon di-oxide were equivalent to 352.57 parts barium oxide, and that 34 parts sulphuric anhydride were equivalent to 66 parts of barium

oxide. If C = 75.4, and S = 200, the value follows. (*Phil. Trans.*, 104, 1814, 20.)

J. J. Berzelius: 136.79 (O = 16).

100 parts of barium chloride gave 138.08 and 138.06 parts argentic chloride. [If Ag = 107.93, and Cl = 35.457, the above value follows.] Berzelius also determined barium from the sulphate; 100 parts barium chloride gave 112.17 and 112.18 parts sulphate. Calculated for S = 200.75 this determination is almost identical with the other; Berzelius, however, expressly adopts the former. [Calculated for S = 32.0742, it gives 135.74.] (Poggend. Annal., 8, 1826, 189, and Lehrbuch der Chemie, 5th ed., 3, 1229.)

E. Turner: 137.4 (O = 16).

Turner determined the chlorine contents of barium chloride at 34.016 per cent. by precipitation with silver. This number was the mean of the best two experiments made, and the value follows from it on the assumption that Cl = 35.42. The barium chloride was prepared from native carbonate by solution in chlorhydric acid, precipitation of impurities with barium oxide, ignition of the chloride, treatment with alcohol, and recrystallization. (*Phil. Trans.*, 119, 1829, 291.*)

T. Thomson: 136 (O = 16); 850 (O = 100).

Thomson had formerly determined this atomic weight at 875 by mixing potassic sulphate with barium chloride in such proportions that the supernatant liquid contained no sensible amount of either sulphuric acid or barium. Turner having shown the fallacy of this method, Thomson substituted ammonium sulphate, and also sulphuric acid for the potassium salt, and found 9.5006 barium oxide equivalent to 5.00 sulphuric anhydride. He also analyzed the chloride with argentic nitrate, assuming silver = 1375, and chlorine = 450, and reached the same conclusion with reference to barium. (Thomson's System of Chemistry, 7th ed., 1, 1831, 426.)

^{*}Turner made the discovery in the course of this investigation that barium sulphate carries down other salts, such as potassic sulphate, which cannot be extracted from the precipitate by any degree of washing, and that determinations, with barium sulphate, are consequently unreliable. Although Berzelius drew attention to the importance of the observation, and Thomson was obliged to acknowledge errors in his work from this cause, the fact was for a long time nearly forgotten, as can readily be proved from the contents of this digest.

—. SALVETAT: 136 (O = 16); 850 (O = 100).

Determined from the loss of weight ensuing on the decomposition of barium carbonate by sulphuric acid. Details not given. (Paris Comptes Rendus, 17, 1843, 318.)

J. Pelouze: 137.28 (O = 16); 858.01 (O = 100).

Into a nitric acid solution of a known weight of perfectly pure silver, a known and slightly more than equivalent weight of barium chloride was introduced. The excess was titrated with decimal silver solution. The value is the mean result of three experiments, which give an extreme difference of 0.22 for O=100. The barium chloride was purified by recrystallizations continued till determinations gave a constant result, and was dessicated in part at 200° , and in part at a temperature just below redness. Pelouze took Ag=1349.01, and Cl=443.2. (Paris Comptes Rendus, 20, 1845, 1047.)

C. Marignac: 137.08 (O = 16); 856.77 (O = 100).

Determined by six experiments on the equivalence of silver and barium chloride performed by Pelouze's method, (vide supra.) 100 silver were found equivalent to 96.365 barium chloride; extreme difference, 0.038; hence the value taken. Marignac takes Ag = 1349.01, and Cl = 443.2. The barium chloride was purified as follows: Commercial chloride was crystallized from boiling aqueous solution; the crystals were heated to redness, dissolved in boiling water, treated with carbon di-oxide, filtered and crystallized, and these crystals were washed with alcohol and again recrystallized. Determinations were made at each stage and the purification was continued until constant results were obtained. (Liebig, Annal., 68, 1848, 214; Bibl. Univ., Arch. des Sciences, 8, 265.)

H. STRUVE: 136.26 (O = 16).

100 parts of barium chloride gave 112.0938 parts of sulphate as a mean of two experiments; extreme difference, 0.005. 8 = 32; Cl = 35.4624. (Liebig, Annal., 80, 1851, 204; Oefversigt af Kongl. vet. Acad. Foehr., 6, 165.)

T. Andrews: 137.578 (O = 16).

Andrews obtained this number from two nearly coincident experiments of which he gives no details. (Brit. Assoc. Rep., 1852, pt. 2, 33.)

C. Marignac: 137.16 (O = 16).

Three experiments were made on the titration of air-dried barium chloride in crystals by Pelouze's method, (vide supra.) Five grammes of the salt required for precipitation (1) 4.4205; (2) 4.4195; (3) 4.4210 grammes silver. Three experiments were made on the conversion of the same barium chloride into sulphate. Ten grammes of the salt gave (1) 9.543; (2) 9.544; (3) 9.542 grammes sulphate. In each of the latter experiments the water was determined, and was found to vary no more than 0.0005 grammes. Comparison of the two series gives for Ag = 108, S = 16, and O =8; barium equal to (1) 68.57; (2) 68.61; (3) 68.55; in mean 68.58, or one-half of 137.16. This result is independent of the possible trace of water the chloride might have contained. In another series of three experiments the water was driven off at a low red heat and determined, and the salt analyzed by Pelouze's method. It was proved that barium chloride is not decomposed at the temperature employed. (1) gave 68.61; (2) 58.59; and (3) 68.55, or a mean of 68.583. The salt for the experiments marked (1) was prepared by recrystallization and precipitation with alcohol; that for (2) by a repetition of the same process, and for (3) by resolution of (2) and precipitation with chlorhydric acid gas. Marignac proved that the precipitated argentic chloride contained entirely insignificant traces of barium salt. Cl = 35.5. (Bibl. Univ., Archives des Sciences, Nouv. Série., 1, 1858, 209.)

J. Dumas: 137 (O = 16).

Determined by fifteen experiments on the titration of barium chloride with argentic nitrate, which give a general average of 68.516 with an extreme difference of 0.11. The barium chloride was prepared from pure nitrate and pure carbonate, and from commercially pure chloride after it had been freed from lead by precipitation with barium sulphide. The chloride was precipitated from solution by chlorhydric acid gas and melted in a current of chlorine to prevent oxidation. Ag = 108; Cl = 35.5. (Annales de Chimie et de Physique, (3,) 55, 1859, 137.)

BERYLLIUM.

The atomic heat of beryllium has been determined by J. Emerson-Reynolds by direct comparison with that of silver.

In a calorimetric apparatus constructed for the purpose, the amount of heat given off during cooling by 108 parts of silver heated to 100° was found to be equal to that communicated by a little more than 9.2 parts of beryllium under the same conditions. Assuming the atomic weight of the metal to be 9.2, the atomic heat found would be 5.91. The smallness of this number the observer accounts for by supposing that there was a trace of platinum present introduced by the use of platinum vessels in the course of reduction. (*Phil. Mag.*, (5,) 3, 1877, 38.)

J. J. Berzelius: 14.5 (O = 16).

Berzelius analysed the salt formed by saturating dilute sulphuric acid with beryllium oxide. From the amount of barium sulphate obtained he inferred that the atomic weight of beryllium was 331.261 on the supposition that the oxide was Be, + O₈ and that the salt was neutral. Berzelius took O = 100; S = 200.75, and Ba = 855.29. [Awdejew having discovered that this salt is basic, this value is reduced to 90.63; or, for O = 16, to 14.5.] Berzelius accepted Awdejew's determination in preference to his own. (Poggend. Annal., 8, 1826, 187; and Lehrbuch der Chemie, 5th ed., 3, 1225.)

T. Thomson: 36 (O = 16).

Experiments not given. The value is four times nine, and may have arisen from a mistake as to saturation. (System of Chem. 7 ed., 1, 1831, 459.)

—. Awdejew: 13.85 (O = 16); 86.58 (O = 100).

Beryllium sulphate, in chlorhydric acid solution, was decomposed with barium chloride. In the filtrate the excess of barium chloride was precipitated with sulphuric acid, and the beryllium oxide thrown down with ammonia, dried, heated, and weighed. The beryllium sulphate was prepared from pure carbonate by treatment with sulphuric acid and precipitation with alcohol. It was purified by recrystallization. Four experiments were made, the mean of which calculated for S = 201.165, gave Be = 58.084 with an extreme difference of 1.955. (Poggend. Annal., 56, 1842, 106.) Weeren recalculated these analyses for S = 200 and got 57.72, [or $\frac{2}{3}$ of 86.58.] (Poggend. Annal., 92, 1824, 124.)

J. Weeren: 13.83 (O = 16); 86.46 (O = 100).

Weeren followed the same method as Awdejew, except that he precipitated the beryllium with ammonium sulphide, the oxide being soluble in excess of ammonia. The mean of four experiments gave 57.64, the extreme difference being 1.52 for O = 100, [57.64 is $\frac{3}{4}$ of 86.46.] Weeren took S = 200. (Poggenii. Annal., 92, 1854, 124.)

G. Klatzo: 13.89 (O = 16).

Klatzo made five analyses of the sulphates containing seven and four molecules of water, precipitating the sulphuric acid as barium sulphate, and the beryllium as oxide by means of ammonia. From a comparison of the sum of the oxide found in all the analyses with the total amount of barium sulphate found, Klatzo deduces Be = 9.227, for Ba = 137, and S = 32. [If Ba is taken equal to 137.16, and S = 32.07, and if each of the analyses is calculated for itself, Be = 13.89. The extreme difference is 0.45.] The sulphates were purified by recrystallization, and treatment with alcohol. (Erdmann's Jouin. für Prak. Chemie, 106, 1868, 227; Klatzo, Ueber die Constitution der Beryllerde, Dorpat, 1868.)

L. F. Nilson and O. Pettersson have redetermined the specific heat of beryllium within a few weeks. They find the specific heat 0.4079, corresponding to a trivalent metal and a sesqui-oxide. The investigation seems to have been made with great care, while that of Emerson-Reynolds was merely preliminary. (Berlin, Bericht der chem. Ges., 11, 1878, 386.)

BISMUTH.

Dulong and Petit, Regnault, and Kopp, have determined the specific heat of Bismuth. It corresponds to an atomic weight of about 210. (Gmelin-Kraut, l. c.)

P. LAGERHJELM: 212.86 (O = 16); 1330.377 (O = 100).

Metallic Bismuth was oxidized in a weighed vessel by nitric acid, and the nitric acid expelled by heat. 10 grammes of bismuth gave 11.1275 oxide. (Berzelius' Lehrbuch der Chemie, 5th ed., 3, 1216; Stockholm, Akad. Handl., 34, 1813, 219.)

19

R. Schneider: 208 (O = 16); 1299.98 (O = 100).

Determined by eight experiments on the conversion of metallic bismuth into oxide by solution in nitric acid and decomposition of the nitrate in the same vessel. The escaping gases were led through nitric acid, and the bismuth caught in this way was separately converted into oxide and weighed. In four experiments the bismuth was prepared by the reduction of basic nitrate, and for the other four by the reduction in hydrogen of the oxide formed in those which preceded. 100 bismuth oxide were found to contain a mean of 89.655 metal; extreme difference, 0.048. (Poggend. Annal., 82, 1851, 303.)

J. Dumas: 210.44 (O = 16).

Determined by seven experiments on bismuth chloride, which was decomposed in solution by sodium carbonate, and the sodium chloride thus formed titrated with silver solution. The value taken is the mean result. The extreme difference is 1.12. Dumas takes Ag = 108, and Cl = 35.5. The bismuth chloride was prepared by the action of chlorine on bismuth, and was purified by fractional distillation over bismuth. That employed in the experiments was colorless. (Annal. de Chimie et de Physique, (3,) 55, 1859, 176.)

BORON.

The specific gravities of a number of volatile compounds of boron have been determined by Dumas, Woehler and Deville, and others, and correspond to an atomic weight of about 11. (Gmelin-Kraut, l. c.; L. Meyer, l. c.)

H. F. Weber has discovered that the specific heat of boron rises rapidly with the temperature, becoming nearly constant at 600°. Above this temperature its specific heat is 0.5, and its atomic heat 5.5. (*Poggend. Annal.*, 154, 1875, 575.)

J. J. Berzelius: 11.01 (O = 16).

Davy's investigations having shown that boracic acid contains about 68 per cent. oxygen, and having thus established the formula of borax, Berzelius determined the atomic weight from the water contents of that salt. He found in three experiments, without variation, 47.1 per cent. Gmelin-Kraut recalculates this composition with Stas' atomic

weights, and gets the value given. (Poggend. Annal., 8, 1826, 19.)

A. LAURENT: 10.86 (O = 16).

Laurent found that borax retains some water even when melted, which, however, can be expelled by the addition of iceland spar. By repeating Berzelius' experiments, and adding a known weight of spar, he found the water contents in two experiments 47.15 and 47.20. He did not regard the experiments as accurate. Gmelin-Kraut recalculates these data with Stas' atomic weights, and gets B = 10.91 and 10.81. (Paris Comptes Rendus, 29, 1849, 5.)

Worhler and Deville: 10.87 (O = 16).

These chemists titrated the bromide and the chloride of boron with argentic nitrate. They do not offer the analyses as atomic weight determinations, but Dumas applies the data to this object. Taking Ag = 108, and Cl = 35.5, Dumas calculates from the analysis of the chloride prepared by the action of H Cl on B, B = 11; from the analysis of the chloride prepared by the action of Cl on B, B = 10.6; from the analysis of the bromide prepared by the action of bromine on boron, B = 11. (Annal. de Chimie et de Physique, (3,) 52, 1858, 88; 65, 1859, 129.)

T. Thomson: 10.67 (O = 16).

Thomson supposed boracic acid to be composed of one atom of boron and two of oxygen, and concluded from Davy's and his own experiments that the atom of B was exactly equal to that of O. For the correct composition of the acid his value 'must be reduced one-third. (System of Chem., 7th ed., 1, 1831, 214.)

BROMINE.

Mitscherlich determined the vapor density of bromine, and Regnault the specific heat in a solid condition at very low temperatures. Both of these constants correspond to an atomic weight of 80. (Gmelin-Kraut, l. c.; L. Meyer, l. c.)

A. J. Balard: 75 (O = 16); 468.85 (O = 100).

1.27 potassium bromide decomposed with sulphuric acid

gave a residue of 0.973 potassic sulphate. [If this analysis is calculated with Stas' atomic weights, it gives Br = 74.65.] In another experiment 100 parts of argentic bromide reduced with zinc, the excess of which was extracted with sulphuric acid, gave 58.9 parts silver. [Calculated with Stas' data this gives Br = 75.3.] Balard mentions no special precautions in the preparation of his salts for this determination. (Annal. de Chimie et de Physique, 32, 1826, 357, 362.)

J. Liebig: 75.29 (O = 16); 470.55 (O = 100).

2.521 potassic bromide precipitated with argentic nitrate gave 4.041 argentic bromide. The potassic bromide was obtained by adding potassic hydrate to an alcoholic solution of bromine until the solution began to lose color. (Annal. de Chimie et de Physique, 33, 1826, 331.)

J. J. Berzelius: 78.264 (O = 16); 489.15 (O = 100).

Berzelius suspected that insufficient precautions had been taken in the preceding determinations to get rid of chlorine. He washed bromine for a long time, and converted it into zinc bromide and ammonium bromide. These salts he partially precipitated with argentic nitrate to get rid of chlorine. From the filtrate he precipitated argentic bromide which he washed, dried, and melted. 7.202 of this bromide, decomposed in a current of chlorine, yielded 5.546 argentic chloride; 7.8805 bromide gave 6.069 chloride. If Ag = 1351.607, and Cl = 442.652, the mean value of Br is as above; difference, 0.09. (Poggend. Annal., 14, 1828, 565; Kongl. vet. Akad. Handl., 1828.)

C. Loewig: 75.76 (O = 16).

According to Gmelin-Kraut, Handbuch der Chemie, the determination was published in a treatise entitled Brom und Seine Chemische Verhältnisse, Heidelberg, 1829.

C. Marignac: 79.957 (O = 16).

In three experiments a known weight of silver was dissolved in nitric acid, precipitated with potassium bromide, and the argentum bromide dried at 200° and weighed. [For Ag = 107.93 these experiments give Br = 79.938, with an extreme difference of 0.018.] In vacuo this result is, according to Stas, 79.968. In seven experiments a known weight of silver was precipitated by a determinate amount of potassic bromide by titration. [If K = 39.137, and Ag = 107.93,

this gives bromine = 79.924 with an extreme difference of 0.046.] In vacuo this becomes, according to Stas, 79.945. In four experiments potassium bromate was decomposed by heat, and the potassic bromide weighed. [For K = 39.137 these experiments give bromine at 80.11 with an extreme difference of 0.56. These latter are evidently much less accurate than the preceding, and I have therefore averaged the first and second series in vacuo.] The KBr was prepared by heating bromate purified by recrystallization. (Berzelius' Lehrbuch der Chemie, 5th ed., 3, 1194; Bibl. Univ., 46, 1843, 357.)

W. Wallace: 79.74 (O = 16).

Determined by analysis of arsenic ter-bromide, by titration with argentic nitrate, according to the method of Pelouze, (see arsenic, Pelouze's determination.) Three experiments were made, giving a mean of 79.738; extreme difference, 0.051. As = 75; Ag = 107.97. The arsenic and bromine were directly combined, and the compound was purified by fractional distillation and recrystallization. (Phil. Mag., (4,) 18, 1859, 279.)

J. Dumas: 80 (O = 16).

Determined by three experiments on the conversion of argentum bromide into chloride in a current of dry chlorine. The mean is 80.03; the extreme difference is 0.18. Silver is taken at 108, and chlorine at 35.5. The argentum bromide was prepared with bromine free from iodine, and was purified from chlorine by digestion with argentum bromide. (Annal. de Chemie et de Physique, (3,) 55, 1859, 162.)

J. S. Stas: 79.952 (O = 16).

Four complete syntheses (the weight of each of the constituents, and of the compound being determined) were made of argentum bromide, a known weight of silver being converted into sulphate, and precipitated with a known weight of bromine which had been converted into hydrobromic acid. The mean result was that 100 Ag = 74.0805 Br; with an extreme difference of 0.004. Two analyses of argentic bromate, made by reducing the salt in suspension with sulphurous acid, gave for the molecular weight of the bromide 187.84, and 187.90, mean 187.87. A comparison of these data gives Br = 79.940. [This, I think, must be a misprint for 79.949.] Fourteen experiments were made on the equivalence of KBr and Ag by Pelouze's method, (see

As, Pelouze's determination.) The mean result was that 100 Ag = 110.345 KBr; extreme difference, 0.029. This gives Br = 79.958 for Ag = 107.93, and K = 39.137. The bromate of silver was prepared from potassic bromate and silver salts. For the preparation of Ag see Stas' determination of it. The potassic bromate was prepared by the action of chlorine on a mixture of KBr and KHO. The bromide was prepared by the action of heat on bromate, by treating bromine with KHO, and in other ways. No reagents were probably ever prepared with such care as those employed in this and the accompanying determinations. The weights are all in vacuo. (Stas, Untersuch. über Chem. Proport., Leipzig, 1867.)

CADMIUM.

Regnault, Kopp, and Bunsen have determined the specific heat of cadmium, which corresponds to an atomic weight of 112. Deville and Troost determined the density of cadmium vapor at above 1000°. It answers to an atomic weight of 114. (Gmelin-Kraut, l. c.; L. Meyer, l. c.)

F. STROMEYER: 111.48 (O = 16); 696.767 (O = 100).

Stromeyer found that 100 parts of cadmium combine with 14.352 parts of oxygen to form the oxide. (Berzelius' Lehrbuch der Chemie, 5th ed., 3, 1219; Schweigger's Journ., 22, 1818, 362.)

T. Thomson: 112 (O = 16); 700 (O = 100).

Thomson says that he has shown this to be the true value by analysis of the sulphate in two different states. (System of Chem., 7th ed., 1, 1831, 555.)

K. von Hauer: 112 (O = 16); 700 (O = 100).

Determined by nine experiments on the reduction of cadmium sulphate to sulphide in a current of hydrogen sulphide under pressure. The mean of the experiments gave Cd = 55.999; extreme difference, 0.16. Von Hauer took S = 16. The sulphate was purified by repeated recrystallizations and by conversion into oxide. It was dried at 200°. The sulphide was in each case carefully examined for undecomposed sulphate. (Erdmann's Journ. für Prak. Chem., 72, 1857, 346.)

J. Dumas: 112.24 (O = 16).

Determined by six experiments on the titration of cadmium chloride with argentic nitrate. The mean of all the experiments was Cd = 56.12; extreme difference, 0.49. The third experiment varies considerably from the rest, and Dumas seems inclined to omit it in the average. If it is left out, the mean becomes 56.06; extreme difference, 0.29. Dumas takes Cl = 35.5; Ag = 108. The cadmium chloride was prepared in two lots by solution of cadmium in chlorhydric acid, evaporation and melting for several hours in a current of chlorhydric acid gas. (Annal. de Chimie et de Physique, (3,) 65, 1859, 158.)

E. Lenssen: 112.06 (O = 16).

Three experiments were made on the decomposition of cadmium oxalate, the salt and the resulting oxide being weighed. The mean result was Cd = 56.03; extreme difference, 0.19. C = 6. The oxalate was prepared from pure chloride by precipitation with oxalic acid, washing and drying at 150°. It was carefully tested, and was found to be anhydrous. (Erdmann's Journ. für Prak. Chemie, 79, 1860, 281.

CÆSIUM.

The great similarity between cæsium and the other alkaline metals renders the deduction of its atomic weight from its equivalent sufficiently certain.

Kirchhoff and Bunsen: 123.35 (O = 16).

Determined by three experiments on the analysis of the chloride with argentic nitrate. The value is the mean; extreme difference, 0.13. The cæsium was separated from the other alkalies by extracting a mixture of oxides and carbonates with alcohol. It was converted into chloride by precipitation with platinum chloride, reduction of the double chloride in hydrogen and solution. These operations were repeated until the cæsium salt gave sensibly the same results after successive purifications. Its purity was also tested spectroscopically. Silver was taken at 107.94, and chlorine at 35.46. (Poggend. Annal., 113, 1861, 363.)

Johnson and Allen: 138 (O = 16).

Determined by four experiments on the precipitation of cæsium chloride with argentic nitrate. The mean result was Cs = 133.036; the extreme difference, 0.842. Ag = 107.94; Cl = 35.46. Cæsium and rubidium were separated by partial crystallization of their bitartrates. The cæsium bitartrate was converted into chloride by precipitation with platinum chloride, reduction and solution. The nitrate formed on the precipitation of the cæsium chloride with silver was reconverted into cæsium chloride and redetermined, and so on. The purity of the salt was tested spectroscopically. (Silliman's Amer. Journ., (2,) 35, 1863, 96.)

R. W. Bunsen: 133 (O = 16).

Determined by three experiments on the precipitation of cæsium chloride with argentic nitrate. The mean result was 132.99; extreme difference, 0.02. Ag = 107.94; Cl = 35.46. In order to prepare pure chloride, a mixture of cæsium and rubidium salts was converted into carbonates, a little more tartaric acid was added than was necessary to form acid tartrate with the rubidium and neutral tartrate with the cæsium, and the mixture was exposed on a filter to the action of a saturated atmosphere of aqueous vapor. The cæsium salt is deliquescent, and gradually passes through the filter, while the rubidium salt is unaffected. The cæsium tartrate was turned into pure chloride by repeated precipitation with platinum chloride, reduction in hydrogen and solution. The determinations were made on the product of successive purifications, and only those were taken into consideration which were made after analysis showed a constant composition. The spectroscope was employed to test the purity of the salt. (Poggend. Annal., 119, 1863, 5.)

—. Mercer: 133 (O = 16).

The fact of this determination, without details, is mentioned by Frankland. (Chem. News, 8, 1863, 18.)

R. Godeffroy: 132.557 (O = 16).

Derived from the mean of four analyses of cæsium chloride with argentic nitrate, the extreme difference being 0.185. Cl = 85.5; Ag = 108. The cæsium was separated from the other alkalies by the fractional crystallization of their alums continued until the cæsium compound was

spectroscopically pure. The aluminium was removed with ammonia, the sulphuric acid with barium chloride and traces of barium with ammonium carbonate. The cæsium chloride, which was not deliquescent, was dried at 150°. (Liebig's Annal., 181, 1876, 185.)

CALCIUM.

Bunsen has determined the specific heat of calcium. It corresponds to an atomic weight of 40. (Gmelin-Kraut, l. c.)

F. H. WOLLASTON: 40.736 (O = 16); 254.6 (O = 100).

Wollaston found that 43.7 parts of carbon di-oxide saturated 56.3 parts of lime. If C = 75.4, the value follows. (*Phil. Trans.*, 104, 1814, 20.)

J. J. Berzelius: 40.32 (O = 16); 252.075 (O = 100).

301 parts of anhydrous calcium chloride gave 775 parts argentic chloride. If Cl = 443.28 and Ag = 1349.66 the value follows. This analysis, made in 1818, was erroneously calculated from a mistake in setting down its results and the atomic weight of Ca was taken at 256.019. (Poggend. Annal., 8, 1826, 189; and Lehrbuch der Chemie, 5th ed., 3, 1227.)

J. Dumas: 40 (O = 16).

Three experiments were made on the calcination of calcium carbonate which contained 0.03 per cent. of ferric oxide and silicic acid. The weight of the residue was in mean 56.07, or, subtracting 0.03, 56.04, with an extreme difference of 0.08. These figures give almost exactly 40. The weighings are reduced to vacuum. (Paris Comptes Rendus, 14, 1842, 537.)

—. SALVETAT: 40 (O = 16); 250 (O = 100).

It is to be inferred from the context that this determination was made from the loss of weight ensuing on the decomposition of calcium carbonate by heat or sulphuric acid. (Paris Comptes Rendus, 17, 1843, 318.)

C. Marignac: 40.208 (O = 16); 251.3 (O = 100).

Determined by precipitating calcium chloride with argentic nitrate; Ag = 1349.01; Cl = 443.2. Marignac laid no weight on this determination finding it impossible to prepare calcium chloride which did not show an alkaline reaction. The presence of caustic lime would make the result erroneously high; no doubt Berzelius' early analysis was defective from the same cause. (Berzelius' Jahresbericht, 24, 1844, 103; Bibl. Univ., 46, 1843, 367.)

ERDMANN and MARCHAND: 40.007 (O = 100).

Four experiments were made on the calcination of calcium carbonate enclosed in a double platinum crucible in a wind-furnace, till the weight was constant. A mean of 56 per cent. calcium oxide was found with an extreme difference of 0.05. This gives Ca = 40 for C = 12. Two experiments were made by decomposing calcium carbonate by sulphuric acid. These gave a mean of 43.99 carbonic acid; difference, 0.02. The value taken is the mean of all experiments. The carbonate was prepared by precipitating calcic chloride with ammonium carbonate, and drying at 160° to 180°. Confirmatory experiments were made on iceland spar. The weighings are reduced to vacuum. (Erdmann's Journ. für Prak. Chem., 26, 1842, 472.)

Berzelius maintained that Erdmann and Marchand employed material containing water, chlorine and magnesium. Erdmann and Marchand answered that there could be no magnesium and was no chlorine but that they had convinced themselves that spar is the only compound of certain and constant composition. Berzelius replied that they then admitted that their carbonate contained water. and Marchand appealed to their experiments on spar, upon which Berzelius made experiments showing that spar, too, retains water at 200°. This Erdmann and Marchand denied and finally assert that all the carbonic acid is not driven off at any attainable temperature, and that their results were therefore too high instead of being too low. The error they estimate to exactly cover the difference between their averages and 40. (Erdmann's Journ. für Prak. Chem., 31, 1844, 257; 37, 1846, 75; 50, 1850, 287.)

ERDMANN and MARCHAND: 40.062 (O = 16); 250.39 (O = 100).

The spar experiments referred to above. Six analyses were made as before, giving a mean of 56.028 oxide; extreme

difference 0.047. (Erdmann's Journ. für Prak. Chemie, 31, 1844, 268.) Another experiment, in which the absence of water was proved, gave 56.03 lime. The weighings are reduced to vacuum. (Erdmann's Journ. für Prak. Chem., 37, 1846, 77.)

J. J. Berzelius: 40.264 (O = 16); 251.651 (O = 100).

Five experiments were made on the conversion of caustic lime into sulphate. The value is the mean for 8 = 200.75; extreme difference 0.962 for 0 = 100. The lime was carefully purified and burnt, but Berzelius says nothing of testing it for carbonic acid, upon which Erdmann and Marchand found an objection. Berzelius expresses himself ill satisfied with the results. (Liebig's Annal., 46, 1843, 241; also Lehrbuch der Chemie, 5th ed., 3, 1228.)

J. Dumas: 40.02 (O = 16).

Five experiments were made on the titration of calcium chloride with argentic nitrate. They give a mean of 20.065, but Dumas considers only three of them as entitled to a voice. These give 20.01; extreme difference, 0.03. The calcium chloride was prepared by dissolving marble in chlorhydric acid, digestion with lime water, filtration, evaporation, treatment with chlorhydric acid and heating in a current of chlorine. For the three experiments averaged the chloride was kept melted in the current of gas for from 8 to 10 hours. Ag = 108; Cl = 35.5. (Annal. de Chimie et de Physique, (3,) 55, 1859, 129.)

CARBON.

The specific gravity of gaseous carbon compounds shows that the atomic weight must be nearly 12. (Gmelin-Kraut, l. c.) Weber has shown that the specific heat of carbon at high temperatures obeys Dulong and Petit's law.

F. H. Wollaston: 12.064 (O = 16); 75.4 (O = 100).

Biot and Arago found the specific gravity of carbon dioxide 1.5196, and that of oxygen, 1.1036. Calculation from these data gives the value. (*Phil. Trans.*, 104, 1814, 20.)

J. J. Berzelius: earlier determinations.

In 1817 Berzelius attempted to determine the atomic weight of carbon by two analyses of plumbic carbonate. These analyses calculated for Pb = 206.926 (Stas,) give C = 11.998 and 11.984, or 74.99 and 74.90.] Considering the difference too great, he calculated the atomic weight from Biot and Arago's determination of the specific gravities of carbon di-oxide and oxygen, 1.10359 and 1.51961. lius gives 75.33 as the result; [I make it 75.394.] Subsequently, (1819,) Berzelius and Dulong determined these specific gravities more accurately at 1.524 and 1.1026 whence he calculated C = 76.437. This number was accepted until Dumas showed it to be false, although in the mean time carbon di-exide had been shown to be a condensible gas. According to Dumas, Berzelius at one time accepted a value 76.52 of which I have found no account. In Berzelius' Lehrbuch, 3, 1174, 76.48 is a misprint for 76.437. (Berzelius' Lehrbuch der Chemie, 5th ed., 3, 1197, et passim.)

T. Thomson: 12 (O = 16); 75 (O = 100).

Thomson found the specific gravity of carbon di-oxide 1.52673. Assuming the specific gravity of oxygen at 1.1111, chiefly to accord with the supposition that air is a compound containing 20 per cent. of oxygen, he calculates the atomic weight of carbon at 75. (Erdmann's Journ. für Prak. Chem., 8, 1836, 372; Records of General Science, by R. D. Thomson, 1836, 179.)

J. Dumas: about 12.16 (O = 16); 76 (O = 100).

From analysis of well crystallized naphthaline, Dumas infers that the atomic weight of carbon cannot be so high as 76.44, and must be nearly as above. (*Poggend. Annal.*, 44, 1838, 110.)

J. J. Berzelius: 12.23 (O = 16); 76.458 (O = 100).

One experiment was made on the decomposition of plumbic carbonate by heat, which gave C = 76.405. [If Pb = 206.926, the data give C = 12.185, or 76.157.] Another experiment was made on the oxalate, which gave C = 76.511. Berzelius regards these results as confirmatory of the value 76.438. The plumbic carbonate was prepared by precipitating the nitrate with ammonium carbonate. The oxalate was obtained by decomposing the acetate with oxalic acid. (Liebig's Annal., 30, 1839, 241.)

G. Fownes: 12.12 (O = 16).

Determined by three analyses of naphthaline with cupric oxide, the usual precautions being observed. The value is the mean; extreme difference, 0.14. The naphthaline was purified by slow sublimation in a florence flask, and was brilliantly white. Fownes does not regard his results as conclusive as to the exact value. (*Phil. Mag.*, (3,) 15, 1839, 62.)

E. MITSCHERLICH: 12.016 (O = 16); 75.1 (O = 100).

Experiments made on the analysis of naphthaline by the ordinary method of organic analysis gave never more than 75.2, and those which seemed most accurate very nearly 75. (Mitscherlich's Lehrbuch der Chemie, 4th ed., 1, 1844, 595.)

Dumas and Stas:
$$12 (O = 16)$$
; 75 $(O = 100)$.

Determined by fourteen experiments on the combustion of carbon in oxygen, the resulting carbon di-oxide being weighed. In five cases natural graphite was employed, and in four graphite from charcoal pig-iron. Both were purified by treatment with acid and heating in chlorine. necessary oxygen was developed in the combustion-tube from potassic chlorate and cupric oxide. In five experiments diamond was employed, and the oxygen was furnished The oxygen was displaced by air, espefrom a gasometer. cially purified from carbon di-oxide by milk of lime. products of combustion were collected in tubes filled with pumice stone moistened with sulphuric acid, Liebig potashbulbs and tubes filled with dry potash. The mean of the experiments on graphite gave C = 74.982; those on diamond gave 75.005; the extreme difference was 0.238. observers point out that the result would not be affected by reduction to vacuum. (Annal. de Chimie et de Physique, (3,) 1, 1841, 5.)

Liebig thinks that potash must have been volatilized, and says that there is no assurance that the oxygen was completely expelled by air. He also points out that the analyses of camphor and benzoic acid, accompanying the investigation, show an excess of carbon for C = 75. (Liebig's Annal., 38, 1841, 195.)

ERDMANN and MARCHAND: 12.009 (O = 16); 75.054 (O = 100).

Erdmann and Marchand repeated Dumas' and Stas' experiments. Five experiments on diamond gave C = 75.028;

31

extreme difference, 0.38. Three experiments on natural and one on artificial graphite gave C = 75.087; extreme difference, 0.13. The number is the mean of all experiments. Erdmann and Marchand adopt 75. Calcium chloride was used in these experiments instead of sulphuric acid to avoid objections as to the possible volatility of the acid. (Erdmann's Journ. für Prak. Chem., 23, 1841, 159.)

Berzelius and Liebig and Redtenbacher: 12.119 (O = 16); 75.741 (O = 100).

Five analyses by Berzelius of the tartrate of lead, the decomposition being effected by heat, gave 62.7431 per cent. plumbic oxide; extreme difference, 0.045. Several analyses of plumbic racemate gave a mean of 62.75 per cent. oxide; extreme difference, 0.05. The salts were prepared by fractional precipitation of plumbic acetate with tartaric and racemic acids respectively. They were dried at 100°. (Poggend. Annal., 19, 1830, 306.) From the analyses of the tartrate Liebig and Redtenbacher calculate C = 75.771, and from the racemate 75.711, taking Pb = 1294.489 and H = 6.2394. (Liebig's Annal., 38, 1841, 137.)

LIEBIG and REDTENBACHER: 12.137 (O = 16); 75.854 (O = 100).

Determined by decomposing known weights of organic salts of silver in a covered crucible by heat and weighing the silver. Five analyses of each of the following salts showed that 18.6113 Ag = 28.8098 acetate; 9.6171 Ag =16.223 tartrate; 16.2641 Ag = 27.438 racemate; 16.0596Ag = 25.9019 malate. If Ag = 1351.607 and H = 6.2394, the above value for C follows, with an extreme difference for the 20 analyses of 0.765, (O = 100.) The figures are all calculated for vacuum: [If Ag = 107.93 and H = 1.0025, the average number obtained from the mean of each set of analyses gives C = 12.06865 or 75.429.] The acetate was prepared by partially neutralizing pure acetic acid with ammonia, precipitating with argentic nitrate and recrystallizing the salt from hot aqueous solution. The crystals were dried at 103°. The acetic acid was prepared from plumbic acetate. The tartrate was prepared by adding tartrate of sodium and potassium to a hot (80° to 85°) dilute solution of argentic nitrate till a small permanent precipitate was formed, and cooling the solution. The racemate was prepared from pure acid racemate of ammonium like the tartrate. The malate was prepared from calcium

malate and argentic nitrate. The salt thus obtained was dissolved in nitric acid, and reprecipitated with ammonia added drop by drop, not to complete neutralization, washed and dried. (*Liebig's Annal.*, 38, 1841, 139.)

A. Strecker recalculated Liebig and Redtenbacher's analyses given above, independently of the atomic weight of silver, from the difference in their atomic composition, employing the method of least squares. He found $C=75.415\pm0.061$, or 12.066 ± 0.01 . In the same way, and from the same analyses he calculated the atomic weight of silver at 1348.79, or 107.9032. [The close coincidence between this result and Stas', is certainly worthy of remark.] (Liebig's Annal., 59, 1846, 280.)

Marignac repeated Liebig and Redtenbacher's experiments and got almost the same results, but, by varying the method so as to preclude loss by spirting, different ones.

(Liebig's Annal., 59, 1846, 287.)

Stas had the same experience as Marignac, and also ascribes Liebig and Redtenbacher's high results to loss by spirting. (Bulletin de l'Acad. Roy. des Sciences de Belgique, 16, 1849, 9.)

C. Marignac: 11.986 (O = 16).

Determined by three analyses of the acetate of silver. The salt was decomposed by heat in a tube in such a way that the products of decomposition were forced to pass through porous silver, and loss by spirting was impossible. 100 parts of the salt were found to contain a mean of 64.664 silver, with an extreme difference of 0.005 in vacuo. [If Ag = 107.93, these figures give the above value.] Marignac regards the analysis as a confirmation of Dumas and Stas' determination. The acetate was prepared by solution of argentic carbonate in acetic acid and successive recrystallizations. (Liebig's Annal., 59, 1846, 287; Bibl. Univ., Arch. des Sciences, 1. 1846.)

Strecker believes that the silver in Marignac's determination must have retained carbon. (Ibid. 284.)

F. von Wrede: 12.019 (O = 16); 75.12 (O = 100).

Von Wrede determined the specific gravity of carbon dioxide, taking into consideration its variation from the law of Marriotte. He found it equal to $1.52037 \frac{1+0.0049.p}{1+at}$. He also found the specific gravity of oxygen 1.1052 and

CARBON. 33

that of carbonic oxide 0.96779. Comparison gives C = 75.11 to 75.14. (Berzelius' Jahresbericht, 22, 1842, 72.) Berze-

lius adopted this determination.

According to Gmelin-Kraut, I, (2,) 70, Regnault's value for the specific gravity of oxygen combined with von Wrede's for carbon di-oxide gives C = 12.0037, and with that for carbonic oxide C = 12.0105.

J. S. Stas: 12.005 (O = 16); 75.029 (O = 100).

Determined by passing carbonic oxide over a known weight of pure cupric oxide, and weighing the carbon dioxide formed. Stas got from eight experiments C = 74.993The number taken is the mean of the results, to 75.055. which is misprinted in Stas' paper 75.039.] The carbonic oxide was prepared from oxalic acid by the action of sulphuric acid. It was purified from carbon di-oxide by passing through potash tubes, and from oxygen by passing over hot copper filings, and was kept in a gasometer over water, in which was dissolved a solution of stannous oxide in pot-The cupric oxide was prepared by igniting pure cupric The carbonic acid formed in the experiments was caught in potash and sulphuric acid tubes. The amount of carbon di-oxide weighed was from 23 to 67 grammes. The weighings are reduced to vacuum. (Bulletin de l'Acad. Roy. des Sciences de Belgique, 16, 1849, 9.)

GRAPHON.

B. C. Brodie: 33 (O = 16).

By the action of potassic chlorate and nitric acid on graphite, Brodie obtained a compound of carbon, oxygen and hydrogen containing 11 atoms of carbon, and by the action of heat on this substance two others containing, respectively, 22 and 66 atoms. The first of these is analogous to the hydrated oxide of silicon obtained by Buff and Woehler, if Si = 21. From this fact, and the specific heat of graphite, Brodie concludes that the atomic weight of the graphitic form of C is 33. (*Phil. Trans.*, 149, 1859, 249.) Graham-Otto points out that if Si = 28, graphon must be 44, and that, in that case, the argument from the specific heat loses its applicability.

CERIUM.

The specific heat of metallic cerium, as determined by W. F. Hillebrand, is 0.04479, and the atomic heat 6.18 if the atomic weight is 138. (*Poggend. Annal.*, 158, 1876, 86.)

It is well known that cerium is always accompanied in nature by lanthanium and didymium. The former was discovered in 1839, and the latter in 1843, both by Mosander.

W. Hisinger: 137.93 (O = 16).

According to Hisinger, as reported by Berzelius, the lower oxide of cerium contains 14.821 O per 100 Ce, giving the atomic weight at 574.718 for O = 100, if the lower oxide is regarded as a protoxide. (*Poggend. Annal.*, 8, 1826, 186.)

T. Thomson: 150 (O = 16).

Thomson analysed the sulphate and obtained for cerium the value 625, (O = 100.) [He probably took barium = 70.] (System of Chem., 7th ed., 1, 1831, 466.)

F. J. Otto: 138.91 (O = 16).

According to Gmelin, Otto found in an approximate determination Ce = 578.8, and recorded it in his revised translation of Graham's Chemistry, 1, 1840, 222.

A. Beringer: 138.48 (O = 16).

[Three analyses of cerous chloride with silver give the atomic weight of cerium at 576.875, or 92.22, if Ag = 107.93, and Cl = 35.457. Inconsistent results are given for an analysis of the sulphide.] Three analyses of the sulphate in which the oxide was determined, gave 57.4717 per cent. so-called protoxide, [or Ce = 576.31, or 92.21, if S = 82.0742.] Analysis of the formate gave Ce = 577.04 for C = 75.85. The material for the preparations was ceric oxide obtained from cerite, and purified from lanthanium by digestion with very dilute nitric acid. The lower oxide was assumed to be Ce O. (Liebig's Annal., 42, 1842, 184.)

R. Hermann: 138 (0 = 16).

The lower oxide was assumed to be Ce O. 23.523 parts of anhydrous cerous sulphate gave 29.160 parts of barium sulphate, giving Ce = 575, for O = 100, Ba = 856.88, and S = 201.16. The salt was obtained by precipitating basic

35

sulphate from a sulphuric solution of the cerite oxides, and converting this precipitate into the neutral salt. (Erdmann's Journ. für Prak, Chem., 30, 1843, 184.)

CERIUM.

C. Rammelsberg: 137.93 (O = 16).

Hermann states that Rammelsberg experimented on cerium salts free from lanthanium, and got Ce = 574.7, the lower oxide being supposed to contain one atom of oxygen. [I cannot find the original paper.] (Erdmann's Journ. für Prak. Chem., 30, 1843, 184.)

C. Marignac: 141.79 (O = 16).

The result of seven experiments on the titration of cerous sulphate, prepared from basic sulphate, with barium chloride. (Erdmann's Journ. für Prak. Chem., 48, 1849, 406; Bibl. Univ. Arch des Sciences, 8, 265.) Marignac subsequently made experiments which showed these results to be too high from the impurity of the barium sulphate precipitate, (see note to Turner's determination of Barium,) and that the number 575 (for O = 100 and cerous oxide Ce O) was more probable. (Annal. de Chimie et de Physique, (3,) 38, 1853, 148.)

T. Kjerulf: 174.56.

Kjerulf obtained, by three organic analyses of cerium oxalate, Ce = 727.33 on the protoxide theory, O = 100. The salt was prepared by dissolving cerium oxide in oxalic acid. (Liebig's Annal., 87, 1853, 12.) Bunsen points out that this must have been a basic salt. (Ibid, 105, 1858, 50.)

R. Bunsen and J. Jegel: 138.192 (O = 16).

The lower oxide was presumed to contain one atom of oxygen. In two experiments cerous sulphate was decomposed with ammonium oxalate. The sulphuric acid thus liberated was determined with barium sulphate; the cerium oxalate precipitate was decomposed by heat with the formation of ceric oxide, which was weighed and the additional oxygen, introduced by heating, determined by iodometric titration. The salt was not anhydrous; the water contents was estimated by difference. The experiments gave respectively 57.49 and 57.46 per cent cerous oxide in the anhydrous salt, or Ce = 576.3 and 575.25 if S = 200. One experiment was made on hydrous cerium oxalate. The cerous oxide was found as before; the water was determined and the

oxalic acid was estimated by difference. This gave 60.02 per cent. cerous oxide, calculated for the anhydrous salt, or Ce = 575.65. The salts were prepared from cerite as follows: the mineral was digested with sulphuric acid, the sulphates formed were leached with water and with dilute nitric acid; this solution was treated with hydrogen sulphide, chlorhydric acid was added and cerium oxalate was precipitated. The oxalate was heated with magnesia to convert the cerium into the higher oxide, which was dissolved in concentrated nitric acid. After diluting the solution, chemically pure basic sulphate was precipitated. In the preparation of cerous sulphate and oxalate oxidation was prevented by the action of sulphurous acid. (Liebig's Annal., 105, 1858, 45.)

C. Rammelsberg: 138.216 (O = 16).

One experiment on the organic analysis of cerium oxalate by heating in a current of oxygen behind copper oxide gave Ce = 575.9, (O = 100), or 92.144, (O = 16), cerous oxide being regarded as Ce O. Rammelsberg does not adopt his own, but Hermann's determination. (*Poggend. Annal.*, 108, 1859, 44.)

C. Wolf: 136.992 (O = 16).

Determined from experiments on the sulphate, prepared and analyzed as by Bunsen and Jegel. Wolf purified the basic sulphate by solution in nitric acid and reprecipitation in hot water, aided by recrystallizations. He found that the oftener these processes were repeated the smaller was the atomic weight resulting from the analysis. The purifications were repeated until the salt was spectroscopically free from didymium, and was perfectly white, (that employed by other investigators had been yellowish or buff.) The value taken, 45.664, [or \(\frac{1}{2} \) of 136.992,] was the smallest and last value reached. The investigation was made in Bunsen's laboratory. (Silliman's Am. Journ., (2,) 46, 1868, 53.)

C. H. Wing: 137.01 (O = 16).

Two experiments were made on the decomposition of hydrous cerium sulphate with oxalic acid, the cerium oxalate being converted into ceric oxide by heat. The amount of cerous oxide in the ceric oxide was calculated according to Wolf's results, giving for the atomic weight of cerium 45.64 and 45.69, 8 being 32. The cerium was six times recon-

verted into basic sulphate, and repeated recrystallizations were made. The salt was white and spectroscopically pure. The determination was made in Gibbs' laboratory. (Silliman's Amer. Journ. (2,) 49, 1870, 356.)

D. Mendelejeff: 138 (O = 16).

Mendelejeff first suggested raising the atomic weight of cerium from 92 to 138. His reasons were a specific heat determination which he had made with very impure metal, and the fact that the supposed sesquioxide had never been shown to exist. He believes that the atomic weight will be found somewhat below 138, because that is the atomic weight of barium. (Liebig's Annal., suppl., 8, 1871, 186.)

H. Buehrig: 140.648 (0 = 16).

Determined from ten analyses of the hydrous oxalate performed by combustion in a current of pure oxygen behind copper oxide. The water was collected in tubes filled with calcic chloride, and the carbonic acid in potash. Five experiments in which the cerium oxide was not determined gave a mean of 94.1304, on the supposition that cerous oxide contains 1 atom of oxygen and that O = 15.96, with an extreme difference of 0.0445. Five determinations in which the cerium was determined as ceric oxide gave 94.2260, with an extreme difference of 0.0431. Carbon was taken at 11.97. The mean result is Ce = 94.1782 for the above mentioned assumptions, [or 140.648 for O = 16, and on the supposition that cerous oxide is a sesqui-compound. The oxalate was prepared from basic nitrate purified by Gibbs' method of oxidation with minium and nitric acid. The salt was spectroscopically pure. (Erdmann's Journ. für Prak. Chem., 120, 1873, 222.)

CHLORINE.

The density of chlorine gas and the specific heat of chlorine compounds leave no doubt that the atomic weight of this element is nearly 35.5. (Gmelin-Kraut, l. c.)

MARCET, BERZELIUS, WOLLASTON: 35.28 (O = 16).

Marcet, by experimenting on the calcination of pure marble, and on the saturation of chlorhydric acid with lime, found as the mean of many trials, that 50.77 calcic carbonate are equivalent to 56.1 calcic chloride. Wollaston, taking the equivalent of calcic carbonate at 630, and that of calcium at 255, calculates the equivalent of chlorine at 441 for O = 100. Wollaston cites Berzelius as having obtained the same number by the conversion of plumbic carbonate into chloride. (*Phil. Trans.*, 97, 1807, 301; 104, 1814, 20.)

J. J. Berzelius: 35.412 (O = 16); 221.327 (O = 100).

The molecular weight of potassium chloride was ascertained from four experiments on the decomposition of potassium chlorate, which on being heated lost 39.15 per cent. oxygen. This gives for the chloride 932.567, (O = 100.) 100 parts of potassium chloride were further found equivalent to 192.4 parts argentic chloride, and 100 parts of silver to 132.75 argentic chloride. The value follows. Berzelius in his Lehrbuch accepts Marignac's determination and ascribes the error of the value he had obtained to the imperfect decomposition of that portion of the chlorate which was carried off as dust during the experiment. (Poggend. Annal., 8, 1826, 17; also Lehrbuch der Chemie, 3, 1189, 1191.)

E. Turner: 35.42 (O = 16).

Turner made two experiments on the decomposition of plumbic chloride with argentic nitrate. Assuming the atomic weight of lead at 103.6, and that 100 silver = 132.8 chloride, these analyses gave Cl = 35.48 and 35.48. Turner also decomposed corrosive sublimate with calcic oxide neutralized with nitric acid and precipitated with argentic If mercury = 201, these analyses give a maximum of 35.28, and a minimum of 35.21, of which Turner selects the largest. From calomel treated in the same way, he arrived at the value 35.35. From his experiments on the composition of argentic chloride (and apparently comparison with potassic chloride and chlorate) Turner got 35.45. The mean of the other experiments was 35.35, but Turner considers 35.42 as being the most likely value. The plumbum chloride was prepared from the carbonate, and was purified by recrystallization, as was also the corrosive sublimate. The calomel was "prepared by Mr. Howard," and retained traces of moisture at 300°, which would make the atomic weight derived from its analysis too small. values are for vacuum. (Phil. Trans., 123, 1833, 529.)

F. Penny: 35.454 (0 = 16).

Six experiments on the conversion of silver into nitrate gave 100 Ag = 157.441 nitrate; extreme difference, 0.028. Twelve experiments by three different methods on the conversion of silver into chloride gave 100 Ag = 132.837 chloride. Four series of experiments on the interconversion of potassic chloride, chlorate and nitrate gave for the difference between the molecular weights of the chloride and the nitrate 26.56. Corresponding experiments with sodium salts gave the same difference 26.568. The mean combined with the data for the silver salts gives the molecular weight of argentic chloride at 143.424, and Cl = 35.454. For further details see Penny's determinations of potassium, sodium, nitrogen and silver. The weighings were calculated for vacuum. (Phil. Trans., 129, 1839, 32.*)

R. PHILLIPS: 35.688 (O = 16).

In order to avoid the error possibly incurred by the melting of argentic chloride, etc., Phillips mixed known and nearly equivalent quantities of silver dissolved in nitric acid, or of crystallized argentic nitrate, with ammonium chloride; filtered, washed, and precipitated the comparatively minute amount of chlorine in the filtrate with silver solution. The fusion of this small quantity could cause no loss of importance. Phillips confesses that his ammonium chloride was acid and the only conclusions he draws are that Cl = 36, N = 14, O = 8 and H = 1 may be taken without considerable error if silver is 108. [The method seems to have been original and is nearly that afterwards adopted by Pelouze. The acidity of the ammonium chloride would of course give Cl too high.] (*Phil. Trans.*, 129, 1839, 35.)

C. Marignao: 36.001 (O = 16); 225.007 (O = 100).

Determined by passing chlorhydric acid gas over hot cupric oxide and condensing the water formed. The mean of three experiments was Cl = 450.013; the extreme difference is 0.2 for O = 100. The gas was made from recrys-

^{*}This is one of the most elegant investigations of the kind to be found in chemical literature, though it scarcely receives a mention except from Stas, who accords to it the praise it deserves. Stas' wonderfully exhaustive researches were necessary to prove beyond question that chemistry has a mathematical basis, and that the atomic weights of the elements are incommensurate. Penny's investigation, taken in connection with Stas', shows that the highest degree of accuracy is not incompatible with the simplest means when they are applied with the care and acumen, without which exact results cannot, under any circumstances, be obtained.

tallized sea-salt and concentrated sulphuric acid and was dried by passing through nine tubes filled with sulphuric acid and pummice stone and with calcium chloride. The water was collected in a condenser to which drying tubes were appended. (Paris Comptes Rendus, 14, 1842, 570.)

A. LAURENT: 35.468 (O = 16); 221.672 (O = 100).

Determined by three analyses of chloronaphthalintetrachloride, which he found to contain 58.22; 58.29; 58.28; per cent. Cl. The mean is 58.27 from which the value follows. (Paris Comptes Rendus, 14, 1842, 456.)

According to Maumené, Laurent confessed that his salt was impure, containing chlorose compounds, in Gerhardt's Comptes Rendus, 1845, 108. (Annal. de Chimie et de Physique, (3,) 18, 1846, 45.)

C. Marignac: 35.37 (O =16); 221.07 (O = 100).

One synthesis of argentic chloride showed that 100 silver equals 32.74 chlorine. Berzelius had found 32.75, which Marignac adopts. Marignac found by six experiments on the decomposition of potassic chlorate by heat, that the molecular weight of potassic chloride was 932.14. He tested the equivalence of potassic and argentic chlorides by precipitating the former with argentic nitrate, filtering without the use of paper through a funnel with a capillary neck. The precipitate was dried and weighed, then melted and reweighed, no loss being observable. 100 potassium chloride gave 192.33 and 192.34 argentic chloride in two experiments, or reduced to vacuum, 192.26. Hence the atomic weight is 442.13. The potassic chloride was prepared by heating chlorate which had been purified by repeated recrystallizations. (Liebig's Annal., 44, 1842, 23.)

C. Marignac: 35.456 (O = 16); 221.6 (O = 100).

In accordance with Pelouze's suggestion, Marignac repeated his determination of the composition of argentic chloride and of the equivalence of potassic and argentic chlorides, retaining the molecular weight of potassic chloride mentioned in the last paragraph. That value was obtained from the mean of six experiments on the decomposition of the chlorate which gave the percentage of oxygen at from 39.155 to 39.167; mean 39.161. Pelouze had got, as the mean of three experiments, 39.157. (Paris Comptes Rendus, 15, 1842, 959.) Marignac made eleven experiments on the equivalence of silver and potassium chloride by Pelouze's

method, a known weight of silver being dissolved in nitric acid and added to a known and nearly equivalent amount of potassic chloride in solution, after which the excess was titrated with decimal standard solution. 100 parts of silver were precipitated by from 69.049 to 69.067, in mean by 69.062 chloride. 100 parts of chloride were precipitated by from 192.33 to 192.37, in mean by 192.348 silver. experiments were made on the composition of argentic chloride by dissolving silver in nitric acid, with precautions against loss by spirting, precipitation with chlorhydric acid, washing, drying, melting and weighing in the same vessel. 100 parts of silver gave from 132.825 to 132.844 chloride, mean 132.84. Calculation from these data gives in vacuo Ag = 1349.01; K = 488.94; Cl = 443.20; for O = 100 [or Ag = 107.921; K = 39.115; Cl = 35.456, for O = 16.7(Berzelius' Jahresbericht, 24, 1844, 58; Bibl. Univ., 46, 1843, 350.)

C. Gerhardt: 36 (O = 16).

By heating potassic chlorate in a current of oxygen Gerhardt got, when he took precautions against loss by spirting, a mean of 60.949 chloride, from which he deduces 36 for chlorine without giving further data. (Paris Comptes Rendus, 21, 1845, 1280.) Marignac shows that no data have ever been published which, in connection with Gerhardt's experiments, would give this value for chlorine. He adds further experiments of his own which, without aiming to establish more exactly the true atomic weight, prove it less than 36 (Liebig's Annal., 59, 1846, 284; Bibl. Univ., Arch. des Sciences, 1, 1846.)

E. J. Maumené: 35.462 (O = 16).

Maumené made seven analyses of argentic chloride by reduction in a current of pure hydrogen. Five of these experiments were made with quantities less than 10 grammes, and gave a mean of 100 silver = 32.736 Cl. Two experiments were made with about 30 grammes each, and gave 100 silver equal to 32.86 and 32.853 chlorine. Maumené prefers the latter, and deduces from them for chlorine the value 43.67 or 35.494 taking silver according to his own experiments at 1350.32. [If silver is taken at 107.93 (Stas) the same analyses give 35.462.] (Annal. de Chimie et de Physique. (3,) 18, 1846, 41.)

A. LAURENT: 35.5 (O = 16); 221.88 (O = 100).

A single experiment was made as follows: pure silver was weighed off and placed in a matrass, nitric and chlorhydric acids were added, the liquid was evaporated and the chloride melted. An empty test was carried on at the same time to act as tare. Silver was taken at 1350. (Paris Comptes Rendus, 20, 1849, 5.)

J. Dumas: 35.5 (O = 16).

Determined by chloridizing different weights of pure silver by heating the metal in a current of chlorine. Experiments on 10 grammes and 20 grammes gave a mean of 35.5055, the difference being 0.013, for chlorine, if silver is 108. (Annal. de Chimie et de Physique, (3,) 55, 1859, 135.)

J. S. Stas: 35.457 (O = 16).

Stas found the atomic weight of chlorine by three inde-

pendent methods:

- (1.) From analysis of argentic chlorate and synthesis of argentic chloride. A known weight of the chlorate was dissolved in water, precipitated with sulphuric acid to secure advantageous division of the salt, and reduced while in suspension by a slow stream of sulphurous anhydride. chloride was washed, dried, and weighed in the flask in which it was produced. The minute amount of chloride present in the chlorate was collected and taken into consideration, and the wash-water was carefully examined for silver. Two analyses (of about 140 and 260 grammes) gave for the molecular weight of the chloride 143.383 and 148.407, mean 143.895. A variety of syntheses of argentic chloride in the wet and in the dry way showed that 100 parts silver combined with nearly 32.850 parts chlorine. Stas assumes that none of his syntheses can possibly have given too much chloride and accepts the relation stated. These data give Cl = 35.458.
- (2.) From the mutual relations of potassic chlorate and chloride and argentic chloride, combined with the composition of the last. The chlorate was decomposed either by gentle heat or in the wet way with chlorhydric acid. 100 parts of chlorate were found to contain 60.846 parts chloride as the mean of eight experiments; extreme difference, 0.012, which gives the molecular weight of potassic chloride at 74.59. The relation between potassic and argentic chloride was ascertained by Pelouze's method, (see Marignac's

determination above.) Twenty experiments on quantities of 32 grammes, and less, of silver gave 100 parts Ag = 69.103 parts KCl; extreme difference, 0.008. These data combined with the composition of argentic chloride given

above, indicate for chlorine 35.460.

(3.) The composition of argentic nitrate was determined, and the difference between the atomic weights of nitrogen and chlorine. In two experiments silver was dissolved in nitric acid, the solution evaporated to dryness, and the nitrate kept melted until there was no further loss of weight. The result obtained was that 100 silver = 157.484 nitrate; difference, 0.008. From series of experiments on the relation of the chlorides of potassium, sodium, lithium and silver to the nitrates, Stas found the difference between a chloride and a nitrate from 26.586 to 26.591; mean 26.588. These data show that the atomic weight of chlorine lies between 35.455 and 35.460, and confirm the mean of all the determinations of Penny, Marignac, and Stas, 35.457. silver for this investigation was either distilled or compared with distilled silver; it was found impossible to reduce the amount of silica in the alkaline salts below 0.002 of one per cent., it was therefore determined and allowed for; every possible method of purification by recrystallization and otherwise was resorted to to ensure purity. The weighings are all reduced to vacuum. (Stas, Unters. über Chem. Proport., Leipzig, 1867.)

CHROMIUM.

The specific heat of chromium, as determined from that of the oxide by Kopp, Regnault, and Neumann, corresponds to an atomic heat of from 5.4 to 5.98, if the atomic weight is taken at 52.4. (Gmelin-Kraut, l. c.)

J. J. Berzelius: 56.29 (O = 16); 351.819 (O = 100).

100 parts of plumbic nitrate, on precipitation with potassic chromate, gave 98.772 parts plumbic chromate. The value follows for Pb = 1294.498, and N = 88.518. (Poggend. Annal., 8, 1826, 22.)

T. Thomson: 64 (O = 16); 400 (O = 100).

3.14 grains of metallic chromium, converted into chromic acid by heating with potash and nitre, gave a precipitate of 16.23 grains plumbic chromate. (*Phil. Trans.*, 117, 1827, 159.)

E. Peligor: 52.48 (O = 16); 328 (O = 100).

Peligot reached this value by a careful carbon determination of chromous acetate, produced by precipitating a dilute solution of chromium protochloride with sodium acetate, C = 75. Peligot does not regard the experiment as definitive, the salt possessing but little stability. (Annal. de Chimie et de Physique, (3,) 12, 1844, 527.)

N. J. Berlin: 62.54 (O = 16); 328.39 (O = 100).

Five experiments were made on the decomposition of argentic chromate with chlorhydric acid and alcohol. silver chloride was washed in the flask in which it was precipitated, treated with aqua regia, melted and weighed without removal. The decanted fluid and the wash-water were evaporated to dryness with excess of ammonia, treated with water and the chromium oxide filtered off, heated to redness and weighed. [Nothing is said of the recovery of any argentic chloride that might have been removed by the decantation.] The value taken is calculated from the comparison of the amounts of argentic chloride and of chromium oxide obtained, $Ag = \overline{1349.66}$; Cl = 443.28. extreme difference is 1, for O = 100. The argentic chromate was prepared by adding nitrate to a solution of potassic chromate. (Erdmann's Journ. für Prak. Chem., 38, 1846, 145.)

V. A. JACQUELIN: 50.08 (O = 16); 313 (O = 100).

By washing and purifying violet chromium chloride, Jacquelain obtained a substance which he took to be the pure chloride and which was more soluble than the unpurified salt. He analysed it by melting with soda, and arrived at the above number. (Liebig's Annal., 64, 1847, 275; Revue Scient., 14, 198.)

A. Moberg: 53.563 (O = 16); 334.769 (O = 100).

Moberg made twelve experiments on the decomposition of chromium salts by heat. In two cases the sulphate dried at a low red heat was decomposed by strong ignition in a platinum crucible; the results being, 335.65 and 335.29 for chromium. Ten experiments were made on the decomposition of ammonium-chromium-alum which had been dried in a pulverized state for a long time. These determinations gave from 333.965 to 335.739. The value taken is the mean. The alum employed was prepared from pure material, and was repeatedly recrystallized. S = 200; N = 87.5. (Erdmann's Journ. für Prak. Chem., 43, 1848, 115.)

J. Lefort: 52.97 (O = 16).

Determined by fourteen experiments on the precipitation of barium with sulphuric acid from a nitric acid solution of barium chromate. The barium chromate was prepared by precipitating potassium chromate with barium nitrate and drying the precipitate at 250°. [If these analyses are calculated for barium = 137 and S = 32, they give 100 barium chromate = 60.244 barium oxide, extreme difference, 0.26, and the atomic weight as above. Lefort seems to have taken Ba = 136.72. Berlin points out the correction which I have verified.] (Erdmann's Journ. für Prak. Chem., 51, 1850, 261; Journ. de Pharm. et de Chim., 18, 27.)

R. WILDENSTEIN: 53.485 (O = 16).

Determined by thirty-two experiments on the precipitation of barium chloride, desiccated at a red heat, by pure, neutral potassic chromate. The mean of these analyses gave 100 barium chromate = 81.70 barium chloride; extreme difference 0.35. Wildenstein calculates 384.48 without giving the assumption for chlorine. [If Cl = 35.457; Ba = 137, the value follows.] (Erdmann's Journ. für Prak. Chem., 59, 1853, 28.)

F. Kessler: 52.3 (O = 16).

Kessler reached this value by comparing the oxidizing action of potassic chromate with that of potassic chlorate on arsenious acid. Six experiments were made on the oxidizing power of the chromate and twelve on that of the chlorate by a method of titration. By combining the maximum of one with the minimum of the other series, Kessler finds the atomic weight of chromium between 25.93 and 26.40; in mean 26.15, K being = 39.12 and Cl = 35.45. Confirmatory experiments were made on the oxidation of ferrous chloride in the same way. These gave a mean of 26.1. (Poggend. Annal., 113, 1861, 137; 95, 1855, 208.)

M. SIEWART: 52.094 (O = 16).

Determined from the amount of chlorine in sublimed violet chromium chloride. Siewart criticises Kessler's determination and deduces from the latter's data a value 25.02. (Kopp's Jahresbericht, 14, 1861, 240; Halle, Zeitschr. für die Gesammt. Naturwis., 17, 530.)

Kessler points out that the number 25.02 is a misprint in the Jahresbericht, and that Siewart's paper ascribes to him the value 26.02. (Poggend. Ann., 117, 1862, 352.)

COBALT.

The atomic heat of cobalt as determined by Regnault is 6.27 if the atomic weight is assumed at 58.8. (Gmelin-Kraut, l. c.)

E. Rothoff: 58.98 (O = 16); 368.65 (O = 100).

269.2 parts of cobalt oxide converted into neutral cobaltous chloride and precipitated with argentic nitrate gave 1029.9 argentic chloride, according to Berzelius' report. (Poggend. Annal., 8, 1826, 185.) Berzelius recalculates this analysis for Cl = 221.64 and Ag = 1349.66, and gets the value taken. (Berzelius' Lehrbuch, 3, 1220.)

R. Schneider: 60.006 (O = 16); 375.04 (O = 100).

Determined from four analyses of the oxalate. The carbon was determined as in organic analysis; the metal by heating a known weight of the salt first in a current of air, then in one of oxygen, and by reduction of the oxide in hydrogen. The mean of the four analyses gave cobalt at 80.003, with an extreme difference of 0.026 for C=6. The oxalate was prepared by converting the chemically pure cobalt of commerce into roseo-cobaltic chloride, from which the metal was again reduced, then dissolved in chlorhydric acid and carbonate precipitated, which was digested with oxalic acid. (*Poggend. Annal.*, 101, 1857, 398.)

Marignac objects to this determination that the oxalate, being insoluble, may very likely have retained portions of the carbonate which could not be removed by washing.

(Bibl. Univ., Arch. des Sciences, (2,) 1, 1858, 372.)

Schneider answers that he obtained nearly identical results from lots prepared at different times, and that he believes that he has convinced himself that the oxalate contained no carbonate. (*Poggend. Annal.*, 107, 1859, 610.)

Gibbs, reporting Schneider's determination, remarks: "Very numerous and carefully made analyses of the ammonium-cobalt bases, executed in my laboratory, indicate 29.5 as the true equivalent of cobalt." (Silliman's Amer. Journ., (2,) 25, 1858, 438.)

C. Marignac: about 59 (O = 16).

Five experiments were made on the decomposition of cobalt sulphate by heat. This salt can be readily dried without decomposition, and the acid is completely driven off by heat, but the resulting protoxide contains a slight

COBALT. 47

excess of oxygen. In order to remove this excess it was melted under a known weight of an acid silicate of lead. The results for cobalt varied from 29.32 to 29.38. The sulphate was purified by recrystallization. Marignac also experimented on the chloride. The weight of this salt varies greatly with the moisture of the atmosphere when crystallized, and attempts to desiccate it usually result in the formation of some insoluble compound. Three analyses of chloride appearing to contain one molecule of water, and dried at 100°, performed by titration with silver solution, gave cobalt at 29.42 to 29.51. Five experiments were made in the same way on chloride either melted in a current of chlorine or of chlorhydric acid gas, or calcined with ammonium chloride. These determinations gave from 29.36 to 29.42. (Bibl. Univ., Arch. des Sciences, (2,) 1, 1858, 374.) [Marignac, in another investigation in the same volume, takes Ag = 108; Cl = 35.5.

J. Dumas: 59 (O = 16).

Determined by five experiments on the titration of cobalt chloride with silver. The mean result for cobalt was 29.542; extreme difference 0.09; Ag = 108; Cl = 35.5. The chloride was prepared by dissolving pure cobalt in aqua regia, evaporating in the presence of excess of chlorhydric acid and heating to redness in a current of chlorhydric acid gas. In two of the determinations cobalt from a different lot, which had been heated in a vacuum was employed. (Annal. de Chimie et de Physique, (3,) 55, 1859, 148.)

W. J. Russell: 58.74 (O = 16).

Determined by fifteen experiments on the reduction of cobalt oxide in hydrogen. The value is the mean; the extreme difference is 0.19. To obtain pure cobalt oxide Claudet's salt was prepared, purified by recrystallization, etc., reduced in hydrogen, the metal dissolved in nitric acid and the resulting salt decomposed by heating in a stream of carbon di-oxide. (Chem. Soc. Journ., (2,) 1, 1863, 57.)

Schneider considers that no sufficient precautions were taken to exclude air in these experiments, and that higher oxides were formed. (*Poggend. Annal.*, 130, 1867, 810.)

E. von Sommaruga: 60 (O = 16).

Determined by seven experiments on the reduction of purpureocobaltic chloride in a current of hydrogen. The mean of the experiments is 29.965; four of them give a

mean 29.996. The extreme difference is 0.098. The salt was prepared by solution of the carbonate in chlorhydric acid, addition of ammonia in excess, exposure to the air, washing of the precipitate with acidulated, then with pure water and drying at 110°. A special examination showed it free from other metals. Sommaruga took Cl = 35.5; N = 14. (Erdmann's Journ. für Prak. Chem., 100, 1867, 118; Sitz.-Bericht der k. k. Akad., 1866.)

C. Winkler: 59 (O = 16).

This value is derived from the mean of five experiments on the precipitation of gold from a solution of neutral crystallized chloride of gold and sodium. The metallic cobalt employed was prepared by the reduction of purpureocobaltic chloride. The latter was made from oxide, and was purified by recrystallization. Gold was assumed at 196. The mean of the results was 29.496; extreme difference, 0.071. (Fresenius' Zeitschr. für Anal. Chem., 6, 1867, 22.)

P. Welesky: 58.98 (O = 16).

Determined from the analysis of cobalti-cyanides, performed by drying the salt at 100°, and heating to redness, first in a current of oxygen then of hydrogen. Four experiments with phenylammonium-cobalti-cyanide gave cobalt at from 29.38 to 29.59. Two experiments with ammonium-cobalti-cyanide gave from 29.46 to 29.55. Mean, 29.48; extreme difference, 0.21. A single experiment by Winkler's method gave 29.42. (Berlin, Bericht der Chem. Ges., 2, 1869, 592.)

W. J. Russel: 58.76 (O = 16).

Determined by the amount of hydrogen set free by the solution of cobalt in hydrochloric acid. The value is the mean of 2 (or 4?) trials. The cobalt employed was that reduced by Russel in his former experiments on the same atomic weight. (Chem. News, 20, 1869, 20.)

R. H. Lee: 59.10 (O = 16).

Determined by analysis of cobalti-cyanide salts. They were decomposed in a crucible by heating from above. The carbon separated was burned off in air and then in oxygen, and the metallic oxide reduced in hydrogen. Six experiments on the strychnine salt gave a mean of 59.05. Six experiments on the brucine salt gave 59.15. Six experi-

ments, made with especial care, on the reduction of purpureo-cobaltic chloride by hydrogen gave 59.09. (Reported by Gibbs. Berlin, Bericht der Chem. Ges., 4, 1871, 789.)

COPPER.

Regnault, Kopp, and others have determined the specific heat of copper. It corresponds to an atomic heat of about 6 if the atomic weight is taken at 63.8. (Gmelin-Kraut, l. c.)

R. CHENEVIX: F. H. WOLLASTON: 64 (O = 16); 400 (O = 100.)

Chenevix found 20 parts of oxygen equivalent to 100 parts of copper, whence Wollaston deduces the atomic weight. (Phil. Trans., 104, 1814, 21.)

J. J. Berzelius: 63.296 (O = 16); 395.6 (O = 100).

Determined by two experiments on the reduction of cupric oxide with hydrogen, which gave 395.695 and 395.507. The water was not weighed. (*Poggend. Annal.*, 8, 1826, 182; and *Lehrbuch*, 3, 1216.)

ERDMANN and MARCHAND: 63.456 (O = 16); 396.6 (O = 100.)

Determined by four experiments on the reduction of large quantities of cupric oxide in a current of hydrogen. The hydrogen was displaced by air after the completion of the reduction. The weight of the oxide and of the copper were reduced to vacuum, but not that of the weights employed. To obtain pure cupric oxide, pure vitrol was prepared and electrolytically decomposed. The copper thus obtained was dissolved in nitric acid, and the nitrate decomposed by heat. The value is the mean; the extreme difference is 0.056 for O = 8, or 0.112 for O = 16. (Erdm. Journ. für Prak. Chem., 31, 1844, 389.)

Berzelius points out that these analyses vary among themselves much more than his own. He makes the difference somewhat greater than it really is by neglecting the reduction to vacuum. (*Ibid.*, 37, 1846, 72.)

Hampe shows that these analyses, correctly calculated, give Cu = 63.46. (Zeitschr. für Berg Hütten-und-Sal-Wesen im Preus. St., 21, 1873, 261.)

J. Dumas: 63.5 (O = 16).

Dumas says that experiments on the reduction of cupric oxide and on the sulphidation of copper have shown him that the atomic weight of copper lies between 31.5 and 32, near 31.75, but that his experiments cannot be regarded as decisive. (Annal. de Chimie et de Physique, (3,) 55, 1859, 129.)

MILLON and COMMAILLE: 63.128 (O = 16); 394.55 (O = 100).

These (three) experiments were in most respects a repetition of Erdmann and Marchand's. The value is the mean; the extreme difference is 0.49 for O = 100, or 0.0784 for O = 16. The sulphate was prepared free from iron or zinc by dissolving copper in ammoniacal sulphate or nitrate. The oxide was obtained by heating the nitrate. (Paris Comptes Rendus, 56, 1863, 1249; and 57, 1863, 145.)

Fresenius sees no reason for preferring this to Erdmann and Marchand's value. (Fresenius' Zeitschr. für Anal. Chem., 2, 1863, 474.)

W. Hampe: 63.3296 (O = 16).

In three experiments cupric oxide was reduced in a current of hydrogen with all possible precautions. The hydrogen was displaced by air before weighing, though it was shown by experiment that porous copper does not condense hydrogen. The metal was heated till incipient melting was The reduction and melting were repeated without altering the weight. Hampe attempted to control his results by reconverting the metal into oxide, but was unable to effect complete oxidation. The water produced by the reduction was found to be perfectly pure. The mean result was Cu = 31.6696, maximum, 31.6729, minimum, 31.6648. The oxide was prepared from metallic copper. To obtain pure metallic copper, sulphate free from bismuth was electrolytically decomposed, the finely divided metal well washed, then melted, first in a current of carbon di-oxide, afterwards in hydrogen, and then again in carbon di-oxide. From the metal, basic nitrate was formed and from this salt, by heating first in air and then in oxygen, oxide. In two experiments the atomic weight of copper was determined by decomposing cupric sulphate by electrolysis, and weighing the metal. The residual fluid was evaporated, and a minute amount of copper, which had escaped decomposition, was

recovered and determined as sulphide. For S = 16.037 and 0 = 8, these experiments gave Cu = 31.6577 and 31.66. The value taken is the mean of the two series. All weighings were reduced to vacuum. (Zeitschr. für Berg Hüttenwad Sal.-Wesen im Preus. St., 21, 1873, 260.)

DIDYMIUM.

W. F. Hillebrand found the specific heat of this metal 0.04563, which corresponds to an atomic heat of 6.60 for an atomic weight of 144.78. (*Poggend. Annal.*, 158, 1876, 78.)

C. Marignac: 148.8 (O = 16); 930 (O = 100).

Determined by decomposing disulphate with barium chloride. Assuming the lower oxide as a prot-oxide, he calculated the atomic weight at 620. As Marignac was not confident of the purity of his salt, and subsequently became certain that the method was untrustworthy, details are unnecessary. (Liebig's Annal., 71, 1849, 313.)

C. Marignac: 143.81 (O = 16); 898.8 (O = 100).

Five experiments were made on the sulphate by decomposition with ammonium oxalate. The didymium oxalate was heated to redness, and the resulting oxide weighed. On the assumption that the oxide was protoxide, these determinations gave a mean of 598.2 for Di, with an extreme difference of 2.5. Three experiments were made on the chloride, the insoluble oxychloride, which is unavoidable in drying the salt, being separated. The chlorine was determined with silver, and the Di as in the previous experiments. These determinations gave Di at 600.2, with an extreme difference of 5.2 for Cl = 443.2 and S = 200. salts were prepared from cerite. The cerium was extracted by treatment at first with dilute and afterwards with concentrated nitric acid. The sulphates of Di and La were separated by partial precipitation with oxalic acid and by partial recrystallization. (Annal. de Chimie et de Phys., (3,) 38, 1853, 148.)

R. Hermann: 142.44 (O = 16); 890.25 (O = 100).

In one experiment sulphate which had been heated to a low red heat, was dissolved, decomposed with ammonium

oxalate, the precipitate incinerated and the oxide weighed. The result was Di = 594.46, on the prot-oxide hypothesis, for S = 200. In one experiment the chloride was decomposed with argentic nitrate, oxychloride being filtered off and allowed for, and the argentic chloride weighed. This experiment gave Di = 592.54 for Cl = 443.2. For the preparation of the salt see Lanthanium. (Erdmann's Journ. für Prak. Chem., 82, 1861, 387.)

H. ZSCHIESCHE: About 144 (O = 16).

In five experiments the sulphate was exposed to a white heat until the weight became constant and the oxide on being tested showed no traces of sulphur. The results varied from Di = 46.585 to 48.08, probably, Zschiesche thinks, on account of the presence of La. S = 16. Di was separated from La by the partial precipitation of the nitrates with oxalic acid, the first portion falling being redissolved, and the partial precipitation repeated twenty times. (Erdmann's Journ. für Prak. Chem., 107, 1869, 74.

C. Erk: 142.695 (O = 16).

The sulphate was decomposed with ammonium oxalate, the oxalate incinerated and the oxide weighed. The sulphuric acid was also precipitated as barium salt, and weighed. Three experiments gave a mean of Di = 95.13, on the prot-oxide hypothesis, with an extreme difference of 0.78. The Di salt was found to contain yttrium which was removed by repeated fractional precipitation with sodium sulphate. This re-agent precipitates a double salt of Di and sodium. The purification was continued until the atomic weight became constant. (Kopp's Jahresbericht, 1870, 819, Jena'sche Zeitschr, für Med. und Nat., 6, 299.)

Casselmann thinks that the salt may still have retained yttrium, and Fresenius objects to the barium sulphate determination on the well-known grounds. (Fresenius' Zeitschr, 10, 510.)

D. Mendelejeff: 138 (O = 16).

From the analogy between Di and cerium and other elements, and from the fact that it forms two oxides, Mendelejeff believes that its lower oxide is a sesqui-oxide, and its atomic weight 138. Mendelejeff points out that an error is to be apprehended in the received values from the fact that we have no guarantee of the pureness of Di salts except recrystallization. (Liebig's Annal. Suppl. 8, 1871, 190.)

P. T. CLEVE: 147.01 (O = 16).

Determined by the conversion of didymium oxide into sulphate. The number is the mean of six experiments; extreme difference 0.58. The Di was separated from lanthanium by repeated precipitations of basic nitrate from nitric acid solution, conversion into formate and decomposition of this salt by heat. (Kopp's Jahresbericht, 1874, 259. Bulletin Soc. Chimique, (2,) 21, 246.)

W. F. HILLEBRAND: 144.78 (O = 16).

Determined by one experiment on the conversion of metallic Di into nitrate, and then, by heat, into oxide. The impurities were determined. The metal was reduced electrolytically from the chloride. (*Poggend. Annal.*, 158, 1876, 78.)

ERBIUM.

The physical and chemical analogies of the salts of this element have led Mendelejeff (Liebig's Annal., Suppl. 8, 1871, 195,) and P. T. Cleve (Kopp's Jahresbericht, 1874, 260; Bulletin Soc. Chimique, (2,) 21, 344,) to regard it as triatomic, and its atomic weight as about 170.

M. Delafontaine: 113.04 (O = 16).

M. Delafontaine investigated gadolinite by Mosander's method, and obtained besides yttrium, two substances which he regarded as erbium and terbium. From the sulphates, in which he supposed the metals to exist as protoxides, he determined erbium at 496 and terbium at 471 for O = 100. Popp (Liebig's Annalen, 131, 189,) and Bunsen and Bahr (Ibid, 137, 1,) have shown that Mosander's method gives only mixtures. Delafontaine's terbium is thought to have been chiefly the erbium of other chemists. (Liebig's Annal., 134, 1865, 108.)

Bahr and Bunsen: 168.9 (O = 16).

A known weight of erbium oxide was treated with a very slightly excessive quantity of sulphuric acid; the solution evaporated and the excess of acid driven off at as low a temperature as possible. The increase of weight indicates 112.6 for S = 32. The oxide was prepared from gadoli-

nite. The mineral was decomposed with chlorhydric acid, and the earths precipitated with oxalic acid. The oxalates were converted into nitrates, the cerium metals separated with potassic sulphate, and calcium and magnesium with ammonia. If the nitrates of yttrium and erbium are dissolved in boiling water, basic erbium nitrate with some yttrium crystallizes out, leaving yttrium nitrate with some erbium in solution. The process of partial crystallization was continued as long as the atomic weight of the erbium salt continued increasing. Bahr and Bunsen believe, however, that the atomic weight may be some hundredths higher. The salt was spectroscopically free from didymium. (Liebig's Annal., 137, 1866, 2.)

P. T. Cleve and O. M. Hoeglund: 170.55 (O = 16).

Determined from four syntheses of the sulphate, giving 113.7 on the diatomic hypothesis. The oxide was purified by heating the nitrates, etc., according to Berlin. (Blomstrand in Berlin, Ber. der Chem. Ges., 1873, 1467; Bull. Soc. Chimique, 1873, 193 and 289.)

FLUORINE.

Dumas and Peligot and others have determined the vapordensity of a number of fluorine compounds. They correspond to an atomic weight of about 19. (L. Meyer, l. c.)

H. DAVY: 18.86 (O = 16).

Determined by the conversion of Derbyshire spar into sulphate. 100 parts of spar gave a maximum of 175.2 parts calcic sulphate. [If S = 32; Ca = 40; the value follows.] (*Phil. Trans.*, 104, 1814, 64.)

J. J. Berzelius: 18.85 (O = 16).

Determined by conversion of calcic fluoride into sulphate. 100 parts fluoride gave, in mean of three experiments, 175 parts sulphate; extreme difference, 0.2. [If S = 32; Ca = 40; the value follows.] (Poggend. Annal., 8, 1826, 18, and Lehrbuch, 3, 1196.)

P. LOUYET: 19 (O = 16).

Determined by six experiments on the conversion of fluor-spar into calcic sulphate. The mean result was 100 parts spar equal 174.36 sulphate, with an extreme difference of 0.3. Spar from Derbyshire was pulverized, digested with chlorhydric acid, and the foreign matter removed by lutration in water. It was completely dissolved in sulphuric acid, the excess of which was driven off by heat continued till a constant weight was obtained. S = 200; Ca = 250. (Erdmann's Journ. für Prak. Chem., 47, 1849, 104; Annal. de Chim. et de Phys., (3,) 25, 1849, 291.)

E. FREMY.

This chemist says that his analyses essentially confirm Berzelius' determination. (Annal. de Chimie et de Phys., (3,) 47, 1856, 27.)

J. Dumas: 19 (O = 16).

Determined by the conversion of fluorides into sulphates. A single experiment on the conversion of calcic fluoride gave 18.96; two experiments on sodic fluoride, 19.06; and two on potassic fluoride, 18.99. The mean is 19.01; extreme difference, 0.12. Ca = 20; Na = 23; K = 39; S = 16. The alkaline salts were well crystallized and were fused before use. (Annal. de Chim. et de Phys., (3,) 55, 170.)

S. DE LUCCA: 18.96 (O = 16).

Determined by four experiments on the conversion of a pure spar from Gerfalco into sulphate. The extreme difference was 0.15. The decomposition was very difficult. The loss on ignition and the residue left on evaporation of the acid employed were taken into consideration. [S apparently = 16; Ca = 20.] (Paris Comptes Rendus, 51, 1860, 299.)

GALLIUM.

Berthelot has determined the specific heat of gallium at 0.079 corresponding to an atomic heat of 5.52, if the atomic weight is 69.9. (Paris Comptes Rend., 86, 1878, 786.)

L. DE BOISBAUDRAN: 69.9 (O = 16).

This chemist "has prepared several chlorides, [samples of chloride?] several bromides, and several anhydrous iodides of gallium. He has determined the atomic weight of gallium, and found it 69.9, (mean of two experiments.)" (Paris Comptes Rend., 86, 1878, 756.)

GOLD.

Dulong and Petit and Regnault have determined the specific heat of gold. It corresponds to an atomic weight of about 200. (Gmelin-Kraut, l. c.)

J. J. Berzelius: 196.4 (O = 16).

Determined by the amount of mercury necessary to precipitate a known weight of gold from solution of chloride. 142.9 mercury were found equivalent to 93.55 gold. [If Hg = 200, this gives Au = 196.397.] (Poggend. Annal., 8, 1826, 178.)

T. Thomson: 200 (O = 16).

This value is derived from a somewhat inaccurate experiment on the reduction of auric chloride by ferrous sulphate. (Edinb. Trans. Roy. Soc., 11, 1831, 26.)

J. J. Berzelius: 196.73 (O = 16).

Determined by five experiments on the relative amount of gold and of potassic chloride in the residue obtained by heating the double chloride of the two metals in an atmosphere of hydrogen. [Calculated for KCl = 74.594, (Stas,) these experiments give a maximum of 196.79, minimum of 196.63 and a mean of 196.727. The atomic weight derived from the first experiment is misprinted in the Lehrbuch, as is the mean in the Jahresbericht.] (Berzelius' Jahresbericht, 25, 1846, 41; and Lehrbuch, 3, 1845, 1212.)

A. Levol: 196.26 (O = 16).

A known weight of gold was converted into chloride, and this salt decomposed in boiling solution by a current of pure, washed sulphurous acid. The sulphuric acid formed was precipitated as barium salt, and the atomic weight calculated by comparison of the gold employed and the barium sulphate obtained. 1000 gold gave 1782 sulphate. [If the atomic weight of S=32.0742, and that of Ba=137.08, the above value follows.] (Annal. de Chimie et de Phys., (3,) 30, 1850, 355.)

HYDROGEN.

The density of hydrogen as determined by a great number of investigators, especially Regnault, is about $\frac{1}{16}$ of that of oxygen. If oxygen is 16, the atomic weight of hydrogen

is consequently about 1.

The atomic weights of the elements are compared either with that of oxygen or with that of hydrogen. The main advantage of assuming hydrogen as unity is the simplicity of the approximate values expressed in terms of the atomic weight of this element. The hypothesis of Prout has also had much influence in giving currency to this unit. The advantages of oxygen as a standard of comparison consist in the fact that it combines with all the elements, except fluorine, and in the superior accuracy of the determination of its specific gravity. The percentage variation between Regnault's determinations of the specific gravity of hydrogen was thirty-six times as great as occurred in his experiments on oxygen. Unnecessary complication in the approximate values of the atomic weights is as well avoided by assuming oxygen at 16 as by taking hydrogen at 1.

These reasons for the adoption of the atomic weight of oxygen as a standard of comparison appear to me conclusive, and accordingly all values in this paper have been re-

duced to O = 16.

F. H. WOLLASTON: 1.06 (O = 16); 6.64 (O = 100).

Gay-Lussac and Humboldt having shown that two volumes of hydrogen and one of oxygen form water, and Biot and Arago having determined the specific gravity of these gases, Wollaston calculated the above atomic weight. (Phil. Trans., 104, 1814, 20.)

BERZELIUS and DULONG: 0.9984 (O = 16); 6.24 (O = 100).

Determined by three experiments on the reduction of cupric oxide by hydrogen. The hydrogen was made from

pure materials, and passed through a solution of litharge in potash, and over a coarse powder of caustic potash before use. The resultant water was caught in calcic chloride and weighed. The determination was also confirmed by experiments on the specific gravity of oxygen and hydrogen. The minimum result for hydrogen was 0.9934, the maximum 1.0086. (Thomson's Annals of Phil., 2, 1821, 48.)

T. Thomson: 1 (O = 16); 6.25 (O = 100).

Thomson found the Sp. Gr. of H = 0.0694. Taking that of O as 1.1111 on theoretical grounds (the supposed compound nature of air, etc.,) he calculates the above value. (Erdmann's Journ. für Prak. Chem., 8, 1836, 374; Records of Gen. Sci., R. D. Thomson, 1836, 179.)

J. Dumas: 1.0012 (O = 16); 6.2575 (O = 100).

Determined by nineteen experiments on the reduction of cupric oxide with pure hydrogen. The gas was made from pure materials and was passed through solutions of plumbic nitrate and argentic sulphate, and over potash, and dried with cold sulphuric acid or with phosphoric acid. The weighings of the oxide and of the reduced copper were made in vacuo. [Dumas corrected the results obtained for the air contained in the sulphuric acid, but does not explain how he estimated it, while certain other possible corrections are not mentioned.] The mean of the corrected results is 12.515. The extreme difference is 0.09 for O = 100. Without the correction for absorbed air the mean is 12.533, [or 1.00264]; maximum 12.583; minimum 12.481. (Paris Comptes Rend., 14, 1842, 537.)

ERDMANN and MARCHAND: 1.0016 (O = 16); 6.26 (O = 100).

Determined by eight experiments on the reduction of cupric oxide with hydrogen, the number is the mean of the results. In four of the experiments the correction for vacuum was calculated. These gave H = 12.548; extreme difference, 0.067. In four experiments the weighings were made in vacuo. These gave a mean of 12.492, with an extreme difference of 0.015. The oxide employed was either copper scale or was produced from cupric nitrate. The hydrogen was made from pure zinc and sulphuric acid, and was purified with potash in solution and in lumps, mercuric chloride, sulphuric acid, and chloride of calcium. In the

last five experiments the gas was also passed over red-hot copper to remove traces of oxygen.) (Erdmann's Journ. für Prak. Chem., 26, 1842, 461.)

J. S. Stas: 1.0025 (O = 16).

From all the investigations that have been made on the specific gravity of the gases, the composition of water, etc., Stas is inclined to believe that the atomic weight of hydrogen cannot be less than above. Stas found that 100 silver were equivalent to 49.5973 ammonium chloride. [If N=14.044, and Cl=35.457, this relation would give H=1.0074.] (Stas, Untersuch. über. Chem. Prop., Leipzig, 1867.)

J. Thomsen: 1.0025 (O = 16).

Thomsen made three experiments on the oxidation of a known volume of hydrogen by cupric oxide, and five experiments on the combustion of a known volume of hydrogen in oxygen, which proved that 2 litres of hydrogen gave 1.6082 grammes of water under normal conditions, and at latitude 45° . According to Regnault, 1 litre of oxygen and 2 litres of hydrogen would weigh 1.6084 grammes. Hence 1 volume oxygen and 2 volumes hydrogen form water; and if H = 1, O = 15.96, [or if O = 16, H = 1.0025.] (Berlin, Ber. der Chem. Ges., 3, 1870, 928.)

INDIUM.

Bunsen found the specific heat of In 0.565 and 0.574, which correspond to an atomic weight of about 114. (Poggend. Annal., 141, 28.)

F. Reich and T. Richter: 111.39 (O = 16).

In one experiment pure indium was dissolved in nitric acid, the oxide precipitated with ammonia and weighed. This experiment gave In = 463.4 for O = 100, and on the supposition that the metal was di-atomic. In a second experiment indium sulphide was dissolved in nitric acid, and the resulting sulphuric acid precipitated with barium chloride. This gave In = 464.9. The number taken is the mean. S = 200. The metal was prepared from the oxide. After the removal of lead, etc., with hydrogen sulphide, the oxides

of iron and indium were precipitated with ammonia, the precipitate dissolved in acetic acid and impure indium sulphide reprecipitated. This operation was repeated, and the last traces of iron were removed by partial precipitation with ammonia. (Erdmann's Journ. für Prak. Chem., 92, 1864, 484.)

C. Winkler: 107.754 (O = 16).

Determined by decomposing the nitrate by heat, and weighing the resulting oxide. The mean result of three experiments was In = 35.918 for O = 8, and assuming the univalence of the metal. Extreme difference, 0.079. Metallic indium was prepared by solution of the impure sulphide in chlorhydric acid, precipitation of indium by barium carbonate, solution in sulphuric acid, and precipitation by ammonia of the oxide which was reduced by hydrogen. [This indium seems to have contained iron.] (Erdmann's Journ. für Prak. Chem., 94, 1865, 1.)

C. Winkler: 113.439 (O = 16).

In two experiments the double chloride of gold and sodium was decomposed by pure indium, giving 37.73 and 37.80 for O = 8, and assuming univalence for the metal. In two experiments the nitrate was decomposed by heat, giving In = 37.845 and 37.879. In one experiment the oxide was precipitated from nitric acid solution by ammonia. This experiment gave In = 37.811. The number taken is the mean. The impure indium sulphide was purified as in Winkler's former determination with barium carbonate, but this process requires to be repeated several times. The reduction of the oxide was performed with sodium, the excess of which was removed from the regulus by cupellation in soda. (Erdmann's Journ. für Prak. Chem., 102, 1867, 282.)

R. Bunsen: 113.76 (O = 16).

Determined by converting metallic indium into oxide by means of nitric acid and heat. He seems to regard the experiment only as confirmatory of Winkler's. The metal was the same which served for the determination of the specific heat, and was carefully tested for all impurities. (Poggend. Annal., 141, 1870, 28.)

IODINE. 61

IODINE.

Dumas determined the specific gravity of iodine vapor. It answers to an atomic weight of about 127. (Annal. de Chim. et de Phys., 33, 1826, 337.)

L. J. GAY-LUSSAC: 123.9 (O = 16).

100 parts of iodine were found equivalent to 26.225 parts of zinc. [If Zn = 65, these figures give the atomic weight at 123.9.] (Poggend. Annal., 14, 1828, 559; Annal. de Chimie, 91, 1814, 5.)

W. Prout: 126 (O = 16).

Prout found 100 parts of iodine equivalent to 25.8 parts of zinc. [If Zn = 65, this gives I = 125.97.] (Thomson's Annals of Phil., 6, 1815, 323.)

T. Thomson: 124 (O = 16); 775 (O = 100).

Thomson found 20.5 potassic iodide = 19.75 zinc iodide, = 20.75 plumbic nitrate. [If K = 39.1, and plumbic nitrate = 331, the relation given leads to an atomic weight of 124.41.] Thomson thinks that his iodine may have been somewhat impure, as he purified it only by sublimation. (Thomson's System of Chem., 7th ed., 1, 1831, 81.)

J. Dumas: 126.13 (O = 16).

Dumas determined the density of iodine vapor at 8.716 for air = 1. [Referred to the molecular weight of oxygen, this density gives the above number for the atomic weight.] Dumas thinks it probable that it can be more accurately determined by analysis. (Annal. de Chim. et de Phys., 33, 1826, 837.)

J. J. Berzelius: 126.26 (O = 16); 789.14 (O = 100).

Determined by decomposing a known weight of argentic iodide in a current of chlorine, melting the chloride and expelling free chlorine by atmospheric air. The number is the mean of two experiments; difference, 0.01. Ag = 1851.607; Cl = 442.653. The iodide was prepared by precipitation from a solution of potassic iodide with argentic nitrate. The first portion of the precipitate was set aside as possibly contaminated with chlorine. (Poggend. Ann., 14, 1828, 562.)

C. Marignac: 126.844 (O = 16).

In five experiments a known weight of silver was dissolved in nitric acid and precipitated by a known amount of potassic iodide according to Pelouze's modification of Gay-Lussac's method. The mean result was 100 Ag = 153.74 KI in air; extreme difference, 0.14. Stas has recalculated this result for Ag = 107.93, and K = 39.137. The atomic weight so found is, in vacuo, 126.847. In three experiments a known weight of silver was dissolved and precipitated as iodide; mean result, 100 Ag = 217.511 iodide. Extrème difference, 0.04. From these data Stas gets I = 126.84. The iodine was purified by recrystallization as potassic iodate. The methods employed by previous experimenters were ineffectual. (Berzelius' Jahresbericht, 24, 75; Bibl. Univ. de Genève, 46, 1842, 367; also, Stas, Untersuch. über Chem. Prop., 153.)

E. MILLON: 126.07 (O = 16); 787.915 (O = 100).

Three experiments were made on the decomposition of potassic iodate. The mean loss of oxygen was 22.473 per cent; extreme difference, 0.03. If K = 488.94, this gives I = 1580.93. In three experiments argentic iodate, which had been dried for a long time at 200°, was employed, which lost 17.0467 per cent. oxygen; extreme difference, 0.03. If Ag = 1349.01, these data give I = 1570.73. [Berzelius cites this as an atomic weight determination; Millon, however, seems to have regarded it only as a confirmation of Berzelius' number.] Millon prepared pure iodine by passing a current of chlorine through a solution of KI till the precipitated I was redissolved, and reprecipitating with an excess of KI. (Annal. de Chim. et de Phys., (3,) 9, 1843, 407.)

V. A. JACQUELIN: 125.6 (O = 16); 785 (O = 100).

Determined by the analysis of iodic acid with silver. The acid was prepared by the oxidation of iodine with nitric acid of sp. gr. 1.5. The purity of the preparation does not seem to have been tested. Ag = 1351. (Erdmann's Journ. für Prak. Chem., 51, 1850, 458; Annal. de Chim. et de Phys., (3,) 30, 1850, 332.)

J. Dumas: 127 (O = 16).

Determined by the conversion of argentic iodide into chloride in a current of dry chlorine. Two experiments gave 127.04 and 127.01 for Ag = 108; Cl = 35.5. In Gmelin-Kraut's Handbuch these data are recalculated for Ag = 108

IODINE. 63

107.93 and Cl = 35.457, giving I = 126.941 and 126.928. The argentic iodide used was prepared from zinc iodide which had been prepared from iodine in large crystals. The argentic iodide was fused. (Annal. de Chim. et de Phys., (3,) 55, 1859, 163.)

J. S. STAS: 126.851.

Stas ascertained the molecular weight of argentic iodide as follows:

In two complete analyses, a known weight of argentic iodate was decomposed by heat in a current of pure, dry nitrogen. The oxygen set free was caught by hot copper and weighed, as well as the residual argentic iodide. In one experiment argentic iodate was dissolved in ammonia, precipitated by sulphuric acid, (to secure advantageous division of the salt,) and reduced while in suspension by a slow current of sulphurous acid. The mean molecular weight reached was 234.779; extreme difference, 0.063. The samples of iodate employed were prepared: (1.) From argentic sulphate and potassic iodate, mixed boiling, the latter in excess, thorough washing and drying in air freed from organic particles; (2.) By the reaction of potassic iodate on argentic hyposulphite. The purity of the salt was carefully tested.

Stas ascertained the composition of argentic iodide as follows:

(I.) A known weight of argentic nitrate was precipitated by hydro-iodic acid and the argentic iodide washed, dried, and weighed in the same vessel. (2.) A known weight of Ag was dissolved in nitric acid, converted into sulphate, dissolved in very dilute sulphuric acid, and precipitated with hydro-iodic acid. The precipitate was washed at temperatures increasing up to 90°. (3.) A known weight of argentic sulphate was allowed to react on a known and nearly equivalent weight of iodine in an aqueous solution of sulphurous and sulphuric acids at 10°, and in the dark, till all the iodine was taken up. The excess of iodine was titrated with silver solution, and the iodide weighed. This method was employed in two experiments. (4) differed from (3) mainly in the conversion of the iodine into ammonium iodide before bringing it into contact with argentic sulphate. Four experiments were made by the last method.

The mean composition of the iodide, as derived from all the experiments, is 100 Ag = 117.5343 iodine. From these data Stas calculates the atomic weight of I at 126.857, and

that of silver at 107.928. The sum of these weights is not the molecular weight, and this, as well as recalculation of the data, shows that the number is a misprint for 126.851. Stas' results are, therefore, even closer to Marignac's than

his memoir would indicate.]

Most of the experiments were made with iodine prepared by the decomposition of nitric iodide decomposed in a large volume of water at 65°. The iodine was further purified by distillation over barium oxide and by other means. the preparation of silver see that metal. All possible precautions were observed in the preparation of all reagents and in the conduct of the experiments. (Stas, Untersuch. über Chem. Prop. Leipzig, 1867.)

IRIDIUM.

Regnault determined the specific heat of iridium. It corresponds to an atomic weight of about 198. (Gmelin-Kraut, *l. c.*)

J. J. Berzelius: 197.19 (O = 16).

Berzelius determined this value from analysis of potassium chloro-iridiate. This salt reduced in hydrogen lost 29 per cent., the same quantity lost by the corresponding platinum salt, (vide platinum.) Berzelius originally calculated the atomic weight of the platinum metals both from the loss of chlorine of these double salts and from the relation between the metal and the potassic chloride left after reduc-In his Lehrbuch he points out the impossibility of complete desiccation, and resorts exclusively to the latter method of calculation. With respect to iridium he merely asserts that its atomic weight is the same as that of platinum, without there, or elsewhere, giving data as to the amounts of iridium and potassic chloride found in the reduced salt. It is, therefore, open to question whether he assumed the identity from the loss on reduction or not. If Pt = Ir, and if KCl = 74.594, the value fellows; see platinum.] Osmium and iridium were separated by fusion with nitre, solution, and distillation. The residue was fused with potassic chloride and sodium carbonate. On solution the iridium remains behind. This residue was repeatedly roasted and reduced to drive off osmium compounds. The potassium chloro-iridiate was formed from the pure metal. (Poggend. Ann., 13, 1828, 468; Kongl. Vet. Acad. Handl., 1828.)

C. E. CLAUS: W. M. WATTS: 197.6 (O = 16).

Watts recalculated two analyses of potassium chloroiridiate by Claus from the loss in reduction, and for Cl = 35.457, (Stas.) From one analysis he finds K = 39.87, and Ir = 198.56; from the other K = 39.93, and Ir = 196.62. (Chem. News, 19, 1869, 302.)

IRON.

Regnault, Kopp and others have determined the specific heat of this metal. It corresponds to an atomic weight of about 56. (Gmelin-Kraut, l. c.)

L. J. Thenard: F. H. Wollaston: 66.2 (O = 16); 345 (O = 100).

Thenard determined the composition of the oxide at 22.5 0 and 77.5 Fe, whence Wollaston calculates the value. (Phil. Trans., 104, 1814, 21.)

J. J. Berzelius: 64.27 (O = 16); 839.218 (O = 100).

Determined by repeated experiments on the oxidation of iron, such as is used for piano wire, with nitric acid. The carbon was determined and allowed for. Berzelius in his Lehrbuch shows that the error in this determination lay in the unsuspected presence of soluble silica and on reanalysis he found enough of it to correct the number when taken into account. (Poggend. Ann., 8, 1826, 185.)

G. Magnus: 54.25 (O = 16); 339.06 (O = 100).

Magnus' experiments were made by reducing ferric oxide in a current of hydrogen at about the temperature of boiling mercury. He regarded them simply as comfirmatory of Berzelius' number. (Poggend. Ann., 3, 1825, 84.)

F. Stromeyer: 55.6 (O = 16).

Determined by reducing ferric oxide at a red heat by hydrogen. The oxide is reduced only with great difficulty at a lower temperature. The mean of the experiments gave the oxygen contents at 30.15 per cent., [whence I have calculated the value.] (Poggend. Ann., 6, 1826, 475.)

H. Capitaine: 51.36 (O = 16); 321 (O = 100).

Determined by the peroxidation of galvanically reduced iron and by measuring the hydrogen evolved on the solution of iron in sulphuric acid. (Annal. de Chim. et de Phys., (8,) 2, 1841, 126.)

H. Wackenboder: 66.48 (O = 16).

Wackenroder helped Stromeyer in his reduction of ferric oxide, of which he gives the details. He also describes five experiments of his own, which gave the oxygen contents of ferric oxide at from 30.01 to 30.38. He took no precautions to purify his hydrogen and thinks that the loss of oxygen may have been apparently reduced. [30.195 oxygen corresponds to the above value for Fe.] (Berzelius' Jahresbericht, 24, 1844, 121; Archiv. der Pharm., 36, 1844, 22.)

SVANBERG AND NORLIN: 55.97 (O = 16); 349.809 (O = 100).

In seven experiments a known weight of iron was dissolved in nitric acid and the salt decomposed by heat. The operation was performed in a glass flask. The mean result in vacuo, was 349.104; extreme difference, 0.803. In seven experiments ferric oxide was reduced with purified hydrogen. The mean was Fe = 350.514; extreme difference, 0.735. The number taken is the mean of all the experiments, in vacuo. Berzelius in his Lehrbuch cites the experiments and, by neglecting the reduction to vacuum, gets a slightly different number. He also expresses a preference for the experiments by reduction: (Berzelius' Jahresbericht, 24, 1844, 121; and Poggend. Ann., 62, 1844, 270.)

J. J. Berzelius: 66.05 (O = 16); 350.32 (O = 100).

Berzelius, as a check on the last determination, made two experiments on the oxidation of iron by nitric acid with special precautions against partial reduction. The number is the mean; difference, 0.101. The iron was melted down with glass and magnetic oxide. In his *Lehrbuch* he adopts the mean of these experiments and Svanberg and Norlin's reduction determinations. (*Poggend. Ann.*, 62, 1844, 270.)

ERDMANN AND MARCHAND: 56.016 (O = 16); 350.1 (O = 100).

Erdmann and Marchand made eight experiments on the reduction of ferric oxide in a carefully purified current of

hydrogen. The weighings of the metal were made in vacuo to avoid possible reoxidation in displacing the gas by air. The number is the mean of the experiments; extreme difference, 1.4 for O = 100. The ferric oxide was prepared by incineration of the oxalate, moistening the residue with nitric acid and reheating. (Erdmann's Journ. für Prak. Chem., 33, 1844, 1.)

L. E. RIVOT: 64.25 (O = 16); 339.01 (O = 100).

Determined by two experiments on the reduction of pure ferric oxide in a current of hydrogen. 100 parts of oxide gave 69.31 and 69.35 parts metallic iron. (Annal. de Chim. d de Phys., (3,) 30, 1850, 188.)

E. Maumené: 56.0016 (O = 16); 350.01 (O = 100).

Maumené made six experiments by dissolving iron wire in aqua regia, precipitating with ammonia, heating the precipitate to redness and weighing. The number is the mean; extreme difference, 0.34. Maumené had convinced himself by analysis of the extreme purity of the wire. (Erdmann's Journ. für Prak. Chem., 51, 1850, 350.)

J. Dumas: 56.2 (O = 16).

Two experiments on the precipitation of ferric chloride by argentic nitrate gave each 28.1. A single experiment by the same method on ferrous chloride which was slightly yellow, gave 28.1. An experiment made on ferrous chloride, which had been heated in a current of hydrogen and of HCl and was colorless, but contained metallic iron, gave when the admixture was determined, 27.99. Dumas takes Ag = 108; Cl = 35.5, (Annal. de Chim. et de Phys., (3,) 55, 1859, 157.)

LANTHANIUM.

W. F. Hillebrand has determined the specific heat of metallic lanthanium. It corresponds to an atomic heat of 6.23, if the atomic weight is taken at 139. (*Poggend. Ann.*, 158, 1876, 82.)

Several investigations on the atomic weight of lanthanium were made previous to Mosander's announcement of the discovery of didymium. F. J. Otto found it 108.41 shortly after its discovery, and announced it in his translation of

Graham's chemistry. (Gmelin's Handbuch, 5th ed., 1, 46.) Choubine, from analysis of the chloride and of the sulphate, arrived at 108.45. (Erdmann's Journ. für Prak. Chem., 26, 1842, 443.) Rammelsberg determined it from the sulphate, which was rose-colored, at 133.17. (Poggend. Ann., 55, 1842, 65.) R. Hermann found La = 144 from rose-colored sulphate. (Erdmann's Journ. für Prak. Chem., 30, 1843, 198.)

C. G. Mosander: 139.2 (O = 16); 870 (O = 100).

Mosander says that his experiments show the true value to be in the neighborhood of 680, (the metal being assumed bivalent,) but that his salts were not pure, and the determination of little value. (*Poggend. Ann.*, 60, 1843, 301.)

C. Marignac: 141.12 (O = 16); 882 (O = 100).

Eleven experiments were made on the decomposition of the sulphate by barium chloride. The results vary greatly. Marignac wrote later (Annal. de Chim et de Phys., (3,) 38, 1853, 148) that experiment had convinced him of the incorrectness of this determination, and that the true value is about 575. (La bivalent.) (Liebig's Ann., 71, 1849, 306.)

M. Holzmann: 139.22 (O = 16); 870.15 (O = 100).

In three experiments La sulphate was decomposed by ammonium oxalate. In the filtrate from the precipitated oxalate the sulphuric acid was determined as barium salt. The oxalate was decomposed by heat, and the lanthanium oxide weighed. These experiments gave a mean of 580; extreme difference, 5.6; for bivalent lanthanium. experiments the iodate was decomposed by oxalic acid, the oxide determined as before, and the iodine titrated by Bunsen's method. These experiments gave a mean of 580.2; extreme difference, 5.3. 8 = 200; Ba = 855. In the prepare ration of the salts analyzed the cerium was separated by peroxidation with magnesium oxide and precipitation as basic sulphate. After the removal of yttrium by potassic sulphate, the lanthanium and didymium salts were separated, by making a saturated solution of the sulphates at a temperature of three or four degrees, and gradually raising the temperature. Lanthanium salt then crystallizes out nearly The purification was repeated until the salts were not discolored when heated in an open crucible over the glass-blower's lamp. Bunsen assisted at this investigation. (Erdmann's Journ. für Prak. Chem., 75, 1858, 343.)

C. CZUDNOWICZ: 140.3 (O = 16); 876 (O = 100).

Czudnowicz especially disclaims making this as an atomic weight determination and he adopts Holzmann's value. The salt analysed was the sulphate, and the method the same as that employed by Holzmann. (Erdmann's Journ. für Prak. Chem., 80, 1860, 31.)

R. Hermann: 139.32 (O = 16); 870.75 (O = 100).

Hermann analyzed the carbonate by decomposing it over mercury by sulphuric acid, and measuring the carbon dioxide liberated. The residue was heated to redness and This experiment gave La = 580.4, the metal being assumed as bivalent. The carbonate was prepared by precipitating the sulphate with sodium bicarbonate. three experiments the sulphate was decomposed by ammonium oxalate and the oxide, obtained by incinerating the oxalate, weighed. These analyses gave La = 580.7. In one experiment the chloride was analysed with argentic nitrate, giving La = 580.4. The number taken is the mean; extreme difference 2.3. In the preparation of the salts, cerium was separated as basic sulphate, La and didymium were partially separated by crystallization after which a portion of the nearly pure sulphate was precipitated by ammonia, and this precipitate digested with the mother liquor. Didymium sulphate is by this means completely precipitated. B = 200; Cl = 443.2; C = 75. Hermann remarks that his former determination was made with impure material. (Erdmann's Journ. für Prak. Chem., 82, 1861, 395.)

H. ZSCHIESCHE: 135.27 (O = 16).

Determined by six experiments on the sulphate. The water was driven off at 230°, and the anhydrous salt exposed to a white heat until the weight became constant, and on being tested, showed no sulphur. The mean result was La = 45.09; extreme difference, 1.15. In preparing the salt from cerite, the cerium was peroxidized with red lead and nitric acid and was precipitated as basic nitrate. The didymium was separated by partial precipitation with oxalic acid and concentration, these operations being repeated as long as the absorption lines of Di were perceptible in the spectroscope. A correction was made for the loss of weight of the crucible, and there was no dust upon its sides. 8 = 16. (Erdmann's Journ. für Prak. Chem., 104, 1868, 174; 107, 1869, 72.)

C. Erk: 135.39 (0 = 16).

Determined by analysis of the sulphate by the method employed by Holzmann. The bases were separated by the methods which Hermann used. Yttrium was also eliminated. Fresenius in his Zeitschrift, 10, 509, objects to the details of the Erk's manipulation of barium sulphate. (Kopp's Jahresbericht, 1870, 319; Jena Zeitschr. für Med. und Nat., 6, 1870, 299.)

D. Mendelejeff: 180 (O = 16).

As La forms but one oxide, the salts of which are not, according to Marignac, isomorphous with those of the lower oxide of didymium, Mendelejeff concludes that it belongs to the same group, but that its oxide is a binoxide, and its atomic weight 180. (Liebig's Ann., Suppl., 8, 1871, 190.)

C. Marignac: 138.75 (O = 16).

By heating the sulphate till all acid was expelled, Marignac, in two experiments, determined La (bivalent) at 92.52 and 92.56; by precipitation with ammonia and heating at 92.24 and 92.48. The sulphate was purified by a great number of partial recrystallizations, and showed only doubtful traces of didymium in the spectroscope. S=16. (Annal. de Chim. et de Phys., (4,) 30, 1873, 67.)

P. T. CLEVE: 139.15 (O = 16).

Determined by the conversion of lanthanium oxide into sulphate. The number is the mean; extreme difference 0.55. The oxide was purified from didymium by repeated partial precipitation from nitric acid solution with ammonia, basic didymium nitrate going down. The lanthanium was finally precipitated with oxalic acid. The oxide was found to be spectroscopically pure by Thalén. (Kopp's Jahresbericht, 1874, 257; Paris Bull. de la Soc. Chim., 21, 196, 246, 344.)

LEAD.

Regnault, Kopp and others have determined the specific heat of lead. It answers to an atomic weight of about 207. (Gmelin-Kraut, l. c.)

71

J. J. Berzelius and F. H. Wollaston: 207.4 (O = 16); 1295 (O = 100).

Berzelius found 16.5 parts carbon di-oxide equal to 83.5 lead oxide, whence the value, if C = 75.4. [If C = 12, these figures give lead at 206.67.] Berzelius also determined the composition of the oxide at 7.15 oxygen and 92.85 lead, giving Pb = 207.52 or 1297. (Phil. Trans., 104, 1814, 20.)

J. J. Berzelius: 207.12 (O = 16); 1294.498 (O = 100).

Determined by the reduction of a known weight of oxide of lead by hydrogen and the weight of the resultant lead; mean of four nearly coincident experiments. (*Poggend. Ann.*, 8, 1826, 184.

- Longchamp is credited in some books with an atomic weight determination of lead. He made none, but only speculated on the composition of minium, taking Berzelius' determination as a basis. (Annal. de Chim. et de Phys., 34, 1827, 105.)
 - J. J. Berzelius: 207.078 (O = 16); 1294.242 (O = 100).

This value is the mean of six experiments on the reduction of the oxide in a current of hydrogen. The oxide was produced by the decomposition of the nitrate by heat. As this compound reacts upon Pt, the crucible was lined out with a coating of a very basic nitrate, which prevented the lumps of neutral salt from coming in contact with the crucible. The glass in which the oxide was reduced was not attacked. [The third analysis is miscalculated. It should show an atomic weight of 1295.595. The mean is, therefore, as above, and the extreme difference 2.421.] (Poggend. Ann., 19, 1830, 314.)

J. J. Berzelius: 207.14 (O = 16); 1294.645 (O = 100).

In his Lehrbuch, Berzelius selects five analyses made by the above method, three of them the same. These give the above mean, with an extreme difference of 0.704 for O = 100. (Lehrbuch, 3, 1219.)

E. Turner: 207.3 (O = 16).

Determined by experiments on the conversion of metallic lead and of oxide of lead into the sulphate by solution in

nitric acid and evaporation with sulphuric acid. In three experiments, Turner found 100 lead = 146.401 sulphate; extreme difference 0.055. Berzelius had found 100 Pb = 146.419 sulphate; extreme difference 0.078. Turner takes the mean of his own and Berzelius' determinations, 146.41. In one experiment Turner found 100 oxide = 135.92 sulphate. Combination of these results gives Pb = 103.6 [or more accurately 103.65.] The oxide was prepared from subnitrate. The lead was prepared from plumbic acetate which was converted into carbonate, then into nitrate, in which form it was recrystallized, then again into carbonate, and reduced by black flux. On testing, it was found perfectly pure. Weighings reduced to vacuum. (Phil. Trans., 123, 1833, 524.)

C. Marignao: 207.04 (O = 16).

Marignac made four experiments on plumbic chloride by Pelouze's modification of the silver titration method. He found Pb = 108.57-.49-.55-.46. The number taken is the mean. The salt was titrated cold, argentic chloride being soluble in hot solutions of plumbic nitrate. The plumbic chloride was purified by recrystallization, and, after being pulverized, was dried at about 200° . According to Marignac there is no difficulty in desiccating it completely at this temperature. Ag = 108; Cl = 35.5. Marignac found it impossible to convert the chloride into the sulphate completely. (Bibl. Univ., Arch. des Sciences, (2,) 1, 1858, 223.)

J. Dumas: 207.1 (O = 16).

From a single experiment on the precipitation of the chloride with argentic nitrate. The chloride used was heated for twelve hours in a current of dry HCl, and the amount of water retained determined. Dumas found it impossible entirely to desiccate the salt without decomposition, drying at 250° does not desiccate it. Cl = 35.5; Ag = 108. (Annal. de Chim. et de Phys., (3,) 55, 1859, 129.)

J. S. Stas: 206.926 (O = 16).

According to the mean of 10 syntheses of plumbic nitrate, 100 lead = 159.9703 nitrate; extreme difference, 0.023. If N = 14.044, this relation gives Pb = 206.918. Stas also made six syntheses of the sulphate, which gave in mean 100 Pb = 146.4275 sulphate; extreme difference, 0.024. If 8 = 82.0742, this relation gives Pb = 206.934. The syntheses were made in the same way as in the determination of the

73

atomic weight of silver. The drying of the nitrate could be accomplished only in vacuo and at about 155°. The weighings are for vacuum. The lead used was prepared from commercial acetate by precipitation with metallic lead, of copper, etc., conversion into sulphate, then into carbonate and reduction by potassic cyanide or black flux. (Stas, Untersuch. über Chem. Prop. Leipzig, 1867, 824.)

LITHIUM.

Regnault has determined the specific heat of lithium. It corresponds to an atomic weight of about 7. (Gmelin-

Kraut, l. c.)

The earliest determinations of this constant seem to have been made with a double salt of lithium and potassium, at all events with a very impure material. According to Arfvedson, 420.4 lithium chloride give 1322.4 argentic chloride, whence he deduces as the atomic weight the number 127.757 [or 10.22.] (Poggend. Ann., 8, 1826, 189.) L. N. Vauquelin found 430 lithium sulphate equivalent to 875 barium sulphate. [If S = 32; Ba = 137.08, this relation gives Li = 9.27.] Vauquelin does not describe the preparation of his salt. (Annal. de Chem. et de Phys., 7, 1818, 287.) C. G. Gmelin found Li = $191.21 \, \text{for } 7.65. \, \text{]}$ (*Poggend*. Ann., 15, 480; Gilbert's Ann., 62, 1819, 399.) Kralovanszky by two analyses of the sulphate with barium chloride got Li at from 10.096 to 10.168 (Liebig's Ann., 121, 94; Schweigger's Journ., 54, 1828, 231.) Thomson and Stromeyer also each got similar values. (Thomson's System of Chem., 7th ed., 1, 1881, 420.)

R. Hermann: 6.085 (O = 16); 38.03 (O = 100).

Experiments were made on the carbonate by decomposing it with acid over mercury, and measuring the resultant di-oxide. For C = 75.33, these determinations give Li = 38. Several experiments were also made by analyzing the sulphate with barium chloride. For S = 201.06 and Ba = 856.88, these give Li = 38.05. Hermann precipitated lithium carbonate with ammonium carbonate, and subsequently converted it into sulphate. The chloride was prepared from the phosphate by Berzelius' method. (Poggend. Ann., 15, 1829, 480.)

J. J. Berzelius: 6.533 (O = 16); 40.83 (O = 100).

Berzelius found that 1.874 lithium sulphate gave 3.9985 barium sulphate, and calculated this relation for 8 = 200.75; Ba = 855.29. He also found 4.4545 melted carbonate = 6.653 sulphate, but rejected the analysis. (Lehrbuch, 3, 1229, and Jahresbericht, 10, 1830, 96.)

R. HAGEN: 6.67 (O = 16).

Hagen precipitated lithium sulphate with barium chloride, and found that 0.852 dry lithium sulphate gave 1.8195 barium sulphate whence he calculates Li = 6.493. [If Ba = 137.08; S = 32; this relation gives Li = 6.57.] (Poggend. Ann., 48, 1839, 363.)

J. W. MALLET: 6.95 (O = 16); 86.89 (O = 100).

In two experiments a known weight of lithium chloride was precipitated by argentic nitrate, and the argentic chloride weighed. In one experiment lithium chloride was titrated with argentic nitrate by Pelouze's method. The number is the mean; the extreme difference is 0.18 for O = 100. Mallet takes Ag = 1849.66; Cl = 443.28. The alkalis were separated from the lithium salt by repeated treatment with ether and alcohol. The salt was examined for impurities, and was fused with a little ammonium chloride to prevent the formation of oxy-chloride. (Silliman's Amer. Journ., (2,) 22, 1856, 849.)

L. Troost: 6.5 (O = 16).

Troost found this number from analysis of the carbonate which had been crystallized from water containing carbon di-oxide and dried at 200°, but does not regard it as definitive. (Annal. de Chim. et de Phys., (8,) 51, 1857, 111.)

J. W. MALLET: 7 (O = 16).

Troost having objected to Mallet's former method of determination, he redetermined it by precipitating the sulphate with a standard solution of barium chloride, the precipitating power of which had been tested on the sulphates of magnesium and sodium. This method was adopted to avoid the well-known imperfections of the sulphur determination. Compared with sodium sulphate the atomic weight of Li was found = 6.92 and 6.95. Compared with magnesium sulphate it was found = 7.07 and 7.09.

LITHIUM. 75

Mg = 24; Na = 23. The sulphate was prepared from carbonate, and dried somewhat below a red heat. (Silliman's Amer. Journ., (2,) 28, 1859, 349.)

K. Diehl: 7.026 (O = 16).

Determined by analysis of lithium carbonate with Bunsen's apparatus and in his laboratory. Four experiments; extreme difference, 0.024. C=12. The salt was purified from alkalis by precipitation as carbonate, resolution in acid and reprecipitation, repeated until the sodium line was no longer visible. Diehl found that precipitation of the sulphate with barium chloride gave a nearly constant error on account of the retention of lithium in the precipitate, and led to nearly the same results as Berzelius got. (Liebig's Am., 121, 1862, 93.)

L. TROOST: 7 (O = 16).

Troost found 1.309 grammes lithium chloride = 4.42 argentic chloride, and 2.75 lithium chloride = 9.3 argentic chloride. From these analyses he deduces the values 7.03 and 6.99. By decomposing the carbonate, dried at 100°, with silicic acid, he found 0.97 carbonate = 0.577 carbon di-oxide and 1.782 carbonate = 1.059 di-oxide, and infers for Li 7 and 7.02. In one experiment the carbonate was converted into sulphate. 1.217 carbonate gave 1.808 sulphate. Troost calculates Li = 7.06. [If Cl = 35.457; Ag = 107.93; C = 12; S = 32; these determinations give, in the same order as above, 7.01; 6.94; 6.98; 7.02; 7.07.] The carbonate was purified by solution in water containing carbon di-oxide, and reprecipitation by boiling, the operation being repeated until the salt was spectroscopically pure. (Paris Comptes Rend., 54, 1862, 366.)

J. S. Stas: 7.022 (O = 16).

According to the mean of three determinations, 100 parts of silver = 89.358 lithium chloride; extreme difference, 0.005. If Ag = 107.93; Cl = 35.457; this ratio gives Li = 7.022. This value is confirmed by experiments on the conversion of the chloride into the nitrate, the results of which give Li = 7.018. The comparison with silver was made according to Pelouze's modification of the silver titration method. The chloride was purified from alkalis, after preliminary treatment with ether and alcohol, by pouring the dissolved salt into a boiling solution of ammonium car-

bonate containing ammonia in excess. All weighings reduced to vacuum. (Stas, Untersuch. über Chem. Prop., Leipzig, 1867.)

MAGNESIUM.

Regnault and Kopp have each determined the specific heat of this metal. It answers to an atomic weight of about 24. (Gmelin-Kraut, l. c.)

J. J. Berzelius: 25.3 (O = 16); 158.139 (O = 100).

Determined by dissolving magnesium oxide in dilute sulphuric acid, evaporating and heating to incipient redness. 100 oxide were found = 293.985 sulphate. The sulphate was perfectly soluble in water and had therefore lost none of its acid. The oxide was purified by solution in an aqueous solution of carbon di-oxide and reprecipitated by boiling. S = 200.75. (Poggend. Ann., 8, 1826, 188; and Lehrbuch, 3, 1227.)

Marchand and Scheerer recalculated this analysis for S = 200 and reached the value 157.74. They assert that the oxide may have contained alkalis and that the sulphuric acid carries off magnesium sulphate in volatilizing. (Erdmann's Journ. für Prak. Chem., 50, 1850, 392.)

W. Henry: F. H. Wollaston: 23.36 (O = 16); 146 (O = 100).

Henry found that magnesium sulphate contained 88 per cent. magnesium oxide. If S=200 the value follows. (Phil. Trans., 104, 1814, 21.)

— Longchamp: 15.35 (O = 16).

In two experiments, Longchamp found that five parts of crystallized magnesium sulphate are equivalent to 4.91 barium sulphate. [If Ba = 137.08; S = 32, the number follows.] Longchamp found 53 per cent. water which is much too high. According to Marchand and Scheerer, the data for the anhydrous salt give Mg = 97.37, for S = 200; Ba = 856.8, [or 15.74.] (Annal. de Chim. et de Phys., 12, 1819, 265.)

L. J. GAY-LUSSAC: 23.62 (O = 16).

16.205 grammes crystallized sulphate were found equal to 15.845 barium sulphate, and 19.895 magnesium sulphate

to 18.8455 barium sulphate. Calculating from the anhydrous salt Gay-Lussac found from these experiments respectively Mg =147.23 and Mg = 148.09 for Ba = 856.8; S = 200. The salt was found to contain 51.43 water. [Calculated from the anhydrous salt these data give Mg = 23.55 and 23.68. Calculated from the hydrous salt (7 molecules water) the numbers give 24.14 and 24.41, if S = 32; Ba = 137.08.] Gay-Lussac remarks that the sulphate is partially decomposed at a red heat. (Annal. de Chim. et de Phys., 13, 1820, 808.)

T. Scheere: 24.16 (O = 16); 150.97 (O = 100).

Mean of six experiments on the precipitation of the sulphate with barium chloride. Extreme difference, 0.79. S = 200.75; Ba = 855.29. After weighing, the barium sulphate was treated with dilute HCl and the chloride thus extracted allowed for. (*Poggend. Ann.*, 69, 1846, 535.)

T. Scheerer: 24.21 (O = 16); 151.33 (O = 100).

Barium sulphate formed as in the last determination was fused with soda, the barium carbonate dissolved in HCl, and reprecipitated as sulphate. In the filtrate additional magnesia was found. If the error in the former determination was the same, its corrected value would be as above. (Poggend. Ann., 70, 1847, 407.)

SVANBERG and Nordenfeldt: 24.72 (O = 16); 154.504 (O = 100).

Four experiments were made on the calcination of the oxalate, and three on the conversion of the magnesia so obtained into sulphate. The oxalate was dried at from 100° to 105° and heated to redness until the weight was constant. The oxide was dissolved in sulphuric acid, evaporated and the excess driven off by heat. The oxalate was prepared from the sulphate by precipitation with sodium carbonate and digestion with oxalic acid. The number is the mean of all experiments; extreme difference, 0.514. S = 200.75; C = 75.12; H = 12.48. (Erdmann's Journ. für Prak. Chem., 45, 1848, 473.)

According to Marchand and Scheerer, the data give Mg = 154.27 for S = 200; H = 12.5; C = 75.

MARCHAND and SCHEERER: 24.03 (O = 16); 150.19 (O = 100).

Eleven experiments were made in calcining massive magnesium carbonate from Frankenstein, and weighing the

caustic magnesia formed. The carbonate was dried at 800° , and the carbon di-oxide, which passes off above 280° , was caught by caustic baryta solution and determined. The traces of carbon di-oxide not expelled by a yellow heat were set free by solution in chlorhydric acid and also determined as barium carbonate. The silicic acid, etc., were also determined. The mean in air is 150.84; in vacuo as above. Extreme difference, 0.57. C = 75. Eleven other experiments were made with comparatively impure material and less precaution, tending to confirm the above. (Erdmann's Journ. für Prak. Chem., 50, 1850, 409.)

T. SCHEERER: 24 (O = 16); 150 (O = 100).

By separating the neutral sulphates of magnesium and calcium by means of alcohol, Scheerer found that the magnesites used by Marchand and himself contained from one-fourth to one-half per cent. calcium oxide. This correction would make their determination almost exactly 250 or 24. (Liebig's Ann., 110, 1858, 236.)

V. A. JACQUELIN: 24.408 (O = 16); 152.55 (O = 100).

Anhydrous, neutral magnesium sulphate, obtained by solution of the oxide in sulphuric acid and heating to redness, gave 33.56 per cent. pure oxide. The method adopted is not described. This oxide by treatment with sulphuric acid gave the original amount of sulphate. If S = 200, the number follows. (Annal. de Chim. et de Phys., (3,) 32, 1851, 195.)

A. MACDONNELL: 23.9 (O = 16).

Determined from analyses of anhydrous and of crystaltized magnesium sulphate. (*Brit. Assoc. Rep.*, 1852, part 2, 36; and *Kopp's Jahresbericht*, 5, 364.)

J. F. Bahr: 24.77 (O = 16); 154.842 (O = 100).

A known weight of purified magnesium oxide was dissolved in sulphuric acid, evaporated and heated to redness till the weight was constant. The number is the mean of three experiments; extreme difference, 0.515. The oxide was obtained from meteoric olivin. After removal of the heavy metals, the solution was evaporated to dryness with soda, washed and heated to redness. The oxide so obtained was dissolved in acetic acid, oxalic acid was added, the

solution evaporated nearly to dryness, and the oxalate thoroughly washed. Bahr says that the presence of alkalis could not be suspected. S=200. (Erdmann's Journ. für Prak. Chem., 56, 1852, 310; Efversigt af Akad. Færh., 1851, 303.)

Scheerer says that oxide so prepared retains carbonic acid, that sulphate is carried off in heating the sulphate to redness, and that the presence of alkalis is to be suspected. (Erdmann's Journ. für Prak. Chem., 56, 1852, 489.)

J. Dumas: 24.6 (O = 16).

Dumas made eleven experiments on the titration of magnesium chloride with argentic nitrate. He found great difficulty in preparing pure chloride, and does not feel confident of his results. The number is the mean; extreme difference, 0.28. Ag = 108; Cl = 35.5. The chloride was prepared from various salts, but was in all cases finally heated in an atmosphere of HCl. Dumas points out, however, that this process does not remove oxide if present. (Annal. de Chim. et de Phys., (8,) 55, 1859, 129.)

MANGANESE.

Regnault has determined the specific heat of manganese. It corresponds to an atomic weight of about 55. (Gmelinkraut, l. c.)

J. J. Berzelius: 56.93 (O = 16); 855.787 (O = 100).

By dissolving manganese in nitric acid, evaporating and heating to a low red, Berzelius found 100 Mn = 142.16 oxide. It was not known at the time that the oxide might be partially reduced by this process. (*Poggend. Ann.*, 8, 1826, 185; and *Jahresbericht*, 9, 136.)

J. A. Arfvedson: 56.25 (O = 16); 351.56 (O = 100).

From 1.508 chloride Arfvedson obtained 3.408 argentic chloride. If Ag = 1351.607; Cl = 221.325; the number follows. (Berzelius' Jahresbericht, 9, 1829, 136; Afhandl. i. Fysik., 6, 236.)

E. Turner: 54.9 (O = 16).

Turner analyzed the carbonate in an apparatus similar to Bunsen's. He found 34.72 per cent. carbon di-oxide and 8.427 water. For C = 6, he calculates Mn = 28.024. By dissolving the protoxide in sulphuric acid, evaporating and heating to redness, he found 9 oxide = 19.01 sulphate. If S = 16, this gives Mn = 27.96. A second experiment gave 27.93. From 12.47 Mn chloride he obtained 28.42 argentic chloride. [If Cl = 85.5, Ag = 108; this gives Mn = 54.9.] The carbonate was obtained by precipitation with potassium carbonate. The protoxide was obtained by reduction of higher oxides in hydrogen. The chloride was melted in a current of HCl gas. (Edinb. Trans., 11, 1831, 143.)

J. J. Berzelius: 66.34 (O = 16); 845.9 (O = 100).

Berzelius repeated Turner's experiments, taking larger quantities. From the chloride he got from 845.84 to .96; from the sulphate from 846.08 to .29. Ag = 1351.607; Cl = 221.825; S = 201.165. (Berzelius' Jahresbericht, 9, 1830, 136.)

J. J. Berzelius: 55.14 (O = 16); 344.684 (O = 100).

In his Lehrbuch he apparently takes the analyses of the chloride above cited, recalculated for Cl = 221.64; Ag = 1849.66. (Lehrbuch, 3, 1224.)

R. Brandes: 67.06 (O = 16); 356.602 (O = 100).

Determined by analysis of crystallized chloride. The chlorine was determined by precipitation with silver. The Mn was precipitated as carbonate, and converted into oxide by heat. The water was determined by difference, and the composition of the oxide was assumed to be as given by Berzelius, (!) whose values for Ag and Cl were also taken. (Poggend. Ann., 22, 1831, 256.)

K. VON HAUER: 54.98 (O = 16); 343.632 (O = 100).

Determined by nine experiments on the reduction of the sulphate to sulphide by heating the salt in a current of hydrogen sulphide. The reduction was performed in a porcelain tube enclosed in a charcoal fire. Number, mean; extreme difference, 0.84, for 0 = 16. The sulphate was prepared from a pyrolusite containing only silica, iron, and barium. It was reduced to protoxide, dissolved in sulphuric acid, oxidized with nitric acid, precipitated with oxalic

acid, converted into red oxide, dissolved in chlorhydric acid and alcohol, precipitated with ammonium carbonate, dissolved in sulphuric acid, repeatedly heated to redness and recrystallized, and was dried at 300°. Accurate experiments on the reduction of the red oxide proved impracticable on account of the hydroscopicity of the compound. Two experiments on the oxidation of the protoxide, undertaken as a check on the other method, gave 27.486 and 27.527 for O = 8; S = 16. (Erdmann's Journ. für Prak. Chem., 72, 1857, 361; Sitz.-Bericht der k. k. Akad., 1857.)

J. Dumas: 54.96 (O = 16).

Determined by the decomposition of the chloride with argentic nitrate. The number is the mean of five experiments; extreme difference, 0.1 for O = 16. Cl = 35.5; Ag = 108. Dumas had previously made experiments on the reduction of the hyperoxide to protoxide by hydrogen. These gave the atomic weight at from 25.99 to 26.09 for O = 8. Dumas believes that a part of the oxide was reduced to metal. The peroxide was prepared from nitrate of the protoxide. (Annal. de Chim. et de Phys., (3,) 55, 1859, 150.)

- RAWACK: 54.02 (O = 16).

Determined, in Schneider's laboratory, by reducing a known weight of red oxide to protoxide in a current of dry hydrogen, and weighing the water formed. The number is derived from the mean of six selected experiments. The extreme difference is 0.22 for O = 16. (Poggend. Ann., 107, 1859, 607.)

R. Schneider: 54.038 (O = 16).

The mean result of four analyses of the oxalate by the ordinary method of organic analysis. Extreme difference, 0.04 for O = 16. C = 12. The oxalate was prepared from chemically pure sulphate by precipitation with sodium carbonate, digestion with oxalic acid, and drying over sulphuric acid. (*Poggend. Ann.*, 107, 1859, 613.)

MERCURY.

The specific heat of mercury in the solid state, as observed by Regnault, and the vapor density, as determined by Dumas, correspond to an atomic weight of slightly above 200. (Gmelin-Kraut, l. c.; L. Meyer, l. c.)

Fourceov and Thenard, Davy, Wollaston: 200.8 (O = 16); 1255 (O = 100).

Fourcroy and Thenard found 80 = 100 Hg. Davy found 80 = 880 Hg, giving Hg = 1266. The latter also found 134 Cl = 880 Hg, which for Cl = 441, gives Hg = 1254. (*Phil. Trans.*, 104, 1814, 21.)

N. G. Sefstroem: 202.53 (O = 16); 1265.822 (O = 100).

Determined by three analyses of the oxide according to which 100 Hg = 7.89, 7.9, and 7.97 O. (Berzelius' Lehrbuch, 3, 1215.)

E. Turner: 200.72 (O = 16).

Turner made a number of determinations of this atomic weight but regarded the value he adopted, 202, only as an approximation. From the oxide, prepared from nitrate, he got 200.77 and 199.97. The compound was decomposed by heat, and the products carried over silver and gold in a narrow tube. Four experiments were made on mercuric chloride which was decomposed by pure calcic oxide, and the Cl precipitated with argentic nitrate. [These analyses recalculated for the Stas' atomic weights of Ag and Cl give 202.079, 201.701, 201.815.] Turner also made two experiments on the reduction of the chloride with stannous chloride, the Hg being collected, dried and weighed. [These experiments recalculated give 199.423 and 199.289.] The mercuric chloride was purified by recrystallization. Weighings reduced to vacuum. (Phil. Trans., 123, 1833. **535.**)

ERDMANN AND MARCHAND: 200.14 (O = 16); 1250.6 (O = 100).

Determined from the mean of four experiments on the reduction of the oxide in a current of carbon di-oxide. Copper, carbon (from sugar) oxide, and carbon, were introduced in successive layers in a combustion tube. Dry carbon di-oxide was passed through and the mercuric oxide heated. The metal was collected in a receiver to which a tube filled with gold foil was appended. The metal was perfectly clean. Moisture was removed by a stream of dry air after distillation. The oxide was purified by heating it to incipient decomposition the metallic fumes being removed

by a current of dry air. It was tested before being analysed. The extreme difference in the results was 0.8 for O=100. All weighings in vacuo. (Erdmann's Journ. für Prak. Chem., 31, 1844, 392.)

E. MILLON: 199.94 (O = 16); 1249.63 (O = 100).

Millon made two experiments by heating mercuric chloride with calcic oxide in a current of hydrogen and condensing the metal. The experiments gave 73.87 and 73.82 per cent. mercury. If Cl = 442.64, the value follows. The chloride was dissolved in ether and sublimed. It was perfectly soluble in ether and alcohol, and was well crystallized. (Paris Comptes Rend., 20, 1845, 1291.)

L. Svanberg: 200 (O = 16); 1250 (O = 100).

Svanberg made three experiments by the same method employed by Millon. The mean result was 1248.47; extreme difference, 0.94; but Svanberg shows that there was probably loss, and that the larger the quantity of chloride employed the higher the result. He regards Erdmann and Marchand's result as most probable, but in need of confirmation. Cl = 448.28. (Erdmann's Journ. für Prak. Chem., 45, 1843, 468; Kongl. Vet. Akad. Handl., 1845, 135.)

MOLYBDENUM.

Regnault determined the specific heat of molybdenum. It answers to an atomic weight of about 96. (Gmelin-Kraut, l. c.)

J. J. Berzelius: 96.36 (O = 16); 596.1 (O = 100).

One hundred parts of anhydrous plumbic nitrate, dissolved and precipitated with neutral ammonium molybdate, gave 110.68 parts plumbic molybdate. If Pb = 1294.645, N = 87.53, the value follows. Berzelius expresses himself dissatisfied with the accuracy of the determination. (Poggend. Ann., 8, 1826, 23; and Lehrbuch, 3, 1208.)

SVANBERG AND STRUVE: 92.13 (O = 16); 575.829 (O = 100).

After trying various methods without getting accordant results, these chemists made ten experiments on the sul-

phide by roasting it first in a current of moist, and then of dry air. Three experiments were excluded as imperfect. The remainder gave a mean of 89.7523 molybdic acid from 100 sulphide; extreme difference, 0.22. The value follows for S = 200. Objections have been made (*Liebig's Ann.*, 68, 211) that the difference in weight between the acid and the sulphate is too small for the purpose of the determination, and that the different analyses give very different atomic weights. The sulphide was prepared by melting together molybdic acid, sulphur, and caustic potash, and leaching the product with water and chlorhydric acid. The sulphide was dried in a current of hydrogen. The molybdic acid was dissolved in ammonia to prove the absence of sulphide. (*Erdmann's Journ. für Prak. Chem.*, 44, 1848, 315.)

N. J. Berlin: 91.96 (O = 16); 574.75 (O = 100).

Determined by four analyses of the double mono-sesquimolybdate of ammonium by heating gently with nitric acid in a platinum crucible until only molybdic acid was left. Extreme difference, 8.32 for O = 100; N = 175; H = 12.5. The preparation of the salt is not given. (*Erdmann's Journ.* für Prak. Chem., 49, 1850, 446.)

J. Dumas: 96 (O = 16).

Dumas made five experiments on the reduction of molybdic acid (prepared from the natural sulphide) by means of hydrogen. The reduction was begun at a low temperature in a glass tube, and completed in an unglazed porcelain tube in a reverberatory furnace, where it was kept till several hours heating produced no further alteration in weight. The molybdenum did not assume a metallic appearance. The number is the mean; extreme difference, 0.8 for O = 16. (Annal. de Chim. et de Phys., (3,) 55, 1859, 142.)

M. Delafontaine: 92 (O = 16); 575 (O = 100).

This chemist made many experiments in various ways without being able to reach constant results, and only remarks that his experiments indicate Svanberg and Struve's value as the best. (Erdmann's Journ. für Prak. Chem., 95, 1865, 137; Bibl. Univ., Arch. des Sciences, 23, 1865.)

H. Debray: 95.94 (O = 16).

Debray made three experiments on the reduction of molybdic acid. The acid was first converted into the red

oxide in platinum, and at a low temperature, and the small portion of the acid volatilized during this operation was caught and determined. The reduction was completed in a porcelain tube at a white heat. Debray gives his results at 48.03; 48.04; and 47.84. [The analytical data, recalculated, give 95.30; 95.55; 95.73; perhaps on account of misprints. Reduction to vacuum would still further reduce the numbers. The acid was purified by sublimation in platinum, conversion into ammonium salt, and regeneration by heat. In two experiments ammoniacal solution of molybdic acid was evaporated in the dark with excess of argentic nitrate, the argentic molybdate dissolved out and the excess of silver determined. Debray found 5.510 acid = 7.657 silver, and 7.236 acid = 10.847 silver. Hence he calculates M=48 and 47.98. [A little calculation shows that the first data are misprinted. They should read 5.11 acid = 7.657 silver. The corrected data give for Ag = 107.93; M = 96.06 and 95.99. The mean of the recalculated analyses is 95.73.] (Paris Comptes Rend., 66, 1868, 732.)

L. Meyer: 96.10 (O = 16).

Calculated from three analyses of the dichloride, two analyses of the tetrachloride, and two analyses of the pentachloride, made by Leichte and Kempe in Meyer's laboratory. The dichloride was analyzed by heating in a current of hydrogen sulphide, and subsequently in a current of hydrogen. Molybdenum disulphide is the residue. The HCl formed was caught in ammonium hydrate and precipitated by argentic nitrate, after the hydrogen sulphide had been driven off by boiling in a flask provided with a condensing drip-tube. The tetra and pentachloride were decomposed with nitric acid, excess of ammonium hydrate was added, and molybdenum trisulphide precipitated with ammonium sulphide. A weighed portion of the dry precipitate was converted into disulphide by heating in a current of hydrogen. The chlorine of the higher chlorides was determined in the filtrate after precipitation of the trisulphide. By comparing the amount of chloride analyzed with the amount of argentic chloride obtained, Meyer finds in mean M = 95.92; extreme difference, 1.87 for O = 15.96. By comparing the amount of disulphide with that of argentic chloride, M = 95.75; extreme difference, 1.35. By comparing the amount of chloride analyzed with the amount of disulphide obtained for one analysis of tetrachloride and two analyses of pentachloride, he gets M = 95.94; extreme difference, 2.15. The general mean is M = 95.86; extreme

difference, 2.15. Ag = 107.66; S = 31.98; Cl = 35.37; O = 15.96. The specific gravities of the chlorides not having been determined, the weighings are not reduced to vacuum. The pentachloride was prepared from M by heating it in a current of Cl entirely free from air. The metal had been freed from oxide by heating in an atmosphere of HCl. By moderate heating of the pentachloride in dry H, and by distilling pentachloride over the product in dry carbon di-oxide, the trichloride is obtained. The trichloride heated in carbon di-oxide is decomposed into tetrachloride and di-chloride, which latter must be purified with warm dilute nitric acid. (Liebig's Ann., 169, 1874, 360, 344.)

NICKEL.

Regnault has determined the specific heat of nickel. It corresponds to an atomic weight of about 59. (Gmelin-Kraut, l. c.)

E. Rothoff: 59.09 (O = 16); 369.333 (O = 100).

Rothoff converted 188 parts of oxide into chloride, a neutral solution of which gave 718.2 parts argentic chloride. If Cl = 221.64, Ag = 1349.66, the value follows. (Berzelius' Lehrbuch, 3, 1221.)

P. BERTHIER.

Lassaigne having announced the atomic weight of nickel at 500, (Schweigger's Jahrbuch, 9, 108,) Berthier re-examined the subject and found Rothoff's number confirmed. (Berzelius' Jahresbericht, 5, 1825, 148; Annal. de Chim. et de Phys., 25, 1824, 148.)

ERDMANN AND MARCHAND: 58.2 (O = 16); 365.9 (O = 100).

Determined "with all precaution" by the reduction of the oxide with hydrogen. The results varied from 29.1 to 29.8, but Erdmann has reason to believe the smaller number the more accurate. (Erdmann's Journ. für Prak. Chem., 65, 1852, 202.)

87

H. SAINTE-CLAIRE DEVILLE:

100 parts fused nickel, containing three-tenths per cent. silicon and one-tenth per cent. copper, gave 262 parts anhydrous, yellow nickel sulphate, "corresponding to the atomic weight as given by Berzelius." (Annal. de Chim. et de Phys., (3,) 46, 1856, 182.)

R. Schneider: 58.05 (O = 16); 362.8 (O = 100).

Determined from four analyses of the oxalate. The carbon determinations were made by the ordinary method of organic analysis, because some hydrocarbon forms when the salt is decomposed by heat alone. The metal was determined by heating a known weight of the salt first in air and then in a current of oxygen, and subsequent reduction by hydrogen. In the preparation of the salt the usual precipitate with ammonium sulphide was washed with dilute chlorhydric acid, and the cobalt separated with barium carbonate and chlorine. From the nickel salt obtained the oxalate was precipitated with oxalic acid. The number is the mean of four analyses; extreme difference, 0.082 for O = 8. (Poggend. Ann. 101, 1857, 396.)

C. Marignac: 59 (O = 16).

Marignac made two analyses of the sulphate by decomposing the salt by heat. The decomposition is perfect. To avoid errors arising from possible reduction of a portion of the oxide, it was moistened with nitric acid, and recalcined at a moderate temperature. The results obtained were Ni = 29.2 and 29.5. The sulphate was purified by recrystallization. He also made experiments on the chloride by titration with argentic nitrate, according to Pelouze's modification of Gay-Lussac's method. Three such analyses gave from 29.4 to 29.5. In one experiment he also evaporated the nickel nitrate, after filtering off the argentic chloride, and converted it into oxide by heat. This experiment gave Ni = 29.64. The chloride, whether it be distilled or calcined with ammonium chloride, is apt to leave an insoluble residue the weight of which must be deducted. S = 16; Ag = 108; Cl = 35.5. (Bibl. Univ. Arch. des Sciences, (2,) 1, 1858, 375.)

J. Dumas: 59.028 (O = 16).

The number is the mean result of five experiments on the titration of the chloride with argentic nitrate; extreme difference 0.08. Ag = 108; Cl = 35.5. In three cases the nickel chloride was prepared by dissolving fused nickel in aquia regia, repeated evaporation to dryness with HCl, and heating for from twelve to twenty-four hours in a current of HCl gas. In two cases it was produced by passing a current of chlorine over spongy nickel. The chloride analyzed was crystalline and volatile without residue. (Annal. de Chim. et de Phys., (3,) 55, 1859, 149.)

R. Schneider: 58.058 (O = 16).

In consequence of Marignac's criticism (that as nickel oxalate is insoluble it cannot be purified by recrystallization) Schneider repeated his former determination, making special tests for oxalic acid, sodium, and chlorine, with the above result. (*Poggend. Ann.*, 107, 1859, 616.)

W. J. Russell: 58.738 (O = 16).

Determined from the mean of thirteen experiments on the reduction of the oxide in hydrogen. Extreme difference, 0.12 for O = 16. The oxide was prepared from three specimens of commercial nickel, which were first converted into pure oxalate and then into nitrate. The oxide was obtained by decomposing the nitrate by a very strong heat. (Journ. Chem. Soc., (2,) 1, 1863, 61.)

Schneider remarks that a portion of the oxide analyzed may have been reduced during the process of decomposing the nitrate. (*Poggend. Ann.*, 130, 1867, 310.) Marignac points out the same danger. (*Bibl. Univ.*, Arch. des Sciences, (2,) 1, 374.)

E. von Sommaruga: 58.026 (O =16).

Determined from the amount of barium sulphate obtained by precipitating the double sulphate of nickel and potassium with barium chloride. The number is the mean of six experiments; extreme difference, 0.168 for O = 8, S = 16; Ba [no doubt] = 68.5; K = 89.2. The salt was prepared by solution of commercial nickel in sulphuric and nitric acid, adding potassic sulphate to the solution, and repeatedly recrystallizing the double sulphate. (Erdmann's Journ. für Prak. Chem., 100, 1867, 115; Sitz.-Ber. der k. k. Akad., 1866.)

C. Winkler: 59.05 (O = 16).

Determined by the amount of gold precipitated from a solution of neutral crystallized potassium chloro-aurate by

89

a known weight of nickel. The number is the mean of four experiments; extreme difference, 0.186 for O=16, Au=196. The nickel was prepared as follows: commercial nickel carbonate was dissolved in chlorhydric acid, cobalt was repeatedly precipitated with sodium hypochlorite, copper, etc., were removed with hydrogen sulphide, the nickel was precipitated with sodium carbonate, the precipitate dissolved in chlorhydric acid, the chloride sublimed and reduced in a current of hydrogen. (Fresenius' Zeitsch., 6, 1867, 22.)

W. J. Russell: 58.76 (O = 16).

Determined by the amount of hydrogen set free by solution of nickel in chlorhydric acid. The nickel was that obtained in Russell's former determination of the atomic weight of nickel. (Chem. News, 20, 1869, 20.)

R. H. Lee: 58.01 (O = 16).

Determined by analyses of nickel cyanide salts. They were decomposed in a platinum crucible by heat from above. The carbon separated out was burned off first in air and then in oxygen. The metallic oxide was reduced in a current of hydrogen. The mean of six experiments on the strychnine salt gave Ni = 58.04. The mean of six experiments on the brucine salt gave Ni = 57.98. The salts were purified by recrystallization. (Berlin. Bericht der Chem. Ges., 4, 1871, 790.)

NIOBIUM.

The vapor density of the chloride and of the oxychloride, as determined by Deville and Troost, places the atomic weight at about 94. (*Paris Comptes Rend.*, 56, 1863, 891.)

H. Rose: 122 (O = 16).

Rose deduced the atomic weight of niobium from analyses of what he supposed to be the tetrachloride, determining the niobium as niobic acid, and the chlorine as argentic chloride. The results, which varied greatly, indicated the value 97.64. [Marignac having proved that the salt is a pentachloride, this number becomes 122.] Marignac showed

that Rose dealt with a compound containing a large amount of the corresponding tantalium chloride. (*Poggend. Ann.*, 104, 1858, 439.)

Rose; Rammelsberg: 94 (O = 16).

Rose analysed the oxychloride, but did not recognize it as an oxychloride. Rammelsberg calculated the atomic weight from Rose's figures and found that the highest chlorine contents corresponds to an atomic weight of 94. Rose's salt must have been nearly pure as there is no corresponding tantalium compound. (Poggend. Ann., 136, 1869, 353.)

R. Hermann: 104.8 (O = 16).

Hermann deduces this value from analyses of a number of chlorides and sodium salts. The formulas which he gives these compounds are complicated, unlikely, and unsupported by evidence. Marignac has shown that Hermann's salts contained tantalium. (Erdmann's Journ. für Prak. Chem., 68, 1856, 73.)

O. W. Blomstrand: 95 (O = 16).

Blomstrand made three determinations of the chlorine contents of the pentachloride, getting 64.712 per cent., extreme difference, 0.32. He also made eleven determinations of the niobium in the same compound, weighing it as niobic acid. 100 chloride gave in mean 49.794 acid. The atomic weight calculated from the chlorine contents is 96.67; from the niobic acid, 96.16. Blomstrand also made experiments on sodium niobate which led him to the conclusion that the most probable number is 95. (Gmelin-Kraut, 2, part 2, 78; Acta Univ. Lund., 1864.)

C. Marignao: 94 (O = 16).

Determined from a number of analyses of potassium fluoniobate containing two atoms of potassium. The compound was decomposed by sulphuric acid with which it was evaporated to dryness. The residue was leached with water, the filtrate evaporated and the potassic sulphate melted and weighed. The sulphuric acid remaining with the niobic acid was driven off by heat and the acid weighed. The salt being readily soluble and crystallizing well, can easily be purified from all substances except titanium which Marignac knows no way of separating or determining.

The larger the amount of titanium present, the lower will the atomic weight be; Marignac therefore takes the highest value. (Liebig's Ann., S. 4, 334, 288, 338; Bibl. Univ., Arch. des Sciences, 23, 1865, 25, 1866.)

NITROGEN.

Regnault has determined the specific gravity of nitrogen. It indicates an atomic weight slightly above 14. (Gmelin-Kraut, l. c.)

BIOT and ARAGO; WOLLASTON: 14.03 (O = 16); 87.7 (O = 100).

Biot and Arago found the specific gravities of N and H 0.96918 and 0.07321. If H = 13.27 the value follows. [This very accurate value is of course the result of two compensating errors.] (*Phil. Trans.*, 104, 1814, 20.)

J. J. Berzelius; 14.163 (O = 16); 88.518 (O = 100).

Calculated from the specific gravity as determined by Berzelius and Dulong, compared with that of oxygen. By decomposing the nitrate of lead by heat, Berzelius also found N = 88.61 for Pb = 1294.498. (Poggend. Ann., 8, 1826, 14.)

E. Turner: 14.15 (O = 16).

Determined by experiments on the nitrates of lead, silver, and barium, which were precipitated with sulphuric and hydrochloric acids, and gave respectively N = 14.201; 14.09; 14.17; if Pb = 103.6; Ba = 68.7; Cl = 35.42; S = 16.085; the weighings being reduced to vacuum. The salts were purified by recrystallization. Turner recommends more direct methods. (*Phil. Trans.*, 123, 1838, 537.)

T. Thomson: 14 (O = 16).

From the hypothesis that air is a compound containing four parts of N and one part oxygen, and from the average of various selected determinations of the specific gravity of oxygen, Thomson concludes the specific gravity of oxygen is 1.1111, and that of N 0.9722. These numbers stand

to one another as 16 to 14. (Erdmann's Journ. für Prak. Chem., 8, 1836, 375; Records of General Science, by R. D. Thomson, 1836, 179.)

F. Penny: 14.018 (O = 16).

From the mean of three series of experiments (vide Penny's determination of potassium) it follows that 100 potassic chloride = 135.636 potassic nitrate. Penny found the molecular weight of KCl = 74.527; hence the difference between a chloride and a nitrate is 26.560. Similar experiments were made on the sodium salts. In four experiments 100 sodium chlorate were found = 54.930 chloride; extreme difference, 0.02. In three experiments, 100 sodium chlorate were found = 79.882 sodium nitrate; extreme difference, 0.015. In six experiments 100 sodium nitrate were found = 68.771 chloride; extreme difference, 0.013. seven experiments 100 chloride were found = 145.416sodium nitrate; extreme difference, 0.016. These data give sodium chloride = 58.5, and the nitrate = 85.068, or the difference between a chloride and a nitrate = 26.568. Penny found Cl = 35.454. If $NO_3 - Cl = 26.564$, N =14.018. Weighings for vacuum. (Phil. Trans., 129, 1839, 25.)

L. Svanberg: 13.91 (O = 16).

Determined by four experiments on the decomposition of plumbic nitrate by heat which gave a mean of 67.4016 per cent. oxide; extreme difference, 0.0087. [If Pb = 206.926 (Stas) the value follows.] (Berzelius' Jahresbericht, 22, 1842, 38.)

C. Marignac: 14.02 (O = 16); 87.625 (O = 100).

Marignac made five experiments by dissolving a known weight of silver in nitric acid and melting and weighing the nitrate formed. The silver carried out of the retort by the vapors was precipitated and determined. The mean result was that 100 silver = 157.430 nitrate; extreme difference, 0.046; or, if Ag = 1349.01, N = 87.585. Six experiments were made by the decomposition of a known weight of argentic nitrate with a known weight of potassic chloride by Pelouze's method. Mean, 100 KCl = 227.986 argentic nitrate; extreme difference, 0.18. This gives N = 87.685 if K = 488.94 and Cl = 443.2. Seven experiments by the same method showed that 100 silver dissolved in nitric acid = 49.522 ammonium chloride; extreme differ-

ence, 0.063; Hence N = 87.655. The weighings are reduced to vacuum. (Berzelius' Jahresbericht, 24, 1842, 44; Bibl. Univ. de Genève, 46, 1842, 363.)

T. Anderson: 13.95 (O = 16); 87.204 (O = 100).

Determined by four experiments on the decomposition of plumbic nitrate by heat at a sufficiently low temperature to permit of complete decomposition. The number is the mean; extreme difference, 0.198 for O = 100. Pb = 1294.5. (Annal. de Chim. et de Phys., (3,) 9, 1848, 254.)

J. Pelouze: 14.014 (O = 16); 87.59 (O = 100).

A known weight of argentic nitrate was brought in contact with a known and slightly excessive weight of ammonium chloride and the excess titrated with silver solution. One experiment gave N = 175.58; a second gave N = 174.78. Ag = 1349.01; Cl = 443.2. The ammonium chloride was purified by sublimation and recrystallization. (Paris Comptes Rend., 20, 1845, 1047.)

P. EINBRODT: 14 (O = 16); 87.5 (O = 100).

Experiments on the decomposition of plumbic nitrate by heat gave N = 87.5 plus a vanishing quantity if Pb = 1294.2239. (Leibig's Ann., 70, 1849, 286.)

J. Dumas: 14 (O = 16).

Determined by experiments on the combustion of ammonia and cyanogen. Particulars not given. C = 6; H = 1. (Annal. de Chim. et de Phys., (3,) 55, 1859, 134.)

J. S. Stas: 14.044 (O = 16).

Stas made seven determinations of the relation between silver and its nitrate by dissolving pure silver in nitric acid, evaporating to dryness and keeping the salt melted until there was no further loss of weight. In two of these experiments the salt was melted in vacuo. The mean result was that 100 Ag = 157.472 nitrate; whence N = 14.040. Later Stas made two more experiments by the same method with all possible precautions to secure accuracy. These gave 100 Ag = 157.484 nitrate and N = 14.042. By the conversion of the chlorides of potassium, sodium, lithium and silver into nitrates, Stas found the difference between a chloride and a nitrate 26.5882. This gives N = 14.045. The weigh-

ings are reduced to vacuum. Cl = 35.457; Ag = 107.93. (Stas, Unters. über Chem. Prop. Leipzig, 1867.)

OSMIUM.

Regnault has determined the specific heat of osmium. It corresponds to an atomic weight of about 199. (Gmelin-Kraut, l. c.)

J. J. Berzelius: 199.04 (O = 16).

Berzelius analyzed potassium chloro-osmate by reduction in a current of hydrogen and solution of the potassium chloride from the residue. 1.3165 grammes of the double salt lost 0.8805 in reduction and the residue was composed of 0.401 potassium chloride and 0.585 osmium. The atomic weight may be calculated either from the chlorine lost or from the relation of the chloride to the metal in the residue. Berzelius preferred the latter as more probably accurate. [If K = 39.137; Cl = 35.457 (Stas;) this relation gives 199.04.] According to W. M. Watts, (Chem. News, 19, 802) the loss of chlorine gives for Stas's values 0s = 199.42. Hyperosmic acid was separated from iridium compounds by distilling at a gentle heat. The first portion is perfectly pure. The metal was precipitated from chlorhydric acid solution of hyperosmic acid by mercury and subsequently purified by heating in a current of hydrogen. Potassium chloro-osmate was formed by heating comminuted metal and KCl in a current of chlorine. (Poggend. Ann., 13, 1828, 580; Kongl. Vet. Acad. Handl., 1828.)

E. Fremy: 199.65 (O = 16); 1247.8 (O = 100).

Pure osmium was burned in a current of oxygen and the fumes led over potassic hydrate, by which they are absorbed. An additional potash tube did not increase in weight. Corks were avoided. Number of experiments not given. (Erdmann's Journ. für Prak. Chem., 33, 1844, 409; Journ. de Pharm. et Chim., 1844, 241.)

DEVILLE and DEBRAY: 198 (O = 16).

These chemists determined the vapor density of hyperosmic acid by Dumas' method, finding it 8.89 at 246°,

and 8.87 at 286°. They hence consider it probable that the atomic weight of osmium is the same as that of platinum. The acid was very pure and was prepared by the combustion of metallic osmium in oxygen. (Paris, Comptes Rend., 44, 1857, 1101.)

OXYGEN.

The atomic weight of oxygen is assumed at 16 for the reasons stated under hydrogen, q. v. If hydrogen is taken as unity, 0 = 15.96.

PALLADIUM.

Regnault determined the specific heat of palladium. It corresponds to an atomic weight of about 106. (Gmelin-Kraut, l. c.)

J. J. Berzelius; 106.51 (O = 16).

In his earliest determinations of this constant, Berzelius saturated the metal with sulphur, getting about 711 for S = 201.165; and decomposed the chloride with mercury, getting 704. [711 appears to be a misprint for 714.618 the number given with corresponding data at Poggend., 8, 180.] In this investigation a known weight of potassium chloropalladate was reduced in a current of hydrogen, the weight of the residue determined, the potassium chloride leached from the residue and the metallic palladium weighed. The double salt was strongly heated, but not to fusion, in a current of dry air before weighing. It being impossible to desiccate this and the similar platinum-metal salts completely without decomposition, the atomic weight was calculated from the relation between the metal and the KCl. Berzelius found 0.575 Pd = 0.809 KCl, and 0.851 Pd = 1.192 KCl. [If KCl] = 74.594 (Stas) the former gives Pd = 106.036, the latter 106.509.7 Berzelius had reason to consider the latter analysis the more accurate. (Poggend. Ann., 13, 1828, 454; Kongl. Vet. Acad. Handl., 1828.)

PHOSPHORUS.

The specific heat of this element, as well as the density of phosphorus and its numerous volatile compounds in the gaseous state, corresponds to an atomic weight slightly above 31. (Gmelin-Kraut, l. c.)

V. Rose; F. H. Wollaston: 35.1 (O = 16).

Wollaston adopted the analysis of Rose, who found that phosphoric anhydride contained 53.28 per cent. oxygen and 46.72 per cent. phosphorus. [This relation gives the above value.] (*Phil. Trans.*, 104, 1814, 20.)

J. J. Berzelius: 31.325 (O = 16).

Berzelius made experiments on the reduction of auric chloride and of argentic sulphate by phosphorus. His results were 0.8115 P = 13.98 Ag; 0.829 P = 8.714 Au; 0.754 P = 7.93 Au. [The first of these analyses is misprinted in the original memoir (Gilbert's Ann., 53, 433).] In the Lehrbuch it is miscalculated as Ruecker has shown. zelius preferred deducing the atomic weight of P from that of silver, because the atomic weight of the latter was more accurately known than that of gold. [If Ag = 107.93, the data give P = 31.825, for Au = 196.67 the latter analyses give P = 31.176 and 81.165.] In all the experiments the precipitated metal was boiled with the solution when the reduction was nearly complete. A trace of gold was observed to precipitate after the experiments were over. The silver was heated to redness before weighing. Cooke, Jr., has shown (atomic weight of antimony) that silver is volatile at a red heat. Berzelius must therefore have got too large a result.] The phosphorus was distilled, melted in a glass tube and cooled very slowly, to permit traces of oxides to rise to the surface, and the lower portion of the tube with the phosphorus broken off and instantly (Gilbert's Ann., 53, 1816, 433, and Lehrbuch 3, weighed. 1188.)

J. Pelouze: 32.024 (O = 16); 200.15 (O = 100).

A known weight of argentic nitrate was brought in contact with a known and slightly excessive weight of phosphorous chloride and the excess titrated. The number of experiments is not given. Ag = 1349.01; Cl = 443.2.

The terchloride was prepared by chloridizing finely divided P with dry chlorine, adding finely divided P, decanting, agitation with tin amalgam and rectification over the same. The fluid was colorless and did not give any precipitate with water. (Paris, Comptes Rend., 20, 1845, 1047.)

 ∇ . A. JACQUELIN: 29.83 (O = 16); 186.438 (O = 100).

Determined by experiments on the chlorides of phosphorus with argentic nitrate and plumbic oxide. The results are utterly discordant. (*Paris, Comptes Rend.*, 33, 1851, 698.)

A. Schroetter: 31.0274 (O = 16).

Determined by burning perfectly pure amorphous phosphorus in dry oxygen and weighing the phosphoric anhydride. The number is the mean of 10 experiments; extreme difference, 0.1242. Previous to burning, the phosphorus was heated for a long time in carbon di-oxide or hydrogen. It was burned not in powder but in lumps. (Erdmann's Journ. für Prak. Chem., 53, 1851, 435; Sitz.-Bericht der k. k. Akad., 1851.)

B. C. Brodie: 81.81 (O = 16).

Three experiments made by oxidation of phosphorus with aqua regia and determination as magnesium pyrophosphate gave this mean. Brodie seems to regard these determinations only as evidence that the atomic weight needs redetermination. (Journ. Chem. Soc., 5, 1852, 295.)

J. Dumas: 31.03 (O = 16).

Determined by five experiments on the titration of the terchloride with argentic nitrate. The chloride was prepared by the action of dry chlorine on amorphous phosphorus and distillation after the chlorine had been displaced by carbon di-oxide. The portion distilling between 76° and 78° only was used. The number is the mean of the results; extreme difference, 0.08. Ag = 108; Cl = 35.5. (Annal. de Chim. et de Phys., (3,) 55, 1859, 172.)

PLATINUM.

Regnault and Kopp have determined the specific heat of platinum. It answers to an atomic weight of about 197. (Gmelin-Kraut, l. c.)

J. J. Berzelius: 197.19 (O = 16).

Determined by the same method as osmium, q. v., from a single experiment on potassium auroplatinate. 2.135 potassium chloride accompanied 2.822 platinum. [If KCl = 74.594 (Stas,) this gives the above value.] The salt was prepared by precipitating an alcoholic solution of platinum chloride with potassium chloride, washing with alcohol and heating to redness in a current of chlorine. Berzelius remarks that the metal used in his former determinations was impure. (Poggend. Ann., 13, 1828, 468, and Lehrbuch, 3, 1213.)

T. Andrews: 197.88 (O = 16).

Determined by three experiments on potassium chloroplatinate. The salt was dried at 105° in vacuo, decomposed by zinc, the excess of zinc removed by acetic acid, the solution filtered off, and the chlorine titrated. The number is the mean; extreme difference, 0.22. The values assumed for Ag and Cl are not given. They were most likely Marignac's. (Brit. Assoc. Rep., 1852, part 2, 33.)

J. S. Stas made preparations for determining the atomic weight of platinum, but not being able to produce potassium chloroplatinate entirely free from water, and being unacquainted with Bunsen's method of accomplishing this end, resigned the attempt. He made, indeed, three analyses by the same method employed by Berzelius, but unfortunately does not communicate the results. (Stas, Untersuch. über Chem. Prop., Leipzig, 1867, 265.)

POTASSIUM.

Regnault determined the specific heat of potassium. It corresponds to an atomic weight of about 39. (Gmelin-Kraut, l. c.)

M. H. KLAPROTH; F. H. WOLLASTON: 39.517 (O = 16).

Klaproth found that 441 Cl = 591 potassium oxide. Hence Wollaston deduced the value 491 (O = 100) for K. [If Cl = 35.457, this relation gives K = 39.517.] (Phil. Trans., 104, 1814, 20.)

J. J. Berzelius: 39.193 (O = 16); 244.958 (O = 100).

Berzelius found that 100 KCl = 192.4 Ag Cl. If Ag = 1351.607; Cl = 442.65; the above value follows. (*Poggend. Ann.*, 8, 1826, 190.)

F. Penny: 39.073 (O = 16).

Penny made six experiments on the conversion of the chlorate into the chloride. Potassic chlorate was dried at about 105°, dissolved in a flask with HCl, evaporated, dried and weighed. The cake contained some free HCl. It was broken up, pulverized, and a known quantity heated to dull redness but not to fusion, and the HCl expelled allowed for. The mean result was that 100 KCl O₂ = 60.823 KCl; extreme difference, 0.015. This relation gives KCl = 74.527and if Cl = 35.454 (Penny,) the value for K follows. Numerous experiments were also made on the introconversion of the nitrate, the chloride and the chlorate, which established the difference between a chloride and a nitrate, besides confirming the value of K. The salts were purified by recrystallization and were carefully tested for impurities. The weighings are all for vacuum. (Phil. Trans., 129, 1839, 18.)

C. Marignao: 39.2 (O = 16); 245 (O = 100).

By six experiments on the decomposition of the chlorate by heat, 100 chlorate were found to lose 39.161 oxygen; extreme difference 0.012; hence KCl = 932.14. By comparing this value with the molecular weight and the composition of argentic chloride, Cl was calculated at 442.13, leaving for K the number 490. Confirmatory experiments were made on potassic perchloride. The chlorate was purified by recrystallization. The weighings are for vacuum. (Liebig's Ann., 44, 1842, 23.)

C. Marignac: 39.115 (O = 16); 244.47 (O = 100).

Having determined the atomic weight of chlorine from syntheses of argentic chloride, and found it 443.2, the molecular weight of KCl in the last determination, gives K = 244.47, for vacuum. Berzelius, by rejecting some analyses and the correction for vacuum, deduces the value 244.429. (Berzelius' Jahresbericht, 25, 1845, 31; Bibl. Univ. de Genève, 46, 1842, 350.)

J. Pelouze: 39.144 (O = 16); 244.65 (O = 100).

A known weight of KCl was brought into contact with a known amount of silver dissolved in nitric acid, the chloride being slightly in excess. This excess was titrated with a decimal solution of silver. The number is the mean of three experiments. Ag = 1349.01; Cl = 443.2. The chloride was prepared by heating the chlorate and recrystallizing the residue. (*Paris Comptes Rend.*, 20, 1845, 1047.)

According to Pelouze, Levol found the molecular weight of KCl 466.245, which combined with Marignac's value of Cl would give K = 244.645 or 39.143. (Ibid.)

E. J. MAUMENÉ: 38.96 (O = 16); 243.502 (O = 100.)

The mean of three experiments on the decomposition of KCl with an excess of argentic nitrate showed that 100 KCl = 192.75 AgCl. If Ag = 1850.82 and Cl = 443.67, according to Maumené's determinations, the number follows. The KCl was prepared from the chlorate by heat. To confirm his values for K and Cl, he made seven experiments on the decomposition of the chlorate by heat, and found that 100 chlorate gave 60.791 chloride. An unaccounted for increase in the weight of the flask occurred in these experiments. (Annal. de Chim. et de Phys., (8,) 18, 1846, 41.)

J. S. Stas: 39.187 (O = 16).

According to the mean of seven determinations, 100 parts of KCl dissolved in nitric acid, and evaporated to dryness give 185.6428 parts of nitrate; extreme difference, 0.017. If Cl = 35.457; N = 14.044; the value follows. This value is confirmed by previous experiments which gave 39.130. Potassic chloride, by whatever means it is prepared, still retains silica. Stas, therefore, determined

the amount of silica in the KCl and allowed for it. Weighings for vacuum. (Stas, Untersuch. über Chem. Prop.,

Leipzig, 1867.)

Stas mentions that Dumas, who was the first to place K at 89, afterwards became convinced that this number was too low. (*Ibid*, page 318.)

RADCLIFFE.

RHODIUM.

Regnault has determined the specific heat of rhodium. It corresponds to an atomic weight of about 104. (Gmelin-Kraut, l. c.)

J. J. Berzelius: 104.3 (O = 16).

Berzelius made two analyses of dipotassic chlororhodiate. This salt can be completely desiccated in a current of chlorine at a red heat without decomposition. 3.146 grammes gave on reduction in a current of hydrogen 0.930 Cl, and the residue contained 1.304 KCl and 0.912 metallic rhodium. [If KCl = 74.594, Cl = 35.457, (Stas,) the atomic weight of the salt calculated from the Cl contents is 359.881, and that of Rh 104.272. The relation between the Rh and the Cl gives Rh=104.312. The relation between the KCl and the Rh gives Rh = 104.340. The mean is 104.308.] Berzelius made a second analysis of the crystallized salt in which he determined the water of crystallization. [Under the same suppositions and in the same order, the resulting values for Rh are 106.279; 104.762; 103.708.] In the Lehrbuch only the former analysis is used to deduce the atomic weight. Rhodium was separated from other metals by its insolubility in aqua regia. The double salt was formed by heating finely pulverized Rh in mixture with KCl in a current of chlorine. The double salt was dissolved in water, precipitated with alcohol, washed with alcohol and dried. (Poggend Ann., 13, 1828, 487; Kongl. Vetens. Akad. Handl., 1828.)

In his earlier determination (Rh = 750.68 for O = 100) Berzelius mistook an hydrated oxide for a chloride. (*Ibid.*)

RUBIDIUM.

Kopp determined the specific heat of rubidium chloride. It corresponds to an atomic weight of about 85. (Gmelin-Kraut, l. c.)

Kirchhoff and Bunsen: 85.36 (O = 16).

Determined from the mean of four experiments on the precipitation of the chloride with argentic nitrate. extreme difference was 0.24. Ag = 107.94; Cl = 35.46. An impure mixture of rubidium and potassium chlorides, nearly free from lithium and the earths, was partially precipitated with platinum chloride and the precipitate freed from KCl by repeated boiling with water. The residue was reduced in a current of hydrogen, the rubidium chloride extracted with water, and reprecipitated with platinum chloride. This process was repeated until the potassium lines in the spectrum disappeared. The rubidium was then converted into a mixture of carbonate and oxide, and the caesium separated by extraction with alcohol. The amount of silver precipitated was also tested from time to time and the purification continued till this became constant. (Poggend. Ann., 113, 1861, 339.)

J. PICCARD: 85.41 (O = 16).

Determined by four analyses of rubidium chloride with argentic nitrate. The number is the mean; extreme difference, 0.09. The separation of potassium from rubidium was effected for the different analyses by 6, 7, and 8 successive partial precipitations with platinum chloride, and the separation of caesium by thirty successive extractions of the anhydrous carbonates with warm absolute alcohol. The salt analysed was spectroscopically pure. Ag = 107.94; Cl = 85.46. The experiments were made with Bunsen's assistance. (Erdmann's Journ. für Prak. Chem., 86, 1862, 449.)

L. Grandeau, who is sometimes credited with making a determination of Rb, expressly disclaims doing so. He mentions Bunsen's value as the true atomic weight and says that his analyses of the sulphate, undertaken to test its purity, led him to adopt the number 85.4; apparently for brevity's sake. (Annal. de Chim. et de Phys., (3,) 67, 1863, 227.)

R. Godeffroy: 85.476 (O = 16).

Determined by four analyses of rubidium chloride prepared and analysed exactly as Godeffroy determined cæsium, q. v.; extreme difference, 0.04. Cl = 35.5; Ag = 108. (Liebig's Am., 181, 1877, 189.)

RUTHENIUM.

Bunsen has determined the specific heat of ruthenium. It corresponds to an atomic weight of about 104. (Gmelin-Kraut, l. c.)

C. E. CLAUS: 104.57 (O = 16).

Determined from three analyses of potassium chlororutheniate by the same method Berzelius had employed for other platinum metals. Claus found an average of 28.783 per cent. Ru; extreme difference 0.48, and 41.063 KCl; extreme difference, 0.51. [If K = 39.137, Cl = 35.457; this composition gives Ru = 104.57. The weighings as given in the memoir are misprinted.] Claus also determined the chlorine with silver; the results were such as to show that the salt was not anhydrous, though it had been dried at 200° in an atmosphere of Cl. The salt was prepared by the evaporation of a solution of ruthenium and potassic hydrate in aqua regia, solution of other chlorides of Ru in dilute HCl, and removal of basic compounds by mechanical concentration in water. Claus later takes the atomic weight of Ru = 104. In this memoir he puts it at 651.387 (0 = 100,) 104.22 (O = 16,) without mentioning the values of K and Cl. (Poggend. Ann., 65, 1845, 218.)

SELENIUM.

Regnault determined the specific heat of selenium, which accords with an atomic weight of about 79. (Gmelin-Kraut, l. c.)

J. J. Berzelius: 79.23 (O = 16).

Berzelius found that 100 Se absorb 179 dry chlorine gas, and that the product was exactly decomposed by water into chlorhydric acid and selenious acid. [If Cl = 35.457 (Stas) the value follows.] (*Poggend. Ann.*, 8, 1826, 21.)

F. SACC: 78.55 (O = 16); 490.93 (O \doteq 100).

Sacc's experiments are very discordant. He made three experiments on the reduction of a known weight of selenious acid with ammonium bisulphite and chlorhydric acid. The mean result was Se = 490.38; extreme difference, 5.5. In four experiments barium seleniate was decomposed by heating to redness with sulphuric acid in excess. was found to contain 41.95 selenious acid; extreme difference 0.04. For Ba = 856.877 the resulting value is 491.49. The selenium was purified by solution in nitric acid, evaporation and sublimation, and by reduction with HCl and ammonium bisulphite. Selenious acid was prepared by oxidation with nitric acid. Barium seleniate was prepared by precipitation of barium nitrate with sodium seleniate and heating to redness. Sacc regards 490.3 as the most probable value of Se. (Annal. de Chim. et de Phys., (8,) 31, 1851, 119.)

A. Schroetter: 78.6 (O = 16).

Details not given. (Kopp's Jahresbericht, 4, 1851, 318; Sitz.-Bericht der k. k. Acad., 6, 1851, 214.)

ERDMANN AND MARCHAND: 78.6 (O = 16); 492.5 (O = 100).

Determined from experiments on mercuric selenide by the same methods employed for the determination of S, q. v. Three experiments gave 71.726, 71.731, 71.741, per cent. mercury. (Erdmann's Journ. für Prak. Chem., 55, 1852, 202.)

J. Dumas: 76.46 (O = 16).

Determined by seven experiments on the chloridation of selenium. The chloride was condensed in a tube cooled to -20° , after which the escaping gases were led through other tubes filled with asbestos. The extreme difference in the results was 0.46. Cl = 35.5. (Annal. de Chim. et de Phys., (3,) 55, 1859, 129.)

O. Pettersson and G. Ekman: 79.08 (O = 16).

Determined by five analyses of selenious acid. A warm solution of the acid was acidified with chlorhydric acid, precipitated with sulphurous acid and the selenium collected on a glass filter. Many precautions are necessary in the precipitation and drying. The value is the mean; extreme difference, 0.04. (Berlin, Bericht der Chem. Gesell., 9, 1876, 1212; in extenso in the Acta of the Scientific Soc. of Upsala.)

SILICON.

The vexed question of the composition of silicic acid has been settled by H. F. Weber, who found that the specific heat of this element becomes nearly constant above 200° and that the atomic heat is 5.8 for Si = 28. (Poggend. Ann., 154, 1875, 575.)

J. J. Berzelius: 29.63 (O = 16); 185.19 (O = 100).

100 parts of silicon, which had been heated to redness, and freed from silicic acid by hydrofluoric acid, gave 208 parts silicic acid, whence the value. Berzelius also made analyses of barium fluosilicide from which he calculated the oxygen contents of the acid at 51.975. This gives for the atomic weight of Si 29.58. (Poggend. Ann., 8, 1826, 20; and Lehrbuch, 3, 1200.)

J. Pelouze: 28.46 (O = 16); 177.88 (O = 100).

A known weight of perfectly pure silver, dissolved in nitric acid, was brought in contact with a known and slightly excessive weight of silicon tetrachloride and the excess titrated with decimal silver solution. The value is derived from the mean of two experiments; difference 0.76 for O = 100; Cl = 448.2, Ag = 1849.01. The chloride was prepared by Ebelmen; it was perfectly transparent, volatilized without residue, and had been dried for a long time in a vacuum. (Paris, Comptes Rend., 20, 1845, 1047.)

J. Dumas: 28.02 (O = 16).

Determined from the mean of two experiments on the tetrachloride which was weighed off in a glass bulb and

introduced, so enclosed, into a vessel centaining water. The bulb was broken and the chlorine contents titrated with argentic nitrate. The difference between the experiments was 0.2 for O = 16, Ag = 108, Cl = 35.5. The chloride was repeatedly rectified; its boiling point was 59° . (Annal. de Chim. et de Phys., (3,) 55, 1859, 129.)

J. Schiel: 28.01 (O = 16).

Determined by two analyses of the tetrachloride. The salt was decomposed with a slight excess of ammonium hydrate and the chlorine titrated with argentic nitrate. The atomic weights of Cl and Ag used are not stated. Schiel found 0.6738 silicic chloride = 2.277 argentic chloride, and 1.3092 silicic chloride = 4.418 argentic chloride. [For Ag = 107.93, Cl = 35.457, these data give Si = 28.13, and 27.89.] (Liebig's Ann., 120, 1861, 94.)

SILVER.

Dulong and Petit, Regnault and others have determined the specific heat of silver and found it in accord with an atomic weight of about 108. (Gmelin-Kraut, l. c.)

MARCET; DAVY; WENZEL; WOLLASTON.

Wollaston in his table of equivalents mentions that Marcet found 441 Cl = 1350 silver, and Davy the same quantity of chlorine = 1360 silver. Wenzel found 200 sulphur = 1360 silver. (*Phil. Trans.*, 104, 1814, 21.)

J. J. Berzelius: 108.129 (O = 16); 675.804 (O = 100).

Berzelius found that 100 silver gave 132.75 argentic chloride. Taking Cl = 221.325 he calculates Ag = 1351.607. He expresses uncertainty whether or no this value should not be reduced to one half. (*Poggend. Ann.*, 8, 1826, 180.)

E. Turner: 108 (O = 16).

Turner determined the composition of argentic chloride at 100 silver to 132.8 chloride. These numbers are for

SILVER. 107

vacuum. If Cl = 35.42 (Turner) the value follows. (Phil. Trans., 123, 1833, 536.)

F. Penny: 107.97 (O = 16).

Penny made six experiments on the conversion of silver into nitrate. The silver was dissolved in cold nitric acid, the solution evaporated, and the nitrate fused all in one flask and with precautions against loss by spiriting. found 100 Ag = 157.441 nitrate; extreme difference, 0.028. In five experiments the nitrate from the preceding determinations was converted into chloride, by means of chlorhydric acid, in the same flask, dried, fused, and weighed. Penny could detect no decomposition in fusion. He found 100 Ag = 132.8372 chloride; extreme difference, 0.01. two experiments silver was dissolved in nitric acid, precipitated with chlorhydric acid, evaporated and fused, giving 132.830 and 132.838. The mean of all seven experiments is 132.836. Penny takes 132.837. From the relations of the chlorides, chlorates, and nitrates of potassium and sodium, Penny had determined the difference between the atomic weights of a chloride and a nitrate at 26.565. gives the molecular weight of argentic chloride at 143.424 and Ag = 107.97. The silver used, as well as the water and the acids, were carefully tested for impurities and a minute amount of solid residue in the twice distilled water and in the acids was allowed for. The weighings were all reduced to vacuum. (Phil. Trans., 129, 1839, 27.)

C. Marignac: 108 (O == 16); 675 (O == 100).

Silver was dissolved in nitric acid and precipitated with chlorhydric acid. One experiment, reduced to vacuum, gave 100 silver = 132.74 chloride, which Marignac considered confirmatory of Berzelius' value, 182.75. He therefore adopted the latter number. 100 potassic chloride were found to produce 192.26 argentic chloride, in two experiments, the difference between which was 0.01. By analysis, by means of heat, of potassic chlorate, Marignac had found the molecular weight of the chloride 932.14, these relations give the molecular weight of argentic chloride at 1792.13 and the atomic weight of silver at 1350. The potassic chloride was prepared by heating the chlorate and cooling the resulting chloride over sulphuric acid. (Liebig's Ann., 44, 1842, 23.)

C. Marignao: 107.922 (O = 16); 674.505 (O = 100).

Marignac redetermined the relation between silver and potassic chloride by Pelouze's method. He found 100 Ag = 69.062 KCl in six experiments, the extreme difference between which was 0.018. In five experiments he found 100 KCl = 192.348 Ag; extreme difference 0.04. He also redetermined the composition of argentic chloride. silver was dissolved in a long-necked flask and the fumes passed into a second flask containing water. Solution being effected, the water from the second flask was added to the contents of the first, and the whole precipitated with HCl. The chloride was washed, dried, melted and weighed in the same flask. The result was 100 Ag = 132.84 chloride; extreme difference 0.019. Combination of these data with Marignac's old value for the molecular weight of KCl, 932.14, gives Ag = 1349.01. All weighings reduced to vacuum. Berzelius revised the result by throwing out one experiment and by rejecting the correction for vacuum. He thus got Ag = 1849.66. (Berzelius' Jahresbericht, 24, 58; 25, 31; Bibl. Univ. de Genève, 46, 1842, 350.)

In opposition to Prout's hypothesis, Marignac cites his analyses of argentic acetate, in which the escaping gases were forced to pass over porous silver. They gave in three experiments 64.664 silver from 100 acetate; extreme difference 0.005. If C = 75, this gives Ag = 1349.6. He also found 100 Ag = 157.455 nitrate. [If N = 87.5, this gives Ag = 1348.88.] He also found 100 Ag = 49.556 ammonium chloride. (*Liebig's Ann.*, 59, 284; *Bibl. Univ. de Genève*, 1846.)

LIEBIG and REDTENBACHER; STRECKER: 107.903 (O = 16); 674.395 (O = 100).

Strecker recalculated Liebig and Redtenbacher's analyses of argentic acetate, tartrate, racemate and malate by the method of least squares, and from the difference in the atomic composition of these salts. He obtained for Ag the value 1348.79. Vide Carbon. (Liebig's Ann., 59, 1846, 280.)

E. J. MAUMENÉ: 108.026 (O = 16); 675.16 (O = 100).

In four experiments argentic oxalate was mixed with sand in a flask and decomposed by heat in a current of air. The

109

products of decomposition were passed over cupric oxide, and through drying tubes and potash tubes. In five experiments the acetate was treated in the same way, but not mixed with sand. The mean result was Ag = 1350.32; extreme difference 0.77. Maumené found it very difficult to purify the oxalate, which showed traces of nitric acid after 100 washings. (Annal. de Chim. et de Phys., (3,) 18, 1846, 41.)

SILVER.

J. S. Stas: 107.93 (O = 16).

Thirteen syntheses of argentic iodide, performed by bringing hydroiodic acid in contact with argentic sulphate or nitrate, gave 100 Ag = 117.5848 iodine. Three analyses of argentic iodate, performed by decomposition by heat in a current of nitrogen or by reduction of the salt, while in suspension, by a current of sulphurous anhydride, gave Hence Ag = 107.928. Agl = 234.779.Four syntheses of the bromide, performed by bringing hydrobromic acid in contact with argentic sulphate, gave 100 Ag = 74.0805 Br. Two analyses of argentic bromate, by reduction while in suspension with sulphurous anhydride, gave Ag Br = 187.87. Hence, Ag = 107.921. Seven syntheses of argentic chloride, three of them by combustion of silver in chlorine, three by precipitation with HCl, and one by precipitation with ammonium chloride, gave 100 Ag = 32.8445 Cl.Stas adopts the number 32.85 on the supposition that no excess of chlorine was possible. The chloride was fused. Two analyses of the chlorate, accomplished by heat or by evaporation with chlorhydric acid, gave Ag Cl = 143.395. Hence Ag = 107.987. Five syntheses of the sulphide, performed by heating silver in a current of sulphur vapor, or of hydrogen sulphide, gave 100 Ag = 114.8522 argentic sulphide. Six analyses of the sulphate by reduction in a current of hydrogen, showed that 100 sulphate contained 69.203 silver, hence Ag = 107.920, [107.926? vide Sulphur.] From analysis of potassium chlorate, Stas had determined the molecular weight of KCl at 74.59. By twenty-four determinations he found 100 Ag = 69.103 KCl, hence Ag = 107.943. The silver was prepared either by Levol's method or by decomposing an ammoniacal solution of argentic nitrate with a mixture of ammonium sulphite and a copper salt. The metal was heated to the boiling point until the sodium line disappeared and the metallic fumes were a pale blue. To test its purity, it was compared with distilled silver. See Stas's determinations of Cl, Br, I, S,

and K. All weighings reduced to vacuum. (Stas, Untersuch. über Chem. Prop., Leipzig, 1867.)

SODIUM.

The specific heat of sodium has been determined by Regnault and indicates an atomic weight of about 23. (Gmelin-Kraut, l. c.)

H. DAVY; F. H. WOLLASTON: 23.28 (O = 16); 145.5 (O = 100).

Davy found that 134 Cl combine with 88 Na to form sodium chloride. If Cl = 441, the value follows. (*Phil. Trans.*, 104, 1814, 20.)

J. J. Berzelius: 23.164 (O = 16).

Berzelius found that 100 Na Cl = 244.6 Ag Cl. [If Ag Cl = 143.387, (Stas,) the value follows.] (*Poggend. Ann.*, 8, 1826, 189.)

F. Penny: 23.046 (O = 16).

Penny made four experiments on the conversion of the chlorate into the chloride by means of HCl. A known weight of the salt was dissolved in a flask in the acid and evaporated, dried and weighed without removal. The sodium chloride was not fused. The mean result was that 100 chlorate equals 54.930 chloride; extreme difference, 0.02. This relation gives the molecular weight of the chloride at 58.5. Penny had found the atomic weight of Cl = 35.454; hence the value for Na. [If Cl = 35.457 (Stas,) Na = 23.043. Stas himself found 23.043.] The sodium chlorate was prepared by precipitating potassium chlorate with sodium bitartrate, and purifying the sodium chlorate by recrystallization. The weighings are for vacuum. (Phil. Trans., 129, 1839, 25.)

J. Pelouze: 22.97 (O = 16); 143.59 (O = 100).

A known weight of perfectly pure silver was dissolved in nitric acid, and brought in contact with a known and slightly excessive weight of sodium chloride, and the excess titrated with decimal silver solution. The mean result of three experiments was that 100 Ag = 51.141 Na Cl; extreme difference, 0.033. The value follows for Ag = 1349.01; Cl = 443.2. The sodium chloride was prepared either from sodium sulphate and barium chloride, or from sodium carbonate and chlorhydric acid, or from a very pure rock salt. It was repeatedly recrystallized and was dried at 200° or melted. (Paris Comptes Rend., 20, 1845, 1047.)

J. Dumas: 23.011 (O = 16).

Determined from the mean of seven experiments on the titration of sodium chloride with argentic nitrate; extreme difference, 0.09. Ag = 108; Cl = 35.5 [Dumas gives the mean as 23.014 instead of 23.0114.] For five experiments Na Cl recrystallized ten times and melted was employed. For two experiments (giving an average of 23.036) the residue from the incineration of the acetate was used to prepare Na Cl, which was recrystallized four times and melted. (Annal. de Chim. et de Phys., (3,) 55, 1859, 129.)

J. S. Stas: 23.043 (O = 16).

According to the mean of 10 determinations 100 Ag = 54.2078 Na Cl; extreme difference 0.0038. The sodium chloride was found to contain a minute quantity of silicic acid which reduces the result from Na = 28.049 to 23.045 for Ag = 107.93; Cl = 35.457. According to the mean of five determinations 100 Na Cl = 145.4526 sodium nitrate; extreme difference 0.025. If N = 14.044 this gives Na = 23.045. The lowest determination gives Na = 23.042. The sodium chloride was purified by recrystallization and in part by conversion into sodium chloroplatinate. The weighings are for vacuum. (Stas, Untersuch. über Chem. Prop., Leipzig, 1867.)

STRONTIUM.

Regnault determined the specific heat of strontium chloride. It corresponds to an atomic weight of about 87.5. (Gmelin-Kraut, l. c.)

M. H. KLAPROTH; F. H. WOLLASTON: 94.4 (O = 16); 590 (O = 100).

Klaproth found 42 sulphuric anhydride = 58 strontium oxide; whence the value for S = 200. (*Phil. Trans.*, 104, 1814, 20.)

F. Stromeyer; 87.34 (O = 16); 545.929 (O = 100).

According to Berzelius, Stromeyer found that 100 strontium chloride = 181.25 argentic chloride; whence the value, for Ag = 1849.66; Cl = 221.64. (Berzelius' Lehrbuch, 3, 1229.) In Gilbert's Ann., 54, 1816, 251, Stromeyer refers to this analysis as by V. Rose. Stromeyer himself found 0.5 grm. carbonate = 75.5394 c. c. carbon di-oxide [which gives Sr = 88.26 if 1000 c. c. carbon di-oxide weigh 1.96483 grm.] Stromeyer calculated Sr = 552.28 for O = 100.

—. SALVETAT: 88 (O = 16); 550 (O = 100).

Determined from the loss of weight of strontium carbonate by calcination and on driving off carbon di-oxide with sulphuric acid. Details not given. (*Paris Comptes Rend.*, 17, 1848, 318.)

J. Pelouze: 87.68 (O = 16); 548.02 (O = 100).

A known weight of perfectly pure silver was brought in contact with a known and slightly excessive amount of strontium chloride and the excess titrated with decimal silver solution. The number is the mean of two experiments; extreme difference, 0.2. Ag = 1849.01; Cl = 443.2. The chloride was purified by recrystallization and was dried at 200° or below redness. (Paris Comptes Rend., 20, 1047.)

C. Marignao: 87.54 (O = 16).

Marignac made experiments on three different preparations of strontium chloride, (1,) (2,) (8.) Compared with silver by Pelouze's method it was found that ten grammes strontium chloride = (1) 8.103; (2) 8.099; (3) 8.101 silver. The same strontium chloride converted into sulphate gave (1) 6.887; (2) 6.8855; (3) 6.884 sulphate. In both these series of experiments the strontium was weighed as airdried, hydrous, crystalline chloride. Comparison gives Sr

=(1) 43.79; (2) 43.82; (8) 43.77. In each experiment of the latter series the water was determined by driving it off at a red heat. It was proved that the chloride does not undergo decomposition at this temperature, and the water contents was found to vary no more than 0.0005 of the total weight. In three more experiments the water was determined, and the anhydrous salt analysed by Pelouze's method giving (1) 43.77; (2) 43.74; (3) 43.76. Ag = 108; Cl = 35.5; S = 16. The chloride was prepared (1) from the chemically pure chloride of commerce by precipitating barium with sulphuric acid, separation of lime by precipitation of the strontium chloride by HCl gas and washing with chlorhydric acid. The purity was tested by the solubility of a portion converted into sulphate. The chloride was finally redissolved and precipitated with alcohol. was prepared from (1) by a repetition of the same process. (3) was prepared from (2) by recrystallization. (Bibl. Univ., Arch. des Sciences, (2,) 1, 1858, 220.)

J. Dumas: 87.52 (O = 16).

Determined from the mean of six experiments on the analysis of strontium chloride with argentic nitrate. The extreme difference was 0.14, Cl = 35.5; Ag = 108. The salt was purified by boiling with sulphuric acid, and precipitation with and recrystallization from chlorhydric acid. These processes were in some cases several times repeated. The pure salt was fused in a current of HCl gas. (Annal. de Chim. et de Phys., (3,) 55, 1859, 129.)

SULPHUR.

Deville and Troost and others have determined the density of sulphur in the gaseous form. It corresponds to an atomic weight of about 32. The specific heat of sulphur also agrees moderately well with this value. (Gmelin-Kraut, l. c.; L. Meyer, l. c.)

J. J. Berzelius; F. H. Wollaston: 32 (O = 16); 200 (O = 100).

According to Wollaston, Berzelius found that plumbic sulphide was composed of 86.64 lead and 13.36 S. Hence the value, for lead = 1295. (Phil. Trans., 104, 1814, 20.)

J. J. Berzelius: 32.19 (O = 16) 201.165 (O = 100).

A known weight of lead was dissolved in pure nitric acid, precipitated with sulphuric acid and evaporated. The mean result of four experiments was that 100 Pb = 146.44 sulphate. The variation was only in the fifth figure. If lead = 1294.498 the value follows. [If this relation is recalculated with Stas's atomic weight of lead, S = 32.096.] (Poggend. Ann. 8, 1826, 16.)

E. Turner: 32.17 (O = 16).

Determined from syntheses of plumbic and baric sulphates. The former gave 16.083, the latter, 16.087. Ba = 68.7, Pb = 103.6. The numbers are for vacuum. Vide Barium and Lead. (Phil. Trans., 123, 1833, 539.)

T. Thomson: 32 (O = 16); 200 (O = 100).

This chemist found the specific gravity of sulphurous acid in mean of two experiments, 2.22216, almost exactly double 1.1111 which he takes (on utterly untenable grounds) for the specific gravity of oxygen. (Erdmann's Journ. für Prak. Chem., 8, 1836, 370; Records of General Science by R. D. Thomson, 1836, 179.)

ERDMANN and MARCHAND:
$$32.004$$
 (O = 16); 200.026 (O = 100).

Determined by four experiments on the decomposition of mercuric sulphide by copper, in a current of carbon dioxide, the mercury being caught in a cold receiver. The mean composition was found to be for vacuum 86.211 mercury and 13.789 sulphur, extreme difference, 0.017 Hg. If Hg = 1250.6, the value follows. In purifying the sulphide it was first heated to drive off excess of sulphur and then sublimed three times, the first and last portions of the sublimate being rejected. (Erdmann's Journ. für Prak. Chem., 31, 1844, 396.)

J. J. Berzelius: 32.12 (O = 16); 200.75 (O = 100).

Berzelius' former value, 201.165, is changed by the new value for lead, 1294.645 to 200.8017. Three new experiments were made by gently heating argentic chloride in a current of hydrogen disulphide. The mean of three experiments gives S = 200.706; extreme difference 0.11. Cl = 443.38, Ag = 1349.66. (Berzelius' Jahresbericht, 25, 1845, 37, and Lehrbuch, 3, 1185.)

H. STRUVE: 32.002 (O = 16).

Determined by six experiments on the reduction of a known weight of argentic sulphate in a current of hydrogen. The number is the mean; extreme difference, 0.146. Ag = 108. The sulphate was prepared by precipitating the nitrate with an excess of sulphuric acid, and drying at a high temperature. (Liebig's Ann., 80, 1851, 203; Berzelius' Jahresbericht, 30, 20.)

J. Dumas: 32.0196 (O = 16).

Determined by five experiments on the combustion of silver in sulphur vapor. The number is the mean; extreme difference, 0.054. Ag = 108. The sulphur was purified by repeated distillation. The silver was heated to redness in a current of sulphur vapor, the excess of sulphur being afterwards distilled off in a current of carbon di-oxide. (Annal. de Chim. et de Phys., (3,) 55, 1859, 147.)

J. S. Stas: 32.0742 [?] (O = 16).

According to the mean of six analyses of argentic sulphate by decomposition in a current of hydrogen at as low a temperature as possible, 100 sulphate yield 69.203 [more exactly 69.20317] silver; extreme difference, 0.012. Five syntheses of the sulphide, performed by heating silver in a current of sulphur vapor or hydrogen disulphide, showed that 100 silver = 114.8522 sulphide; extreme difference, 0.005. By comparing these figures, which are for vacuum, Stas deduces S = 32.0742; Ag = 107.920. [There seems to be a trifling error in this calculation. The weighings seem to be correct, for the means correspond to the details given. As given, the numbers indicate S = 32.058; Ag = 107.926. The latter is almost identical with Stas's mean value, 107.930.] The sulphate was prepared by the action of sulphuric acid on argentic nitrate, or by solution of silver in sulphuric acid. The salt was heated above the boiling point of sulphuric acid. (Stas, Unters. über Chem. Prop., Leipzig, 1867.)

TANTALIUM.

Deville and Troost have determined the vapor density of tantalium chloride. It agrees with an atomic weight of 182. (Paris Comptes Rend., 64, 1867, 294.)

J. J. Berzelius: 167.74 (O = 16).

Berzelius decomposed the sulphide in dry chlorine gas and decomposed the resulting chloride with water. 99.75 parts sulphide yielded 89.35 tantalic acid. On the supposition that the acid contains three atoms of oxygen Berzelius calculates the atomic weight at 1148.365 for S = 200.75. [If the acid contains five atoms of oxygen the value becomes 167.74.] (Poggend. Ann., 4, 1825, 14, and Lehrbuch, 3, 1209.)

Rose denies that the sulphide formed, as Berzelius prepared it, by heating tantalium in carbon disulphide vapor is a constant compound. (*Poggend. Ann.*, 99, 580.) Marignac, however, shows that Berzelius, Rose and Hermann, obtained constant results from its analysis, from 89.50 to 90 acid from 100 sulphide. If Ta = 182, the sulphide would give 90.24 acid. (*Liebig's Ann.*, S, 4, 1866, 358.)

H. Rose: 172 (O = 16).

Out of twelve analyses of the chloride, in which both the chlorine and the tantalic acid were determined, Rose selected two in which the agreement was best. [These analyses calculated for Ag = 107.93, Cl = 35.457, give Ta = 171.96.] The chloride was prepared from tantalic acid especially freed from tungsten and tin by mixing with carbon, drying in carbon di-oxide, and heating in a current of chlorine in which the salt was allowed to cool. Excess of chlorine was expelled by dry air, and the salt was hermetically sealed in glass. Rose supposed the acid to contain two atoms of oxygen and therefore deduces the value 859.81. (O = 100). (Poggend. Ann., 99, 1856, 75.)

Marignac seems to prove that the material with which Rose dealt contained niobium. He states that the chlorides of the two elements cannot be separated from one another, and that there are no characteristics by which their purity can be decided. (*Liebig's Ann.*, S, 4, 1866, 352.)

R. HERMANN:

Hermann made many analyses of tantalium salts to which, however, he ascribes quite incomprehensible formulas. Marignac has shown that his methods were utterly inadequate to produce pure preparations. He assumes two atoms of tantalium and three atoms of oxygen in the acid and gives the atomic weight as 645. (O = 100.) (Erdmann's Journ. für Prak. Chem., 70, 1857, 193.)

C. Marignac: 182 (O = 16).

Berzelius', Rose's and Marignac's analyses of the double fluoride of tantalium and potassium show that the fluorine is combined with Ta and potassium in proportions of two to five. The salt has also exactly the crystal form of the niobium salt. Hence the acid is a ditantalic pentoxide. Four experiments were made on this salt by drying at 100°, moistening with sulphuric acid and heating gradually till the excess of acid was driven off. The potassic sulphate was leached out, evaporated, melted and weighed, and the tantalic acid heated to redness and weighed. The mean potassic sulphate contents was found to be 44.29 per cent; extreme difference, 0.15. The mean amount of tantalic acid obtained was 56.59; extreme difference, 0.25. If K = 39, these data give Ta = 182.3. Four analyses were also made of the ammonium salt. This contained traces of potassium which were determined and allowed for in each case. The mean amount of tantalic acid obtained was 65.25 per cent; extreme difference, 0.34. This gives Ta = 182, the number which Marignac adopts. The salts were obtained by dissolving tantalic acid, which had not been heated to redness, in fluohydric acid, adding potassic or ammonic hydrate and purifying by recrystallization. These salts are much less soluble than the corresponding niobium and titanium salts. (Liebig's Ann., S. 4, 1866, 284.)

TELLURIUM.

Regnault and Kopp have each determined the specific heat of tellurium and found it in accord with an atomic weight of about 128. (Gmelin-Kraut, l. c.)

J. J. Berzelius 129.03 (O = 16); 806.452 (O = 100).

A known weight of metallic tellurium was oxidized with nitric acid, the excess of acid being driven off by heat. It was found that 100 Te gave 124.8 tellurious acid. (*Poggend. Ann.*, 8, 1826, 24.)

J. J. Berzelius: 128.28 (O = 16); 801.76 (O = 100).

Determined as before but with purer material. Three experiments were made, which gave 802.838, 801.786, 801.74. Berzelius took the mean of the latter two. The tellurium was prepared from tetradymite by heating with potassium carbonate and olive oil in a closed crucible, dissolving the potassium telluride so formed in water free from air, precipitating the tellurium by a current of air and distilling it in a current of hydrogen. (Poggend. Ann., 32, 1834, 16.)

K. von Hauer: 128.06 (O = 16).

Determined from the mean of five experiments on the precipitation of bromine with argentic nitrate from the double bromide of potassium and tellurium. The bromine contents was found to be 69.9236 per cent., for Ag = 108.1; Br = 80; extreme difference 0.172. If K = 39.2, the value follows. The salt was prepared by mixing tellurium and potassic bromide in atomic proportions, adding water and bromine, heating to drive off excess of bromine and repeated recrystallization. (Erdmann's Journ. für Prak. Chem., 73, 1858, 98; Sitz-Bericht der k. k. Acad.)

J. Dumas: 129 (O = 16).

No details are given. (Annal. de Chim. et de Phys., (3,) 55, 1859, 129.)

THALLIUM.

Regnault determined the specific heat of thallium. It agrees with an atomic weight of 204. (Gmelin-Kraut, l. c.)

A. Lamy: 204 (O = 16).

Three analyses of the chloride with argentic nitrate gave

A mean of 204; extreme difference 1.2. An experiment on the precipitation of the sulphate with barium nitrate gave 204.3. [The atomic weights used were probably those accepted by Dumas.] The salts were purified by recrystallization. (Annal. de Chim. et de Phys., (3,) 67, 1863, 411.)

W. Crookes: 202.96 (O = 16).

These determinations were made from the sulphate, which was prepared with great care. By decomposing the sulphate with potassic iodide and weighing the thallic iodide formed, the atomic weight was found at 202.73; by precipitation with barium nitrate, 203.55; with chlorhydric acid and alcohol, thallic chloride being weighed, 201.85; from the amount of sulphate produced from a known weight of metal, 203.1; by precipitation with platinum chloride, 203.56. The values taken for Cl, I, etc., are not given; [they were probably those accepted by Dumas.] (Erdmann's Journ. für Prak. Chem., 92, 1864, 277; Chem. News.)

H. Werther: 204 (O = 16).

In five experiments Werther decomposed thallic iodide with potassic hydrate and zinc, both perfectly pure, and precipitated the iodine with silver. The mean result of these experiments was Tl = 204.4; extreme difference 1.7. [The value assumed for I is not stated. One experiment, which gave exactly 204, according to Werther, recalculated for Ag = 107.93; I = 126.85 gives Tl = 203.63.] Three experiments were made by decomposing the iodide with ammoniacal solution of argentic nitrate and weighing the argentic iodide formed. These determinations gave Tl = 203.47; extreme difference 0.3. The preparation of the iodide is not given. (Erdmann's Journ. für Prak. Chem., 92, 1864, 136.)

M. Hebberling: 203.94 (O = 16).

Hebberling made three experiments on the sulphate by precipitation with barium chloride, which gave in mean Tl = 204.13; extreme difference 0.2. He also made two experiments on the chloride by precipitation with argentic nitrate. These gave 203.8 and 203.5. The atomic weights assumed are not stated. [If Ag = 107.93; Cl = 35.457; the first analysis of the chloride gives Tl = 203.44. The data for the second analysis are misprinted. If a probable correction of a single figure is made, the data give Tl = 100.000

203.026.] The salts were purified by recrystallization. (Liebig's Ann., 134, 1865, 11.)

W. Crookes: 204.155 (O = 16).

Determined by experiments on the solution of metallic thallium in nitric acid and evaporation to dryness. The number is the mean of ten experiments; extreme difference, 0.038. The balance stood in a partial vacuum, and the weighings were made at two different pressures and calculated for vacuum. Very elaborate precautions were taken throughout. Crookes also mentions determinations made with barium nitrate, but gives no data. The thallium was prepared in seven different lots by the reduction of as many different salts which had been purified by recrystallization &c. The metal was fused in lime. The reagents were expecially prepared by methods similar to those of Stas. Crookes took N = 14.009, O = 15.96, and calculated for Tl the value 203.642. [If O = 16, the value becomes 204.155.] (Phil. Trans., 163, 1873, 277.)

THORIUM.

From the isomorphism existing between thorium, tin, and titanium, and from the similarity of thorium to zirconium, Delafontaine and Marignac believe the oxide to contain two atoms of oxygen. (Liebig's Ann., 131, 100.) Neither the specific heat of this element nor the vapor density of any of its compounds has been determined so far as I know.

J. J. Berzelius; 238 (O = 16); 1887.72 (O = 100).

From the sulphate, precipitated by heating a solution of the salt and redissolved in cold water, Berzelius got the values 748.493 and 735.713 by precipitating with barium chloride. He also analysed the double sulphate of potassium and thorium. From the relation between the sulphuric acid and the thorium oxide found, the atomic weight would seem to be 750.63, while the relation between the potassic sulphate obtained, and the amount of oxide gives 740.6. These numbers are calculated on the supposition that the oxide contains a single atom of oxygen. Ba =

855.29, 8 = 200.75, K = 488.856. (Poggend. Ann., 16, 1829, 398, and Lehrbuch, 3, 1224.)

J. J. Chydenius: 236.64 (O = 16).

This chemist analysed the sulphate, the double sulphate of potassium and thorium, the oxalate, the acetate and the formate, getting results which vary from 228.52 to 243.76. He averages with his own results analyses made by Berzelius and by Berlin, which, however, alter the result inappreciably. According to Delafontaine, the methods employed for purification are ineffectual. Chydenius assumes a single atom of oxygen in the oxide. (Poggend. Ann., 119, 1863, 55.)

N. J. Berlin: 231.64 (O = 16).

Chydenius reports two analyses of the oxalate by Berlin which gave for thorium 57.87 and 57.95, or 231.48 and 231.80. (Poggend. Ann., 119, 1863, 56.)

M. Delafontaine: 231.5 (O = 16).

Determined from analyses of the sulphate. Fourteen experiments on the decomposition of this salt, by the heat of a strong double-draught lamp, gave a mean of 52.51 per cent. oxide; extreme difference, 0.83. In three experiments the sulphur contents of the salt was determined by precipitation with barium chloride after the sulphate had been decomposed with ammonium oxalate. The mean amount of sulphuric anhydride so found was 31.92 per cent.; extreme difference, 0.78. Three experiments on the water contents gave 15.68 per cent; extreme difference, 0.21. The sum of these means is 100.11. The value of thorium was calculated from the relation of the oxide to the sulphuric anhydride for S = 32, Ba = 137. The salt was prepared from thorite and from orangite by decomposition with sulphuric acid and recrystallization of the sulphate with the help of heat. The purification was continued until the crystals and the mother liquor had exactly the same composition. Marignac assisted at this investigation. (Liebig's Ann., 131, 1864, 100.)

P. T. CLEVE: 233.88 (O = 16).

Cleve made six analyses of the anhydrous sulphate, getting in mean Th = 233.8; extreme difference, 1.36. From

analyses of the oxalate he got 283.97; extreme difference, 0.6. (Kopp's Jahresbericht, 1874, 261; Bull. Soc. Chim., (2,) 21, 116.)

TIN.

Regnault and Kopp have each determined the specific heat of tin. It agrees with an atomic weight of about 118. Dumas, Cahours and others have determined the vapor density of volatile tin compounds with a similar result. (Gmelin-Kraut, l. c.; L. Meyer, l. c.)

J. J. Berzelius; 117.647 (O = 16); 735.294 (O = 100).

Berzelius determined this value by oxidizing pure tin foil by means of nitric acid and weighing the oxide. He found 100 tin = 127.2 stannic acid. (*Poggend. Ann.*, 8, 1826, 184.)

G. J. MULDER: 116.112 (O = 16); 725.7 (O = 100).

Two experiments were made by oxidizing tin with nitric acid, evaporating, drying, and heating to redness. They gave each 100 tin = 127.56 stannic acid; whence the value. All possible precautions are said to have been taken. The metal was prepared by the reduction of pure oxide with soot and a flux. (Erdmann's Journ. für Prak. Chem., 48, 1849, 85; Scheikundige Onderzoek., 5. Deel, 260.)

C. L. VLAANDEREN: about 118. (O = 16).

Determined from experiments on the oxidation and reduction of tin and stannic acid in vessels of various materials. The experiments regarded as the most accurate were made on the reduction of the acid in a current of hydrogen in porcelain vessels. The acid had been heated in platinum. These experiments gave 59.04 and 59.12. Stannic acid heated in glass or porcelain was found to retain nitric acid. (Kopp's Jahresbericht, 11, 1858, 138; Mulder, Scheikundige Verh. en Onderzoek., 2. Deel, 150.)

J. Dumas: 118.08 (O = 16).

Two experiments were made on the oxidation of pure tin by nitric acid. The stannic acid being heated white hot in platinum vessels gave for the atomic weight 59.1, and 58.96. The tin employed was prepared from pure chloride. Two experiments on the titration of the chloride with argentic nitrate gave 59.06 and 59.03. Ag = 108, Cl = 35.5. (Annal. de Chim. et de Phys., (3,) 55, 1859, 156.)

TITANIUM.

The specific heat of titanic acid has been determined by Regnault and by Kopp, and indicates an atomic weight of about 50. Dumas determined the vapor density of the tetrachloride at 6.836. [If the molecular weight of O = 32, and if Cl = 35.457, this gives Ti = 56.025.] (Gmelin-Kraut, l. c., and Poggend. Ann., 9, 1827, 441.)

H. Rose: 61.17 (0 = 16).

Determined by roasting titanium sulphide and weighing the titanic acid formed. The highest result obtained was 1.017 sulphide from 0.757 acid. This result Rose adopted on the supposition that an excess was impossible. For S=201.16 these data give Ti=62.25 (O=16); 389.1 (O=100.) [If S=32, Ti=61.17.] The sulphide was prepared by heating titanic acid in a current of carbon disulphide. (Gilbert's Ann., 73, 1828, 135.)

Rose subsequently expressed the opinion that the sulphide employed in this analysis was impure, and contained undecomposed titanic acid, but afterwards came to the conclusion that it was perfectly pure, accounting for the variation of the results from those he obtained later by the theory that the sulphide and the oxide of this element, like those of tantalium, were entirely dissimilar compounds. Marignac has shown that tantalium sulphide is of normal constitution. (Poggend. Ann., 99, 1856, 576.)

H. Rose: 48.28 (O = 16).

Titanium chloride was decomposed with water, titanic acid precipitated by ammonic hydrate, and the chlorine precipitated from the filtrate with argentic nitrate. Taking Ag=1351.607, Cl = 221.325; Rose calculated the chlorine contents in four experiments at from 74.43 to 74.53 per

cent; mean 74.46 and Ti at 303.686. According to Gmelin-Kraut, these analyses recalculated for Stas's values give Ti = 48.28. The chloride was prepared by the action of chlorine on a mixture of titanic acid and carbon, and was rectified four or five times over potassium and mercury. It was clear and developed no chlorine on decomposition with water. (Poggend. Ann., 15, 1829, 145.)

C. G. Mosander: 47.33 (O = 16); 295.81 (O = 100).

Mosander determined the oxygen contents of titanic acid at from 39.83 to 40.82 per cent.; mean 40.427. Mosander never described the method of analysis. [The oxygen contents was probably determined from the chloride, for the above data give Ti = 294.7, while Berzelius records the determination as having given 295.81.] (Poggend. Ann., 19, 1830, 212, and Berzelius' Lehrbuch, 3, 1211.)

J. Pierre: 50.36 (O = 16).

Determined by three experiments on the titration of the chloride with argentic nitrate by Pelouze's method. Pierre does not give the values taken for Cl and Ag. He calculates the atomic weight of Ti at 314.69. [If Ag = 107.93, Cl = 35.457; his data give Ti = 314.75 (O = 100); 50.36 (O = 16), with an extreme difference in the latter case of 0.08.] He made two other determinations giving lower results, but it was found that the chloride employed was slightly decomposed by contact with air. The chloride was prepared from artificial titanic acid which was free from iron, and was further purified by fractional distillation. (Annal. de Chim. et de Phys., (3,) 20, 1847, 257.)

• A. Demoly: 56.512 (O = 16).

Determined by experiments on the tetrachloride. The salt was decomposed with water, the titanic acid precipitated by ammonic hydrate, and the chlorine precipitated in the filtrate, after the excess of ammonic hydrate had been volatilized and the solution acidified. Both precipitates were weighed. Demoly calculates the atomic weight of Ti at 350, without mentioning what values he accepted for silver and chlorine. [If Ag = 107.93, Cl = 35.457; the atomic weight, calculated from the argentic chloride, is 353.2 (O = 100); or 56.512 (O = 16), with an extreme difference in the three experiments of 0.88 for O = 16.] The chloride was prepared from rutile by preliminary conver-

sion into nitride, &c. It was purified by rectification over mercury and potassium. (Liebig's Ann., 72, 213; Laurent and Gerhardt, Comptes Rend., 1849, 325.)

TUNGSTEN.

Regnault has determined the specific heat of tungsten, and Roscoe the vapor density of the chloride. These experiments place the atomic weight of tungsten at about 184. (Gmelin-Kraut, l. c.; L. Meyer, l. c.)

J. J. Berzelius: 189.26 (O = 16); 1183.355 (O = 100).

A weighed quantity of tungstic acid was reduced in a current of hydrogen, again weighed, then re-oxidized and reweighed. The number is the mean result of the two operations. The number is given in Berzelius' Lehrbuch as 1188.36 with the data, which are also given in Poggend. Ann., 8, 23. It is pointed out in Graham-Otto that this value must be misprinted, an observation which I have verified. (Poggend. Ann., 4, 1825, 152.)

Berzelius made an earlier determination than the foregoing by the oxidation of the sulphide, getting 1207. He points out the source of error in this experiment arising from the formation of irreducible sulphate. (Berzelius' Jahresbericht, 5, 1825, 121.)

R. Schneider: 184.12 (O = 16); 1150.78 (O = 100).

Schneider made five experiments on the reduction of tungstic acid with hydrogen in a porcelain tube heated by a charcoal fire. These analyses gave the mean contents of the acid at 79.316 tungsten per hundred; extreme difference, 0.096. This composition corresponds to an atomic weight of 1150.39. He also made three experiments on the combustion of tungsten, getting a mean of 79.327 tungsten per 100 acid; extreme difference, 0.005, or an atomic weight of 1151.17. The value taken is the mean. The tungstic acid was prepared by decomposing ammoniotungstic sulphide with chlorhydric acid, washing the precipitate with acid, solution in ammonia, reprecipitation with chlorhydric acid, and so on until a perfectly pure product was obtained. The tungstic acid was finally dried and

heated to redness. (Erdmann's Journ. für Prak. Chem., 50, 1850, 163.)

R. F. MARCHAND: 184.1 (O = 16); 1150.6 (O = 100).

Determined from two experiments on the reduction of tungstic acid in a current of hydrogen, and two experiments on the combustion of tungsten. These determinations were made in the same manner as and at the same time with Schneider's. The extreme difference was 3.5 for O = 100. (Liebig's Ann., 77, 1851, 263.)

J. B. von Borck: 183.816 (O = 16); 1148.85 (O = 100).

Determined by seven experiments on the reduction of tungstic acid at a white heat by hydrogen, and by two experiments on the combustion of tungsten. The number is the mean; extreme difference, 10.38 for O = 100. The tungstic acid was prepared from Wolframite by fusing the mineral with potassium carbonate, solution in water containing alcohol, precipitation with calcic chloride and decomposition of the calcic tungstate with chlorhydric acid. The tungstic acid so produced was converted into ammonium salt which, on decomposition, yields a compound free from iron and manganese. (Erdmann's Journ. für Prak. Chem., 54, 1851, 254.)

A. RICHE: 174 (O = 16).

This value was reached by five determinations of the amount of water produced by the reduction of tungstic acid in a current of hydrogen, which gave a mean of 87.07; extreme difference, 1.78. The tungstic acid was obtained by heating the ammonium salt, or by the decomposition of the oxychloride produced by heating tungstic acid and carbon in a current of chlorine. (Annal. de Chim. et de Phys., (3,) 50, 1857, 10.)

J. Dumas: 184 (O = 16).

Dumas made six experiments on the reduction of tungstic acid in hydrogen at a high temperature in a nacelle of unglazed porcelain, and two experiments on the titration of the chloride with argentic nitrate. The extreme difference between the results was 0.69 for O = 8. The acid was pre-

pared by gently heating the ammonium salt in a muffle. (Annal. de Chim. et de Phys., (3,) 55, 1859, 144.)

F. A. Bernoulli: 186.8 (O = 16); 1167.5 (O = 100).

Berneulli made five experiments on the reduction of tungstic acid by hydrogen in a porcelain tube at a very high temperature, two experiments on the amount of water formed in reduction, and four experiments on the oxidation of tungsten. The mean result was W = 93.41; extreme difference, 0.75. [If experiment 9, in which oxidation seems to have taken place, is left out, the mean becomes 93.35; extreme difference, 0.18.] The tungstic acid was prepared from ammonium tungstate which had been boiled for several days with pitric acid. The tungstic acid was One part of it was green, another part. heated to redness. The determinations from the different colored acids did not differ, and Bernoulli considers them isomeric modifications of the same compound. There appear to be misprints in the data given. (Poggend. Ann., 111, 1860, 599.)

C. Schefbler: 184 (O = 16).

Scheibler reached this value by five determinations of the water contents (9 molecules) of barium metatungstate. From determinations of the barium and the tungsten in the same compound Scheibler reached other values, but he regards the water determination as the most trustworthy. (Erdmann's Journ. für Prak. Chem., 83, 1861, 328.)

E. Zettnow: 183.952 (O = 16).

Determined from analyses of ferrous tungstate and argentic tungstate. A known weight of ferrous tungstate was melted with sodium carbonate and the mass dissolved. The ferric hydrate was thoroughly washed, dissolved in chlorhydric acid, reduced to ferrous chloride with zinc of known composition, and titrated with potassic permanganate in several measured portions. Four such series of experiments were made, and gave a mean of 92.038 for W; extreme difference, 0.33. The ferrous tungstate was prepared by melting pure anhydrous sodium tungstate with ferrous chloride and sodium chloride, dissolving, separating impurities, crystallizing, washing the crystals with water, chlorhydric acid and sodium carbonate. The argen-

tic tungstate was decomposed with nitric acid and titrated with sodium chloride or decomposed with hot sodium chloride solution, the argentic chloride being weighed. Five experiments gave a mean of 91.915 for W; extreme difference, 0.13. The argentic tungstate was prepared by the precipitation of sodium tungstate with argentic nitrate, thorough washing and drying in yellow light. The permanganate solution was prepared according to Mohr and tested with ammonio-ferrous sulphate. Fe = 28, Ag = 108. (Poggend. Ann., 130, 1867, 30.)

H. E. Roscoe: 184.04 (O = 16).

Determined by reducing tungstic acid in a current of hydrogen, by reoxidizing the metal, and by reducing the chloride in a current of hydrogen, the chlorhydric acid being condensed and estimated as argentic chloride. In the experiments on the acid, that compound was reduced, and reoxidized three times with almost identical results. The mean of the second and third reductions of the same sample gave W = 183.84. In the experiments on the chloride, the chlorine and the tungsten were each determined, and gave a mean of 184.25 for Cl = 35.5. The tungstic acid was prepared by the decomposition of the chloride, washing and heating to redness in a platinum vessel. It was canary yellow. The chloride was prepared from pure tungsten. (Liebig's Ann., 162, 1872, 366.)

URANIUM.

No certainty exists as to the relation between the equivalent and the atomic weight of uranium. The latter is commonly accepted as about 120. Mendelejeff gives grounds for supposing it to be 240, (Liebig's Ann., S. 8, 1871, 178,) and L. Meyer regards it as probably 180, a value which accords well with the specific heat of the black oxide as observed by Regnault. (Gmelin-Kraut, l. c.) For the purposes of this paper it seems best to retain the customary value.

J. A. Arfvedson: 128.6 (O = 16).

Determined by experiments on the reduction of uranoso-

uranic oxide and on the oxidation of uranous oxide. By combustion of uranous oxide in oxygen he found in two experiments that 100 oxide combined with 3.695 and with 3.73 oxygen. From the reduction of the green oxide he found that 100 uranous oxide combine with 3.67 oxygen. He deduces as the mean 3.688. Regarding uranous oxide as the metal, Arfvedson calculated the atomic weight at 2711.36. [If the lower oxide is a protoxide, the data give 128.6 for O = 16.] The uranous oxide was prepared from pitchblende by solution in aqua regia, precipitation of heavy metals with hydrogen sulphide, precipitation with ammonic hydrate, solution in ammonium carbonate to remove iron, reprecipitation, heating to redness, washing with chlorydric acid to remove impurities, and reduction in hydrogen. (Poggend. Ann., 1, 1824, 254.)

E. Peligot: 119.128 (O = 16).

In two experiments the amount of carbon in the acetate was found to be 11.27 and 11.3; mean 11.285. In one experiment the uranic oxide was determined at 67.8 per cent. [From these data the above value follows.] Peligot takes 120 or 750, C = 75. The preparation of the salt is not given. Peligot mentions the oxalate and gives analyses, but does not deduce an atomic weight from them. (Annal. de Chim. et de Phys., (3,) 5, 1842, 39.)

J. J. EBELMEN: 118.86 (O = 16); 742.875 (O = 100).

Ebelmen made six experiments on the reduction of the oxalate to uranous oxide by hydrogen and heat. The value follows with an extreme difference of 0.65 for C = 75; H = 12.5. All the weighings were reduced to vacuum. To obtain pure oxalate, the nitrate was precipitated by oxalic acid and this preparation decomposed by heat. The oxide thus obtained was digested with chlorhydric acid, washed, dissolved in nitric acid, recrystallized, and precipitated with oxalic acid. The oxalate was dried at 100°. According to Rammelsberg the reduction of the oxalate is accompanied by the separation of carbon which remains with the oxide. (Annal. de Chim. et de Phys., (8,) 5, 1842, 189.)

Berzelius, Arfvedson, Marchand: 128.4 (O = 16); 802.49 (O = 100).

While Arfvedson was making his first determination, Berzelius also made an experiment on the combustion of ura-

nous oxide getting 103.685 uranic from 100 uranous oxide. Marchand (Erdmann's Journ. für Prak. Chem., 23, 1841, 498) got in the same way 103.668. The average of the combustion experiments of all three chemists is 103.694, whence Berzelius calculates the value. (Berzelius' Jahresbericht, 22, 1842, 113.) Peligot and Rammelsberg, as well as Marchand, point out faults in this method, such as the probable condensation of hydrogen in the protoxide and the tendency to form higher oxides. (Poggend. Ann., 59, 1843, 4.)

C. RAMMELSBERG.

This chemist made experiments on the reduction by hydrogen of the green oxide, prepared in various ways, and got results varying from 580.4 to 767.6 for O = 100. (Poggend. Ann., 59, 1843, 9.) By precipitation of uranous chloride with silver he reached the number 787.5 for Cl = 442.65. The chlorine contents found varies in three experiments from 73.89 to 74.46. The chloride was prepared by heating uranous oxide in an atmosphere of chlorine. (Poggend. Ann., 55, 1842, 321.)

J. Wertheim: 119.42 (O = 16); 746.36 (O = 100).

Determined by three experiments on the decomposition of the double acetate of uranium and sodium. The mean loss of acetic acid by heating the salt to redness was 32.477 per cent.; extreme difference, 0.036. The number follows for C = 75, H = 6.25, Na = 390.9. [In Poggend. Ann., 57, 484, an abstract is given of a paper read before the academy (of Berlin?) by Mitscherlich, in which he states that Wertheim's experiments above described give 740.512. Berzelius in his Jahresbericht, 23, 137, makes or quotes the same statement, so also does Rammelsberg, Poggend. Ann., 59, 4, and it has been repeated elsewhere. I have recalculated the data given by Wertheim and find the results correctly deduced in his own report. For Na = 23.043 (Stas); the data give U = 119.53.] The salt was prepared from uraninite by solution in nitric acid, precipitation with hydrogen sulphide, evaporation of the filtrate to dryness, solution in hot water, crystallization and recrystallization, heating the crystals to drive off nitric acid, solution in acetic acid, digestion with sodium carbonate and recrystallization. (Erdmann's Journ. für Prak. Chem., 29, 1843, 209.)

C. Rammelsberg: about 120 (O = 16).

Determined in six experiments, undertaken at Berzelius' suggestion, by treating uranous oxide with nitric acid and sulphuric acid and weighing the sulphate. It is very difficult to weigh the uranous oxide which constantly increases in weight. Two experiments were made on the green oxide, which was prepared either by heating uranous oxide, or the nitrate, in air. Two experiments were made on magnesium uraniate by dissolving the compound in nitric scid and heating to redness. The compound was found unstable in character. One experiment was made by heating the double acetate of uranium and sodium and three experiments by heating the double acetate of barium and The results obtained varied from 633.17 to uranium. Rammelsberg considers the determinations confirmatory of Wertheim's and Ebelmen's. (Poggend. Ann., 66, 1845, 95.)

E. Peligor: 120 (O = 16); 750 (O = 100).

Determined by combustion of the oxalate in a current of air, both the carbonic acid and the green oxide of uranium being weighed. At first Peligot got only 730 as the atomic weight by this process, but by repeating the recrystallization of the salt until determinations gave constant results, he got a mean of 750. He says that he came to the same value by comparing the amount of uranic oxide obtained from the acetate with the weight of the salt employed. (Paris Comptes Rend., 22, 1846, 487.)

VANADIUM.

Roscoe has determined the vapor density of vanadium chloride. It agrees with an atomic weight of about 51. (L. Meyer, l. c.)

J. J. Berzelius: 52.47 (O = 16).

Berzelius made four experiments on the relation between the higher and the lower oxides of vanadium, three by reduction with hydrogen at a very high temperature and one by oxidation. He supposed the higher oxide to have the formula VO₈, and the lower VO, and consequently got for the atomic weight the number 855.84 (O = 100). R. Schneider has shown that the data as given by Berzelius are discordant, (Poggend. Ann., 88, 319,) a fact of small importance in view of the succeeding investigation. The higher oxide analyzed by Berzelius was produced by gently heating the ammonium salt. (Poggend. Ann., 22, 1831, 14; Kongl. Vet. Akad. Handl., 1831.)

Roscoe examined some ammonium vanadate which Berzelius had sent Faraday and found that it contained phos-

phorus. (*Liebig's Ann.*, S, 6, 1868, 93.)

H. E. Roscor: 51.33 (O = 16).

Roscoe made four experiments on the reduction of vanadic acid $(\nabla_2 O_5)$ in carefully purified hydrogen. The acid was prepared from ammonium vanadate. To free this compound from phosphorus and silicic acid it was powdered, decrepitated with sodium in an iron crucible, washed with water and with chlorhydric acid, re-oxidized with nitric acid, chloridized in a current of chlorine, the chloride rectified and decomposed with water. The acid so obtained was dried, moistened with sulphuric acid, exposed to the fumes of fluohydric acid for ten days and melted. This pure acid was first heated for several hours in dry air and afterwards in hydrogen. The mean result of four experiments was V = 51.371; extreme difference, 0.228. Nine experiments were made on the titration of the chloride by Pelouze's method. Eight experiments were also made on the analysis of the chloride with argentic nitrate by the ordinary method. The mean of the seventeen experiments on the chloride gives the contents in chlorine at 61.276 per cent.; extreme difference, 0.69. This composition indicates an atomic weight of 51.29. Roscoe takes Cl = 35.457, Ag =107.93. The vanadium chloride was purified by rectification over sodium in a current of carbon di-oxide. The reagents were prepared according to Stas. (Liebig's Ann., S, 6, 1868, 86.)

Roscoe mentions atomic weight determinations by Czudnowicz as giving 55.35. This chemist, however, did not calculate an atomic weight from his analyses, but used that obtained by Berzelius. (*Poggend. Ann.*, 120, 1863, 17.)

YTTRIUM.

The composition of yttrium oxide is not definitely settled. Mendelejeff concludes from the general behavior of its compounds that it is a sesqui-oxide. As, however, all the chemists who have made atomic weight determinations of this element have considered it a prot-oxide, I shall assume it to be so and the atomic weight, therefore, about 60.

J. J. Berzelius: 64.29 (O = 16); 401.84 (O = 100).

This determination was made before the discovery of erbium and can scarcely be correct. The value was reached by analysis of the sulphate with barium chloride. Ba = 856.88, S = 201.165. (Poggend. Ann., 8, 1826, 186; 10, 1827, 341.)

N. J. Berlin: 59.7 (O = 16).

According to Blomstrand in Berlin, Ber. der Chem. Ges., 1873, 1467. I can find no other record of this determination which probably appeared in the Forhandl. ved de Skandinaviske Naturforsk, 1860, 448.

O. Popp: 68 (O = 16),

The mean of four analyses of the sulphate showed that 40.15 oxide were equivalent to 38.23 sulphuric anhydride, giving a molecular weight for the oxide of 42.015; extreme difference, 0.018. The yttrium was precipitated with sublimed oxalic acid, the free acid being afterwards neutralized with ammonia. The sulphuric acid was precipitated with barium chloride in the filtrate with precautions. Popp, who denies the existence of erbium and terbium, separated yttrium from the cerite oxides by precipitation with barium carbonate, yttrium remaining in solution, S = 16, Ba = 68.5. (Liebig's Ann., 131, 1864, 183.)

M. Delafontaine: about 64 (O = 16).

Delafontaine does not pretend that this number is exact. It is derived from analyses of the sulphate. His method of separation was essentially Mosander's, which was proved by Popp and by Bunsen and Bahr to give impure salts. (Liebig's Ann., 134, 1865, 108.)

Bahr and Bunsen: 61.7 (O = 16).

Determined by saturating the oxide with sulphuric acid as in the determination of erbium, q.v. Partial recrystallization does not produce pure yttrium nitrate, but only concentrates traces of didymium in the salt. Didymium must be separated with potassic sulphate. Erbium nitrate is more easily decomposed by heat than yttrium nitrate. The nitrates were therefore partially decomposed, yttrium nitrate dissolved out and the process repeated until there was no trace of erbium or didymium visible in the spectroscope. The mean of two determinations gave Y = 30.85; difference, 0.1. S = 16. (Liebig's Ann., 137, 1866, 21.)

M. Delafontaine: 58.5 (O = 16).

Determined by three experiments on the sulphate which gave in mean 48.28 per cent. oxide for S=32. [In the Jahresbericht this determination is reported as giving Y=74.5. Yttrium is apparently a misprint for yttrium oxide.] The yttrium salt seems to have been prepared according to the method of Bahr and Bunsen. (Kopp's Jahresbericht, 1866, 184; Bibl. Univ., Arch. des Sciences, (2), 25, 1866, 112.)

P. T. CLEVE AND O. M. HOEGLUND: 59.7 (O = 16).

Determined by analysis of the sulphate. The oxide was purified by heating the nitrates, etc., according to N. J. Berlin. (Blomstrand, in Berlin, Bericht der Chem. Ges., 1873, 1467; Bihang till Vet. Akad. Handl, 1873, B. 1, 3, No. 8.)

ZINC.

The specific heat of zinc has been determined by Regnault and others. The vapor density of volatile organic compounds has been determined by Frankland and others. These experiments agree in placing the atomic weight at about 65. (Gmelin-Kraut, l. c.; L. Meyer, l. c.)

GAY-LUSSAC, BERZELIUS, WOLLASTON: 65.547 (O = 16); 409.67 (O = 100).

In his experiments on the oxidation of zinc Gay-Lussac

ZINC. 135

found that 100 Zn = 24.41 oxygen. This value is repeatedly cited in his memoir. (Gilbert's Ann., 30, 1811, 297; Mêmoire D'Arceuil, 2, 174.) Wollaston gives the same figures on Gay-Lussac's authority. (Phil. Trans., 104, 1814, 21.) Wollaston calculates from these data Zn = 410, (O = 100.) Berzelius in each of two experiments got 100 Zu = 124.4oxide. (Gilbert's Ann., 37, 1811, 460.) In Poggend. Ann., 8, 1826, 184, as well as in his Lehrbuch, Berzelius cites Gay-Lussac as having found 100 Zn = 24.8 oxygen. states that his own determinations were in perfect accordance with these figures, and calculates from them the atomic weight of zinc at 403.226 or 64.52, and this was the accepted value for many years. I cannot find any other determinations by either of these chemists, and am obliged to suppose that there was a mistake made in recording the data from which Berzelius made his calculations; if so, it is remarkable that neither Berzelius nor the other chemists who determined this value perceived it; for the question was reopened during Berzelius' life, and A. Erdmann made his determination at Berzelius' request.

V. A. JACQUELIN: 66.24 (O = 16); 414 (O = 100).

This number was reached by measuring the amount of hydrogen developed by a known weight of zinc from sulphuric acid on the supposition that the specific gravity of hydrogen is 0.0624. The results seem to have been inconsistent. Subsequently Jacquelain arrived at the same number by oxidizing an impure zinc of known composition. (Paris Comptes Rend., 14, 1842, 636; and Annal. de Chim. et de Phys., (3,) 7, 1843, 204.)

P. A. FAVRE:
$$66$$
, $(O = 16)$; 412.5 $(O = 100)$.

Favre made four experiments on the combustion of zinc oxalate, the carbon di-oxide being collected and its weight compared with that of the oxide. The mean result was Zn = 412.66; extreme difference, 1.11. C = 75. He also made three experiments by passing the hydrogen developed by a known weight of zinc over cupric oxide, the water being caught. These experiments gave in mean Zn = 412.16; extreme difference, 0.65 for H = 12.5. (Annal. de Chim. et de Phys., (3), 10, 1844, 163.)

A. Erdmann; 65.05 (O = 16); 406.591 (O = 100). Determined by oxidizing pure zinc with nitric acid, and

driving off the acid by heating the salt in a porcelain crucible. Platinum is attacked. The number is the mean of four experiments; extreme difference, 0.698. The zinc was prepared by mixing pure oxide with carbon, and distilling in a current of hydrogen. (Berzelius' Jahresbericht, 24, 1844, 132; Efversigt af Kongl. Vet. Akad. Handl., 1, 3.)

ZIRCONIUM.

Deville and Troost have determined the vapor density of the chloride. It agrees with an atomic weight of about 90. (L. Meyer, l. c.)

J. J. Berzelius: 89.6 (O = 16).

In one experiment the sulphate was decomposed with ammonic hydrate, the oxide weighed and the sulphuric acid precipitated with barium chloride. In five experiments the sulphate was decomposed at a white heat, ammonium carbonate being added at the close of the operation. The mean result was that 100 parts of sulphuric anhydride unite with 75.853 parts of zirconium oxide; extreme difference, 0.23. Berzelius deduces the value 840.08 for O = 100, S = 201.165; on the supposition that the oxide contains three atoms of oxygen. [Being a binoxide, this relation gives Zr = 89.6 for O = 16.] The sulphate seems to have been prepared by dissolving the oxide in sulphuric acid and expelling the excess of acid by heat. (Poggend. Ann., 4, 1825, 126.)

R. HERMANN:

This chemist made some experiments on the chloride getting in three determinations a mean of 839.45 for O = 100 and on the tri-oxide supposition. The extreme difference was 20.1. Cl = 443.65. The chloride was produced by heating the oxide with carbon in a current of chlorine. Hermann adopts not his own but Berzelius' determination. (Erdmann's Journ. für Prak. Chem., 31, 1844, 77.)

C. Marignac: 90 (O = 16).

Determined from analyses of potassium fluo-zirconiate. The salt was decomposed with sulphuric acid, the excess

of acid driven off by heat, the residue weighed, the potassic sulphate leached out with water, and the residue again weighed. Marignac does not pretend that the determination is accurate. The results gave from 45.01 to 45.48. He thinks that some potassic sulphate may have escaped solution, and therefore takes the minimum. K = 39, S = 16.

According to Marignac, Deville also found the atomic weight of zirconium somewhat higher than Berzelius by analysis of the chloride with which he determined the vapor density. (Annal. de Chim. et de Phys., (3,) 60, 1860, 257.)

•				
		•		
		•		
	•		ł	
·				
				1
				ı
	•			
				•
			•	

APPENDIX.

DETERMINATIONS BY T. THOMSON.

In Thomson's Annals of Philosophy, volumes 16 and 17, 1820-21, Thomson published a series of papers descriptive of experiments undertaken for the purpose of verifying Prout's hypothesis. His method consisted in mixing reagents in what he considered equivalent proportions, and after precipitation examining portions of the supernatant liquid for an excess of each of the salts supposed to neutralize one another. In all except four cases, either the salt analyzed was a sulphate and the precipitant barium chloride, or the determination was dependent upon such an analysis; yet although Thomson took barium = 70, in no instance was he able to detect either barium or sulphuric acid in the residual solution when the quantity of the re-agents corresponded to the atomic weights which he adopts. Comparison of his results with those reached by more accurate experimenters will make this exact neutralization appear impossible, nor were his contemporaries able to repeat his experiments successfully. Thomson's determinations are, as such, utterly valueless, yet as they were for many years extensively accepted in English and American scientific literature they are inserted here for reference. In the following table Thomson's numbers are multiplied, when necessary, for the sake of comparison with the values now accepted.

DETERMINATIONS INVOLVING BARIUM = 70.

Arsenic76	Magnesium24
	Manganese56
Bismuth216	Nickel52
	Nitrogen14
Carbon12	Phosphorus32
Chlorine36	Potassium40
	Silver110
Cobalt52	Sodium24
Copper64	Strontium88*
Iron56	Sulphur32
Lead208	Zinc68
	189

THOMSON FURTHER DETERMINED-

Antimony at 132 by oxidation.

Boron at 12 from analysis of borax.

Mercury at 200 by conversion of the oxide into chloride.

Tin at 116 by oxidation with nitric acid.

REDUCTION OF WEIGHINGS TO VACUUM.

In discussing the analyses recorded in the foregoing pages, or in reconciling atomic weight determinations by various chemists, it may be found convenient to employ the following table. The maximum error involved is less than 0.01 per cent. or 0.1 milligram per gram.

GRAM WEIGHTS BEING OF BRASS, FRACTIONS OF PLATINUM.

For substances the sp. gr. of which exceeds 6.1; no correction is necessary.

For substances the sp. gr. of which is less than 6.1:—

To correct the entire grams; multiply their number by the correction in the table opposite the sp. gr. of the substance, found in the first column, and add the product to the observed

weight.

To correct the fractions of a gram, multiply the correction opposite the sp. gr. of the substance, found in the third column of the table, by the first two decimal figures of the observed weight, if the sp. gr. of the substance is less than 3, and by the first decimal only, if the sp. gr. exceeds 3, and add the product to the observed weight.

ALL WEIGHTS USED BEING OF PLATINUM.

For substances the sp. gr. of which exceeds 7.8, no correc-

tion is necessary.

For substances the sp. gr. of which is less than 7.8:—Multiply the correction opposite the sp. gr. of the substance, found in the third column, by the number of grams, tenths and hundredths observed, if the sp. gr. falls short of 3, or by the number of grams and tenths, if the sp. gr. exceeds 3, and add the product to the observed weight.

The table shows within what limits it is necessary to know

the sp. gr.

	Brass) for ity between—	Correction per Gram $Error < \frac{1}{30} Mg$.	Specific Grave	
	d 11.064	-0.000 067 gram		
11.064	6.904	0.000 000	51.766 and	
6.904	5.019	+0.000 067	18.568	7.807
5.019	8.948	0.000 188	7.807	5.480
8.94 8	8.247	0 000 200	5.480	4.222
3.247	2.759	0.000 267	4.222	8.4 33
2.759	2.899	0 000 888	8.488	2.893
2.899	2.122	0.000 400	2.898	2.500
2,122	1.903	0.000 467	2.500	2.201
1.903	1.724	0.000 588	2.201	1.965
1.724	1.576	0.000 600	1.965	1.776
1.576	1.452	0.000 667	1.776	1.619
1.452	1.877	0.000 788	1.619	1.488
1.877	1.254	0 000 800	1.488	1.877
1.254	1.174	0.000 867	1.877	1.281
1.174	1.103	0.000 988	1.281	1.197
1.103	1.041	0.001 000	1.197	1.124
1.041	0.985	0.001 067	1.124	1.059
- • • • •	*****	0.001 188	1.059	1.002
		0.001 200	1.002	0.950

(Sill. Amer. Jour., 16, 1878, 265; Liebig's Ann., 195, 1879, 222.)



INDEX TO AUTHORITIES.

Acta Universitatis Lundensis, 90.

Afhandlingar i Fysik, etc., 79.

Annales de Chimie et de Physique, 8, 10, 12, 18, 15, 19, 20, 21, 22, 24, 28, 30, 85, 40, 41, 42, 48, 47, 50, 51, 55, 56, 61, 62, 68, 65, 66, 67, 71, 72, 78, 74, 77, 78, 79, 81, 84, 86, 87, 88, 98, 97, 100, 102, 104, 106, 109, 111, 118, 115, 118, 119, 128, 124, 126, 127, 129, 185, 187.

Berlin, Bericht der Deutschen Chemischen Gesellschaft, 18, 48, 49, 54, 59, 89, 105, 184.

Berzelius' Lehrbuch der Chemie, 7, 12, 14, 17, 18, 22, 28, 26, 27,28, 29, 88, 46, 49, 54, 56, 71, 74, 78, 80, 82, 88, 86, 98, 105, 112, 114, 116, 120, 121, 124.

Berzelius' Jahresbericht über die Fortschritte der Chemie, etc., 8, 41, 56, 62, 65, 66, 74, 78, 79, 80, 86, 92, 98, 100, 108, 114, 115, 125, 130, 186.

Bibliothèque Universelle de Genève, Archives des Sciences, 15, 16, 22, 27, 82, 41, 46, 47, 62, 72, 84, 87, 88, 91, 98, 100, 108, 118, 184.

Bihang till Vetenskaps Akademien Handlingar, 184. (See Kongliga Vetenskaps Akademien.)

British Association Reports, 15, 19.

Bulletin de la Société Chimique, 58, 54, 122.

Bulletin de l'Académie Royale des Sciences, etc., de Belgique, 82, 88.

Chemical News, 25, 48, 65, 89.

Edinburgh Royal Society Transactions, 56, 80.

Erdmann's Journal für Praktische Chemie, 18, 28, 24, 27, 28, 29, 81, 85, 87, 44, 45, 48, 49, 52, 55, 58, 59, 60, 62, 67, 68, 69, 76, 77, 78, 79, 81, 88, 86, 88, 90, 92, 94, 96, 102, 104, 114, 117, 119, 122, 126, 127, 180, 186.

Fresenius' Zeitschrift für Analytische Chemie, 48, 50, 52, 89.

Forhandlinger ved de Skandinaviske Naturforskeres, 188.

Gilbert's Annalen der Physik, etc., 96, 128, 135.

Gmelin-Kraut, Handbuch der Chemie, 7, 9, 12, 18, 18, 19, 20, 21, 28, 26, 37, 48, 46, 49, 56, 62, 64, 65, 68, 78, 76, 79, 81, 88, 86, 90, 91, 94, 95, 96, 98, 101, 102, 108, 106, 110, 111, 118, 117, 118, 122, 128, 125, 128, 184.

Halle, Zeitschrift für die Gesammten Naturwissenschaften, 45.

Jenaische Zeitschrift für Medicin und Naturwissenschaft, 52.

Journal de Pharmacie et de Chimie, 94.

Journal of the Chemical Society, 47, 88, 97.

Klatzo, Ueber die Constitution der Beryllerde, 18.

Kongliga Vetenskaps Akademien Handlingar, 21, 83, 94, 95, 101.

Kopp's Jahresbericht über die Fortschritte der Chemie, 45, 52, 58, 78, 104, 122, 184.

Laurent and Gerhardt's Comptes Rendus Mensuels, etc., 125.

Liebig's Annalen der Chemie, etc., 15, 26, 28, 29, 80, 81, 82, 85, 36, 87, 40, 41, 48, 51, 52, 58, 54, 68, 78, 75, 78, 86, 91, 98, 99, 108, 106, 107, 115, 116, 117, 120, 121, 125, 126, 128, 182, 183, 184.

Meyer, L., Moderne Theorien der Chemie, 7, 9, 12, 20, 28, 81, 118, 122.

Mitscherlich's Lehrbuch der Chemie, 80.

Œfversigt af Vetenskaps Akademien Færhandlingar, 79, 186. (See Kongliga Vetenskaps Akademien.)

Otto's German Translation of Graham's Chemistry, 88, 84, 67.

Paris Comptes Rendus, 8, 18, 15, 20, 26, 40, 42, 50, 55, 56, 58, 75, 88, 85, 89, 98, 95, 97, 100, 105, 111, 112, 116, 181, 185.

Pelouze, Traité de Chimie, 8.

Philosophical Magazine, 8, 17, 22, 80.

Philosophical Transactions of the Royal Society, 14, 26, 28, 88, 87, 28, 39, 48, 49, 54, 56, 65, 71, 72, 76, 82, 91, 92, 96, 99, 106, 107, 110, 112, 118, 114, 120.

Poggendorff's Annalen der Physik, etc., 9, 10, 11, 12, 18, 14, 17, 18, 19, 21, 24, 25, 26, 29, 88, 86, 88, 48, 45, 46, 49, 51, 53, 54, 59, 60, 61, 64, 65, 66, 67, 68, 91, 94, 95, 98, 99, 101, 102, 108, 104, 105, 106, 110, 114, 116, 118, 121, 122, 128, 124, 125, 127, 128, 129, 180, 181, 132, 188, 185, 186.

Proceedings of the American Academy of Arts and Sciences, 11, 12.

Schweigger's Journal für Chemie und Physik, 28, 86.

Scheikundige Verhandelingen en Onderzoekingen, 122.

Silliman's American Journal of Science, 25, 26, 87, 46, 74, 75.

Sitzungs-Bericht der k. k. Akademie zu Wien, 48, 81, 88, 97, 104, 118.

Stas, Untersuchungen über Chemischen Proportionen und Atomgewichte, 28, 48, 59, 62, 64, 78, 76, 94, 98, 101, 110, 111, 115.

Stockholm Akademien Handlingar, 18. (See Kongliga Vetenskaps Akademien.)

Thomson's System of Chemistry, 7, 14, 17, 20, 28, 24, 29, 61, 78.

Thomson's Annals of Philosophy, 58, 61, 189.

Thomson, R. D., Records of General Science, 92, 114.

Zeitschrift für Berg Hütten-und Salinen-Wesen im Preussichen Staate, 49, 51.

INDEX TO ATOMIC WEIGHT DETERMINATIONS.

	•	
ALLEN. (JOHNSON and)	BERZELIUS, J. J. Chlorine 37, 38	Borck, J. B. von
Cæsium 25	Chlorine 37, 38	Tungsten126
wantitum	Curomum 43	DOIOH 19, 140
ANDERSON, T.	Copper 49	Brandes, R.
Nitrogen 93	Fluorine 54	Manganese 80
Andrews, T.	Gold 56	Brodie, B. C.
Barium 15	Iodine 61	Graphon 22
Platinum 98	Iridium 64	Phosphorus 97
Antimony Q. 140	Iron 65, 66	Bromine 20
ARAGO. (BIOT and) Carbon 28	Lead 71	
Carbon28	Lithium 74	Cerium 37
Nitrogen 91	Magnesium 76	Bunsen, R. W.
ARFVEDSON, I. A.	Manganese 70.80	Cæsium 25
Lithium 73	Molybdenum 82	Cæsium 25 Indium 60
Manganese 79	Nickel 87	Bunsen, R.W., and J. JEGEL.
Uranium128, 129	Nitmoren of	Cerium 35
Arsenic 12	Oemium 04	Bunsen, R. W. (J. F. Bahr
AWDEJEW.	Palladium 95	and)
Beryllium 17	Phoenhouse of	and) Erbium 53
-crymum 17	Distinum of	Yttrium 134
Ritto T P		
BAHR, J. F.		Bunsen. (Kirchoff and)
Magnesium 78	Rhodium101	Cæsium 24
BAHR, J. F., and R. W.	Selenium104	
Bunsen.	Silicon 105	
Erbium 53	Silver 106	Cadmium 23
Yttrium 134	Sodium110	Cæsium 24
BALARD, A. J.	Sulphur 113, 114	Calcium 26, 139 CAPITAINE, H.
Bromine 20	Tantalium 116	CAPITAINE, H.
Barium 13	Tellurium118	
BERINGER, A.	Thorium120	Carbon 28
Cerium 34	Tin122	Cerium 34
BERLIN, N. J.	Tungsten 125	
Chromium 44	Uranium129	Copper 49
Molybdenum 84	Vanadium131	
Thorium121	Yttrium 133	CHOUBINE.
Yttrium 133	Zinc134	
Bernoulli, F. A.	Zirconium 136	Chromium 43
Tungsten 127		CHYDENIUS, J. J.
Berthier, P.	BERZELIUS and DULONG.	Thorium121
Nickel 86	Hydrogen 57	
Beryllium 16	Nitrogen 91	Iridium 65
Berzelius, J. J.	BERZELIUS and LIEBIG.	Ruthenium103
Aluminium 7		CLEVE, P. T.
	BIOT and ARAGO.	Didymium 53
Arsenic 12		Erbium 53
Barium 14		Lanthanium 70
	Bismuth 18, 139	Thorium121
		CLEVE, P. T., and O. M.
	BLOMSTRAND, C. W. Niobium 90	
Bromine 21	ROTERATION AND TOP	Trkina =
Carbo-	Boisbaudran, L. de	Erbium 54
Carbon 29	Gallium 56	Yttrium134
10		145

Cobalt 46, 139	Dumas, J. Magnesium 79	GAY-LUSSAC, L. J.
COMMAILLE. (MILLON and)	Magnesium 79	Zinc134
Copper50		GERHARDT, C.
Cook 3, J. P., Jr.	Molybdenum 84	Chlorine 41
Antimony II		GIBBS, W.
Copper 49, 139	Nitrogen 93	Cerium37
Crookes, W.	Phosphorus 97	Cobalt 46
Thallium119, 120	Potassium100	
Czudnowicz, C.	Selenium 104	Lithium 73
Lanthanium 69		
Dame II	SodiumIII	Boron19
DAVY, H.	Suonuum113	Carbon 33
Fluorine 54	Sulphur115 Tellurium118	Copperson P
Silver106	Tin	Consistence of
Davy I	Tin122	Pubidium 101
DAVY, J.	Tungsten 126 Dumas and Stas.	Gold re
Debray, H.	Carbon 30	Graphon 33
Molybdenum 84	Carbon30	Grapuon
	FREINEN I I	Hagen, R.
DEBRAY. (DEVILLE and) Osmium 94	EBELMEN, J. J.	Tithinm 74
Delafontaine, M.	FINDRODT P	HAMP W
Erbium 53	Uranium129 EINBRODT, P. Nitrogen 93	Conner to
Molybdenum 84	EKMAN, G. (O. PETTERSSON	HATTER K VON
Thorium121	and)	Cadmium 23
Vttrium 122 124	Selenium tos	Manganese 80
Demoly, A.	Erhium 52	Tellurium 118
Titanium 124	and) Selenium 105 Erbium 53 ERDMANN, A.	HERRELING. M.
Deville, H. Sainte-Claire	Zinc 125	Thallium119
Nickel 87	ERDMANN and MARCHAND.	HENRY. W.
DEVILLE and DEBRAY.	Calcium 27	Magnesium 76
Osmium 94	Carbon 30	HERMANN, R.
DEVILLE. (WOEHLER and)	Copper 49	
Boron19	Hydrogen 58	Didymium 51
DEXTER, W. P.	Iron 66	Didymium 51 Lanthanium 68, 69
Antimony to	Mercury 82	Lithium 73
Didymium51	Nickel 86	Niobium 00
DIEHL, K.	Selenium 104	Tantalium117
Lithium 75	Sulphur114	Zirconium136 HILLEBRAND, W. F.
	Erk, C.	HILLEBRAND, W. F.
Hydrogen 57		Didymium 53
Nitrogen 91	_ *	HISINGER, W.
Dumas, J.	·	Cerium 34
Aluminium 8	Favre, P. A.	Hoeglund, O. M. (P. T.
Antimony 10	Zinc 135	CLEVE and)
Arsenic 12, 13	Fluorine 54	Erbium 54
Barium 16	FOURCROY and THENARD.	Yttrium 134
Bismuth 19	Mercury 82	HOLZMANN, M.
	I ————————————————————————————————————	Lanthanium 68
	Carbon 30	Hydrogen 57
Calcium 26, 28	- · · · · · · · · · · · · · · · · · · ·	
Carbon 29	Fluorine 55	Indium 59
Chlorine 42	Osmium 94	Iodine61
Cobalt 47	FREMY. (PELOUZE and)	Iridium 64
Copper50		
Fluorine 55		ISNARD.
Hydrogen 58	Gallium 55	Aluminium 8
Iodine 61, 62	GAY-LUSSAC, L. J.	T
Iron 67		JACQUELAIN, V. A.
Lead 72	Magnesium 76	Chromium 44


INDEX.

JACQUELAIN, V. A.	LIEBIG, J.	MARIGNAC, C.
Iodine 62	Bromine 21 LIEBIG and REDTENBACHER.	Silver 107, 108
Magnesium 78	LIEBIG and REDTENBACHER.	Strontium112
Phosphorus 97	Carbon 31 Silver 108 LIEBIG. (BERZELIUS and)	Tantalium 11
Zinc 125	Silver 108	Zirconium 12
Ircr I /P Rinery and)	LIPPIC (REPORTING and)	MATURE W W
Comments of	Corbon	Alemainium
Cerium 35	Carbon 31 Lithium 73	Aluminium
JOHNSON and ALLEN.	Lithium 73	MAUMENE, E. J.
Cæsium 25	Loewig, C.	Chlorine 41
	Bromine 21	Iron 67 Potassium 100
Vmm (I manne and)	Longchamp.	Potassium100
KEMPE. (LEICHTE and) Molybdenum 85	Magnesium 76	Silver108
Molybdenum 85	LOUYET, P.	MENDELRIREE D.
Kessler, F.		Cerium
Antimony 10 Arsenic 12	I Terope C mm	T: J
Arsenic 13	LUCCA, S. DE	Didymium 52 Erbium 53 Lanthanium 70
Chromium 45	Fluorine 55	Eroium 53
Venezue - J Demeses		Lanthanium 70
	! MACDONNETT A	l Branium ros
Dakidian 24	Magnesium 78	MERCER.
Kubidium 102	Magnesium 78 Magnesium 76, 139 MAGNUS, G. Iron 65	Cæsium 26
KJERULP, T.	MAGNUS. G	Mercury 81 140
Cerium 35	Iron 65	Meven I
Klaproth, M. H.		,,,
Potassium 99	MALLET, J. W.	molybdenum 85
Strontium 112	Lithium 74	Uranium128
KLAPROTH. (WOLLASTON	Manganese 79, 139	MILLON, E.
and)	Chlorine 37 Silver 106	Mercury 83
Вапит 13	Silver106	MILLON and COMMAILLE.
Klatzo, G.	MARCHAND, R. F.	Copper
Beryllium 18	Tungsten 126	Mitscherlich, E.
KRALOVANSZKY.	Timmium	Carbon
Lithium 73	Uranium129 MARCHAND and SCHEERER.	Manne A
KRAUT. (GMELIN-)	MARCHAND and Scheerer.	MOBERG, A.
Boron 19	Magnesium 76	Chromium 44
Corbon	residential (Dichitalis	Molybdenum 83
Carbon 33	and)	Mosander, C. G. Lanthanium 68
Iodine 62	Calcium 27	Lanthanium 68
İ	Carbon 20	Titanium124
Lagerhjelm, P.	Carbon 30 Copper 49	MULDER, G. I.
Bismuth 18	Hydrogen	Tin
	Hydrogen 58	1 111 122
LAMY, A.	Iron 66	37:4 .1 .00
Thallium118	Mercury 82	Nickel 86, 139
Lanthanium 67		Niobium 89
Lassaigne.	Selenium104	Nitrogen 91
Nickel 86	Sulphur114	NORDENFELDT. (SVANBERG
Laurent, A.	MARIGNAC, C.	and)
Boron 20	Barium 15. 16	Magnesium 77
Chlorine 40, 42	Romine 21	Magnesium 77 Norlin. (Svanberg and) Iron 66
I end to too	Coloium	Iron 66
Lead 70, 139	Carbon 27	11011 00
Lee, R. H.	Carbon 32	O 337
Cobalt 48	Cerium 35	ODLING, W.
Nickel 89	Chlorine 39, 40	Aluminium 8
LEFORT, J.	Cobalt 46	Osmium 94
Chromium 45	Didymium 51	OTTO, F. J.
LEICHTE and KEMPE.	Iodine 62	Cerium 34
Molybdenum 85	Lanthanium 60	Lanthanium 67
	Tand	Ovugan
enssen, E.	Lead 72	Oxygen 95
Cadmium 24	Nickel 87	75 11 11
Levol, A.		Palladium 95
Gold 56		Peligot, E.
Potassium100	Potassium 99, 100	Chromium 44
	1 1 2 2 7 7 7 3 4 1	• •


Peligot, E.	ROSCOR, H. E.	Stas, J. S.
Uranium120, 131	Roscoe, H. E. Tungsten128 Vanadium132	Chlorine 42
Pelouze, J.	Vanadium132	Hydrogen 59
Arsenic 12	I KOSR. H.	l lodine 62
Barium 15	Niobium 80, 90	Lead 72
Chlorine 40		Littlium 75
Nitrogen 93	Titanium123	Nitrogen 93
	Rose, H., and Weber.	Potassium100
Potassium100		
Silicon105		SodiumTII
Sodium IIO	Phosphorus 96 Strontium 112	Sulphur115
Delotte and Frence	ROTHORE F	Carbon 30
Aluminium &	ROTHOFF, E. Cobalt 46 Nickel 86	STRECKED A
PRNNY. F.	Nickel 86	Carbon 32
Chlorine 20	Rubidium102	Silver108
Nitrogen 02	Russell, W. J.	STROMEYER, F.
Potassium oo	Russell, W. J. Cobalt 47, 48	Cadmium 23
Silver107	Nickel 88, 89	Iron 65
Sodium110	Ruthenium103	Lithium 73
PETTERSSON, O., and G.		Strontium112
EKMAN.	SACC, F.	Strontium111, 139
Selenium105	Selenium104	
PHILLIPS, R.	SALVETAT.	Barium 15
Chlorine 39	Barium 15 Calcium 26 Strontium 112	Support 115
Process I	Strontium 772	and)
Ruhidium 102	SCHEERER, T.	Molyhdenum 82
Pierre, J.	Magnesium 77. 78	Sulphur 113, 139
	SCHEERER, T. (MARCHAND	Svanberg, L.
Platinum 98	and)	Mercury 82
Popp, O.	Magnesium 76, 77	Nitrogen 92 SVANBERG and NORDEN-
Yttrium 133	Scheibler, C.	SVANBERG and NORDEN-
Potassium 98, 139	Tungsten127	FELDT.
PROUT, W.	Schiel, J.	Magnesium 77 SVANBERG and NORLIN.
Iodine 61	Silicon106	SVANBERG and NORLIN.
Dansar appea	Schneider, R.	Iron 66
RAMMELSBERG, C.	Riemuth to	SVANBERG and STRUVE.
Cerium 35, 36 Lanthanium 68	Cobalt 46	Molybdenum 83
Niobium 90		Tantalium116
Uranium130, 131	Nickel87. 88	Tellurium117
RAWACK.	Tungsten125	Thalén.
	SCHROETTER, A.	Lanthanium 70
REDTENBACHER. (LIEBIG	SCHROETTER, A. Phosphorus 97	Thallium118
and)	Selenium 104	THENARD, L. J., and F. H.
Carbon 31	SEFSTROEM, N. G. Mercury 82	Wollaston.
	Mercury 82	Iron 65
REGNAULT.	Selenium103	THENARD L. J., (FOURCEOY
Hydrogen 57	SIEWART, M.	and)
Nitrogen 91	Silicon105	Mercury 82
REICH, F., and T. RICHTER. Indium 59	Silver 106 120	Hydrogen so
Riche, A.	Sodium	Hydrogen 59 Thomson, T.
	SOMMARUGA. E. VON.	Aluminium _ 7
RICHTER, T. (F. REICH and)	Sommaruga, E. von. Cobalt 47	Antimony140
Indium 59	Nickel 88	Arsenic139
Rivor, L. E.	Stas, J. S.	Barium 14, 130
Iron 67	Bromine 22	Beryllium 17
RhodiumIOI	Carbon32, 33	Bismuth139
•		

INDEX.

THOMSON, T.	Turner, E.	WOEHLER and DEVILLE.
Boron 20, 140	Nitrogen 91	Boron 19
Cadmium 23	Silver 106 Sulphur 114	Graphon 33
Calcium139	Sulphur 114	Wolf, C.
Carbon29, 139		Cerium 36
Cerium 34		WOLLASTON, F. H.
	Antimony II	
	Uranium128	
Cobalt 139		Chlorine 37
	Vanadium131	Copper 49
	VAUQUELAIN, L. N.	Hydrogen 57
	Lithium 73	Magnesium 76
Iodine 61		Mercury 82
Iron139		Nitrogen 91
Lead139		Phosphorus 96
Lithium 73		Potassium 99
Magnesium 139	WACKENRODER, H.	Silver106
Manganese139		Sodium 110
Mercury140		Strontium112
Nickel 130	Bromine 22	Sulphur113
Nitrogen91, 139	WATTS, W. M.	Zinc134
Phosphorus139	Íridium 65	WOLLASTON and KLAPROTH.
	Osmium 94	Barium 13
Silver139	WEBER. (H. Rose and)	WOLLASTON. (BERZELIUS
	Antimony 9	•
Strontium139	WEEREN, J.	Lead 71
Sulphur 114	Beryllium 17	WOLLASTON. (L. J. THE-
Tin140	WELESKY, P.	NARD and)
Zinc139	Cobalt 48	Iron 65
Thorium120		WREDE, F. VON
Tin122, 140	Silver 106	Carbon 32
Tissier, C.	WERTHEIM, J.	3
Aluminium 8	Uranium130	
Titanium123	WERTHER, H.	Yttrium133
TROOST, L.	ThalliumI19	30
Lithium74, 75	WILDENSTEIN, R.	
Tungsten125	Chromium 45	ZETTNOW, E.
TURNER, E.	Wing, C. H.	Tungsten 1 27
Barium 14	Cerium 36	Zinc134, 139
Chlorine 38	WINKLER, C.	Zirconium 1 36
Lead 71	Cobalt 48	ZSCHIESCHE, H.
Manganese 80	Indium 60	Didymium 52
Manganese 80 Mercury 82	Nickel 88	
microusy		







SMITHSONIAN MISCELLANEOUS COLLECT

CONSTANTS OF NATU

PART V.

A RECALCULATION

-0F-

THE ATOMIC WEIGH

ΒY

FRANK WIGGLESWORTH CLARKE, S. B.,
Professor of Chemistry and Physics in the University of Cincit



WASHINGTON: BMITHSONIAN INSTITUTION. 1882.



•

•

•

•

•

•

•

•

1

ADVERTISEMENT.

The present publication is one of a series devoted to the discussion and more precise determination of various "Constants of Nature;" and forms the *Fifth* contribution to that subject published by this Institution.

The First number of the series, embracing tables of "Specific Gravities" and of Melting and Boiling Points of Bodies, prepared by the same author, Prof. F. W. Clarke, was published in 1873. The Fourth part of the series, comprising a complete digest of the various "Atomic Weight" determinations of the chemical elements published since 1814, commencing with the well-known "Table of Equivalents" by Wollaston, (given in the Philosophical Transactions for that year,) compiled by Mr. George F. Becker, was published by the Institution in 1880. The present work which may be regarded as practically supplementary to that digest, (or perhaps rather as the memoir to which that digest is introductory,) comprises a very full discussion and re-calculation of the "Atomic Weights" from all the existing data, and the assignment of the most probable value to each of the elements.

The manuscript of the work was presented to the Institution in its completed form by Prof. F. W. Clarke, the cost of publication only being at the expense of the Smithsonian fund.

SPENCER F. BAIRD, Secretary of Smithsonian Institution.

WASHINGTON, January, 1882.




TABLE OF CONTENTS.

Introduction	
Formulæ for the Calculation of Probable Error	
1. Oxygen	
2. Silver, Potassium, Sodium, Chlorine, Bromine, Iodine, and Sulp	hur
3. Nitrogen	~
4. Carbon	
5. Barium	
6. Strontium	
7. Calcium	
8. Lead	
9. Fluorine	
Phosphorus	
BI. Boron	÷
12. Silicon	
13. Lithium	
14. Rubidium	
15. Cæsium	
16. Thallium	
17. Glucinum	
18. Magnesium	
19. Zinc	
20. Cadmium	
21. Mercury	
22. Chromium	
23. Manganese	
24. Iron	
25. Copper	
26. Molybdenum	
27. Tungsten	
28. Uranium	
29. Aluminum	
30. Gold	
Nickel and Cobalt	
22 Selenium	
Tellurium	
17-madium	
A	
7. Bismuth	• •
7. Bishida (v	·)

CONTENTS.

		Page.
_	Tin	204
39.	Titanium	207
40.	Zirconium	212
4 I.	Thorium	214
42.	Gallium	218
43.	Indium	219
44.	Cerium	220
45.	Lanthanum	229
	Didymium	236
47.	The Yttrium Group. Scandium, Yttrium, Ytterbium, Erbium, Ter-	_
	bium, Phillipium, Decipium, Thulium, Samarium, etc.	240
48 .	Columbium (Niobium)	247
49.	Tantalum	248
50.	Platinum	249
51.	Osmium	254
	Iridium	255
_	Palladium	256
54.	Rhodium	258
•	Ruthenium	259
	oendix	261

INTRODUCTION.

In the autumn of 1877 the writer began collecting data relative to the determinations of atomic weights, with the purpose of preparing a complete resumé of the entire subject, and of recalculating all the estimations. The work was fairly under way, the material was collected and partly discussed, when I received from the Smithsonian Institution a manuscript by Professor George F. Becker, entitled "Atomic Weight Determinations: a Digest of the Investigations Published since 1814." This manuscript, which has lately been issued as Part IV of the "Constants of Nature," covered much of the ground contemplated in my own undertaking. It brought together all the evidence, presenting it clearly and thoroughly in compact form; in short, that portion of the task could not well be improved ¹¹Pon. Accordingly, I decided to limit my own labors to a critical recalculation of the data; to combine all the figures "Pon a common mathematical basis, and to omit everything *hich could as well be found in Professor Becker's "Digest."

At the very beginning of my work certain questions confronted me. Should I treat the investigations of different individuals separately, or should I combine similar data together in a manner irrespective of persons? For example, ought I, in estimating the atomic weight of silver, to take Stas' work by itself, Marignac's work by itself, and so on, and then average the results together; or should I rather combine all series of figures relating to the composition of Potassium chlorate into one mean value, and all the data concerning the composition of silver chloride into another mean, and, finally, compute from such general means the constant sought to be established? The latter plan was finally adopted; in fact, it was rendered necessary by the method of least squares, which method was alone adequate to supply me with good processes for calculation.

The mode of discussion and combination of results was briefly as follows. The formulæ employed are given in another chapter. I began with the ratio between oxygen and hydrogen; in other words, with the atomic weight of oxygen referred to hydrogen as unity. Each series of experiments was taken by itself, its arithmetical mean was found, and the probable error of that mean was computed. Then the several means were combined according to the appropriate formula, each receiving a weight dependent upon its probable error. The general mean thus established was taken as the most probable value for the atomic weight of oxygen, and, at the same time, its probable error was mathematically asssigned.

Next in order came a group of elements which were best discussed together, namely, silver, chlorine, potassium, sodium, bromine, iodine, and sulphur. For these elements there were data from thirteen experimenters. All similar figures were first reduced to common standards, and then the means of individual series were combined into general Thus all the data were condensed into twenty ratios, from which several independent values for the atomic weight of each element could be computed. The probable errors of these values, however, all involved the probable error of the atomic weight of oxygen, and were, therefore, higher than they would have been had the latter element not entered into consideration. Here, then, we have suggested a chief peculiarity of this whole revision. atomic weight of each element involves the probable errors of all the other elements to which it is directly or indirectly Accordingly, an atomic weight determined by reference to elements whose atomic weights have been defectively ascertained will receive a high probable error, and its weight, when combined with other values, will be relatively For example, an atomic weight ascertained by direct comparison with hydrogen will, other things being equal, have a lower probable error than one which is referred to hydrogen through the intervention of oxygen; and a metal whose equivalent involves only the probable error of oxygen

will be more exactly known than one which depends upon the greater errors of silver and chlorine. These points will appear more clearly evident in the subsequent actual discussions.

But although the discussion of atomic weights is ostensibly mathematical, it cannot be purely so. Chemical considerations are necessarily involved at every turn. In assigning weights to mean values I have been, for the most part, rigidly guided by mathematical rules; but in some cases I have been compelled to reject altogether series of data which were mathematically excellent, but chemically worthless because of constant errors. In certain instances there were grave doubts as to whether particular figures should be included or rejected in the calculation of means; there having been legitimate reasons for either procedure. Probably many chemists would differ with me upon such points of judgment. In fact, it is doubtful whether any two chemists, working independently, would handle all the data in precisely the same way, or combine them so as to produce exactly the same final results. Neither would any two mathematicians follow identical rules or reach identical conclusions. In calculating the atomic weight of any element those values are assigned to other elements which have been determined in previous chapters. Hence a variation in the order of discussion might lead to slight differences in the final results.

As a matter of course the data herein combined are of very unequal value. In many series of experiments the weighings have been reduced to a vacuum standard; but in most cases chemists have neglected this correction altogether. In a majority of instances the errors thus introduced are slight; nevertheless they exist, and interfere more or less with all attempts at a theoretical consideration of the results. For example, they affect seriously the investigation of Prout's hypothesis, and are often great enough to account for seeming exceptions to it. Such questions as these will be considered in the appendix.

Another serious source of error affecting many of the re-

sults was not discovered until recently. A large number of computations had been actually finished, involving, among other things, the greater part of Stas' work, when Dumas published his investigation upon the occlusion of oxygen by silver. Here it was shown that a very great number of atomic weight determinations must have been vitiated by constant errors, which, though constant for each series, were probably of different magnitude in different series, and, therefore, could not be systematically corrected for. At the time of the announcement of this discovery of Dumas my work was so far under way that I thought it best to complete my discussion without reference to it, and then to study its influence in the appendix. In the chapter upon aluminum, however, it will be noted that Mallet eliminated this error in great part from his experimental results.

Necessarily, this work omits many details relative to experimental methods, and particulars as to the arrangements of special forms of apparatus. For such details original memoirs must be consulted. Their inclusion here would have rendered the work unwarrantably bulky. There is such a thing as over-exhaustiveness of treatment, which is equally objectionable with under-thoroughness.

Of course, none of the results reached in this revision can be considered as final. Every one of them is liable to repeated corrections. To my mind the real value of the work, great or little, lies in another direction. The data have been brought together and reduced to common standards, and for each series of figures the probable error has been determined. Thus far, however much my methods of combination may be criticized, I feel that my labors will have been useful. The ground is now cleared, in a measure, for future experimenters; it is possible to see more distinctly what remains to be done; some clues are furnished as to the relative merits of different series of results. I hope to be able, from time to time, as new determinations are published, to continue the task here begun, and perhaps, also, to add, in the near future, some data of my own establishing.

In addition to the usual periodicals the following works

- have been freely used by me in the preparation of this volume:
- Berzelius, J. J. Lehrbuch der Chemie. 5 Auflage. Dritter Band. SS. 1147-1231. 1845.
- Van Geuns, W. A. J. Præve eener Geschiedenis van de Æquivalentgetallen der Scheikundige Grondstoffen en van hare Soortelijke Gewigten in Gasvorm, voornamelijk in Betrekking tot de vier Grondstoffen der Bewerktuigde Natuur. Amsterdam, 1853.
- MULDER, E. Historisch-Kritisch Overzigt van de Bepalingen der Æquivalent-Gewigten van 13 Eenvoudige Ligchamen. Utrecht, 1853.
- MULDER, L. Historisch-Kritisch Overzigt van de Bepalingen der Æquivalent-Gewigten van 24 Metalen. Utrecht, 1853.
- Oudemans, A. C., Jr. Historisch-Kritisch Overzigt van de Bepaling der Æquivalent-Gewigten van Twee en Twintig Metalen. Leiden, 1853.
- STAS, J. S. Untersuchungen über die Gesetze der Chemischen Proportionen über die Atomgewichte und ihre gegenseitigen Verhältnisse. Uebersetzt von Dr. L. Aronstein. Leipzig, 1867.

The four Dutch monographs above cited are especially valuable. They represent a revision of all atomic weight data down to 1853, as divided between four writers.

FORMULÆ FOR THE CALCULATION OF PROBABLE ERROR.

Although the ordinary formula for the probable error of an arithmetical mean is familiar to all physicists, it is perhaps best to reproduce it here, as follows:

(1.)
$$e = \pm .6745 \sqrt{\frac{S}{n (n-1)}}$$

Here n represents the number of observations or experiments in the series, while S is the sum of the variations of the individual results from the mean.

In combining several arithmetical means, representing several series, into one general mean each receives a weight indicated by its probable error; greater as the latter becomes less, and vice versa. Let A, B, C, etc., be such mean results, and a, b, c, their probable errors respectively. Then the general mean is determined by this formula:

(2.)
$$M = \frac{\frac{A}{a^3} + \frac{B}{b^2} + \frac{C}{c^2}}{\frac{1}{a^2} + \frac{1}{b^3} + \frac{1}{c^3}} \dots$$

For the probable error of this general mean we have:

(3.)
$$m = \frac{1}{\sqrt{\frac{1}{a^2} + \frac{1}{b^2} + \frac{1}{c^2}}} \dots$$

In the calculation of atomic and molecular weights the following formulæ have been employed. For assistance in connection with them my thanks are due to Professors H. T. Eddy and E. W. Hyde of the University of Cincinnati.

Using, as before, capital letters to represent known quantities and small letters for their probable errors respectively,

we have for the sum or difference of two quantities, A and B:

$$(4.) \qquad \epsilon = \sqrt{a^1 + b^2}$$

For the product of A multiplied by B the probable error is

(5.)
$$e = \sqrt{(Ab)^2 + (Ba)^2}$$

For the product of three quantities, ABC:

(6.)
$$e = \sqrt{(BCa)^2 + (ACb)^2 + (ABc)^2}$$

For a quotient, $\frac{B}{A}$, the probable error becomes

Given a proportion, A:B::C:x, the probable error of the fourth term is as follows:

(8.)
$$e = \frac{\sqrt{\left(\frac{BCa}{A}\right)^2 + (Cb)^2 + (Bc)^2}}{A}$$

This formula is used in nearly every atomic weight calculation, and is, therefore, exceptionally important. Rarely a more complicated case arises in a proportion of this kind:

$$A:B::C+x:D+x$$

In this proportion the unknown quantity occurs in two terms. Its probable error is found by this expression, and is always large:

(9.)
$$e = \sqrt{\frac{(C-D)^2}{(A-B)^4}} \left(B^2 a^2 + A^2 b^2\right) + \frac{B^2 c^2 + A^2 d^2}{(A-B)^2}$$

When several independent values have been calculated for an atomic weight they are treated like means, and combined according to formulæ (2) and (3.) Each final result

is, therefore, to be regarded as the general mean of all reliable determinations. This method of combination may not be the best one theoretically possible, but it seemed to be the only one practically available. The data are too imperfect to warrant the use of much more elaborate processes of discussion.





RECALCULATION OF THE ATOMIC WEIGHTS.

OXYGEN.

The ratio between oxygen and hydrogen is the foundation upon which the entire system of atomic weights depends. Hence, the accuracy of its determination has, from the beginning, been recognized as of extreme importance. A trifling error here may become cumulative when repeated through a moderate series of other ratios.

Leaving out of account the earliest researches, which have now only a historical value, we find that three methods have been employed for fixing this important constant. First, the synthesis of water, effected by passing hydrogen gas over red hot oxide of copper. Secondly, the exact determination of the relative density of the two gases. Thirdly, by weighing the quantity of water formed upon the direct union of a known volume of hydrogen with oxygen.

The first of these methods has been employed in three leading investigations, namely, by Dulong and Berzelius,* by Dumas, and by Erdmann and Marchand. The essential features of the method are in all cases the same. Hydrogen gas is passed over heated oxide of copper, and the water thus formed is collected and weighed. From this weight and the loss of weight which the oxide undergoes, the exact composition of water is readily calculated. Dulong and Berzelius made but three experiments, with the following results for the percentages of oxygen and hydrogen in in water:

Ο.	H.
88.942	11.058
88.809	11.191
88.954	11.046

Thomson's Annals of Philosophy, July, 1821, p. 50.

These figures, rather roughly determined, and by no means exact enough to meet the requirements of modern science, give a mean value of 16.021 for the atomic weight of oxygen. As the weighings were not reduced to a vacuum, this correction was afterwards applied by Clark,* who showed that these syntheses really make O = 15.894; or, in Berzelian terms, if O = 100, H = 12.583.

In 1842 Dumast published his elaborate investigation upon the composition of water. The first point was to get pure hydrogen. This gas, evolved from zinc and sulphuric acid, might contain oxides of nitrogen, sulphur dioxide, hydrosulphuric acid, and arsenic hydride. These impurities were removed in a series of wash bottles; the H₂S by a solution of lead nitrate, the H_sAs by silver sulphate, and the others by caustic potash. Finally, the gas was dried by passing through sulphuric acid, or, in some of the experiments, over phosphorus pentoxide. The copper oxide was thoroughly dried, and the bulb containing it was weighed. By a current of dry hydrogen all the air was expelled from the apparatus, and then, for ten or twelve hours, the oxide of copper was heated to dull redness in a constant stream of the gas. The reduced copper was allowed to cool in an atmosphere of hydrogen. The weighings were made with the bulbs exhausted of air. The following table gives the results:

Column A contains the symbol of the drying substance. B gives the weight of the bulb and copper oxide. C, the weight of bulb and reduced copper. D, the weight of the vessel used for collecting the water. E, the same, plus the water. F, the weight of oxygen. G, the weight of water formed. H, the crude equivalent of H when O = 10,000. I, the equivalent of H, corrected for the air contained in the sulphuric acid employed. This correction is not explained, and seems to be questionable.

^{*} Philosophical Magazine, 3d series, 20, 341.

[†] Compt. Rend., 14, 537.

-	ï	1 (•	1248.0	1247.3	1240.0	254.	1255.0	'n	\$, vi	1248.9	6	Ö		vi.	લં	3	i	1254.7	8	1251.5
	H.	1	•	1249.0	1248.1	0.0		56.	54.		1258.3	ď	51.	53.	1257.7	200	Š	50	57.		8	1253.3
	త	•	÷	22.905	23.053	64.044	85.960	49.047	39.178	51.623	67.586	58.320	59.078	67.282	668.69	58.360	63.577	41.390	38.458	36.175	34.677	Means
	6	•	-	20.362	20.495	47.004	76.364	43.571	34.811	45.887	60.031	51.838	52.508	59.789	62.090	51.838	56.483	36.789	34.162	13	30.827	
	ĸi		•	_		-	973.291				-	•		· ~	. (N)	*	\sim	O	~	00	912.539	•
	D.	180 80x			. 439.711	_	887.331	867.159	839.304	824.624	822.660	741.095	874.832	931.487	682.374	741.097	1064.762	878.640	887.817	888.662	877.862	·
	Ü	X	j	4	6.17	58.85	728.182	8.15	27.10	56.73	4.61	84.0	35. I	Š	98.7	8.4	81.3	19.	20.0	27.6	16.8	
	Ä,	201 085	204.163	344.548	316.671	624.829	804.546	533.726	661.915	612.625	904.643	642.325	587.645	673.280	660.855	642.325	937.845	756.352	754.162	759.762	747.652	•
	Α.	CS. H		***************************************	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	P.O.	H SO.	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,		P,0,		H*SO*	P.O.		II, SO,		1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	P.O.		***************************************		

In the sum total of these nineteen experiments, 840.161 grammes of oxygen form 945.439 grammes of water. This gives, in percentages, for the composition of water, oxygen 88.864; hydrogen, 11.136. Hence the atomic weight of oxygen, calculated in mass, is 15.9608. In the following column the values are given as deduced from the individual data given under the headings F and G:

15.994 16.014 16.024 15.992 15.916 15.916 15.943 16.000 15.892 15.995 15.984 15.958 15.902 15.987 15.926 15.992 15.904 15.900 16.015

Mean, 15.9607, with a probable error of \pm .0070.

In calculating the above column several discrepancies were noted, probably due to misprints in the original memoir. On comparing columns B and C with F, or D and E with G, these anomalies chiefly appear. They were detected and carefully considered in the course of my own calculations; and, I believe, eliminated from the final result.

The paper by Erdmann and Marchand* followed closely after that of Dumas. The method of research was essentially the same as that of the latter chemist, varying only in points of comparatively unimportant detail. The results are given in two series, in one of which the weighings were

^{*} Journ. f. Prakt. Chem., 1842, bd. 26, s. 461.

5

not actually made in vacuo, but were, nevertheless, reduced to a vacuum standard. The second series represents actual vacuum weighings. The quantity of water formed in each experiment, was from 41.664 to 95.612 grammes. I give below only the percentages of oxygen and hydrogen in water as deduced from Erdmann and Marchand's data:

First Series.

O.	H.
88.836	11.164
88.821	11.179
88.874	11.126
88.868	11.132

Second Series.

0.	H.
88.887	11.113
88.898	11.102
88.895	11.105
88.899	11.101

Hence, the atomic weight of oxygen is, as follows:

Fi	rst Series.	Second Series.
	15.915	15.997
	15.891	16.015
	15.976	16.010
	15.966	16.016
Mean,	$15.9369, \pm .0138$	Mean, 16.0095, ± .0030

The effect of discussing these two series separately is somewhat startling. It gives to the four experiments in Erdmann and Marchand's second group a weight vastly greater than their other four and Dumas' nineteen taken together. For so great a superiority as this there is no adequate reason; and it is highly probable that it is due almost entirely to fortunate coincidences, rather than to greater accuracy of work. We will, therefore, treat Erdmann and Marchand's experiments as one series, giving all equal weight, and then combine them with the results obtained by Dumas. We now have—

```
O = 15.9607, ± .0070

By Erdmann and Marchand O = 15.9733, ± .0113

General mean O = 15.9642, ± .0060
```

In discussing the relative density of oxygen and hydrogen gases we need only consider the more modern researches of Dumas and Boussingault, and of Regnault. As the older work has some historical value, I may in passing just cite its results. For the density of hydrogen we have .0769, Lavoisier; .0693, Thomson; .092, Cavendish; .0732, Biot and Arago; .0688, Dulong and Berzelius. For oxygen there are the following determinations: 1.087, Fourcroy, Vauquelin, and Séguin; 1.103, Kirwan; 1.128, Davy; 1.088, Allen and Pepys; 1.1036, Biot and Arago; 1.1117, Thomson; 1.1056, De Saussure; 1.1026, Dulong and Berzelius; 1.106, Buff; 1.1052, Wrede.*

In 1841 Dumas and Boussingault† published their determinations of gaseous densities. For hydrogen they obtained values ranging from .0691 to .0695; but beyond this mere statement they give no details. For oxygen three determinations were made, with the following results:

1.1055 1.1058 1.1057

Mean, 1.10567, \pm .00006

If we take the two extreme values given above for hydrogen, and regard them as the entire series, they give us a mean of $.0693, \pm .00013$.

This mean hydrogen value, combined with the mean oxygen value, gives for the atomic weight of the latter element the number 15.9538, \pm .031.

Regnault's researches, published four years later,‡ were of

^{*} For Wrede's work, see Berzelius' Jahresbericht for 1843. For Dulong and Berzelius, see the paper already cited. All the other determinations are taken from Gmelin's Handbook, Cavendish edition, v. 1, p. 279.

[†] Compt. Rend., 12, 1005. Compare also with Dumas, Compt. Rend., 14, 537.

[†] Compt. Rend., 20, 975.

7

a more satisfactory kind. Indeed, they are among the classics of physical science; and probably approach as near to absolute accuracy as is possible for experiment.

For hydrogen three determinations of density gave the following results:

For oxygen four determinations were made, but in the first one the gas was contaminated by traces of hydrogen, and the value obtained, 1.10525; was, therefore, rejected by Regnault as too low. The other three are as follows:

Now, combining the hydrogen and oxygen series, we have for the atomic weight of oxygen, 15.9628, $\pm .0044$.*

Upon combining the result of Regnault's work with that from Dumas and Boussingault's we get the following value:

This result, it will be seen, agrees remarkably well with that obtained in the experiments upon the synthesis of water.

^{*}Since these computations were made, Professor John Le Conte has called my attention to the existence of slight numerical errors in Regnault's own reductions. As corrected by Le Conte, Regnault's figures give 1.105612 for the density of oxygen, and 0.069269 for that of hydrogen. Hence the atomic weight of O becomes 15.9611, instead of 15.9628. The difference is slight, but still it ought not to be ignored. All the computations in the body of this work, having been finished before I received Professor Le Conte's figures, must stand, nevertheless, as they are. For further details Le Conte refers to Phil. Mag., (4,) 27, p. 29, 1864; and also to the Smithsonian Report for 1878, p. 428.

The third method indicated at the beginning of this discussion has been recently employed in part by J. Thomsen* of Copenhagen. Unfortunately this chemist has not published the details of his work, but only the end results. These serve to confirm the values for oxygen fixed by other methods, but they cannot well be included in the systematic discussion. Partly by the oxidation of hydrogen over heated copper oxide, and partly by its direct union with oxygen, Thomsen finds that at the latitude of Copenhagen, and at sea level, one litre of dry hydrogen at 0° and 760 mm. pressure will form .8041 gramme of water. According to Regnault, at this latitude, level, temperature, and pressure, a litre of hydrogen weighs .08954 gramme. these data, O = 15.9605. It will be seen at once that Thomsen's work depends in great part upon that of Regnault, and yet that it affords an admirable reinforcement of the latter.

It is now plain, in conclusion, that all the different lines of research point to an atomic weight for oxygen a little below 16.00. Five distinct investigations confirm each other wonderfully. Upon combining the values obtained by the two chief methods we get the following final results:

```
From synthesis of water..... O = 15.9642, \pm .0060
From gaseous densities..... O = 15.9627, \pm .0043
```

In the general mean the atomic weight of oxygen becomes 15.9633, with a probable error of \pm .0035.†

^{*} Ber. d. Deutsch. Chem. Gesellschaft, 1870, s. 928.

[†] Le Conte's correction of Regnault's figures introduced here would make 0 = 15.9622, instead of 15.9633. Difference, .0011.

SILVER, POTASSIUM, SODIUM, CHLORINE, BROMINE, IODINE, AND SULPHUR.

The atomic weights of these seven elements depend upon each other to so great an extent that they can hardly be considered independently. Indeed, chlorine, potassium, and silver have always been mutually determined. From the ratio between silver and chlorine, the ratio between silver and potassium chloride, and the composition of potassium chlorate, these three atomic weights were first accurately fixed. Similar ratios, more recently worked out by Stas and others, have rendered it desirable to include bromine, iodine, sulphur, and sodium in the same general discussion.

Several methods of determination will be left altogether out of account. For example, in 1842 Marignac* sought to fix the atomic weight of chlorine by estimating the quantity of water formed when hydrochloric acid gas is passed over heated oxide of copper. His results were wholly inaccurate, and need no further mention here. A little later Laurent† redetermined the same constant from the analysis of a chlorinated derivative of naphthalene. This method did not admit of extreme accuracy, and it presupposed a knowledge of the atomic weight of carbon; hence it may be properly disregarded. Maumené's‡ analyses of the oxalate and acetate of silver gave good results for the atomic weight of that metal; but they also depend for their value upon our knowledge of carbon, and will, therefore, be discussed further on with reference to that element.

Let us now consider the ratios upon which we must rely for ascertaining the atomic weights of the seven elements in question. After we have properly arranged our data we may then discuss their meaning. First in order we may

^{*}Compt. Rend., 14, 570. Also, Journ. f. Prakt. Chem., 26, 304.

[†] Compt. Rend., 14, 456. Journ. f. Prakt. Chem., 26, 307.

[†] Ann. d. Chim. et d. Phys., (3,) 18, 41. 1846.

conveniently take up the percentage of potassium chloride obtainable from the chlorate.

The first reliable series of experiments to determine this percentage was made by Berzelius.* All the earlier estimations were vitiated by the fact that when potassium chlorate is ignited under ordinary circumstances a little solid material is mechanically carried away with the oxygen gas. Minute portions of the substance may even be actually volatilized. These sources of loss were avoided by Berzelius, who devised means for collecting and weighing this trace of potassium chloride. All the successors of Berzelius in this work have benefitted by his example; although for the methods by which loss has been prevented we must refer to the original papers of the several investigators. In short, then, Berzelius ignited potassium chlorate, and determined the percentage of chloride which remained. Four experiments gave the following results:

60.850 60.850 60.851

Mean, 60.851, with a probable error of ± .0006

The next series was made by Penny,† in England, who worked after a somewhat different method. He treated potassium chlorate with strong hydrochloric acid in a weighed flask, evaporated to dryness over a sand bath, and then found the weight of the chloride thus obtained. His results are as follows, in six trials:

60.825 60.822 60.815 60.820 60.823 60.830

Mean, 60.8225, $\pm .0014$

^{*} Poggend. Annalen, 1826, bd. 8, s. 1.

[†] Phil. Transactions, 1839, p. 20.

In 1842 Pelouze* made three estimations by the ignition of the chlorate, with these results:

60.843 60.857 60.830 Mean, 60.843, ± .0053

Marignac, in 1842,† worked with several different recrystallizations of the commercial chlorate. He ignited the salt, with the usual precautions for collecting the material carried off mechanically, and also examined the gas which was evolved. He found that the oxygen from 50 grammes of chlorate contained chlorine enough to form .003 gramme of silver chloride. Here are the percentages found by Marignac:

In chlorate once crystallized	60.845
In chlorate once crystallized	60.835
In chlorate twice crystallized	60.833
In chlorate twice crystallized	60.844
In chlorate three times crystallized	60.839
In chlorate four times crystallized	60.839
Mean,	$60.8392, \pm .0013$

In the same paper Marignac describes a similar series of experiments made upon potassium perchlorate, KClO₄. In three experiments it was found that the salt was not quite free from chlorate, and in three more it contained traces of iron. A single determination upon very pure material gave 46.187 per cent. of oxygen and 53.813 of residue.

In 1845 two series of experiments were published by Gerhardt.† The first, made in the usual way, gave these results:

60.871 60.881 60.875 Mean, 60.8757, ± .0020

^{*} Compt. Rend., 15, 959.

[†] Ann. d. Chem. u. Pharm., bd. 44, s. 18.

[‡] Compt. Rend., 21, 1280.

In the second series the oxygen was passed through a weighed tube containing moist cotton, and another filled with pumice stone and sulphuric acid. Particles were thus collected which in the earlier series escaped. From these experiments we get—

60.947 60.947 60.952

Mean, 60.9487, ± .0011

These last results were afterwards sharply criticized by Marignac,* and their value seriously questioned.

The next series, in order of time, is due to Maumené.† This chemist supposed that particles of chlorate, mechanically carried away, might continue to exist as chlorate, undecomposed; and hence that all previous series of experiments might give too high a value to the residual chloride. In his determinations, therefore, the ignition tube, after expulsion of the oxygen, was uniformly heated in all its parts. Here are his percentages of residue:

60.788 60.790 60.793 60.791 60.785 60.795

Mean, 60.791, ± .0009

The question which most naturally arises in connection with these results is, whether portions of chloride may not have been volatilized, and so lost.

Closely following Maumené's paper there is a short note by Faget,‡ giving certain mean results. According to this chemist, when potassium chlorate is ignited slowly, we get

^{*} Supp. Bibl. Univ. de Genéve, Vol. I.

[†] Ann. d. Chim. et d. Phys., (3,) 18, 71. 1846.

[‡] Ann. d. Chim. et d. Phys., (3,) 18, 80. 1846.

60.847 per cent. of residue. When the ignition is rapid, we get 60.942. As no detailed experiments are given, these figures can have no part in our discussion.

Last of all we have two series determined by Stas.* In the first series we have the results obtained by igniting the chlorate. In the second series the chlorate was reduced by strong hydrochloric acid, after the method followed by Penny:

```
First Series.
60.8380
60.8395
60.8440
60.8473
60.8450
Mean, 60.84276, ± .0012

Second Series.
60.850
60.853
60.844

Mean, 60.849, ± .0017
```

In these experiments every conceivable precaution was taken to avoid error and ensure accuracy. All weighings were reduced to a vacuum standard; from 70 to 142 grammes of chlorate were used in each experiment; and the chlorine carried away with the oxygen in the first series was absorbed by finely divided silver and estimated. It is difficult to see how any error could have crept in.

Now, to combine these different series of experiments.

```
Berselius, mean result_____ 60.851, ± .0006
           ----- 60.8225, \pm .0014
Penny,
Pelouze.
           \pm .0053
            ----- 60.8392, ± .0013
Marignac,
           ---- 60.8757, ± .0020
Gerhardt, Ist
           ---- 60.9487, \pm .0011
     2d
           ---- 60.791, \pm .0009
Manmené,
           \pm .0012
Stas.
     Ist
     2d
44
             \pm .0017
   General mean, from all nine series, representing forty
      experiments _____ 60.846, ± .00038
```

^{*} See Aronstein's Translation, p. 249.

This value is exactly that which Stas deduced from both of his own series combined, and gives great emphasis to his wonderfully accurate work. It also finely illustrates the compensation of errors which occurs in combining the figures of different experimenters.

Similar analyses of silver chlorate have been made by Marignac and by Stas. Marignac's figures I have not been able to find,* and Stas gives but two experiments. The following are his percentages of oxygen in silver chlorate:†

Mean, 25.0795, \pm .0010

For the direct ratio between silver and chlorine there are seven available series of experiments. Here, as in many other ratios, the first reliable work was done by Berzelius.‡

He made three estimations, using each time twenty grammes of pure silver. This was dissolved in nitric acid. In the first experiment the silver chloride was precipitated and collected on a filter. In the second and third experiments the solution was mixed with hydrochloric acid in a flask, evaporated to dryness, and the residue then fused and weighed without transfer. One hundred parts of silver formed of chloride:

^{*}Since all the calculations were finished I have secured a copy of Marignac's figures. They are as follows: The third column gives the percentage of O in AgClO₂.

24.510 gr	m. AgC	ClO ₃ gave	18.3616 AgCl.	25.103
25.809	44	66	19.3345 "	25.086
30.306	46	44	22.7072 "	25.074
28.358	46	4.6	21.2453 "	25.082
28.287	66	44	21.1833 "	25.113
57.170	"	66	42.8366 "	25.072

Mean, 25.088, ± .0044

The introduction of these figures into the subsequent calculations could not produce any appreciable result. They would practically vanish from the general mean. However, they serve here as confirmation of Stas' work.

[†] Aronstein's Translation, p. 214.

[†] Thomson's Annals of Philosophy, 1820, v. 15, p. 89.

Turner's work* closely resembles that of Berzelius. Silver was dissolved in nitric acid and precipitated as chloride. In experiments one, two, and three the mixture was evaporated and the residue fused. In experiment four the chloride was collected on a filter. A fifth experiment was made, but has been rejected as worthless.

The results were as follows: In a third column I put the quantity of AgCl proportional to 100 parts of Ag.

```
28.407 grains Ag gave 37.737 AgCl.

41.917 " " 55.678 " 132.829

40.006 " " 53.143 " 132.837

30.922 " " 41.070 " 132.818

Mean, 132.832, ± .0038
```

The same general method of dissolving silver in nitric acid, precipitating, evaporating, and fusing without transfer of material was also adopted by Penny.† His results for 100 parts of silver are as follows, in parts of chloride:

```
132.836
132.840
132.830
132.840
132.840
132.830
```

Mean, 132.8363, ± .0012

In 1842 Marignact found that 100 parts of silver formed 132.74 of chloride, but gave no available details. Later,

^{*} Phil. Transactions, 1829, 291.

[†] Phil. Transactions, 1839, 28.

¹ Ann. Chem. Pharm., 44, 21.

See Berzelius' Lehrbuch, 5th Ed., Vol. 3, pp. 1192, 1193.

in another series of determinations, he is more explicit, and gives the following data: The weighings were reduced to a vacuum standard.

79.853 g	rm. Ag gave	106.080	AgCl.	Ratio,	132.844
69.905	66	92.864	66		132.843
64.905	66	86.210	66		132.825
92.362	66	122.693	66		132.839
99.653	66	132.383	66		132.844
					•

Mean, 132.839, ± .0024

The above series all represent the synthesis of silver chloride. Maumené* made analyses of the compound, reducing it to metal in a current of hydrogen. His experiments make 100 parts of silver equivalent to chloride:

132.734 132.754 132.724 132.729 132.741

Mean, 132.7364, ± .0077

By Dumast we have the following estimations:

9.954 Ag gave 13.227 AgCl. Ratio, 132.882 19.976 " 26.542 " 132.869

Mean, 132.8755, ± .0044

Finally, there are seven determinations by Stas,‡ made with his usual accuracy and with every precaution against error. In the first, second, and third, silver was heated in chlorine gas, and the synthesis of silver chloride thus effected directly. In the fourth and fifth silver was dissolved in nitric acid, and the chloride thrown down by passing hydrochloric acid gas over the surface of the solution. The whole was then evaporated in the same vessel, and the chloride fused, first in an atmosphere of hydrochloric acid,

^{*} Ann. d. Chim. et d. Phys., (3,) 18, 49. 1846.

[†] Ann. Chem. Pharm., 113, 21. 1860.

[‡] Aronstein's Translation, p. 171.

and then in a stream of air. The sixth synthesis was similar to these, only the nitric solution was precipitated by hydrochloric acid in slight excess, and the chloride thrown down was washed by repeated decantation. All the decanted liquids were afterwards evaporated to dryness, and the trace of chloride thus recovered was estimated in addition to the main mass. The latter was fused in an atmosphere of HCl. The seventh experiment was like the sixth, only ammonium chloride was used instead of hydrochloric acid. From 98.3 to 399.7 grammes of silver were used in each experiment, the operations were performed chiefly in the dark, and all weighings were reduced to vacuum. In every case the chloride obtained was beautifully white. The following are the results in chloride for 100 of silver:

```
132.841
132.843
132.849
132.846
132.848
122.8417
Mean, 132.8445, ± .0008
```

We may now combine the means of these seven series, representing in all thirty-three experiments. One hundred parts of silver are equivalent to chlorine, as follows:

Here, again, we have a fine example of the evident compensation of errors among different series of experiments. We have also another tribute to the accuracy of Stas, since this general mean varies from the mean of his results only within the limits of his own variations.

The ratio between silver and potassium chloride, or, in other words, the weight of silver in nitric acid solution which can be precipitated by a known weight of KCl, has been fixed by Marignac and by Stas. Marignac,* reducing all weighings to vacuum, obtained these results. In the third column I give the weight of KCl proportional to 100 parts of Ag.

```
4.7238 grm. Ag = 3.2626 KCl.
                                         69.067
            "
                   15.001
                                         69.050
21.725
                   15.028
                                         69.066
21.759
21.909
                   15.131
                                         69.063
                   15.216
                                         69.063
22.032
                                         69.063
25.122
                   17.350
                                  Mean, 69.062, \pm .0017
```

Stas' experiments upon this ratio may be divided into two series.† In the first series the silver was slightly impure, but the impurity was of known quantity, and corrections could therefore be applied. In the second series pure silver was employed. The potassium chloride was from several different sources, and in every case was purified with the utmost care. From 10.8 to 32.4 grammes of silver were taken in each experiment, and the weighings were reduced to vacuum. The method of operation was, in brief, as follows: A definite weight of potassium chloride was taken, and the exact quantity of silver necessary, according to Prout's hypothesis, to balance it was also weighed out. The metal, with suitable precautions, was dissolved in initric acid, and the solution mixed with that of the chloride. After double decomposition the trifling excess of silver remaining in the liquid was determined by titration with a normal solution of potassium chloride. One hundred parts of silver required the following of KCl:

^{*} See Berzelius' Lehrbuch, 5th edition, Vol. 3, pp. 1192, 1193.

[†] Aronstein's Translation, pp. 250-257.

First Series.

```
69.105
       69.104
       69.103
       69.104
       69.102
Mean, 69.1036, \pm .0003
   Second Series.
       69.105
       69.099
       69.107
       69.103
       69.103
       69.105
       69.104
       69.099
       69.1034
       69.104
       69.103
       69.102
       69.104
       69.104
       69.105
       69.103
       69,101
       60.105
       69.103
Mean, 69.1033, \pm.0003
```

Now, combining the three series, with their thirty experiments, we get the following:

```
      Marignac
      69.062, ± .0017

      Stas, 1st series
      69.1036, ± .0003

      Stas, 2d series
      69.1033, ± .0003

      General mean
      69.1032, ± .0002
```

The quantity of silver chloride which can be formed from a known weight of potassium chloride has also been determined by Berzelius, Marignac, and Maumené. Berzelius* found that 100 parts of KCl were equivalent to 194.2 of

^{*} Poggend. Annal., 8, 1. 1826.

AgCl; a value which, corrected for weighings in air, becomes 192.32. This experiment will not be included in our discussion.

In 1842 Marignac* published two determinations, with these results from 100 KCl:

192.33 192.34

Mean, corrected for weighing in air, 192.26, ± .003

In 1846 Marignac† published another set of results, as follows. The weighings were reduced to vacuum. The usual ratio is in the third column.

17.034 g	rm. KCl gave	32.761	AgCļ.	192.327	
14.427	66	27.749	66	192.341	
15.028	66	28.910	66	192.374	
15.131	16	29.102	66	192.334	
15.216	66	29.271	46	192.370	
					
				Maan saa saa	

Mean, 192.349, \pm .006

Three estimations of the same ratio were also made by Maumené,‡ as follows:

```
10.700 grm. KCl gave 20.627 AgCl. 192.776
10.5195 " 20.273 " 192.716
8.587 " 16.556 " 192.803
Mean, 192.765, ± .017
```

The three series of ten experiments in all foot up thus:

These figures show clearly that the ratio which they represent is not of very high importance. It might be rejected altogether without impropriety, and is only retained for the

^{*} Ann. Chem. Pharm., 44, 21. 1842. >

⁺ Berzelius' Lehrbuch, 5th Ed., Vol. 3, pp. 1192, 1193.

[†] Ann. d. Chim. et d. Phys., (3,) 18, 41. 1846.

sake of completeness. It will obviously receive but little weight in our final discussion.

In estimating the atomic weight of bromine the earlier experiments of Balard, Berzelius, Liebig, and Löwig may all be rejected. Their results were all far too low, probably because chlorine was present as an impurity in the materials employed. Wallace's determinations, based upon the analysis of arsenic tribromide, are tolerably good, but need not be considered here. In the present state of our knowledge, Wallace's analyses are better fitted for fixing the atomic weight of arsenic, and will, therefore, be discussed with reference to that element.

The ratios with which we now have to deal are closely similar to those involving chlorine. In the first place there are the analyses of silver bromate by Stas.* In two careful experiments he found in this salt the following percentages of oxygen:

There are also four analyses of potassium bromate by Marignac.† The salt was heated, and the percentage loss of oxygen determined. The residual bromide was feebly alkaline. We cannot place much reliance upon this series. The results are as follows:

28.7016 28.6496 28.6050 28.7460

Mean, 28.6755, $\pm .0207$

When silver bromide is heated in chlorine gas, silver chloride is formed. In 1860 Dumast employed this method

^{*}Aronstein's Translation, pp. 200-206.

[†] See E. Mulder's Overzigt, p. 117; or Berzelius' Jahresbericht, 24, 72. † Ann. Chem. Pharm., 113, 20.

for estimating the atomic weight of bromine. His results are as follows: In the third column I give the weight of AgBr equivalent to 100 parts of AgCl.

2.028 grm.	AgBr gave	1.547	AgCl.	131.092
4-237	66	3.235	66	130.974
5.769	66	4.403	46	131.024

Mean, 131.030, \pm .023

This series is evidently of but little value.

But the two ratios upon which, in connection with Stas' analyses of silver bromate, the atomic weight of bromine chiefly depends are those which connect silver with the latter element directly and silver with potassium bromide.

Marignac,* to effect the synthesis of silver bromide, dissolved the metal in nitric acid, precipitated the solution with potassium bromide, washed, dried, fused, and weighed the product. The following quantities of bromine were found proportional to 100 parts of silver:

74.072 74.055 74.066

Mean, reduced to a vacuum standard, 74.077, ± .003

Much more elaborate determinations of this ratio are due to Stas.† In one experiment a known weight of silver was converted into nitrate, and precipitated in the same vessel by pure hydrobromic acid. The resulting bromide was washed thoroughly, dried, and weighed. In four other estimations the silver was converted into sulphate. Then a known quantity of pure bromine, as nearly as possible the exact amount necessary to precipitate the silver, was transformed into hydrobromic acid. This was added to the dilute solution of the sulphate, and, after precipitation was complete, the minute trace of an excess of silver in the clear supernatant fluid was determined. All weighings were re-

^{*} E. Mulder's Overzigt, p. 116. Berzelius' Jahresbericht, 24, 72.

[†] Aronstein's Translation, pp. 154-170.

duced to a vacuum. From these experiments, taking both series as one, we get the following quantities of bromine corresponding to 100 parts of silver:

	74.0830
	74.0790
	74.0795
	74.0805
	74.0830
	•
Mean,	$74.081, \pm .0006$

Combining this with Marignac's result, 74.077, $\pm .003$, we get as a general mean the value 74.0809, $\pm .0006$.*

The ratio between silver and potassium bromide was first accurately determined by Marignac.† I give, with his weighings, the quantity of KBr proportional to 100 parts of Ag:

2.131 g	rm. Ag	= 2.351	KBr.	110.324
2.559	66	2.823	66	110.316
2.447	66	2.700	44	110.339
3.025	"	3.336	66	110.283
3.946	"	4.353	44	110.314
11.569	66	12.763	46	110.321
20.120	66	22.191	66	110.293

Mean, corrected for weighing in air, 110.343, ± .005

Stas,‡ working in essentially the same manner as when he fixed the ratio between potassium chloride and silver, obtained the following results:

^{*}O. W. Huntington, in his paper upon the atomic weight of cadmium, (Amer. Acad. Proc., 1881,) gives three analyses and three syntheses of silver bromide. These give a mean value of Ag: Br:: 100:74.064. This figure I record here in order that other chemists may not overlook the work of Mr. Huntington, although it came out too late for use in my own calculations.

[†] E. Mulder's Overzigt, p. 116. Berzelius' Jahresbericht, 24, 72.

[‡] Aronstein's Translation, pp. 334-347.

110.361 110.360 110.360 110.346 110.338 110.360 110.336 110.344 110.332 110.343 110.357 110.334

Mean, 110.3463, \pm .0020

Combining this with Marignac's mean result, 110.343, \pm .005, we get a general mean of 110.3459, \pm .0019.

The ratios upon which we must depend for the atomic weight of iodine are exactly parallel to those used for the determination of bromine.

To begin with, the percentage of oxygen in potassium iodate has been determined by Millon.* In three experiments he found:

22.46
22.49
22.47

Mean, 22.473, ± .005

Millon also estimated the oxygen in silver iodate, getting the following percentages:

The analysis of silver iodate has also been performed with extreme care by Stas.† From 76 to 157 grammes were used

^{*} Ann. d. Chim. et d. Phys., (3,) 9, 400. 1843.

[†] Aronsteins' Translation, pp. 179-200.

in each experiment, the weights being reduced to a vacuum standard. As the salt could not be prepared in an absolutely anhydrous condition, the water expelled in each analysis was accurately estimated and the necessary corrections applied. In two of the experiments the iodate was decomposed by heat, and the oxygen given off was fixed upon a weighed quantity of copper heated to redness. Thus the actual weights, both of the oxygen and the residual iodide, were obtained. In a third experiment the iodate was reduced to iodide by a solution of sulphurous acid, and the oxygen was estimated only by difference. In the three percentages of oxygen given below the result of this analysis comes last. The figures for oxygen are as follows:

```
16.976
16.972
16.9761
Mean, 16.9747, ± .0009
```

This, combined with Millon's series above cited, gives us a general mean of 16.9771, $\pm .0009$.

The ratio between silver and potassium iodide seems to have been determined only by Marignac,* and without remarkable accuracy. In five experiments 100 parts of silver were found equivalent to potassium iodide as follows:

```
1.616 grm. Ag = 2.483 KI.

2.503 " 3.846 " " 153.665

3.427 " 5.268 " " 152.720

2.141 " 3.290 " " 153.667

10.821 " 16.642 " " 153.794

Mean, 153.6994, ± .0178
```

The synthesis of silver iodide has been effected by both Marignac and Stas. Marignac, in the paper above cited, gives these weighings. In the last column I add the ratio between iodine and 100 parts of silver:

```
15.000 grm. Ag gave 32.625 AgI. 117.500
14.790 " 32.170 " 117.512
18.545 " 40.339 " 117.519
```

Mean, corrected for weighing in air, 117.5335, ± .0036

^{*} Berzelius' Lehrbuch, 5th Ed., 3, 1196.

Stas* in his experiments worked after two methods, which gave, however, results concordant with each other and with those of Marignac.

In the first series of experiments Stas converted a known weight of silver into nitrate, and then precipitated with pure hydriodic acid. The iodide thus thrown down was washed, dried, and weighed without transfer. By this method 100 parts of silver were found to require of iodine:

117.529 117.536

Mean, 117.5325, \pm .0024

In the second series a complete synthesis of silver iodide from known weights of iodine and metal was performed. The iodine was dissolved in a solution of ammonium sulphite, and thus converted into ammonium iodide. The silver was transformed into sulphate and the two solutions mixed. When the precipitate of silver iodide was completely deposited the supernatant liquid was titrated for the trifling excess of iodine which it always contained. As the two elements were weighed out in the ratio of 127 to 108, while the atomic weight of iodine is probably a little under 127, this excess is easily explained. From these experiments two sets of values were deduced; one from the weights of silver and iodine actually employed, the other from the quantity of iodide of silver collected. From the first set we have of iodine for 100 parts of silver:

> 117.5390 117.5380 117.5318 117.5430 117.5420 117.5300

Mean, 117.5373, \pm .0015

From the weight of silver iodide actually collected we

^{*} Aronstein's Translation, pp. 136, 152.

get as follows. For experiment number three in the above column there is no equivalent here:

117.529
117.531
117.539
117.538
117.530
———
Mean, 117.5334, ± .0014

Now, combining these several sets of results, we have the following general mean:

One other comparatively unimportant iodine ratio remains for us to notice. Silver iodide, heated in a stream of chlorine, becomes converted into chloride; and the ratio between these two salts has been thus determined by Berzelius and by Dumas.

From Berzelius* we have the following data: In the third column I give the ratio between AgI and 100 parts of AgCl.

```
5.000 grm. AgI gave 3.062 AgCl. 163.292
12.212 " 7.4755 " 163.360
Mean, 163.326, ± .023
```

Dumas't results were as follows:

General mean from the combination of both series, 163.733, $\pm .0076$.

We now come to the ratios connecting sulphur with silver

^{*} Ann. d. Chim. et d. Phys., (2,) 40, 430. 1829.

[†] Ann. Chem. Pharm., 113, 28. 1860.

and chlorine. Other ratios have been applied to the determination of the atomic weight of sulphur, but they are hardly applicable here. The earlier results of Berzelius were wholly inaccurate, and his later experiments upon the synthesis of lead sulphate will be used in discussing the atomic weight of lead. Erdmann and Marchand determined the amount of calcium sulphate which could be formed from a known weight of pure Iceland spar; and later they made analyses of cinnabar, in order to fix the value of sulphur by reference to calcium and to mercury. Their results will be applied in this discussion towards ascertaining the atomic weights of the metals just named. For our present purposes only three ratios need be considered.

First in order let us take up the composition of silver sulphide, as directly determined by Dumas, Stas, and Cooke. Dumas'* experiments were made with sulphur which had been thrice distilled and twice crystallized from carbon disulphide. A known weight of silver was heated in a tube in the vapor of the sulphur, the excess of the latter was distilled away in a current of carbon dioxide, and the resulting silver sulphide was weighed.

I subjoin Dumas' weighings, and also the quantity of Ag₂S proportional to 100 parts of Ag, as deduced from them:

```
9.9393 grm. Ag = 1.473 S. Ratio, 114.820

9.962 " 1.4755 " " 114.811

30.637 " 4.546 " " 114.838

30.936 " 4.586 " " 114.824

30.720 " 4.554 " " 114.824
```

Mean, 114.8234, ± .0029

Dumas used from ten to thirty grammes of silver in each experiment. Stas,† however, in his work, employed from sixty to two hundred and fifty grammes at a time. Three of Stas' determinations were made by Dumas' method, while in the other two the sulphur was replaced by pure sulphu-

^{*} Ann. Chem. Pharm., 113, 24. 1860

[†] Aronstein's Translation, p. 179.

retted hydrogen. In all cases the excess of sulphur was expelled by carbon dioxide, purified with scrupulous care. Impurities in the dioxide may cause serious error. The five results come out as follows for 100 parts of silver:

114.854 114.853 114.854 114.851 114.849

Mean, 114.8522, ± .0007

The experiments made by Professor Cooke* with reference to this ratio were only incidental to his elaborate researches upon the atomic weight of antimony. They are interesting however, for two reasons: they serve to illustrate the volatility of silver, and they represent, not syntheses, but reductions of the sulphide by hydrogen. Cooke gives three series of results. In the first the silver sulphide was long heated to full redness in a current of hydrogen. Highly concordant and at the same time plainly erroneous figures were obtained; the error being eventually traced to the fact that some of the reduced silver, although not heated to its melting point, was actually volatilized and lost. The second series, from reductions at low redness, are decidedly better. In the third series the sulphide was fully reduced below a visible red heat. Rejecting the first series we have from Cooke's figures in the other two the subjoined quantities of sulphide corresponding to 100 parts of silver:

```
7.5411 grm. Ag<sub>2</sub>S lost .9773 grm. S. Ratio, 114.889
5.0364
               66
                         .6524
                                                 114.882
2.5815
                                                 114.886
               66
                        ·3345
                         .3387
                                                 114.892
2.6130
                                  "
                                                 114.891
2.5724
                         -3334
                                          Mean, 114.888, \pm .0012
1.1357 grm. Ag<sub>2</sub>S lost .1465 S.
                                          Ratio, 114.810
1.2936
                         .1670 "
                                                 114.823
                                          Mean, 114.8165, \pm .0044
```

^{*} Proc. American Acad. of Arts and Sciences, v. 12. 1877.

Now, combining all four series, we get the following results:

Dumas	
Stas	$114.8522, \pm .0007$
Cooke's 2d	114.888, \pm .0012
" 3d	114.8165, ± .0044
General mean	114.8581. + .0006

Here again we encounter a curious and instructive compensation of errors, and another evidence of the accuracy of Stas.

The percentage of silver in silver sulphate has been determined by Struve and by Stas. Struve* reduced the sulphate by heating in a current of hydrogen, and obtained these results:

```
5.1860 grm. Ag<sub>2</sub>SO<sub>4</sub> gave 3.5910 grm. Ag.
                                                   69.244 per cent.
                                                   69.243
   6.0543
                              4.1922
   8.6465
                                                   69.228
                              5.9858
                              8.0608
  11.6460
                                                   69.215
                              6.3045
                                                   69.212
. 9.1090
   9.0669
                              6.2778
                                                   69.239
                                            Mean, 69.230, \pm .004
```

Stas,† working by essentially the same method, with from 56 to 83 grammes of sulphate at a time, found these percentages:

Combining this mean with that from Struve's series we get a general mean of $69.205, \pm .0011$.

^{*} Ann. Chem. Pharm., 80, 203. 1851.

[†] Aronstein's Translation, pp. 214-218.

The third and last sulphur ratio with which we have now to deal is one of minor importance. When silver chloride is heated in a current of sulphuretted hydrogen the sulphide is formed. This reaction was applied by Berzelius* to determining the atomic weight of sulphur. He gives the results of four experiments; but the fourth varies so widely from the others that I have rejected it. I have reason to believe that the variation is due, not to error in experiment, but to error in printing; nevertheless, as I am unable to track out the cause of the mistake, I must exclude the figures involving it entirely from our discussion.

The three available experiments, however, give the following results: The last column contains the ratio of silver sulphide to 100 parts of chloride.

```
6.6075 grm. AgCl gave 5.715 grm. Ag<sub>2</sub>S. 86.478

9.2323 " 7.98325 " 86.471

10.1775 " 8.80075 " 86.472

Mean, 86.4737, ± .0015
```

We have also a single determination of this value by Svanberg and Struve.† After converting the chloride into sulphide they dissolved the latter in nitric acid. A trifling residue of chloride, which had been enclosed in sulphide, and so protected against change, was left undissolved. Hence a slight constant error probably affects this whole ratio. The experiment of Svanberg and Struve gave 86.472 per cent. of silver sulphide derived from 100 of chloride. If we assign this figure equal weight with the results of Berzelius, and combine, we get a general mean of 86.4733, \pm .0011.

For sodium there are but two ratios of any definite value for present purposes. The early work of Berzelius we may disregard entirely, and confine ourselves to the consideration of the results obtained by Penny, Pelouze, Dumas, and Stas.

^{*} Berzelius' Lehrbuch, 5th Ed., Vol. 3, p. 1187.

⁺ Journ. für Prakt. Chem., 44, 320. 1848.

The percentage of oxygen in sodium chlorate has been determined only by Penny,* who used the same method which he applied to the potassium salt. Four experiments gave the following results:

```
45.060
45.075
45.080
45.067
Mean, 45.0705, ± .0029
```

The ratio between silver and sodium chloride has been fixed by Pelouze, Dumas, and Stas. Pelouze† dissolved a weighed quantity of silver in nitric acid, and then titrated with sodium chloride. Equivalent to 100 parts of silver he found of chloride:

By Dumast we have seven experiments, with results as follows: The third column gives the ratio between 100 of silver and NaCl.

2.0535 gr	m. NaCl :	= 3.788 grm.	Ag.	54.211
2.169	44	4.0095 "	i	54.097
4-3554	46	8.0425 "	i	54.155
6.509	46	12.0140 "	:	54.178
6.413	66	11.8375 "		54.175
2.1746	46	4.012 "	;	54.202
5.113	**	9.434 "		54.187

Mean, 54.172, \pm .0096

Stas, || applying the method used in establishing the similar ratio for potassium chloride, and working with salt from

^{*} Phil. Transactions, 1839, p. 25.

[†] Compt. Rend., 20, 1047. 1845.

[†] Ann. Chem. Pharm., 113, 31. 1860.

^{||} Aronstein's Translation, p. 274.

six different sources, found of sodium chloride equivalent to 100 parts of silver:

```
54.2093
54.2088
54.2070
54.2070
54.2070
54.2060
54.2066
54.2081
54.2083
54.2089

Mean, 54.2078, ± .0002
```

Now, combining these three series, we get the following result:

```
Pelouze_______ 54.141, ± .0063

Dumas ______ 54.172, ± .0096

Stas ______ 54.2078, ± .0002

General mean _____ 54.2076, ± .0002
```

Here the work of Stas is of such superior excellence that the other series might be completely rejected without appreciably affecting our calculations.

We have now before us the data establishing, with greater or less accuracy, twenty different ratios relating to the atomic weights of the seven elements under discussion. In these we are to discuss the results of about two hundred and fifty separate experiments. Before beginning upon our calculations we will tabulate our ratios, and number them for convenient future reference. Of course it will be understood that the probable errors given below relate to the last term of each proportion:

```
(9.) Ag: NaCl:: 100: 54.2076, \pm .0002

(10.) Ag: KCl:: 100: 69.1032, \pm .0002

(11.) Ag: KBr:: 100: 110.3459, \pm .0019

(12.) Ag: KI:: 100: 153.6994, \pm .0178

(13.) Ag: Cl:: 100: 32.8418, \pm .0006

(14.) Ag: Br:: 100: 74.0809, \pm .0006

(15.) Ag: I:: 100: 117.5345, \pm .0009

(16.) Ag: Ag<sub>2</sub>S:: 100: 114.8581, \pm .0006

(17.) KCl: AgCl:: 100: 192.294, \pm .0029

(18.) AgCl: AgBr:: 100: 131.030, \pm .023

(19.) AgCl: AgI:: 100: 86.4733, \pm .0011
```

Now, from ratios 1 to 7 inclusive, we can at once, by applying the known atomic weight of oxygen, deduce the molecular weights of seven haloid salts. Let us consider the first calculation somewhat in detail.

Potassium chlorate yields 39.154 per cent. of oxygen and 60.846 per cent. of residual chloride. For each of these quantities the probable error is \pm .00038. The atomic weight of oxygen is 15.9633, \pm .0035, so that the value for three atoms becomes 47.8899, \pm .0105. We have now the following simple proportion: 39.154:60.846::47.8899:x, = the molecular weight of potassium chloride, = 74.4217. The probable error being known for the first, second, and third term of this proportion, we can easily find that of the fourth term by the formula given in our introduction. It comes out \pm .0164. By this method we obtain the following series of values, which may conveniently be numbered consecutively with the foregoing ratios:

```
(21.) KCl, from (1,) = 74.4217, \pm .0164
(22.) KBr, " (2,) = 119.117, \pm .0962
(23.) KI, " (3,) = 165.210, \pm .0529
(24.) NaCl, " (4,) = 58.366, \pm .0137
(25.) AgCl, " (5,) = 143.062, \pm .0320
(26.) AgBr, " (6,) = 187.453, \pm .0432
(27.) AgI, " (7,) = 234.195, \pm .0530
```

With the help of these molecular weights we are now able to calculate eight independent values for the atomic weight of silver:

```
from (10) and (21,) Ag = 107.696, \pm .024
First.
             (11) " (22,) " = 107.948, \pm .087
Second.
             (12) " (23,) " = 107.488, \pm .037
Third.
Fourth.
             (9) " (24,) " = 107.671, \pm .025
             (13) " (25,) " = 107.694, \pm .024
Fifth.
             (14) " (26,) " = 107.681, \pm .025
Sixth,
Seventh.
             (15) " (27,) " = 107.659, \pm .024
             (8) " (16,) " = 107.712, \pm .025
Eighth,
             General mean, " = 107.675, \pm .0096
```

It is noticeable that six of these values agree very well. The second and third, however, diverge widely from the average, but in opposite directions; they have, moreover, high probable errors, and consequently little weight. Of these two, one represents little and the other none of Stas' work. Their trifling influence upon our final results becomes curiously apparent in the series of silver values given a little further along.

When we consider closely, in all of its bearings, any one of the values just given, we shall see that for certain purposes it must be excluded from our general mean. For example, the first is derived partly from the ratio between silver and potassium chloride. From this ratio, the atomic weight of one substance being known, we can deduce that of the other. We have already used it in ascertaining the atomic weight of silver, and the value thus obtained is included in our general mean. But if from it we are to determine the molecular weight of potassium chloride, we must use a silver value derived from other sources only, or we should be assuming a part of our result in advance. other words, we must now use a general mean for silver from which this ratio with reference to silver has been re-Hence the following series of silver values, which are lettered for reference:

```
A. General mean from all eight______ 107.675, ± .0096
                rejecting the first ..... 107.671, \pm .0105
B.
                           second _____ 107.671, ± .0097
C.
                     "
                           third _____ 107.679, ± .0100
D.
                           fourth _____ 107.675, ± .0104
E.
                           fifth .... 107.671, ± .0105
F.
                     "
                           sixth _____ 107.674, \pm .0104
G.
                            seventh_____ 107.678, ± .0105
                     46
H.
                            eighth _____ 107.679, ± .0104
I.
```

These values are essentially the same, both in magnitude and in weight. For all practical purposes any one of them is as good as any other. Still, on theoretical grounds, it may be well to keep them distinct and separate in the remainder of this discussion.

We are now in a position to determine more closely the molecular weights of the haloid salts which we have already been considering.

For silver chloride, still employing the formula for the probable error of the last term of a proportion, we get the following values:

Subtracting from this the atomic weight of silver, 107.675, \pm .0096, we get for the atomic weight of chlorine, Cl = 35.370, \pm .014.

For silver bromide we have these results:

```
From (6) _____AgBr = 187.453, \pm .043

From (14) and (G) ____ " = 187.440, \pm .018

From (18) and (25) ____ " = 187.454, \pm .053

General mean ___ " = 187.443, \pm .016
```

Hence, using the general mean for silver as above, Br = 79.768, $\pm .019$.

Silver iodide comes out as follows:

```
From (7)------AgI = 234.195, \pm .053

From (15) and (H)----- " = 234.237, \pm .023

From (19) and (25)----- " = 234.240, \pm .054

General mean---- " = 234.232, \pm .019
```

Hence I = 126.557, $\pm .022$.

For the molecular weight of sodium chloride we have:

Hence, if chlorine = 35.370, $\pm .014$, then Na = 22.998, $\pm .011$.

For potassium chloride:

For potassium bromide we get:

And for potassium iodide:

Now, taking the molecular weights of these three potassium salts in connection with the atomic weights just found for chlorine, bromine, and iodine, we get these values for potassium:

Finally, the three sulphur ratios give us three estimates for the atomic weight of sulphur. In the third of these I have applied the "A" value for silver and the general mean for silver chloride:

From (8) and (I) ______S = 31.968,
$$\pm$$
 .014
From (16) and (I) _____ " = 31.995, \pm .032
From (20) ____ " = 32.041, \pm .028
General mean ___ " = 31.984, \pm .012

We may now appropriately compare the results of this

discussion with the atomic weights deduced by Stas from his own experiments only. His values are given under two headings: one for oxygen = 16, the other for O = 15.96. As we have been using the figure 15.9633 for oxygen, here is at the outset a discrepancy. Starting from this value we found:

Ag =
$$107.675$$
, $\pm .0096$
Cl = 35.370 , $\pm .014$
Br = 79.768 , $\pm .019$
I = 126.557 , $\pm .022$
Na = 22.998 , $\pm .011$
K = 39.019 , $\pm .012$
S = 31.984 , $\pm .012$

If we assume 16 to be the true figure for oxygen, we get the following results, which I have placed in a column parallel with the values found by Stas:

The Ne	rw Values.	Stas.	Differences.
Silver 10	7.923	107.930	.007
Chlorine 3	5.451	35-457	.006
Bromine 7	9.951	79.952	100.
Iodine 12	6.848	126.850	.002
Sodium 2	3.051	23.043	.009
Potassium 3	9.109	39.137	.028
Sulphur 3	2.058	32.074	.016

These differences are insignificant. No other criticism could more severely test the character of Stas' work, or more definitely illustrate his magnificent accuracy of manipulation.

NITROGEN.

The atomic weight of nitrogen has been determined from the density of the gas, from the ratio between ammonium chloride and silver, and from the composition of certain nitrates.

Upon the density of nitrogen a great many experiments have been made. In early times this constant was determined by Biot and Arago, Thomson, Dulong and Berzelius, Lavoisier, and others. But all of these investigations may be disregarded as of insufficient accuracy; and, as in the case of oxygen, we need consider only the results obtained by Dumas and Boussingault, and by Regnault.

Taking air as unity, Dumas and Boussingault* found the density of nitrogen to be—

.970 .972 .974

Mean, .972, $\pm .00078$

For hydrogen, as was seen in our discussion of the atomic weight of oxygen, the same investigators found a mean of .0693, \pm .00013. Upon combining this with the above nitrogen mean, we find for the atomic weight of the latter element, $N = 14.026, \pm .0295$.

By Regnault† much closer work was done. He found the density of nitrogen to be as follows:

.97148 .97148 .97154 .97155 .97108

Mean, .97137, ± .000062

^{*} Compt. Rend., 12, 1005. 1841.

[†] Compt. Rend., 20, 975. 1845.

For hydrogen, Regnault's mean value is .069263, ± .000019. Hence, combining as before, N = 14.0244, ± .0039.*

The value found by combining both series of experiments is N = 14.0244, $\pm .0039$.

In discussing the more purely chemical ratios for establishing the atomic weight of nitrogen, we may ignore, for the present, the researches of Berzelius, of Anderson, and of Svanberg. These chemists experimented chiefly upon lead nitrate, and their work is consequently now of greater value for fixing the atomic weight of lead. Their results will be duly considered in the proper connection further on.

The ratio between ammonium chloride and silver has been determined by Pelouze, by Marignac, and by Stas. The method of working is essentially that adopted in the similar experiments with the chlorides of sodium and potassium.

For the ammonium chloride equivalent to 100 parts of silver, Pelouze† found:

49.556 49.517

Mean, 49.5365, \pm .013

Marignact obtained the following results. The usual ratio for 100 parts of silver is given also:

8.063	grm. Ag =	3.992 grn	n. NH ₄ Cl.	49.510
9.402	46	4.656	46	49.521
10.339	46	5.120	46	49.521
12.497	4	6.191	46	49.540
11.337	46	5.617	46	49.546
11.307	46	5-595	46	49.483
4.326	64	2.143	46	49.538

Mean, 49.523, \pm .0055

^{*} Professor Le Conte, in his corrections of Regnault's calculations, already cited in a foot note to the chapter on oxygen, finds for the density of nitrogen the value 0.971346. Hence N = 14.0225. This correction is very slight, but it should be considered in any future revision of the atomic weights.

[†] Compt. Rend., 20, 1047. 1845.

¹ Berzelius' Lehrbuch, 5th Ed., 3d v., 1184, 1185.

But neither of these series can for a moment compare with that of Stas.* He used from 12.5 to 80 grammes of silver in each experiment, reduced his weighings to a vacuum standard, and adopted a great variety of precautions to ensure accuracy. He found for every 100 parts of silver the following quantities of NH₄Cl:

49.600 49.599 49.597 49.598 49.597 49.597 49.597 49.602 49.597 49.598 49.592

Mean, 49.5973, $\pm .0005$

Now, combining these three series, we get:

 Pelouze
 49.5365, ± .013

 Marignac
 49.523, ± .0055

 Stas
 49.5973, ± .0005

 General mean
 49.597, ± .0005

The quantity of silver nitrate which can be formed from a known weight of metallic silver has been determined by Penny, by Marignac, and by Stas. Penny† dissolved silver in nitric acid in a flask, evaporated to dryness without transfer, and weighed. One hundred parts of silver thus gave of nitrate:

157.430 157.437 157.458 157.440 157.430 157.455 Mean, 157.4417, ± .0033

^{*} Aronstein's Translation, pp. 56-58.

[†] Phil. Trans., 1839.

Marignac's* results were as follows. In the third column they are reduced to the common standard of 100 parts of silver:

68.987	grm. Ag gave	108.608 grm.	AgNO ₃ .	157-433
57.844	46	91.047	66	157.401
66.436	66	104.592	66	157-433
70.340	"	110.718	66	157.404
200.000	"	314.894	66	157.447

Mean, 157.4236, \pm .0061

Stas,† employing from 77 to 405 grammes of silver in each experiment, made two different series of determinations at two different times. The silver was dissolved with all the usual precautions against loss and against impurity, and the resulting nitrate was weighed, first after long drying without fusion just below its melting point; and again, fused. Between the fused and the unfused salt there was in every case a slight difference in weight, the latter giving a maximum and the former a minimum value.

In Stas' first series there are eight experiments; but the seventh he himself rejects as inexact. The values obtained for the nitrate from 100 parts of silver are given below in two columns, representing the two conditions in which the salt was weighed. The general mean given at the end I have deduced from the means of the two columns considered separately:

	Unfused.	Fused.
	157.492	157.474
	157.510	157.481
	157.485	157.477
	157.476	157.471
	157.478	157.470
	157.471	157.463
	157.488	157.469
Mean,	157.4857 Mean	1, 157.472
	General mean. 157.474. +	0014

^{*} Berzelius' Lehrbuch, 5th Ed., 3, pp. 1184, 1185.

[†] Aronstein's Translation, pp. 305 and 315.

In the later series there are but two experiments, as follows:

	Unfused.		Fused.
	157.4964		157.488
	157.4940		157.480
Mean,	157.4952	Mean,	157.484
	General	mean, 157.486, ±	.0003

Now, to combine all four sets of results:

Penny	$157.4417, \pm .0033$
Marignac	$157.4236, \pm .0061$
Stas, 1st series	157.4740, ± .0014
Stas, 2d series	
C	
General mean	157.479, 土.0003

For the direct ratio between silver nitrate and silver chloride there are two series of estimations. A weighed quantity of nitrate is easily converted into chloride, and the weight of the latter ascertained. In two experiments Turner* found of chloride from 100 parts of nitrate:

Penny,† in five determinations, found the following percentages:

The general mean from both series is 84.3743, $\pm .0025$.

The ratio directly connecting silver nitrate with ammonium chloride has been determined only by Stas.‡ The

^{*} Phil. Trans., 1833, 537.

[†] Phil. Trans., 1839.

[†] Aronstein's Translation, p. 309.

usual method of working was followed; namely, nearly equivalent quantities of the two salts were weighed out, the solutions mixed, and the slight excess of one estimated by titration. In four experiments 100 parts of silver nitrate were found equivalent to chloride of ammonium as follows:

The similar ratio between potassium chloride and silver nitrate has been determined by both Marignac and Stas.

Marignac* gives the following weights. I add the quantity of KCl proportional to 100 parts of AgNO₈:

1.849 gr	m. KCl	= 4.218 gr	n. AgNO ₃ .	43.836	
2.473	44	5.640	66	43.848	
3.317	46	7.565	. "	43.847	
2.926	66	6.670	66	43.868	
6. 191	66	14.110	44	43.877	
4.351	44	9.918	66	43.870	
			М	ean, 43.858 , \pm .0	044

Stas't results are given in three series, representing silver nitrate from three different sources. In the third series the nitrate was weighed in vacuo, while for the other series this correction was applied in the usual way. For the KCl equivalent to 100 parts of AgNO₃ Stas found:

Fir	st Series.
4	13.878
4	43.875
4	43.875
4	43.874
-	
Mean, 4	$\pm 3.8755, \pm .0005$

^{*} Berzelius' Lehrbuch, 5th Ed., 3d vol., 1184, 1185.

[†] Aronstein's Translation, p. 308.

Second Series.

43.864
43.869
43.876
Mean, 43.8697, ± .0023

Third Series.

43.894
43.878
43.885
Mean, 43.8857, ± .0031

Combining all four series we have:

Marignac 43.858, ± .0044

Stas, 1st series 43.8755, ± .0005

43.8697, ± .0023

43.8857, ± .0031

General mean 43.8715, ± .0004

There have also been determined by Penny and by Stas a series of ratios connecting the alkaline chlorides and chlorates with the corresponding nitrates. One of these, relating to the lithium salts, will be studied further on with reference to that metal.

The general method of working upon these ratios is due to Penny.* Applied to the ratio between the chloride and nitrate of potassium it is as follows: A weighed quantity of the chloride is introduced into a flask which is placed upon its side and connected with a receiver. An excess of pure nitric acid is added, and the transformation is gradually brought about by the aid of heat. Then, upon evaporating to dryness over a sand bath, the nitrate is brought into weighable form. The liquid in the receiver is also evaporated, and the trace of solid matter which had been mechanically carried over is recovered and also taken into account. In another series of experiments the nitrate was taken, and by pure hydrochloric acid converted into chloride; the process being the same. In the following columns of figures I

^{*} Phil. Trans., 1839.

have reduced both series to one standard; namely, so as to express the number of parts of nitrate corresponding to 100 of chloride:

```
First Series. - KCl treated with HNO2.
                135.639
                135.637
                135.640
                135.635
                135.630
                135.640
                135.630
         Mean, 135.636, \pm .0011
Second Series.—KNO<sub>3</sub> treated with HCl.
                135.628
                135.635
                135.630
                135.641
                135.630
                135.635
                135.630
         Mean, 153.633, \pm .0011
```

Stas* results are as follows:

```
135.643
135.638
135.647
135.649
135.640
135.645
135.655
```

These figures by Stas represent weighings in the air. Reduced to a vacuum standard this mean really becomes 135.6423.

Now, combining, we have:

```
Penny, 1st series ______ 135.636, ± .0011

" 2d " ______ 135.633, ± .0011

Stas _____ 135.6423, ± .0014

General mean _____ 135.6363, ± .0007
```

^{*} Aronstein's Translation, p. 270.

By the same general process Penny* determined how much potassium nitrate could be formed from 100 parts of chlorate. He found as follows:

82.505 82.497 . 82.498 82.500 Mean, 82.500, ± .0012

For 100 parts of sodium chlorate he found of nitrate:

For the ratio between the chloride and nitrate of sodium Penny made two sets of estimations as in the case of potassium salts. The subjoined figures give the amount of nitrate equivalent to 100 parts of chloride:

```
First Series.—NaCl treated with HNO<sub>3</sub>.

145.415
145.408
145.420
```

145.424 145.410 145.418

145.420

Mean, 145.4164, \pm .0015

Second Series.—NaNO₃ treated with HCl.

145.419 145.391 145.412 145.415 145.412

Mean, 145.410, \pm .0026

^{*} Phil. Trans., 1839.

Stas* gives the following series:

145.453 145.468 145.465 145.469 145,443

Mean, after reducing to vacuum standard, 145.4526, ± .0030

Combining, we have as follows:

```
Penny, 1st series ______ 145.4164, ± .0015

" 2d " ______ 145.410, ± .0026

Stas _____ 145.4526, ± .0030

General mean _____ 145.4185, ± .0012
```

We have now, apart from the determinations of gaseous density, nine ratios, representing one hundred and fourteen experiments from which to calculate the atomic weight of nitrogen. Let us first collect and number these ratios:

```
(1.) Ag: AgNO<sub>3</sub>:: 100: 157.479, \pm .0003

(2.) AgNO<sub>3</sub>: AgCl:: 100: 84.3743, \pm .0025

(3.) AgNO<sub>3</sub>: KCl:: 100: 43.8715, \pm .0004

(4.) AgNO<sub>3</sub>: NH<sub>4</sub>Cl:: 100: 31.488, \pm .0006

(5.) Ag: NH<sub>4</sub>Cl:: 100: 49.597, \pm .0005

(6.) KCl: KNO<sub>3</sub>:: 100: 135.6363, \pm .0007

(7.) KClO<sub>3</sub>: KNO<sub>3</sub>:: 100: 82.500, \pm .0012

(8.) NaCl: NaNO<sub>3</sub>:: 100: 145.4185, \pm .0012

(9.) NaClO<sub>3</sub>: NaNO<sub>3</sub>:: 100: 79.8823, \pm .0029
```

From these ratios we are now able to deduce the molecular weight of ammonium chloride and of the three nitrates named in them. For these calculations we may use the already determined atomic weights of silver, oxygen, potassium, sodium, and chlorine, and the molecular weights of silver chloride and sodium chloride. These two molecular weights involve, respectively, the most probable values for silver, sodium, and chlorine. We cannot, however, appropriately use the directly determined molecular weight of potassium chloride, since the most probable value for the

^{*} Aronstein's Translation, p. 278.

atomic weight of potassium is only in part derived from that salt. The following are the values which we shall employ:

Ag = 107.675,
$$\pm$$
.0096
K = 39.019, \pm .012
Na = 22.998, \pm .011
Cl = 35.370, \pm .014
O₃ = 47.8899, \pm .0105
AgCl = 143.045, \pm .0108
NaCl = 58.3676, \pm .0052

Now, from ratio number five we can get the molecular weight of ammonium chloride, $NH_4Cl = 53.4048$, $\pm .0048$, and N = 14.0336, $\pm .0153$.

From ratio number four an independent value for nitrogen can be calculated, namely, $N = 14.0330, \pm .015$.

For the molecular weight of silver nitrate three values are deducible, namely:

Hence $N = 13.9840, \pm .0174$.

The molecular weight of potassium nitrate is twice calculable, as follows:

And $N = 13.9774, \pm .0216$.

So also for sodium nitrate we have:

And N = 13.9906, $\pm .0163$.

We have now before us six estimates of the atomic weight of nitrogen. It only remains for us to combine these after the usual method, as follows, in order to obtain the most probable value:

I.	From	specific gravity of $N_{}N = 14.0244$, $\pm .0039$
2.	66	ammonium chloride" = 14.0336, ± .0153
3.	44	ratio number four " = 14.0330, \pm .0150
4.	66	silver nitrate " = 13.9840, \pm .0174
5.	44	potassium nitrate
6.	64	sodium nitrate = 13.9906, ± .0163
		•
	G	eneral mean

If oxygen is 16, this becomes 14.0291. Stas found N = 14.044. The difference is .015, showing a remarkably close agreement.

CARBON.

Although there is a large mass of material relating to the atomic weight of carbon, much of it may be summarily set aside as having no value for present purposes. The density of carbon dioxide, which has been scrupulously determined by many investigators,* leads to no safe estimate of the constant under consideration. The numerous analyses of hydrocarbons, like the analyses of naphthalene by Mitscherlich, Woskresensky, Fownes, and Dumas, give results scarcely more satisfactory. In short, all the work done upon the atomic weight of carbon before the year 1840 may be safely rejected as unsuited to the present requirements of exact science. As for methods of estimation we need consider but three, as follows:

First.—The analysis of organic salts of silver.

Second.—The determination of the weight of carbon dioxide formed by the combustion of a known weight of carbon.

^{*} Notably by Lavoisier, Biot and Arago, De Saussure, Dulong and Berzelius, Buff, Von Wrede, Regnault, and Marchand. For details, Van Geuns' monograph may be consulted.

51

Third.—The method of Stas, by the combustion of carbon monoxide.

The first of these methods, which is also the least accurate, was employed by Liebig and Redtenbacher* in 1840. They worked with the acetate, tartrate, racemate, and malate of silver, making five ignitions of each salt, and determining the percentage of metal. From one to nine grammes of material were used in each experiment.

In the acetate the following percentages of silver were found:

64.615 64.624 64.623 64.614 64.610

Mean, 64.6172, \pm .0018

After applying corrections for weighing in air this mean becomes 64.6065.

In the tartrate the silver came out as follows:

59.297 59.299 59.287 59.293 59.293

Mean, 59.2938, \pm .0014

Or, reduced to a vacuum, 59.2806

In the racemate we have:

59.290 59.292 59.287 59.283 59.284

Mean, 59.2872, ± .0012 Or, corrected, 59.2769

[#] Ann. Chem. Pharm., 38, 137. Mem. Chem. Soc., 1, 9. Phil. Mag., (3,) 19, 210.

And from the malate:

```
61.996
61.972
62.015
62.059
62.011
```

Mean, 62.0106, ± .0096 Or, corrected, 62.0016

Now, applying to these mean results the atomic weights already found for oxygen and silver, we get the following values for carbon:

```
From the acetate _____C = 12.0306, ± .0047

" tartrate ____ " = 12.0356, ± .0064

" racemate ___ " = 12.0413, ± .0063

" malate ___ " = 12.0408, ± .0054

General mean ___ " = 12.0363, ± .0028
```

Now these results, although remarkably concordant, are by no means unimpeachable. They involve two possible sources of constant error, namely, impurity of material and the volatility of the silver. These objections have both been raised by Stas, who found that the silver tartrate, prepared as Liebig and Redtenbacher prepared it, always carried traces of the nitrate, and that he, by the ignition of that salt, could not get results at all agreeing with theirs. In the case of the acetate a similar impurity would lower the percentage of silver, and thus both sources of error would reinforce each other and make the atomic weight of carbon come out too high. With the three other salts the two sources of error act in opposite directions, although the volatility of the silver is probably far greater in its influence than the impurity. Even if we had no other data relating to the atomic weight of carbon, it would be clear from these facts that the results obtained by Liebig and Redtenbacher must be decidedly in excess of the true figure.

A different method of dealing with organic silver salts was adopted by Maumené,* in 1846, for the purpose of estab-

^{*} Ann. d. Chim. et d. Phys., (3,) 18, 41.

CARBON. 53

We will simply reverse his results and apply them to the atomic weight of carbon. He effected the combustion of the acetate and the oxalate of silver, and, by weighing both the residual metal and the carbon dioxide formed, he fixed the ratio between these two substances. In the case of the acetate his weighings show that for every gramme of metallic silver the weights of CO₂ were produced, which are shown in the third column:

8.083 g	rm. Ag =	= 6.585 gr	m. CO ₂ .	•	8147
11.215	44	9.135	66		8136
14.351	44	11.6935	••	•	8148
9.030	66	7.358	46	•	8148
20.227	66	16.475	44	•	8145
				Mean, .	 81448

The oxalate of silver, ignited by itself, decomposes too violently to give good results; and for this reason it was not used by Liebig and Redtenbacher. Maumené, however, found that when the salt was mixed with sand the combustion could be tranquilly effected. The oxalate employed, however, with the exception of the sample represented in the last experiment of the series, contained traces of nitrate, so that these results involve slight errors. For each gramme of silver the appended weights of CO, were obtained:

```
14.299 grm. Ag = 5.835 grm. CO_2.
                                                    .4081
17.754
                                                    .4059
11.550
                   4.703
                                                     .4072
                   4.387
10.771
                                                     .4073
8.674
                   3.533
                                                    .4073
                   4.658
11.4355
                                                     .4073
                                             Mean, .40718
```

Now, one of these salts being formed by a bivalent and the other by a univalent acid, we have to reduce both to a common standard. Doing this, we have the following results for the ratio between the atomic weight of silver and the molecular weight of CO_2 ; if Ag = 1.00,

```
From the acetate, CO_3 = .40724, \pm .000076

" oxalate, " = .40718, \pm .000185

General mean, " = .40723, \pm .000071
```

Here the slight error due to the impurity of the oxalate becomes of such trifling weight that it practically vanishes.

From these data, if Ag = 107.675, $\pm .0096$, CO₂ = 43.8485, $\pm .0086$.

Hence $C = 11.9219, \pm .0111$.

As has already been said, the volatility of silver renders all the foregoing results more or less uncertain. Far better figures are furnished by the combustion of carbon directly, as carried out by Dumas and Stas* in 1840 and by Erdmann and Marchand† in 1841. In both investigations weighed quantities of diamond, of natural graphite, and of artificial graphite were burned in oxygen, and the amount of dioxide produced was estimated by the usual methods. The graphite employed was purified with extreme care by treatment with strong nitric acid and by fusion with caustic alkali. I have reduced all the published weighings to a common standard, so as to show in the third column the amount of oxygen which combines with a unit weight (say one gramme) of carbon. Taking Dumas and Stas' results first in order we have from natural graphite:

With artificial graphite:

```
.992 grm. C gave 3.642 grm. CO<sub>2</sub>.

.998 " 3.662 " 2.6682

1.660 " 6.085 " 2.6654

1.465 " 5.365 " 2.6744

Mean, 2.66985, ± .0013
```

^{*} Compt. Rend., 11, 991-1008. Ann. Chim. Phys., (3,) 1, 1.

⁺ Journ. f. Prakt. Chem., 23, 159.

55

And with diamond:

Mean, 2.6665, $\pm .0007$

Erdmann and Marchand's figures for natural graphite give the following results:

```
1.5376 grm. gave 5.6367 grm. CO<sub>2</sub>. 2.6659
1.6494 " 6.0384 " 2.6609
1.4505 " 5.31575 " 2.6647
```

In one experiment 1.8935 grm. of artificial graphite gave 6.9355 grm. CO₂. Ratio for O, 2.6628. This, combined with the foregoing series, gives a mean of 2.6636, \pm .0007.

With diamond they found:

```
.8052 grm. gave 2.9467 grm. CO<sub>3</sub>. 2.6596
1.0858 " 3.9875 " 2.6632
1.3557 " 4.9659 " 2.6629
1.6305 " 5.97945 " 2.6673
.7500 " 2.7490 " 2.6653

Mean, 2.6637, ±.0009
```

Now, combining all these series, we get the following result:

Hence, if O = 15.9633, $\pm .0035$, C = 11.973, $\pm .0030$.

Another very exact method for determining the atomic weight of carbon was employed by Stas* in 1849. Carefully purified carbon monoxide was passed over a known weight

^{*} Bull. Acad. Bruxelles, 1849, (1,) 31.

of copper oxide at a red heat, and both the residual metal and the carbon dioxide formed were weighed. The weighings were reduced to a vacuum standard, and in each experiment a quantity of copper oxide was taken representing from eight to twenty-four grammes of oxygen. The method, as will at once be seen, is in all essential features similar to that usually employed for determining the composition of water. The figures in the third column, deduced from the weights given by Stas, represent the quantity of carbon monoxide corresponding to one gramme of oxygen:

```
9.265 \text{ grm. O} = 25.483 \text{ CO}_2.
                                            1.75046
 8.327
                  22.900 "
                                             1.75010
                 38.351 "
13.9438
                                             1.75040
11.6124
           "
                 31.935 "
                                             1.75008
18.763
                 51.6055 "
                                             1.75039
                 53.8465 "
19.581
                                             1.74994
                 61.926 "
          46
22.515
                                             1.75043
                 67.003 "
24.360
                                            1.75053
                                      Mean, 1.75029, \pm .00005
```

Hence the molecular weight of carbon monoxide is 27.9404, $\pm .0062$. And C = 11.9771, $\pm .0071$.

Now, in order to complete our discussion, we must combine the four values we have found for carbon:

```
1. By Liebig and Redtenbacher... C = 12.0363, ± .0028
2. By Maumené's figures ____ " = 11.9219, ± .0111
3. By combustion of carbon... " = 11.9730, ± .0030
4. By Stas' method ____ " = 11.9771, ± .0071

General mean ___ " = 12.0021, ± .0019
```

But values one and two are hardly reliable enough to be included in our final estimate. They involve dangerous constant errors, and ought, therefore, to be disregarded. Rejecting them altogether, and taking a general mean from values three and four, we get for the most probable figure for the atomic weight of carbon, C = 11.9736, $\pm .0028$. If oxygen is 16, then carbon becomes 12.0011. In other words, the ratio between oxygen and carbon is almost exactly 16 to 12.

BARIUM.

For determining the atomic weight of barium we have a series of six ratios, established by the labors of Berzelius, Turner, Struve, Pelouze, Marignac, and Dumas. Andrews* and Salvetat,† in their papers upon this subject, gave no details nor weighings; and, therefore, their work may be properly disregarded. First in order in point of importance, if not first chronologically, is the ratio between silver and anhydrous barium chloride, as determined by Pelouze, Marignac, and Dumas.

Pelouze,‡ in 1845, made the three subjoined estimations of this ratio, using his well known volumetric method. A quantity of pure silver was dissolved in nitric acid, and the amount of barium chloride needed to precipitate it was accurately ascertained. In the last column I give the quantity of barium chloride proportional to 100 parts of silver:

```
3.860 grm. BaCl, ppt. 4.002 grm. Ag. 96.452
5.790 " 6.003 " 96.452
2.895 " 3.001 " 96.468

Mean, 96.4573, ± .0036
```

Essentially the same method was adopted by Marignac|| in 1848. His experiments were made upon four samples of barium chloride, as follows. A, commercial barium chloride, purified by recrystallization from water. B, the same salt, calcined, redissolved in water, the solution saturated with carbonic acid, filtered, and allowed to crystallize. C, the preceding salt, washed with alcohol, and again recrystallized. D, the same, again washed with alcohol. For 100 parts of silver the following quantities of chloride were required:

^{*} Chemical Gazette, October, 1852.

[†] Compt. Rend., 17, 318.

¹ Compt. Rend., 20, 1047. Journ. für Prakt. Chem., 35, 73.

^{||} Arch. d. Sci. Phys. et Nat., 8, 271.

Dumas* employed barium chloride prepared from pure barium nitrate, and took the extra precaution of fusing the salt at a red heat in a current of dry hydrochloric acid gas. Three series of experiments upon three samples of chloride gave the following results:

		Se	ries A.	
1.7585 gr	m. BaCl ₂ =	1.826 g	rm. Ag.	Ratio, 96.303
3.842	66	3.988	66	96.339
2.1585	66	2.2405	66	96.340
4.0162	"	4.168	"	96.358
				Mean, 96.3325, ± .0068

^{*} Ann. Chem. Pharm., 113, 22. 1860. Ann. Chim. Phys., (3,) 55, 129.

59

Series B.

1.6625 gr	m. BaCl ₂	= 1.727 gr	m. Ag.	Ratio, 96.265
2.4987	"	2.5946	66	96.304
3.4468	44	3.579	66	96.306 .
4.0822	66	4.2395	66	96.290
4.2062	66	4.3683	66	96.289
4.4564	44	4.629	46	96.271
8.6975	66	9.031	66	96.307
				Mean, 96.2902, ± .0043
		Sei	ries C.	

2.2957 gr	m. BaCl,	= 2.3835	grm. Ag.	Ratio, 96.316
4.1372	44	4.293	et	96.371
4.2662	64	4.430		96.303
4.4764	46	4.647	66	96.329
5.6397	66	5.852	**	96.372
				 Mean. 06.3382. + .0

We have now eight series of experiments upon this ratio, representing thirty distinct estimations. Combining, we get a general mean as follows:

Pelouze	96.4573, ± .0036
Marignac, A	96.3543, ± .0033
" B	96.3540, ± .0013
" C	96.3605, ± .0017
" D	96.3670, ± .0057
Dumas, A	$96.3325, \pm .0068$
" B	96.2902, ± .0043
" C	96.3382, ± .0096
General mean	96.3596, ± .0009

The ratio between silver and crystallized barium chloride has also been fixed by Marignac.* The usual method was employed, and two series of experiments were made; in the second of which the water of crystallization was determined previous to the estimation. Five grammes of chloride were taken in each determination. The following quantities of BaCl₂·2H₂O correspond to 100 parts of silver:

^{*} Journ. f. Prakt. Chem., 74, 212. 1858.

A.	В.
113.109	113.135
113.135	113.122
113.097	113.060
Mean, 113.114, ± .0074	Mean, 113.106, ± .0154

The general mean from both series is 113.113, \pm .0067.

The direct ratio between the chlorides of silver and barium was early established both by Berzelius* and Turner.† Berzelius found that 100 parts of dry barium chloride gave of silver chloride:

Turner made five experiments, with the following results:

137.45 137.54 137.70 137.62 137.64

Of these, Turner regards the fourth and fifth as the most exact. These give a mean of 137.63, $\pm .007$, while the other three are in mean 137.563, $\pm .049$. Combining Berzelius' figures with those of Turner, we get as follows:

Incidentally to some of his other work Marignact determined the percentage of water in crystallized barium chloride. Two sets of three experiments each were made, the first upon five grammes and the second upon ten grammes of salt. The following are the percentages obtained:

^{*} Poggend. Annal., 8, 177.

[†] Phil. Trans., 1829, 291.

[‡] Journ. f. Prakt. Chem., 74, 212. 1858.

A.	В.
14.790	14.80
14.796	14.81
14.800	14.80
Mean, 14.795, ± .0019	Mean, 14.803, \pm .002
General mean of both seri	es, 14.799, ± .0014

The ratio between barium nitrate and barium sulphate has been determined only by Turner.* According to his experiments 100 parts of sulphate correspond to the following quantities of nitrate:

For the similar ratio between the sulphate and the chloride there are experiments by Turner, Berzelius, Struve, and Marignac. Turner† found that 100 parts of chloride ignited with sulphuric acid gave 112.19 parts of sulphate. By the common method of precipitation and filtration a lower figure was obtained, because of the slight solubility of the sulphate. This point bears directly upon many other atomic weight determinations.

Berzelius,‡ treating barium chloride with sulphuric acid, obtained the following results in BaSO₄ for 100 parts of BaCl₂:

Struve, || in two experiments, found:

^{*} Phil. Trans., 1833, 538. † Phil. Trans., 1829, 291. † Poggend. Annal., 8, 177. || Ann. Chem. Pharm., 80, 204. 1851.

Marignac's* three results are as follows:

```
8.520 grm. BaCl<sub>2</sub> gave 9.543 BaSO<sub>4</sub>. Ratio, 112.007
8.519 " 9.544 " 112.032
8.520 " 9.542 " 111.995
Mean, 112.011, ± .0071
```

Rejecting Turner's single result as unimportant, we may combine the other series:

```
Berzelius ______ 112.175, ± .0034
Struve _____ 112.0938, ± .0018
Marignac _____ 112.011, ± .0071
General mean _____ 112.106, ± .0015
```

The data from which we are to calculate the atomic weight of barium may now be tabulated as follows:

```
(1.) Ag_2: BaCl_2:: 100: 96.3596, \pm .0009
(2.) Ag_2: BaCl_2.2H_2O:: 100: 113.113, \pm .0067
(3.) BaCl_2: 2AgCl:: 100: 137.841, \pm .0047
(4.) Per cent. of H_2O in BaCl_2.2H_2O, 14.799, \pm .0014
(5.) BaSO_4: BaN_2O_6:: 112.028, \pm .014
(6.) BaCl_2: BaSO_4:: 100: 112.106, \pm .0015
```

From these ratios, with the aid of the atomic weights already established, we can immediately calculate four independent values for the molecular weight of BaCl₂:

We have here an interesting example of the compensation of constant errors. Ratios (2) and (4) both represent work done by Marignac upon barium chloride containing water of crystallization. If now, as is not improbable, the salt contained a trifling excess of water, the molecular weight of barium chloride as calculated from (2) would come out too high, while on the other hand the result from ratio (4) would err in the opposite direction. In point of fact, the

^{*} Journ. f. Prakt. Chem., 74, 212. 1858.

63

two results in the present calculation nearly compensate each other, and, on account of their relatively high probable errors, they exert but an unimportant influence upon the general mean.

In conclusion, we have three independent values for the atomic weight of barium:

```
From mol. wt. of BaCl<sub>2</sub> .... Ba = 136.765, \pm .031

From ratio (5) .... " = 136.795, \pm .364

From ratio (6) .... " = 136.595, \pm .309

General mean .... " = 136.763, \pm .031
```

If O = 16, then Ba = 137.007. In other words, the ratio between oxygen and barium is almost an exact ratio between two whole numbers.

In the above discussion it will at once be noticed that the second and third values for Ba have very high probable errors, and that they therefore exert almost no influence upon the general mean. This fact by no means renders them worthless however, for, at the lowest estimate, they are useful in confirmation of the better determinations. It is also highly probable that the method of discussion, rigidly carried out, does not do them absolute justice.

STRONTIUM.

The ratios which fix the atomic weight of strontium resemble in general terms those relating to barium, only they are fewer in number and represent a comparatively small amount of work. The early experiments of Stromeyer,* who measured the volume of CO₂ evolved from a known weight of strontium carbonate, are hardly available for the present discussion. So also we may exclude the determination by Salvétat,† who neglected to publish sufficient details.

Taking the ratio between strontium chloride and silver first in order, we have series of figures by Pelouze and by Dumas. Pelouze‡ employed the volumetric method already described under barium, and in two experiments obtained the subjoined results. In another column I append the ratio between SrCl, and 100 parts of silver:

Dumas, || by the same general method, made sets of experiments with three samples of chloride which had previously been fused in a current of dry hydrochloric acid. His results, expressed in the usual way, are as follows:

Series A.

3.137 gm	m. SrCl ₂	= 4.280 g	rm. Ag.	Ratio, 73.2944	
1.982	46	2.705	46	73.2717	
3.041	"	4.142	46	73.4186	
3.099	46	4.219	44	73-4534	
				-	
				Mean, 73.3595, ± .030	3

^{*} Schweigg. Journ., 19, 228. 1816.

[†] Compt. Rend., 17, 318. 1843.

[†] Compt. Rend., 20, 1047. 1845.

[|] Ann. Chim. Phys., (3,) 55, 29. 1859. Ann. Chem. Pharm., 113, 34.

Series R.

3.356 grm.	$SrCl_2 =$	4-574	grm.	Ag.	Ratio,	73-3713	•
6.3645	44	8.667	66			73.4327	
7.131	66	9.712	66			73.4246	•
					Mean,	73.4095,	士 .0130

Series C.

7.213 g	rm. SrCl ₂ =	= 9.811 g	rm. Ag.	Ratio, 73.519	5
2.206	66	3.006	66	73.386	6
4.268	66	5.816	66	73.552	9
4.018	46	5-477	66	73.361	3
				Mean. 73.455	- 1. + .0321

Combining, we have:

The foregoing figures apply to anhydrous strontium chloride. The ratio between silver and the crystallized salt, SrCl₂.6H₂O, has also been determined in two series of experiments by Marignac.* Five grammes of salt were used in each estimation, and, in the second series, the percentage of water was first determined. The quantities of the salt corresponding to 100 parts of silver are given in the last column:

Series A.

5 grm.
$$SrCl_2.6H_2O = 4.0515$$
 grm. Ag. 123.411

" " 4.0495 " 123.472

" 4.0505 " 123.442

Mean, 123.442, \pm .012

Series B.

General mean of both series, 123.470, ± .006

^{*} Journ. Prakt. Chem., 74, 216. 1858.

In the same paper Marignac gives two sets of determinations of the percentage of water in crystallized strontium chloride. The first set, corresponding to "B" above, comes out thus:

In the second set ten grammes of salt were taken at a time, and the following percentages were found:

General mean, from both series, 40.575, ± .0015

The chloride used in the series of estimations last given was subsequently employed for ascertaining the ratio between it and the sulphate. Converted directly into sulphate, 100 parts of chloride yield the quantities given in the third column:

```
5.942 grm. SrCl<sub>2</sub> gave 6.887 grm. SrSO<sub>4</sub>. 115.932
5.941 " 6.8855 " 115.949
5.942 " 6.884 " 115.927
... Mean, 115.936, ± .004
```

Now, to sum up the ratios and calculate the atomic weight of strontium.

```
(1.) Ag: SrCl<sub>2</sub>:: 100: 73.4655, ± .0046

(2.) Ag: SrCl<sub>2</sub>.6H<sub>2</sub>O:: 100: 123.470, ± .006

(3.) Per cent. of H<sub>2</sub>O in SrCl<sub>2</sub>.6H<sub>2</sub>O, 40.575, ± .0015

(4.) SrCl<sub>2</sub>: SrSO<sub>4</sub>:: 100: 115.936, ± .004
```

We now have the molecular weight of SrCl,, as follows:

And for the atomic weight of strontium itself we have two values, as follows:

If O = 16, then Sr = 87.575.

CALCIUM.

For determining the atomic weight of calcium we have sets of experiments by Berzelius, Erdmann and Marchand, and Dumas. Salvétat* also has published an estimation, but without the details necessary to enable us to make use of his results. I also find a reference† to some work of Marignac; which, however, seems to have been of but little importance. The earlier work of Berzelius was very inexact as regards calcium, and it is not until we come down to the year 1842 that we find any material of decided value.

The most important factor in our present discussion is the composition of calcium carbonate, as worked out by Dumas and by Erdmann and Marchand.

In 1842 Dumast made three ignitions of Iceland spar, and determined the percentages of carbon dioxide driven off and of lime remaining. The impurities of the material were also determined, the correction for them applied, and the weighings reduced to a vacuum standard. The percentage of lime came out as follows:

56.12 56.04 56.06 ———— Mean, 56.073, ± .016

^{*} Compt. Rend., 17, 318. 1843.

⁺ See Oudeman's monograph, p. 51.

[‡] Compt. Rend., 14, 537. 1842.

About this same time Erdmann and Marchand* began their researches upon the same subject. Two ignitions of spar, containing .04 per cent. of impurity, gave respectively 56.09 and 56.18 per cent. of residue; but these results are not exact enough for us to consider further. Four other results obtained with artificial calcium carbonate are more noteworthy. The carbonate was precipitated from a solution of pure calcium chloride by ammonium carbonate, was washed thoroughly with hot water, and dried at a temperature of 180°. With this preparation the following residues of lime were obtained:

It was subsequently shown by Berzelius that calcium carbonate prepared by this method retains traces of water even at 200°, and that minute quantities of chloride are also held by it. These sources of error are, however, in opposite directions, since one would tend to diminish and the other to increase the weight of residue.

In the same paper there are also two direct estimations of carbonic acid in pure Iceland spar, which correspond to the following percentages of lime:

In a still later paper† the same investigators give another series of results based upon the ignition of Iceland spar. The impurities were carefully estimated, and the percentages of lime are suitably corrected:

^{*} Journ. für Prakt. Chem., 26, 472. 1842.

[†] Journ. für Prakt. Chem., 31, 269. 1844.

69

4.2134	grm. CaCO _s g	ave 2.3594 gi	m. CaO.	55.997 P	er cent.
15.1385	66	8.4810	66	56.022	44
23.5503	64	13.1958	64	56.031	46
23.6390	46	13.2456	66	56.032	44
42.0295	46	23.5533	66	56.044	44
49.7007	44	27.8536	66	56.042	46
					

CALCIUM.

Mean, 56.028, \pm .0047

Six years later Erdmann and Marchand* published one more result upon the ignition of calcium carbonate. They found that the compound began giving off carbon dioxide below the temperature at which their previous samples had been dried, or about 200°, and that, on the other hand, traces of the dioxide were retained by the lime after ignition. These two errors do not compensate each other, since both tend to raise the percentage of lime. In the one experiment now under consideration these errors were accurately estimated, and the needful corrections were applied to the final result. The percentage of residual lime in this case came out 55.998. This agrees tolerably well with the figures found in the direct estimation of carbonic acid, and, if combined with those two, gives a mean for all three of 56.006, \pm .0043.

Combining all these series we get the following result:

Dumas Erdmann an			
46	66	 56.028,	± .0047
44	44	 56.006,	± .0043
Gene	ral mean	 56,0108	+ .0020

For reasons given above this mean is probably vitiated by a slight constant error, which makes the figure a trifle too high.

In the earliest of three papers by Erdmann and Marchand there is also given a series of determinations of the ratio between calcium carbonate and sulphate. Pure Iceland

^{*} Journ. für Prakt. Chem., 50, 237. 1850.

spar was carefully converted into calcium sulphate, and the gain in weight noted. One hundred parts of spar gave of sulphate:

136.07 136.06 136.02 136.06

Mean, 136.0525, \pm .0071

In 1843 the atomic weight of calcium was redetermined by Berzelius,* who investigated the ratio between lime and calcium sulphate. The calcium was first precipitated from a pure solution of nitrate by means of ammonium carbonate, and the thoroughly washed precipitate was dried and strongly ignited in order to obtain lime wholly free from extraneous matter. This lime was then, with suitable precautions, treated with sulphuric acid, and the resulting sulphate was weighed. Correction was applied for the trace of solid impurity contained in the acid, but not for the weighing in air. The figures in the last column represent the percentage of weight gained by the lime upon conversion into sulphate:

```
      1.80425 grm. CaO gained
      2.56735 grm.
      142.295

      2.50400
      " 3.57050 " 142.592

      3.90000
      " 5.55140 " 142.343

      3.04250
      " 4.32650 " 142.202

      3.45900
      " 4.93140 " 142.567
```

Mean, 142.3998, ± .0518

Last of all we have the ratio between calcium chloride and silver, as determined by Dumas.† Pure calcium chloride was first ignited in a stream of dry hydrochloric acid, and the solution of this salt was afterwards titrated with a silver solution in the usual way. The CaCl, proportional to 100 parts of Ag is given in a third column:

^{*} Journ. für Prakt. Chem., 31, 263. Ann. Chem. Pharm., 46, 241.

[†] Ann. Chim. Phys., (3,) 55, 129. 1859. Ann. Chem. Pharm., 113, 34.

```
2.738 grm. CaCl<sub>2</sub> = 5.309 grm. Ag. 51.573

2.436 " 4.731 " 51.490

1.859 " 3.617 " 51.396

2.771 " 5.3885 " 51.424

2.240 " 4.3585 " 51.394
```

Mean, 51.4554, ± .0230.

We have now four ratios to calculate from, as follows:

- (1.) Per cent. of CaO in CaCO₂, 56.0198, \pm .0029
- (2.) $CaO : SO_1 :: 100 : 142.3998, \pm .0518$
- (3.) $CaCO_8 : CaSO_4 :: 100 : 136.0525, \pm .0071$
- (4.) Ag: CaCl₂:: 100: 51.4554, ± .0230

These give us the subjoined values for calcium:

General mean..... = 39.990, ± .010

If O = 16, then Ca = 40.082.

A glance at the above figures will show that, if, as is probable, the value deduced from the composition of calcium carbonate is a trifle too high, the general mean must be too high also. It is, therefore, interesting to see what result the very latest of Erdmann and Marchand's experiments will lead to. They found, after taking every precaution, in a single experiment that calcium carbonate yielded 55.998 per cent. of lime. From this we get Ca = 39.905; or, if O = 16, Ca = 39.997. It is possible, then, that "Prout's law" may hold good for calcium.

LEAD.

For the atomic weight of lead we have to consider experiments made upon the oxide, chloride, nitrate, and sulphate. The researches of Berzelius upon the carbonate and various organic salts need not now be considered, nor is it worth while to take into account any work of his done before the year 1818. The results obtained by Döbereiner* and by Longchamp† are also without special present value.

For the exact composition of lead oxide we have to depend upon the researches of Berzelius. His experiments were made at different times through quite a number of years; but were finally summed up in the last edition of his famous "Lehrbuch." In general terms his method of experiment was very simple. Perfectly pure lead oxide was heated in a current of hydrogen, and the reduced metal weighed. From his weighings I have calculated the percentages of lead thus found and given them in a third column:

Earlier Results.

8.045 grm.	PbO	gave 7.4675	grm. 1	Pb. 92.8217	per cent.
14.183	44	13.165	66	92.8224	64
10.8645	44	10.084	44	92.8160	66
13.1465	44	12.2045	64	92.8346	66
21.9425	44	20. 3695	4.6	92.8313	66
11.159	44	10.359	46	92.8309	• 6
		Lat	est.		
6.6155	44	6. 141	44	92.8275	44
14.487	66	13.448	46	92.8280	46
14.626	44	13.5775	44	92.8313	d's

Mean, 92.8271, $\pm .0013$

For the synthesis of lead sulphate we have data by Berzelius, Turner, and Stas. Berzelius, | whose experiments

^{*} Schweig. Journ., 17, 241. 1816.

[†] Ann. Chim. Phys., 34, 105. 1827.

[‡] Bd. 3, s. 1218.

^{||} Lehrbuch, 5th Ed., 3, 1187.

73

were intended rather to fix the atomic weight of sulphur, dissolved in each estimation ten grammes of pure lead in nitric acid, then treated the resulting nitrate with sulphuric acid, brought the sulphate thus formed to dryness, and weighed. One hundred parts of metal yield of PbSO₄:

Turner,* in three similar experiments, found as follows:

146.430 146.398 146.375 Mean, 146.401, ± .011

In these results of Turner's absolute weights are implied.

The results of Stas' syntheses,† effected after the same general method, but with variations in details, are as follows. Corrections for weighing in air were applied:

146.443 146.427 146.419 146.432 146.421 146.423 Mean, 146.4275, ± .0024

Combining, we get the subjoined result:

 Berzelius
 146.419, ± .012

 Turner
 146.401, ± .011

 Stas
 146.4275, ± .0024

 General mean
 146.4262, ± .0023

Turner, in the same paper, also gives a series of syntheses of lead sulphate, in which he starts from the oxide instead

^{*} Phil. Trans., 1833, 527-538.

[†] Aronstein's Translation, 333.

of from the metal. One hundred parts of PbO, upon conversion into PbSO₄, gained weight as follows:

35.84 35.71 35.84 35.75 35.79 35.78 35.92

Mean, 35.804, ± .018

These figures are not wholly reliable. Numbers one, two, and three represent lead oxide contaminated with traces of nitrate. The oxide of four, five, and six contained traces of minium. Number seven was free from these sources of error, and, therefore, deserves more consideration. The series as a whole undoubtedly gives too low a figure; and this error would tend to slightly raise the atomic weight of lead.

Still a third series by Turner establishes the ratio between the nitrate and the sulphate; a known weight of the former being in each experiment converted into the latter. One hundred parts of sulphate represent of nitrate:

> 109.312 109.310 109.300

Mean, 109.307, \pm .002

In all these experiments by Turner the necessary corrections were made for weighing in air.

For the ratio between lead chloride and silver we have a series of results by Marignac and one experiment by Dumas. There are also unavailable data by Turner and by Berzelius.

Marignac,* applying the method used in his researches upon barium and strontium, and working with lead chloride which had been dried at 200°, obtained these results.

^{*} Journ. für Prakt. Chem., 74, 218. 1858.

LEAD. 75

The third column gives the ratio between PbCl, and 100 parts of Ag:

```
4.9975 grm. PbCl<sub>2</sub> = 3.8810 grm. Ag. 128.768

4.9980 " 3.8835 " 128.698

5.0000 " 3.8835 " 128.750

5.0000 " 3.8860 " 128.667

Mean, 128.721, ± .016
```

Dumas,* in his investigations, found that lead chloride retains traces of water even at 250°, and is sometimes also contaminated with oxychloride. In one estimation 8.700 grammes PbCl, saturated 6.750 of Ag. The chloride contained .009 of impurity; hence, correcting, Ag: PbCl,:: 100: 128.750. If we assign this figure equal weight with those of Marignac, we get as the mean of all, 128.7266, ± .013. The sources of error indicated by Dumas, if they are really involved in this mean, would tend slightly to raise the atomic weight of lead.

The synthesis of lead nitrate, as carried out by Stas,† gives excellent results. Two series of experiments were made, with from 103 to 250 grammes of lead in each determination. The metal was dissolved in nitric acid, the solution evaporated to dryness with extreme care, and the nitrate weighed. All weighings were reduced to the vacuum standard. In series A the lead nitrate was dried in an air current at a temperature of about 155°. In series B the drying was effected in vacuo. 100 of lead yield of nitrate:

```
A.

159.973
159.975
159.982
159.975
159.968
159.973
Mean, 159.9743, ± .0012
```

^{*} Ann. Chem. Pharm., 113, 35. 1860.

[†] Aronstein's Translation, 326.

B.
159.970
159.964
159.959
159.965

Mean, 159.9645, ± .0015

Mean from both series, 159.9704, \pm .0010

There still remain to be noticed two sets of experiments upon lead nitrate, which were originally intended to establish the atomic weight of nitrogen. Lead nitrate was carefully ignited and the residual oxide weighed. The first series, bearing Svanberg's name,* gives simply the percentage of oxide found, as follows:

67.4030 67.4036 67.4043 67.3956

Mean, 67.4016, $\pm .0014$

The second series is by Anderson,† and gives the weighings upon which the percentages rest. The latter come out thus:

```
5.19485 grm. PbN<sub>2</sub>O<sub>6</sub> gave 3.5017 grm. PbO. 67.4071 per cent.

9.7244 " 6.5546 " 67.4037 "

9.2181 " 6.2134 " 67.4044 "

9.6530 " 6.5057 " 67.3957 "

Mean, 67.4027, ± .0016
```

It will at once be seen that these series are identical; the discordance between the first figures of the two being undoubtedly due to some misprint in the weighings of the Anderson set. How it happens that the same work has been published by two separate authors I will not attempt to explain; neither will I undertake to determine which of the two is really entitled to credit.

^{*} Journ. für Prakt. Chem., 27, 381. 1842.

[†] Ann. Chim. Phys., (3,) 9, 254. 1843.

LEAD. 77

We have now seven ratios upon which to base our computations:

```
(1.) Per cent. of Pb in PbO, 92.8271, ± .0013
```

- (2.) Per cent. of PbO in PbN₂O₆, 67.4016, \pm .0014
- (3.) Pb: PbSO₄:: 100: 146.4262, \pm .0023
- (4.) PbO: $PbSO_4$:: 100: 135.804, \pm .018
- (5.) $PbSO_4: PbN_2O_6:: 100: 109.307, \pm .002$
- (6.) Pb: PbN_2O_6 :: 100: 159.9704, \pm .0010
- (7.) Ag: PbCl₂:: 100: 128.7266, \pm .013

Discussing these separately, we get an equal number of values for the atomic weight of lead:

```
From (1) ______Pb _= 206.587, \pm .059

" (2) ______ " = 207.046, \pm .041

" (3) _____ " = 206.435, \pm .041

" (4) _____ " = 207.131, \pm .118

" (5) _____ " = 204.803, \pm .329

" (6) ____ " = 206.454, \pm .037

" (7) _____ " = 206.473, \pm .042

General mean ____ " = 206.604, \pm .019
```

If O = 16, this becomes Pb = 207.079.

In the above discussion are included several values which diverge widely from this general mean, and which, for other reasons, are probably erroneous. Although but one of these carries much weight, it is as well to exclude them, and to base our computations upon the others. If, now, we reject the second, fourth, and fifth values, we get for the atomic weight of lead, $Pb = 206.471, \pm .021$. If O = 16, this becomes Pb = 206.946.

From the synthesis of the nitrate Stas found 206.918, and from the sulphate, 206.934. The agreement of these values with our own general mean is certainly very close.

FLUORINE.

The atomic weight of fluorine has been determined only by one general method, namely, by the conversion of fluorides into sulphates. Excluding the early results of Davy,* we have only to consider the experiments of Berzelius, Louyet, Dumas, and DeLuca, with reference to the fluorides of calcium, sodium, potassium, barium, and lead.

The ratio between calcium fluoride and sulphate has been determined by the four investigators above named, and by one general process. The fluoride is treated with strong sulphuric acid, the resulting sulphate is ignited, and the product weighed. In order to ensure complete transformation special precautions are necessary; such, for instance, as repeated treatment with sulphuric acid, and so on. For details like these the original papers must be consulted.

The first experiments in chronological order are those of Berzelius,† who operated upon an artificial calcium fluoride. He found, in three experiments, for one part of fluoride the following of sulphate:

1.749 1.750 1.751

Mean, 1.750, ± .0004

Louyet's researchest were much more elaborate than the foregoing. He began with a remarkably concordant series of results upon fluor spar, in which one gramme of the fluoride yielded from 1.734 to 1.737 of sulphate. At first he regarded these as accurate, but he soon found that particles of spar had been coated with sulphate, and had therefore escaped action. In the following series this source of error was guarded against.

^{*} Phil. Trans., 1814, 64.

[†] Poggend. Annal., 8, 1. 1826.

[‡] Ann. Chim. Phys., (3,) 25, 300. 1849.

Starting with fluor spar, Louyet found of sulphate as follows:

1.742 1.744 1.745 1.744 1.7435 1.7435 Mean, 1.7437, ± .0003

A second series, upon artificial fluoride, gave:

1.743 1.741 1.741 Mean, 1.7417, ± .0004

Dumas* published but one result for calcium fluoride. .495 grm. gave .864 grm. sulphate, the ratio being 1: 1.7455.

De Luca† worked with a very pure fluor spar, and published the following results. The ratio between CaSO₄ and one gramme of CaF₂ is given in the third column:

.9305 gm	m. CaF ₂ gav	re 1.630 gr	m. CaSO ₄ .	1.7518
.836	66	1.459	66	1.7452
.502	66	.8755	66	1.7440
.3985	44	.6945	46	1.7428

If we include Dumas' single result with these, we get a mean of 1.7459, $\pm .0011$.

Upon combining all these series, we get as follows:

Berzelius	1.7500,	± .0004
Louyet, 1st series	1.7437,	±.0003
" 2d "	1.7417,	±.0004
De Luca and Dumas	1.7459,	1100. ±
General mean	1.74493,	± .0002

For the ratio between the two sodium salts we have experiments by Dumas and by Louyet.‡ According to Louyet one gramme of NaF gives of Na₂SO₄:

^{*} Ann. Chem. Pharm., 113, 28. 1860.

[†] Compt. Rend., 51, 299. 1860.

[‡] See the papers already quoted.

The weighings published by Dumas are as follows:

The general mean of both series is 1.6863, \pm .0004.

Dumas also gives experiments upon potassium fluoride. The quantity of sulphate formed from one gramme of fluoride is given in the last column:

The ratios for the fluorides of lead and of barium are due entirely to Louyet. One gramme BaF, gave of BaSO₄:

With the lead fluoride a new method of treatment was adopted. The salt was fused, powdered, dissolved in nitric acid, and precipitated by dilute sulphuric acid. The evaporation of the fluid and the ignition of the sulphate was then effected without transfer. Five grammes of fluoride were taken in each operation, yielding of sulphate:

We now have five ratios to calculate from, as follows:

```
(1.) CaF_2: CaSO_4:: 1.0: 1.74493, \pm .0002

(2.) NaF: Na_2SO_4:: 1.0: 1.6863, \pm .0004

(3.) KF: K_2SO_4:: 1.0: 1.4991, \pm .0007

(4.) BaF_2: BaSO_4:: 1.0: 1.3310, \pm .0004

(5.) PbF_2: PbSO_4:: 5.0: 6.1783, \pm .0002
```

From these we get five values for F:

If O = 16, this becomes 19.027.

Before leaving the subject of fluorine we must notice two possible sources of error beyond the always to be considered one of impurities in the materials employed. First, an incomplete conversion of a fluoride into a sulphate would lead to results tending to raise the atomic weight of fluorine. On the other hand, the value for fluorine which has most weight is that derived from calcium fluoride. But it was shown under calcium that the atomic weight determined for that metal was probably a trifle too high. This error, introduced into our fluorine calculations, tends to lower our final results. These two errors, then, if they really exist, will, in part at least, compensate each other.

PHOSPHORUS.

The material from which we are to calculate the atomic weight of phosphorus is by no means abundant. Berzelius, in his Lehrbuch,* adduces only his own experiments upon the precipitation of gold by phosphorus, and ignores all the earlier work relating to the composition of the phosphates. These experiments we will consider with reference to gold.

Pelouze,† in a single titration of phosphorus trichloride with a standard solution of silver, obtained a wholly erroneous result; and Jacquelain,‡ in his similar experiments, did even worse. Schrötter's criticism upon Jacquelain sufficiently disposes of the latter.

There are, in short, but two investigations upon the atomic weight of phosphorus which have any value for present purposes, namely, the researches of Schrötter and of Dumas. These chemists worked with different materials and by different methods, and yet obtained beautifully concordant results.

Schrötters burned pure amorphous phosphorus in dry oxygen, and weighed the pentoxide thus formed. One gramme of P yielded P₂O₅ in the following proportions:

```
\begin{array}{c} 2.28909 \\ 2.28783 \\ 2.29300 \\ 2.28831 \\ 2.29040 \\ 2.28788 \\ 2.28848 \\ 2.28856 \\ 2.28959 \\ 2.28872 \\ \hline \\ \text{Mean, } 2.289186, \pm .00033 \\ \text{Hence P} = 30.9562, \pm .0074. \end{array}
```

```
*5th Ed., 1188.

† Compt. Rend., 20, 1047.

‡ Compt. Rend., 33, 693.

| Journ. für Prakt. Chem., 57, 315.

§ Journ. für Prakt. Chem., 53, 435. 1851.
```

Dumas* prepared pure phosphorus trichloride by the action of dry chlorine upon red phosphorus. The portion used in his experiments boiled between 76° and 78°. This was titrated with a standard solution of silver in the usual manner. Dumas publishes weights, from which I calculate the figures given in the third column, representing the quantity of trichloride proportional to 100 parts of silver:

```
      1.787 grm. PCl<sub>3</sub> = 4.208 grm. Ag.
      42.4667

      1.466 " 3.454 " 42.4435

      2.056 " 4.844 " 42.4443

      2.925 " 6.890 " 42.4528

      3.220 " 7.582 " 42.4690
```

Mean, 42.4553, $\pm .0036$

Hence $P = 31.0314, \pm .0467$.

Now, combining these two values, we have:

If O = 16, this becomes 31.0292.

The fact here noticeable, that Dumas' figures give a value for P slightly higher than that deduced from those of Schrötter, may be accounted for upon the supposition that the phosphorus trichloride contained traces of oxychloride. Such an impurity would tend to raise the apparent atomic weight of phosphorus, and its occurrence is by no means improbable.

^{*} Ann. Chem. Pharm., 113, 29. 1860.

BORON.

The atomic weight of this element has been determined by Berzelius and by Laurent, and calculated by Dumas from some experiments by Deville.

Berzelius* based his determination upon three concordant estimations of the percentage of water in borax. Laurent† made use of two similar estimations, and all five may be properly put in one series, thus:

Hence $B = 10.943, \pm .023$.

Dumas't calculations were based on Deville's analyses of the chloride and bromide of boron, which give the ratios between AgCl and BCl_s, and between AgBr and BBr_s. Reducing the weighings to a common standard, 100 parts of silver chloride correspond to the quantities of boron trichloride given in the third column:

Mean, 47.13, \pm .013

Hence $B = 10.808, \pm .174$.

With the bromide, 2.446 BBr₃ gave 5.496 AgBr. If we assign this experiment equal weight with one in the chloride series, and include the probable error of Br, B = 10.964, $\pm .364$.

The three values combine as follows:

^{*} Poggend. Annal., 8, 1. 1826.

[†] Journ. für Prakt. Chem., 47, 415. 1849.

[†] Ann. Chem. Pharm., 113, 31. 1860.

If O = 16, B = 10.966.

Further investigation of the atomic weight of boron is evidently desirable.

SILICON.

Although Berzelius* attempted to ascertain the atomic weight of silicon, first by converting pure Si into SiO₂, and later from the analysis of BaSiF₆, his results were not satisfactory. We need only consider the estimations of Pelouze, Schiel, and Dumas.

Pelouze,† experimenting upon silicon tetrachloride, employed his usual method of titration with a solution containing a known weight of silver. One hundred parts of Ag gave the following equivalencies of SiCl₄:

Hence Si = 28.408.

Essentially the same method was adopted by Dumas.‡ Pure SiCl₄ was weighed in a sealed glass bulb, then decomposed by water, and titrated. The results for 100 Ag are given in the third column:

2.899 grm.
$$SiCl_4 = 7.3558$$
 grm. Ag. 39.411
1.242 " 3.154 " 39.379
3.221 " 8.1875 " 39.340
Mean, 39.377, \pm .014

Hence Si = 28.117.

^{*} Lehrbuch, 5 Aufl., 3, 1200.

[†] Compt. Rend., 20, 1047. 1845.

[‡] Ann. Chem. Pharm., 113, 31. 1860.

Dumas and Pelouze's series combine as follows:

Hence $SiCl_4 = 169.810, \pm .034$.

Schiel,* also studying the chloride of silicon, decomposed, it by ammonia. After warming and long standing it was filtered, and in the filtrate the chlorine was estimated as AgCl. One hundred parts of AgCl correspond to the quantities of SiCl₄ given in the last column:

Hence $SiCl_4 = 169.437$, $\pm .080$, and Si = 27.957. Combining the values for $SiCl_4$ we have this result:

Pelouze and Dumas_____SiCl₄ = 169.810,
$$\pm$$
 .034
Schiel____ " = 169.437, \pm .080
General mean ___ " = 169.675, \pm .031

Hence Si = 28.195, $\pm .066$; or, if O = 16, Si = 28.260.

It will be observed that all of these determinations rest upon the composition of SiCl₄, a compound for which it would not be easy to guarantee absolute purity. All the errors likely to occur in the determination of the atomic weight would be plus errors, so that the value deduced above is almost certainly too high.

^{*} Ann. Chem. Pharm., 120, 94.

LITHIUM.

The earlier determinations of the atomic weight of lithium by Arfvedson, Stromeyer, C. G. Gmelin, and Kralovanzky were all erroneous, because of the presence of sodium compounds in the material employed. The results of Berzelius, Hagen, and Hermann were also incorrect, and need no further notice here. The only investigations which we need to consider are those of Mallet, Diehl, Troost, and Stas.

Mallet's experiments* were conducted upon lithium chloride, which had been purified as completely as possible. In two trials the chloride was precipitated by nitrate of silver, which was collected upon a filter and estimated in the ordinary way. The figures in the third column represent the LiCl proportional to 100 parts of AgCl:

In a third experiment the LiCl was titrated with a standard solution of silver. 3.9942 grm. LiCl balanced 10.1702 grm. Ag, equivalent to 13.511 grm. AgCl. Hence 100 AgCl = 29.563 LiCl. Mean of all three experiments, 29.581, \pm .0087.

Diehl,† whose paper begins with a good resumé of all the earlier determinations, describes experiments made with lithium carbonate. This salt, which was spectroscopically pure, was dried at 130° before weighing. It was then placed in an apparatus from which the carbon dioxide generated by the action of pure sulphuric acid upon it could be expelled, and the loss of weight determined. From this loss the following percentages of CO₂ in Li₂CO₃ were determined:

59.422 59.404 59.440 59.401 Mean, 59.417, ± .006

^{*}Silliman's Amer. Journal, November, 1856. Chem. Gazette, 15, 7.

[†] Ann. Chem. Pharm., 121, 93.

Diehl's investigation was quickly followed by a confirmation from Troost.* This chemist, in an earlier paper,† had sought to fix the atomic weight of lithium by an analysis of the sulphate, and had found a value not far from 6.5; thus confirming the results of Berzelius and of Hagen, who had employed the same method. But Diehl showed that the BaSO₄ precipitated from Li₂SO₄ always retained traces of Li, which were recognizable by spectral analysis, and which accounted for the error. In the later paper Troos made use of the chloride and the carbonate of lithium, both spectroscopically pure. The carbonate was strongly ignited with pure quartz powder, thus losing carbon dioxide, which loss was easily estimated. The subjoined results were obtained:

This combined with Diehl's mean, 59.417, $\pm .006$, gives a general mean of 59.420, $\pm .0057$.

The lithium chloride employed by Troost was heated in a stream of dry hydrochloric acid gas; of which the excess, after cooling, was expelled by a current of dry air. The salt was weighed in the same tube in which the foregoing operations had been performed, and the chlorine was then estimated as silver chloride. The usual ratio between LiCl and 100 parts of AgCl is given in the third column:

This combined with Mallet's mean, 29.581, \pm .0087, gives a general mean of 59.584, \pm .0075.

Finally, we come to the work of Stas,‡ which was exe-

^{*} Zeit. Anal. Chem., 1, 402.

[†] Annales d. Chim. et d. Phys., 51, 108.

[†] Aronstein's Translation, 279-302.

89

cuted with his usual wonderful accuracy. In three titrations, in which all the weights were reduced to a vacuum standard, the following quantities of LiCl balanced 100 parts of pure silver:

In a second series of experiments, intended for determining the atomic weight of nitrogen, LiCl was converted into LiNO₃. The method was that employed for a similar purpose with the chlorides of sodium and of potassium. One hundred parts of LiCl gave of LiNO₃:

We have now the following ratios from which to deduce the atomic weight of lithium:

- (1.) AgCl: LiCl:: 100: 29.584, \pm .0075 (2.) Ag: LiCl:: 100: 39.358, \pm .001
- (3.) LiCl: LiNO₃:: 100: 162.5953, \pm .0025
- (4.) Per cent. of CO₂ in Li₂CO₃, 59.420, ± .0057

Hence two values for the molecular weight of LiCl:

For lithium itself we get three values:

If O = 16, then Li = 7.0235. Stas himself gives 7.022 as his determination. Difference, .0015.

RUBIDIUM.

The atomic weight of rubidium has been determined by Bunsen, Piccard, and Godeffroy; but only from analyses of the chloride.

Bunsen,* employing ordinary gravimetric methods, estimated the ratio between AgCl and RbCl. His rubidium chloride was purified by fractional crystallization of the chloroplatinate. He obtained the following results, to which, in a third column, I add the ratio between RbCl and 100 parts of AgCl:

```
One grm. RbCl gave 1.1873 grm. AgCl.

" 1.1873 " 84.225

" 1.1850 " 84.388

" 1.1880 " 84.175

Mean, 84.253, ± .031
```

The work of Piccard† was similar to that of Bunsen. In weighing, the crucible containing the silver chloride was balanced by a precisely similar crucible, in order to avoid the correction for displacement of air. The filter was burned separately from the AgCl, as usual; but the small amount of material adhering to the ash was reckoned as metallic silver. The rubidium chloride was purified by Bunsen's method. The results, expressed according to the foregoing standard, are as follows:

```
1.1587 grm. RbCl = 1.372 AgCl + .0019 Ag.
                                                  84.300
                    1.6632
                                  .0030 "
1.4055
                                                  84.303
                    1.1850
                            66
                                   .0024 "
100.1
                                                  84.245
                                   .0018 "
                                                  84.313
1.5141
                    1.7934
                                          Mean, 84.290, ± .0105
```

Godeffroy,‡ starting with material containing both rubidium and cæsium, separated the two metals by fractional

^{*} Zeit. Anal. Chem., 1, 136. Poggend. Annal., 113, 339. 1861.

[†] Journ. für Prakt. Chem., 86, 454. 1862. Zeit. Anal. Chem., 1, 518.

¹ Ann. Chem. Pharm., 181, 185. 1876.

CÆSIUM. 91

crystallization of their alums, and obtained salts of each spectroscopically pure. The nitric acid employed was tested for chlorine and found to be free from that impurity, and the weights used were especially verified. In two of his analyses of RbCl the AgCl was handled by the ordinary process of filtration. In the other two it was washed by decantation, dried, and weighed in a glass dish. The usual ratio is appended in the third column:

```
1.4055 grm. RbCl gave 1.6665 grm. AgCl. 84.338
1.8096 " 2.1461 " 84.320
2.2473 " 2.665 " 84.326
2.273 " 2.6946 " 84.354

Mean, 84.3345, ± .0051
```

Combining the three series, we get the following result:

Hence Rb = 85.251, $\pm .018$. If O = 16, Rb = 85.529.

CÆSIUM.

The atomic weight of cæsium, like that of rubidium, has been determined from the analysis of the chloride. The earliest determination, by Bunsen,* was incorrect, because of impurity in the material employed.

In 1863 Johnson and Allen published their results.† Their material was extracted from the lepidolite of Hebron, Maine, and the cæsium was separated from the rubidium as bitartrate. From the pure cæsium bitartrate cæsium chloride was prepared, and in this the chlorine was estimated as

^{*} Zeit. Anal. Chem., 1, 137.

[†] Amer. Journ. Sci. and Arts, (2,) 35, 94.

silver chloride by the usual gravimetric method. Reducing their results to the convenient standard adopted in preceding chapters, we have, in a third column, the quantities of CsCl equivalent to 100 parts of AgCl:

```
      1.8371 grm. CsCl gave
      1.5634 grm. AgCl.
      117.507

      2.1295 " 1.8111 " 117.580

      2.7018 " 2.2992 " 117.511

      1.56165 " 1.3302 " 117.399

      Mean, 117.499, ± .025
```

Shortly after the results of Johnson and Allen appeared a new series of estimations was published by Bunsen.* His cæsium chloride was purified by repeated crystallizations of the chloroplatinate, and the ordinary gravimetric process was employed. The following results represent, respectively, material thrice, four times, and five times purified:

```
1.3835 grm. CsCl gave 1.1781 grm. AgCl. Ratio, 117.435
1.3682 " 1.1644 " 117.503
1.2478 " 1.0623 " 117.462

Mean, 117.467, ± .013
```

Godeffroy's work† was, in its details of manipulation, sufficiently described under rubidium. In three of the experiments upon cæsium the silver chloride was washed by decantation, and in one it was collected upon a filter. The results are subjoined:

```
1.5825 grm. CsCl gave 1.351 grm. AgCl. Ratio, 117.135
1.3487 " 1.1501 " 117.265
1.1880 " 1.0141 " 117.148
1.2309 " 1.051 " 117.107

Mean, 117.164, ± .023
```

We may now combine the three series to form a general mean:

^{*} Poggend. Annal., 119, 1. 1863. † Ann. Chem. Pharm., 181, 185. 1876.

Hence Cs = 132.583, $\pm .024$; or, if O = 16, Cs = 132.918.

THALLIUM.

The atomic weight of this interesting metal has been fixed by the researches of Lamy, Werther, Hebberling, and Crookes. Lamy and Hebberling investigated the chloride and sulphate; Werther studied the iodide; Crooke's experiments involved the synthesis of the nitrate. The last mentioned work was so thorough and admirable that the other researches are included here only for the sake of historical completeness.

Lamy* gives the results of one analysis of thallium sulphate and three of thallium chloride. 3.423 grammes Tl.SO₄ gave 1.578 grm. BaSO₄; whence 100 parts of the latter are equivalent to 216.920 of the former. In the thallium chloride the chlorine was estimated as silver chloride. The following results were obtained. In the third column I give the amount of TlCl proportional to 100 parts of AgCl:

```
3.912 grm. TlCl gave 2.346 grm. AgCl. 166.752
3.000 " 1.8015 " 166.528
3.912 " 2.336 " 167.466

Mean, 166.915, ± .1905
```

Hebberling's† work resembles that of Lamy. Reducing his weighings to the standards adopted above, we have from his sulphate series, as equivalent to 100 parts of BaSO₄, the amounts of Tl₂SO₄ given in the third column:

^{*} Zeit. Anal. Chem., 2, 211. 1863.

[†] Ann. Chem. Pharm., 134, 11. 1865.

1.4195 grm.	. Tl ₂ SO ₄	gave	.6534	grm. BaSO ₄ .	217.248
1.1924	46		.5507	46	216.5 24
.8560	46	•	-3957	46	216.325
					Mean, 216.699

Including Lamy's single result, as of equal weight, we get a mean of $216.754. \pm .1387.$

From the chloride series we have these results, with the ratio stated as usual:

Lamy's mean was 166.915, $\pm .1905$. Both means combined give a general mean of 166.555, $\pm .0865$.

Werther's* determinations of iodine in thallium iodide were made by two methods. In the first series TlI was decomposed by zinc and potassium hydroxide, and in the filtrate the iodine was estimated as AgI. One hundred parts of AgI correspond to the amounts of TlI given in the last column:

```
.720 grm. TlI gave .51 grm. AgI. 141.176
2.072 " 1.472 " 140.761
.960 " .679 " 141.384
.385 " .273 " 141.026
1.068 " .759 " 140.711

Mean, 141.012, ± .085
```

In the second series the thallium iodide was decomposed by ammonia in presence of silver nitrate, and the resulting AgI was weighed. Expressed according to the foregoing standard the results are as follows:

General mean of both series, 140.648, \pm .016.

^{*} Journ. für Prakt. Chem., 92, 128. 1864.

From the foregoing results three values for the atomic weight of thallium are calculable:

In 1873 Crookes,* the discoverer of thallium, published his final determination of its atomic weight. His method was to effect the synthesis of thallium nitrate from weighed quantities from absolutely pure thallium. No precaution necessary to ensure purity of materials was neglected; the balances were constructed especially for the research; the weights were accurately tested and all their errors ascertained; weighings were made partly in air and partly in vacuo, but all were reduced to absolute standards; and unusually large quantities of thallium were employed in each experiment. In short, no effort was spared to attain as nearly as possible absolute precision of results. The details of the investigation are too voluminous, however, to be cited here; the reader who wishes to become familiar with them must consult the original memoir. Suffice it to say that the research is a model which other chemists will do well to copy.

The results of ten experiments by Professor Crookes may be stated as follows. In a final column we may state the quantity of nitrate producible from 100 parts of thallium. The weights given are in grains:

Thallium.	$TlNO_8 + Glass.$	Glass Vessel.	Ratio.
497-9 729 95	1121.851852	472.557319	130.3875
293.193507	1111.387014	729.082713	130.3930
288. 562777	971.214142	594.949719	130.3926
324.963740	1142.569408	718.849078	130.3900
183.790232	1005.779897	766. 133831	130.3912
190.842532	997.334615	748.491271	130.3920
195.544324	1022.176679	767.203451	130.3915
201.816245	1013.480135	750.332401	130.3897
295.683523	1153.947672	768.403621	130.3908
299.203036	1159.870052	769.734201	130.3917
		Mean.	130,3010. + .00034

^{*} Philosophical Transactions, 1873, p. 277.

Hence, using the atomic weights and probable errors previously found for N and O, Tl = 203.715, $\pm .0365$. If O = 16, Tl = 204.183.

Crookes himself, using 61.889 as the molecular weight of the group NO₃, gets the value Tl = 203.642; the lowest value in the series being 203.628, and the highest 203.666; an extreme variation of 0.038. This is extraordinary accuracy for so high an atomic weight, at least as far as Crookes' work is concerned. But its value depends in reality upon the accuracy of other chemists in fixing the atomic weights of N and O; a slight variation in either of the latter constants producing a large variation here. What Crookes really has done has been to fix with almost absolute certainty the ratio between Tl and NO₃. If the latter group should have the molecular weight 62, in accordance with Prout's hypothesis, then Tl = 204.008. In other words, the ratio thus fixed by Crookes is almost exactly represented by two whole numbers, and supports Prout's hypothesis in a very decided way. Crookes himself seems to have overlooked this fact, for he regards his results as militating against the hypothesis in question.

GLUCINUM.

The atomic weight of glucinum is at present much in doubt; our knowledge of it depending upon the unsettled question whether the oxide is GlO or Gl₂O₃. The formula GlO agrees with Mendelejeff's law, and is advocated by Reynolds,* Lothar Meyer,† and Brauner.‡ The symbol Gl₂O₃, on the other hand, is favored by Nilson and Pettersson,|| and by Humpidge.§ Humpidge, Meyer, and Brauner

^{*} Phil. Mag., (5,) 3, 38. 1877. Chem. News, 42, 273. 1880.

[†] Ber. der Deutsch. Chem. Gesell., 13, 1780. 1880. Also, 11, 576. 1879.

[†] Phil. Mag., (5,) 11. Jan., 1881.

^{||} Berichte, 11, 381 and 906. 1879. Also, 13, 2035. 1880.

[¿]Chem. News, 42, 261. 1880.

offer only theoretical discussions of the subject; Reynolds and Nilson and Pettersson have determined the specific heat of the metal, but give opposed results. In the following calculations the simpler formula will be assumed, not as a finality, but because of its accordance with the system of Mendelejeff.

The data from which we are to calculate the atomic weight of glucinum have been determined by Awdejew, Weeren, Klatzo, Debray, and Nilson and Pettersson. Berzelius'* single experiment on the sulphate may be left out of account.

Awdejew,† whose determination was the earliest of any value, analyzed the sulphate. The sulphuric acid was thrown down as barium sulphate; and in the filtrate, from which the excess of barium had been first removed, the glucina was precipitated by ammonia. The figures which Awdejew publishes represent the ratio between SO₃ and GlO, but not absolute weights. As, however, his calculations were made with SO₃ = 501.165, and Ba probably = 855.29, we may add a third column showing how much BaSO₄ is proportional to 100 parts of GlO:

GlO.	Ratio.
1406	921.242
1420	927.304
2480	915.903
4065	920.814
	1406 1420 2480

Mean, 921.316, ± 1.577

The same method was followed by Weeren and by Klatzo, except that Weeren used ammonium sulphide instead of ammonia for the precipitation of the glucina. Weeren‡ gives the following weights of GlO and BaSO₄. The ratio is given in a third column, just as with the figures by Awdejew:

^{*} Poggend. Annal., 8, 1. † " 56, 106. 1842. † " 92, 124. 1854.

GlO.	$BaSO_4$.	<i>Ratio.</i> 9 27. 031	
.3163 grm.	2.9332 grm.		
.2872 "	2.6377 "	918.419	
.2954 "	2.7342 "	925.592	
.5284 "	4.8823 "	902.946	

Mean, 918.497, \pm 3.624

Klatzo's* figures are as follows, with the third column added by the writer:

GlO.	$BaSO_4$.	Ratio.	
.2339 grm.	2.1520 grm.	920.052	
.1910 "	1.7556 "	919.162	
.2673 "	2.4872 "	930.490	
.3585 "	3.3115 "	923.710	
.2800 "	2.5842 "	922.989	

Mean, 923.281, ± 1.346

Combining these series into a general mean, we get the subjoined result:

Awdejew Weeren Klatzo	$918.497, \pm 3.624$
General mean	922.164. + 0.985

Hence GlO = 25.224, $\pm .269$.

Debray† analyzed a double oxalate of glucinum and ammonium, $Gl(NH_4)_2C_4O_8$. In this the glucina was estimated by calcination, after first converting the salt into nitrate. The following percentages were found:

The carbon was estimated by an organic combustion. I give the weights, and put in a third column the percentages of CO, thus obtained:

^{*} Zeitschrift für Anal. Chem., 8, 523. 1869.

[†] Ann. de Chim. et de Phys., (3,) 44, 37. 1855.

Salt.	CO_2 .	Per cent. CO2.
.600 grm.	.477 grm.	79.500
.603 "	.478 "	79.270
.600 "	. •477 "	79.500
		Mean, 79.423, \pm .052

Calculating the ratio between CO, and GlO, we have for the molecular weight of the latter, GlO = 25.220, \pm .180. The agreement between this result and the one previously deduced from the sulphate is certainly very striking.

Last of all and best of all we come to the determinations recently published by Nilson and Pettersson.* These chemists sought to use the sublimed chloride of glucinum, but found it to contain traces of lime derived from a glass tube. They finally resorted to the sulphate as the most available salt for their purposes. This, which they write $Gl_2(SO_4)_3$ 12H₂O, and which we formulate as $GlSO_4$.4H₂O, yields pure glucina upon strong ignition. The subjoined percentages of glucina were thus obtained:

14.171 14.169 14.160 14.176 Mean, 14.169, ± .0023

Hence GlO = 25.048, and Gl = 9.085, \pm .0055. If O = 16, Gl = 9.106. If SO₈ = 80, then Gl = 9.096.

If the oxide is Gl₂O₃, then the value Gl = 9.085, $\pm .0055$ becomes Gl = 13.628, $\pm .0082$.

It would be easy enough to combine this value for Gl with those derived from the experiments of the investigators previously cited, but it is hardly worth while. All the other estimations have such high probable errors that they would practically vanish from the general mean. Their influence would hardly extend to the third decimal place, and they may therefore be neglected.

^{*} Compt. Rend., 91, 168. 1880.

MAGNESIUM.

There is perhaps no common metal of which the atomic weight has been subjected to closer scrutiny than that of magnesium. The value is low, and its determination should, therefore, be relatively free from many of the ordinary sources of error; it is extensively applied in chemical analysis, and ought consequently to be accurately ascertained. Strange discrepancies, however, exist between the results obtained by different investigators; so that the generally accepted figure cannot be regarded as absolutely free from doubt.

The determinations of Berzelius* and other early chemists need not be here considered. Nor does the estimation made by Macdonnell† deserve more than a passing mention. He puts the atomic weight of magnesium at 23.9, but gives no details concerning his method of determination. The researches which we have to consider are those of Scheerer, Svanberg and Nordenfeldt, Jacquelain, Bahr, Marchand and Scheerer, and Dumas.

Scheerer's method of investigation was exceedingly simple.‡ He merely estimated the sulphuric acid in anhydrous magnesium sulphate, employing the usual process of precipitation as barium sulphate. He gives no weighings, but reports the percentages of SO_3 thus found. In his calculations, O = 100, $SO_3 = 500.75$, and BaO = 955.29. It is easy, therefore, to recalculate the figures which he gives, so as to establish what his method really represents, viz., the ratio between the sulphates of barium and magnesium.

Thus revised, his four analyses show that 100 parts of MgSO₄ yield the following quantities of BaSO₄:

^{*} Lehrbuch, 5 Aufl., Bd. 3, s. 1227.

[†] British Association Report, 1852, part 2, p. 36.

[†] Poggend. Annal., 69, 535. 1846.

	Per cent. SO ₈ .		
193.575	66.573		
193.677	66.608		
193.767	66.639		
193.631	66.592	- ₁	· FFE
			· ••• 🗗 🎜 🖟

Mean, 193.6625, ± .0274

Hence, using the atomic weights deduced in previous chapters for Ba, S, and O, Mg = 24.544, $\pm .0311$. In a subsequent note* Scheerer shows that the barium sulphate of the foregoing experiments carried down with it magnesium salts in such quantity as to make the atomic weight of magnesium 0.39 too low. Corrected, Mg becomes = 24.545.

The work of Bahr, of Jacquelain, and in part that of Svanberg and Nordenfeldt, also relates to the composition of magnesium sulphate. Jacquelain's experiments were as follows.† Dry magnesium sulphate was prepared by mixing the ordinary hydrous salt to a paste with sulphuric acid, and calcining the mass in a platinum crucible over a spirit lamp to constant weight and complete neutrality of reaction. This dry sulphate was weighed and intensely ignited three successive times. The weight of the residual MgO having been determined, it was moistened with sulphuric acid and recalcined over a spirit lamp, thus reproducing the original weight of MgSO₄. Jacquelain's weighings for these two experiments show that 100 parts of MgO correspond to the quantities of MgSO₄ given in the last column:

Jacquelain also made one estimation of sulphuric acid in the foregoing sulphate as BaSO₄. His result, (1.464 grm. MgSO₄ = 2.838 grm. BaSO₄,) reduced to the standard adopted in dealing with Scheerer's experiments, give for 100 parts of MgSO₄, 193.852 BaSO₄. If this figure be given equal weight with a single experiment in Scheerer's series,

^{*} Poggend. Annal., 70, 407.

[†] Ann. d. Chim. et Phys., 3 serie, 32, 202.

and combined with the latter, the mean will be 193.700, \pm .0331. From this the atomic weight of magnesium becomes 24.244, \pm .033. This again, corrected according to Scheerer for the magnesium salts carried down by the barium sulphate, becomes 0.39 higher, or Mg = 24.283. Of course this correction, determined by Scheerer for a single experiment, can only be a rough approximation in a mean like the foregoing. It is better than no correction at all, the character of the error involved being known.

Bahr's* work resembles in part that of Jacquelain. This chemist converted pure magnesium oxide into sulphate, and from the increase in weight determined the composition of the latter salt. From his weighings 100 parts of MgO equal the amounts of MgSO₄ given in the third column:

About four years previous to the investigations of Bahr the paper of Svanberg and Nordenfeldt† appeared. These chemists started with the oxalate of magnesium, which was dried at a temperature of from 100° to 105° until it no longer lost weight. The salt then contained two molecules of water, and upon strong ignition it left a residue of MgO. The percentage of MgO in the oxalate comes out as follows:

```
7.2634 grm. oxalate gave 1.9872 grm. oxide. 27.359 per cent.
6.3795 " 1.7464 " 27.375 "
6.3653 " 1.7418 " 27.364 "
6.2216 " 1.7027 " 27.368 "
Mean, 27.3665, ± .0023
```

In three of these experiments the MgO was treated with H₂SO₄, and converted, as by Jacquelain and by Bahr in their later researches, into MgSO₄. One hundred parts of MgO gave of MgSO₄ as follows:

^{*} Journ. für Prakt. Chem., 56, 310. 1852.

[†] Journ. für Prakt. Chem., 45, 473. 1848.

```
1.9872 grm. MgO gave 5.8995 grm. MgSO<sub>4</sub>. 296.875
1.7464 " 5.1783 " 296.513
1.7418 " 5.1666 " 296.624

Mean, 296.671, ± .072
```

We have now for this ratio between MgO and MgSO₄ three series; not at all concordant. We may combine them, assigning to each of Jacquelain's two results a weight corresponding to one of Bahr's:

```
Jacquelain ______ 297.968, ± .0999

Bahr _____ 296.200, ± .0815

Svanberg and Nordenfeldt ____ 296.671, ± .072

General mean ____ 296.806, ± .0475
```

In 1850 the elaborate investigations of Marchand and Scheerer* appeared. These chemists undertook to determine the composition of some natural magnesites, and, by applying corrections for impurities, to deduce from their results the sought for atomic weight. The magnesite chosen for the investigation was, first, a yellow, transparent variety from Snarum; second, a white opaque mineral from the same locality; and, third, a very pure quality from Frankenstein. In each case the impurities were carefully determined; but only a part of the details need be cited here. Silica was of course easily corrected for by simple subtraction from the sum of all of the constituents; but iron and calcium, when found, having been present in the mineral as carbonates, required the assignment to them of a portion of the carbonic acid. In the atomic weight determinations the mineral was first dried at 300°. The loss in weight upon ignition was then carbon dioxide. It was found, however, that even here a correction was necessary. Magnesite, upon drying at 300°, loses a trace of CO₂, and still retains a little water; on the other hand, a minute quantity of CO2 remains even after ignition. The CO₂ expelled at 300° amounted in one experiment to .054 per cent.; that retained after calcination to .055 per cent. Both errors tend in the

^{*} Journ. für Prakt. Chem., 50, 385.

same direction, and increase the apparent percentage of MgO in the magnesite. On the yellow mineral from Snarum the crude results are as follows, giving percentages of MgO, FeO, and CO₂ after eliminating silica:

CO_2 .		MgO.	FeO.
51.8958		47.3278	.7764
51.8798 -		47-3393	.7809
51.8734		47.3154	.8112
51.8875		47.3372	· 7 753
	Mean.	47.3299 ± .0037	

After applying corrections for loss and retention of CO,, as previously indicated, the mean results of the foregoing series become—

CO_2 .	MgO.	FeO.
51.9931	47.2743	.7860

The ratio between the MgO and the CO₂, after correcting for the iron, will be considered further on.

Of the white magnesite from Snarum but a single analysis was made, which, for present purposes may be ignored. Concerning the Frankenstein mineral three series of analyses were executed. In the first series the following results were obtained:

```
8.996 grm. CO_2 = 8.2245 grm. MgO. 47.760 per cent. MgO. 7.960 " 7.2775 " 47.761 " 9.3265 " 8.529 " 47.767 " 47.775 " 47.775 " 47.775 " 47.775 " 47.775 " 47.775 "
```

This mean, corrected for loss of CO, in drying, becomes 47.681. I give series second with corrections applied:

6.8195	grm. MgCO ₃	gave 3.2500	grm. MgC). 47.658 ₁	per cent.
11.3061	66	5.3849	46	47.628	66
9.7375	66	4.635	46	47.599	46
12.3887	44	5.9033	"	47.650	44
32.4148	44	15.453	44	47.674	44
38.8912	44	18.5366	66	47.663	46
26.5223	16	12.6445	46	47.675	66
				Mean, 47.650,	± .0069

The third series was made upon very pure material, so that the corrections, although applied, were less influential. The results were as follows:

```
4.2913 grm. MgCO<sub>8</sub> gave 2.0436 grm. MgO. 47.622 per cent.
27.8286 " 13.2539 " 47.627 "
14.6192 " 6.9692 " 47.672 "
18.3085 " 8.7237 " 47.648 "

Mean, 47.642, ± .0077
```

In a supplementary paper* by Scheerer, it was shown that an important correction to the foregoing data had been overlooked. Scheerer, re-examining the magnesites in question, discovered in them traces of lime, which had escaped notice in the original analyses. With this correction the two magnesites in question exhibit the following mean composition:

	Snarum.	Frankenstein.
CO,	52.131	52.338
MgO	46.663	47.437
CaO	.430	.225
FeO	.776	•••
	100.000	100.000

Correcting for lime and iron, by assigning each its share of CO_2 , the Snarum magnesite gives as the true percentage of magnesia in pure magnesium carbonate, the figure 47.624. To this, without serious mistake, we may assign the weight indicated by the probable error, \pm .0037; the quantity previously deduced from the percentages of MgO given in the uncorrected analyses.

From the Frankenstein mineral, similarly corrected, the final mean percentage of MgO in MgCO₃ becomes 47.628. This, however, represents three series of analyses, whose combined probable errors may be properly assigned to it. The combination is as follows:

^{*} Ann. d. Chem. und Pharm., 110, 240.

± .0022 ± .0069 ± .0077

Result, ± .0020, probable error of the general mean.

We may now combine the results obtained from both magnesites:

```
Snarum mineral ___ Per cent. MgO, 47.624, ± .0037
Frankenstein mineral __ " 47.628, ± .0020

General mean __ " 47.627, ± .0018
```

The last investigation upon the atomic weight of magnesium which we have to consider is that of Dumas.* Pure magnesium chloride was placed in a boat of platinum, and ignited in a stream of dry hydrochloric acid gas. The excess of the latter having been expelled by a current of dry carbon dioxide, the platinum boat, still warm, was placed in a closed vessel and weighed therein. After weighing, the chloride was dissolved and titrated in the usual manner with a solution containing a known quantity of pure silver. The weighings which Dumas reports give, as proportional to 100 parts of silver, the quantities of MgCl, stated in the third column:

2.203 gm	n. MgCl ₂	= 4.964 g	rm. Ag.	44.380
2.5215	46	5.678	**	44.408
2.363	66	5.325	66	44.376
3-994	66	9.012	66	44.319
2.578	46	5.834	66	44.189
2.872	46	6.502	66	44.171
2.080	66	4.710	44	44.161
2.214	**	5.002	66	44.262
2.086	66	4.722	"	44.176
1.688	66	3.823	66	44.154
1.342	66	3.031	46	44.276

There are now before us the following ratios, from which to deduce the sought-for atomic weight:

Mean, 44.261, ± .020

^{*} Ann. Chem. Pharm., 113, 33. 1860.

```
(1.) MgSO_4: BaSO_4:: 100: 193.700, \pm .0331
```

- (2.) $MgO: MgSO_4:: 100: 296.806, \pm .0475$
- (3.) Per cent. of MgO in oxalate, 27.3665, \pm .0023
- (4.) Per cent. of MgO in carbonate, 47.627, \pm .0018
- (5.) Ag: $MgCl_2$:: 100: 44.261, \pm .020

From these we find three values for the molecular weight of MgO:

We have also three values for the atomic weight of magnesium:

Or, if O = 16, Mg becomes = 24.159.

In this general mean all the determinations are included, good or bad. Dumas' result is unquestionably wrong; the error, probably, being due to the presence of oxychloride in the MgCl, which was used. It is doubtful whether any precautions could have eliminated that error. If we take only Marchand and Scheerer's work on magnesium carbonate as having positive value, we shall get from their analyses the following result, viz: Mg = 23.959, \pm .0046. Or, if O = 16, this becomes 24.014. The atomic weight of magnesium, therefore, varies from the whole number 24, only within the ordinary limits of experimental error.

ZINC.

The several determinations of the atomic weight of zinc are by no means closely concordant. The results obtained by Gay-Lussac* and Berzelius† were undoubtedly too low, and may be disregarded here. We need consider only the work done by Jacquelain, Favre, and Axel Erdmann.

In 1842 Jacquelain published the results of his investigations upon this important constant.‡ In two experiments a weighed quantity of zinc was converted into nitrate, and that by ignition in a platinum crucible was reduced to oxide. In two other experiments sulphuric acid took the place of nitric. As the zinc contained small quantities of lead and iron, these were estimated, and the necessary corrections applied. From the weights of metal and oxide given by Jacquelain the percentages have been calculated:

Nitric Series.

9.917 gm	n. Zn gave	12.3138 grm. ZnO.	80.536 per	r cent. Zn.
9.809	44	12.1800 "	80.534	46
		Sulphuric Series.		
2.398	66	2.978 grm. ZnO.	80.524	44
3.197	44	3.968 "	80.570	44

Mean of all four, 80.541, ± .007

Hence $Zn = 66.072, \pm .028$.

The method adopted by Axel Erdmann || is essentially the same as that of Jacquelain, but varies from the latter in certain important details. First, pure zinc oxide was prepared, ignited in a covered crucible with sugar, and then, to complete the reduction, ignited in a porcelain tube in a current of hydrogen. The pure zinc thus obtained was converted into oxide by means of treatment with nitric acid and sub-

^{*} Mémoire d'Arceuil, 2, 174.

[†] Gilb. Annal., 37, 460.

[‡] Compt. Rend., 14, 636.

^{||} Poggend. Annal., 62, 611. Berz. Lehrb., 3, 1219.

109

sequent ignition in a porcelain crucible. Erdmann's figures give us the following percentages of metal in the oxide:

Hence $Zn = 64.9045, \pm .019$.

If we combine the results of Jacquelain with those of Erdmann, we get a mean percentage of zinc, 80.324, $\pm .0032$; and an atomic weight of Zn = 65.168, $\pm .018$. The reason for the discordance between the two experimenters will be considered further along.

Favre * employed two methods of investigation. First, zinc was dissolved in sulphuric acid, the hydrogen evolved was burned, and the weight of water thus formed was determined. To his weighings I append the ratio between metallic zinc and 100 parts of water:

25.389 grm. Zn gave 6.928 grm. H₂O. 366.469
30.369 " 8.297 " 366.024
31.776 " 8.671 " 366.463
Mean, 366.319,
$$\pm$$
 .088

The second method adopted by Favre was to burn pure zinc oxalate, and to weigh the oxide and carbonic acid thus produced. From the ratio between these two sets of weights the atomic weight of zinc is easily deducible. From Favre's weighings, if CO₂ = 100, ZnO will be as given in the third column below:

7.796 grm.
$$ZnO = 8.365$$
 grm. CO_2 .

7.342 " 7.883 " 93.137

5.2065 " 5.588 " 93.173

Mean, 93.169, \pm .012

Hence $Zn = 65.8395$, \pm .022.

^{*} Ann. Chim. Phys., (3,) 10, 163. 1844.

A fourth combustion of the oxalate is omitted from the above series, having been rejected by Favre himself. In this the oxide formed was contaminated by traces of sulphide.

The four values for zinc now before us are so discordant that a combination of them after the usual method can have only a trifling significance. The following is the result thus obtained:

```
From Jacquelain's figures ___Zn = 66.072, \pm .028

From Favre's water series__ " = 65.803, \pm .020

From Favre's oxalate series__ " = 65.8395, \pm .022

From Erdmann's figures___ " = 64.9045, \pm .019

General mean____ " = 65.557, \pm .011
```

It will be seen that three of these values agree tolerably well, placing the atomic weight of zinc in the neighborhood of 66, while the other is, in round numbers, about a unit lower. This lower figure, however, has the smallest probable error, and it will be found also, upon careful consideration, that it is less likely than the others to be vitiated by experimental inaccuracies. Both chemically and mathematically it is the best.

Upon comparing Erdmann's results with those of Jacquelain two points are worth noticing: first, Erdmann worked with purer material than Jacquelain, although the latter applied corrections for the impurities which he knew were present; secondly, Erdmann calcined his zinc nitrate in a porcelain crucible, while Jacquelain used platinum. In the latter case it has been shown that portions of zinc may become reduced and alloy themselves with the platinum of the crucible. Hence a lower weight of oxide from a given quantity of zinc, a higher percentage of metal, and an increased atomic weight. This source of constant error has undoubtedly affected Jacquelain's experiments, and vitiated his results. In Erdmann's work no such errors seem to be present.

Over Favre's experiments Erdmann's have the important merit of simplicity. In the latter it is difficult to detect sources of error; in the former it is easy. In Favre's water series it was essential that the hydrogen should first be thoroughly dried before combustion, and then that every trace of water formed should be collected. A trivial loss of hydrogen or of water would tend to increase the apparent atomic weight of zinc.

In the combustion of the zinc oxalate equally great difficulties are encountered. Here a variety of errors are possible, such as are due, for example, to impurity of material, to imperfect drying of the carbon dioxide, and to incomplete collection of the latter. It may not be easy to prove that such errors actually did creep into Favre's work, and yet their possibility hinders us from absolutely accepting his results.

All things considered, then, Erdmann's determination of the atomic weight of zinc is the one most entitled to credit, and must be taken for the present in lieu of the general mean deduced from all four of the values. This determination, Zn = 64.9045, $\pm .019$, becomes, if O = 16, 65.054.

CADMIUM.

The earliest determination of the atomic weight of this metal was by Stromeyer, who found that 100 parts of cadmium united with 14.352 of oxygen.* With our value for the atomic weight of oxygen these figures make Cd = 111.227. This result has now only a historical interest.

The more modern estimates of the atomic weight of cadmium are four in number, by v. Hauer, Lenssen, Dumas, and Huntington. Of these that by v. Hauer † comes first in chronological order. He heated pure anhydrous cadmium sulphate in a stream of dry hydrogen sulphide, and weighed the cadmium sulphide thus obtained. His results

^{*} See Berz. Lehrbuch, 5th Ed., 3, 1219.

[†] Journ. sur Prakt. Chem., 72, 350. 1857.

were as follows, with the percentage of CdS in CdSO, therefrom deduced:

7.7650 grm	. CdSO ₄ gav	ve 5.3741 g	rm. CdS.	69.209	per cent.
6.6086	64	4.5746	66	69.222	44
7.3821	66	5.1117	66	69.245	46
6.8377	66	4.7336	46	69.228	66
8.1956	66	5.6736	44	69.227	66
7.6039	66	5.2634	46	69. 22 0	"
7.1415	66	4.9431	"	69.217	66
5.8245	66	4.0335	"	69.251	**
6.8462	44	4.7415	66	69.257	44

Mean, 69.231, \pm .0042

Lenssen* worked upon pure cadmium oxalate, handling. however, only small quantities of material. This salt, upon ignition, leaves the following percentages of oxide:

```
.5128 grm. oxalate gave .3281 grm. CdO. 63.982 per cent.
.6552 " .4193 " 63.996 "
.4017 " .2573 " 64.053 "

Mean, 64.010, ± .014
```

Dumas dissolved pure cadmium in hydrochloric acid, evaporated the solution to dryness, and fused the residue in hydrochloric acid gas. The cadmium chloride thus obtained was dissolved in water and titrated with a solution of silver after the usual manner. From Dumas' weighings I calculate the ratio between CdCl₂ and 100 parts of silver:

```
2.369 \text{ grm. } CdCl_2 = 2.791 \text{ grm. } Ag.
                                                     84.880
                                                     84.892
4.540
                       5.348
                       7.260
                                                     85.803
6.177
                       2.841
                                                     84.618
2.404
                       4.166
3.5325
                                                     84.794
4.042
                       4.767
                                                     84.791
                                              Mean, 84.843, ± .026
```

Latest of all comes Huntington's twork, done under the direction of Professor J. P. Cooke. Bromide of cadmium

^{*} Journ. für Prakt. Chem., 79, 281. 1860.

[†] Ann. Chem. Pharm., 113, 27. 1860.

[†] Proc. Amer. Acad., 1881.

was prepared by dissolving the carbonate in hydrobromic acid, and the product, dried at 200°, was purified by sublimation in a porcelain tube. Upon the compound thus obtained two series of experiments were made.

In one series the bromide was dissolved in water, and a quantity of silver not quite sufficient for complete precipitation of the bromine was then added in nitric acid solution. After the precipitate had settled, the supernatant liquid was titrated with a standard solution of silver containing one gramme to the litre. The precipitate was washed by decantation, collected by reverse filtration, and weighed. To the weighings I append the ratio between CdBr, and 100 parts of silver bromide:

1.5592 gm	n. CdBr, g	ave 2.1529 gr	rm. AgBr.	Ratio, 72.423
* 3. 7 456	66	5.1724	66	72.415
2.4267	**	3.3511	66	72.415
* 3.6645	66	5.0590	"	72.435
* 3.7679	44	5.2016	66	72.437
2.7938	44	3.8583	66	72.410
¥ 1.9225	66	2.6552	66	72.405
3-4473	46	4-7593	66	72.433

Mean, 72.4216, \pm .0028

The second series was like the first, except that the weight of silver needed to effect precipitation was noted, instead of the weight of silver bromide formed. In the experiments marked with an asterisk, both the amount of silver required and the amount of silver bromide thrown down were determined in one set of weighings. The third column gives the CdBr, proportional to 100 parts of silver:

* 3.7456	grm. CdBr ₂ =	2.9715	grm. Ag.	126.051
5.0270	46	3.9874	66	126.072
* 3.6645	44	2.9073	66	126.045
* 3.7679	64	2.9888	66	126.067
* 1.9225	66	1.5248	46	126.082
2.9101	66	2.3079	44	126.093
3.6510	e i	2.8951	66	126.110
3.9782	66	3.1551	66	126.088

Mean, 126.076, ± .0052

From the first seriesCdBr, =
$$271.498$$
, $\pm .032$
From the second series ... " = 271.505 , $\pm .027$
General mean ... " = 271.502 , $\pm .0215$

Hence Cd = 111.966, $\pm .043$.

According to Huntington's own calculations these experiments fix the ratio between silver, bromine, and cadmium as Ag: Br: Cd::108:80:112.31. This result militates strongly against Prout's hypothesis.

Upon combining all the determinations we get the following result:

Or, if O = 16, then Cd = 112.092.

It will be seen that Dumas and Huntington's determinations, both made with haloid salts of cadmium, agree with wonderful closeness, and so confirm each other. On the other hand, v. Hauer's data give a value for the atomic weight of cadmium which is much lower. Apparently, v. Hauer's method was good, and the reason for the discrepancy remains to be discovered. Until it is ascertained I prefer to use the above mean value for Cd, rather than to adopt one investigation and reject the others.

MERCURY.

In dealing with the atomic weight of mercury we may reject the early determinations by Sefström* and a large part of the work done by Turner.† The latter chemist, in addition to the data which will be cited below, gives figures

^{*} Sefström. Berz. Lehrb., 5th Ed., 3, 1215. Work done in 1812.

[†] Phil. Trans., 1833, 531-535.

to represent the percentage composition of both the chlorides of mercury; but these results are neither reliable nor in proper shape to be used.

First in order we may consider the percentage composition of mercuric oxide, as established by Turner and by Erdmann and Marchand. In both investigations the oxide was decomposed by heat, and the mercury was accurately weighed. Gold leaf served to collect the last traces of mercurial vapor.

Turner gives four estimations.* Two represent oxide obtained by the ignition of the nitrate, and two are from commercial oxide. In the first two the oxide still contained traces of nitrate, but hardly in weighable proportions. A comparison of the figures from this source with the others is sufficiently conclusive on this point. The third column represents the percentage of mercury in HgO:

```
      144.805 grains Hg = 11.54 grains O.
      92.619 per cent.

      125.980 " 10.08 " 92.592 "

      173.561 " 13.82 " 92.625 "

      114.294 " 9.101 " 92.620 "

      Mean, 92.614, \pm.0050
```

In the experiments of Erdmann and Marchand† every precaution was taken to ensure accuracy. Their weighings, reduced to a vacuum standard, give the subjoined percentages:

Combining, we have:

```
Turner ______ 92.614, ± .0050

Erdmann and Marchand _____ 92.5996, ± .0015

General mean _____ 92.601, ± .0014
```

^{*} Phil. Trans., 1833, 531-535.

[†] Journ. für Prakt. Chem.; 31, 395. 1844.

With a view to establishing the atomic weight of sulphur Erdmann and Marchand also made a series of analyses of pure mercuric sulphide. These data are now best available for discussion under mercury. The sulphide was mixed with pure copper and ignited; mercury distilling over and copper sulphide remaining behind. Gold leaf was used to retain traces of mercurial vapor, and the weighings were reduced to vacuum:

```
34.3568 grm. HgS gave 29.6207 grm. Hg. 86.215 per cent. Hg. 24.8278 " 21.40295 " 86.206 " 37.2177 " 32.08416 " 86.207 " 86.223 " Mean, 86.2127, ± .0027
```

For the percentage of mercury in mercuric chloride we have data by Turner, Millon, and Svanberg. Turner,* in addition to some precipitations of mercuric chloride by silver nitrate, gives two experiments in which the compound was decomposed by pure stannous chloride, and the mercury thus set free was collected and weighed. The results were as follows:

```
44.782 grains Hg == 15.90 grains Cl.

73.798 per cent.

73.798 per cent.

73.784 "

Mean, 73.791, ± .005
```

Millon† purified mercuric chloride by solution in ether and sublimation, and then subjected it to distillation with lime. The mercury was collected as in Erdmann and Marchand's experiments. Percentages of metal as follows:

```
73.87
73.81
73.83
73.87
Mean, 73.845, ± .010
```

Svanberg,‡ following the general method of Erdmann

^{*} Phil. Trans., 1833, 531-525. † Ann. Chim. Phys., (3,) 18, 345 1846.

[‡] Journ. für Prakt. Chem., 45, 472. 1848.

and Marchand, made three distillations of mercuric chloride with lime, and got the following results:

```
12.048 grm. HgCl, gave 8.889 grm. Hg. 73.780 per cent.
12.529 " 9.2456 " 73.794 "
12.6491 " 9.3363 " 73.810 "

Mean, 73.795, ± .006
```

Combining these series we have:

Svanberg 73.795, \pm .00	Υ .
Millon 73.845, \pm .01	
Turner 73.791, ± .00	_

In this mean Turner's figures undoubtedly receive undue weight, for, on experimental grounds, they are probably inferior to both of the other series. It is better, however, that the general mean should remain as it is, than that I should deal arbitrarily with any of the data.

We now have three figures to calculate from:

These give us three values for the atomic weight of mercury and a general mean as follows:

If O = 16, then this becomes 200.171.

CHROMIUM.

Concerning the atomic weight of chromium there has been much discussion, and many experimenters have sought to establish the true value. The earliest work upon it having any importance was that of Berzelius,* in 1818 and 1826, which led to results much in excess of the correct figure. His method consisted in precipitating a known weight of lead nitrate with an alkaline chromate and weighing the lead chromate thus produced. The error in his determination arose from the fact that lead chromate, except when thrown down from very dilute solutions, carries with it minute quantities of alkaline salts, and so has its apparent weight notably increased. When dilute solutions are used, a trace of the precipitate remains dissolved, and the weight obtained is too low. In neither case is the method trustworthy.

In 1844 Berzelius' results were first seriously called in question. The figure for chromium deduced from his experiments was somewhat over 56; but Peligot† now showed, by his analyses of chromous acetate and of the chlorides of chromium, that the true number was near 52.5. Unfortunately, Peligot's work, although good, was published with insufficient details to be useful here. For chromous acetate he gives the percentages of carbon and hydrogen, but not the actual weights of salt, carbon dioxide, and water from which they were calculated. His figures vary considerably moreover; enough to show that their mean would carry but little weight when combined with the more explicit data furnished by other chemists.

Jacquelain's work we may omit entirely. He gives an atomic weight for chromium which is notoriously too low, and prints none of the numerical details upon which his result rests. The researches which particularly command our attention are those of Berlin, Moberg, Lefort, Wildenstein, Kessler, and Siewert.

Among the papers upon the atomic weight under consideration that by Berlin is one of the most important. His starting point was normal silver chromate; but in one ex-

^{*} Schweigg. Journ., 22, 53, and Poggend. Annal., 8, 22.

[†] Compt. Rend., 19, 609 and 734; 20, 1187; 21, 74.

[†] Compt. Rend., 24, 679. 1847.

^{||} Journ. für Prakt. Chem., 37, 509, and 38, 149. 1846.

periment the anhydrochromate Ag, Cr, O, was used. These salts, which are easily obtained in a perfectly pure condition, were reduced in a large flask by means of hydrochloric acid and alcohol. The chloride of silver thus formed was washed by decantation, dried, fused, and weighed without transfer. The united washings were supersaturated with ammonia, evaporated to dryness, and the residue treated with hot water. The resulting chromic oxide was then collected upon a filter, dried, ignited, and weighed. The results were as follows:

```
4.6680 grm. Ag<sub>2</sub>CrO<sub>4</sub> gave 4.027 grm. AgCl and 1.0754 grm. Cr<sub>2</sub>O<sub>3</sub>.
3.4568 " 2.983 " .7960 "
2.5060 " 2.1605 " .5770 "
2.1530 " 1.8555 " .4945 "
4.3335 grm. Ag<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub> gave 2.8692 " 1.5300 "
```

From these weighings three values are calculable for the atomic weight of chromium. The three ratios upon which these values depend we will consider separately; taking first that between the chromic oxide and the original silver salt. In the four analyses of the normal chromate the percentages of Cr₂O₃ deducible from Berlin's weighings are as follows:

And from the single experiment with Ag, Cr, O, the percentage of Cr, O, was 35.306.

For the ratio between Ag₂CrO₄ and AgCl, putting the latter at 100, we have for the former:

In the single experiment with anhydrochromate 100 AgCl is formed from 151.035 Ag₂Cr₂O₇.

Finally, for the ratio between AgCl and Cr₂O₃, the five experiments of Berlin give, for 100 parts of the former, the following quantities of the latter:

```
26.705
26.685
26.707
26.650
26.662
Mean, 26.682, ± .0076
```

These results will be discussed in connection with the work of other investigators at the end of this chapter.

In 1848 the researches of Moberg* appeared. His method simply consisted in the ignition of anhydrous chromic sulphate and of ammoniacal chrome alum, and the determination of the amount of chromic oxide thus left as residue. In the sulphate, $Cr_2(SO_4)_8$, the subjoined percentages of Cr_2O_3 were found. The brackets indicate two different samples of material, to which, however, we are justified in ascribing equal value:

Mean, 39.1946, ± .0280

From the alum, (NH₄)₂Cr₂(SO₄)₂.24H₂O, we have these percentages of Cr₂O₃. The first series represents a salt long dried under a bell jar at a temperature of 18°. The crystals taken were clear and transparent, but may possibly have lost traces of water,† which would tend to increase the atomic weight found for chromium. In the second series the salt was carefully dried between folds of filter paper, and

^{*} Journ. für Prakt. Chem., 43, 114.

[†] This objection is suggested by Berlin in a short note upon Lefort's paper. Journ. sur Prakt. Chem., 71, 191.

results were obtained quite near those of Berlin. Both of these series are discussed together, neither having a remarkable value:

```
1.3185 grm. alum gave .213 grm. Cr<sub>2</sub>O<sub>2</sub>.
                                                  16.155 per cent.
                                                  16.151
 .7987
                           .129
1.0185
                           .1645
                                                  16.151
1.0206
                66
                           .1650
                                       46
                                                  16.167
                                                  16.201
 .8765
                           .1420
 .7680
                                                  16.172
                           .1242
1.6720
                                       66
                                                  16.190
                           .2707
                           .0875
                                                  16.174
 .5410
                                                  16.153
                                                             66
                           .1940
1.2010
                           .1620
                "
                                       66
                                                  16.184
1.0010
                                                  16.007
                           .1235
 -7715
                                                  16.012
                                                             "
                           .2200
1.374
```

Mean, 16.143, ± .0125

The determinations made by Lefort* are even less valuable than those by Moberg. This chemist started out from pure barium chromate, which, to thoroughly free it from moisture, had been dried for several hours at 250°. The chromate was dissolved in pure nitric acid, the barium thrown down by sulphuric acid, and the precipitate collected upon a filter, dried, ignited, and weighed in the usual manner. The natural objection to the process is that traces of chromium may be carried down with the sulphate, thus increasing its weight. In fact, Lefort's results are somewhat too high. Calculated from his weighings, 100 parts of BaSO₄ correspond to the amounts of BaCrO₄ given in the third column:

1.2615	grm. BaCrO ₄ gave	1.1555	grm.	BaSO ₄ .	109.174
1.5895	46	1.4580		"	109.019
2.3255	66	2.1340		66	108.974
3.0390	66	2.7855		"	109.101
2.3480	66	2.1590		66	108.754
1.4230	66	1.3060		44	108.708
1.1975	66	1.1005		66	108.814
3.4580	44	3.1690		66	109.119
2.0130		1.8430		"	109.224

^{*} Journ. für Prakt. Chem., 51, 261. 1850.

3.5570 grm	. BaCrO ₄ gave	3.2710 gr	m. BaSO ₄ .	108.744
1.6470	66	1.5060	46	109.363
1.8240	44	1.6725	46	109.058
1.6950	66	1.5560	66	108.933
2.5960	66	2.3870	44	108.756

Mean, 108.9815, ± .0369

Wildenstein,* in 1853, also made barium chromate the basis of his researches. A known weight of pure barium chloride was precipitated by a neutral alkaline chromate, and the precipitate allowed to settle until the supernatant liquid was perfectly clear. The barium chromate was then collected on a filter, washed with hot water, dried, gently ignited, and weighed. Here again arises the objection that the precipitate may have retained traces of alkaline salts, and again we find deduced an atomic weight which is too high. One hundred parts BaCrO₄ correspond to BaCl₂ as follows:

81.87	81.57
81.So	81.75
81.61	81.66
81.78	81.83
81.52	81.66
81.84	81.80
81.85	81.66
81.70	81.85
81.68	81.57
81.54	81.83
81.66	81.71
81.55	81.63
81.81	81.56
	•
81.86	81.58
81.54	81.67
81.68	81.84

Mean, 81.702, ± .014

Next in order we have to consider two papers by Kessler, who employed a peculiar volumetric method entirely his own. In brief, he compared the oxidizing power of potassium anhydrochromate with that of the chlorate, and from

^{*} Journ. für Prakt. Chem., 59, 27.

his observations deduced the ratio between the molecular weights of the two salts.

In his earlier paper* the mode of procedure was about as follows: The two salts, weighed out in quantities having approximate chemical equivalency, were placed in two small flasks, and to each was added 100 cc. of a ferrous chloride solution and 30 cc. hydrochloric acid. The ferrous chloride was added in trifling excess, and, when action ceased, the amount unoxidized was determined by titration with a standard solution of anhydrochromate. As in each case the quantity of ferrous chloride was the same, it became easy to deduce from the data thus obtained the ratio in question. I have reduced all of his somewhat complicated figures to a simple common standard, and give below the amount of chromate equivalent to 100 of chlorate:

120.118 120.371 120.138 120.096 120.241 120.181

Mean, 120.191, ± .028

In his later paper Kessler substituted arsenic trioxide for the iron solution. In one series of experiments the quantity of anhydrochromate needed to oxidize 100 parts of the arsenic trioxide was determined, and in another the latter substance was similarly compared with the chlorate. The subjoined columns give the quantity of each salt proportional to 100 of As, O₃:

$K_2Cr_2O_7$.	KClO3.
98.95	41.156
98.94	41.116
99.17	41.200
98.98	41.255
99.08	41.201
99.15	41.086
	41.199
Mean, 99.045, ± .028	41.224

^{*} Poggend. Annal., 95, 208. 1855.

[†] Poggend. Annal., 113, 137. 1861.

From the data given in the earlier paper, if we use our recent values for chlorine, potassium, and oxygen,

K₂Cr₂O₇ = 293.937,
$$\pm$$
 .086
And from the later, " = 294.159, \pm .119
General mean, " = 294.013, \pm .0697

Finally, we come to the determinations published by Siewert,* whose work does not seem to have attracted general attention. He, reviewing Berlin's work, found that upon reducing silver chromate with hydrochloric acid and alcohol, the chromic chloride solution always retained traces of silver chloride dissolved in it. These could be precipitated by dilution with water; but, in Berlin's process, they naturally came down with the chromium hydroxide, making the weight of the latter too high. Hence too large a value for the atomic weight of chromium. In order to find a more correct value Siewert resorted to the analysis of sublimed, violet, chromic chloride. This salt he fused with sodium carbonate and a little nitre, treated the fused mass with water, and precipitated from the resulting solution the chlorine by silver nitrate in presence of nitric acid. The weight of the silver chloride thus obtained, estimated after the usual manner, gave means for calculating the atomic weight of chromium. His figures, reduced to a common standard, give, as proportional to 100 parts of chloride of silver, the quantities of chromic chloride stated in the third of the subjoined columns:

^{*} Zeitschrift Gesammt. Wissenschaften, 17, 530. 1861.

.2367 gm	n. Cr ₂ Cl ₆ g	ave .6396 gr	m. AgCl.	37.007
.2946	66	·7994	44	36.853
.2593	66	.7039	46	36.838
-4935 ·	66	1.3395	66	36.842
.5850	66	1.5884	66	36.830
.6511	66	1.76681	46	36.852
.5503	66	1.49391	66	36.836

Mean, 36.865, $\pm .0158$

The first of these figures varies so widely from the others that we are justified in rejecting it; in which case the mean becomes $36.842, \pm .0031$.

Siewert also made two analyses of silver anhydrochromate by the following process. The salt, dried at 120°, was dissolved in nitric acid. The silver was then thrown down by hydrochloric acid, and, in the filtrate, chromium hydroxide was precipitated by ammonia. Reduced to a uniform standard, we find from his results, corresponding to 100 parts of AgCl, Ag, Cr, O₇, as in the last column:

Giving Berlin's single estimation equal weight with one of these, and combining, we get a general mean of 150.816, $\pm .074$.

Siewert's percentages of Cr₂O₃ obtained from Ag₂Cr₂O₇, are as follows, calculated from the above weighings.

Combining, as before, with Berlin's single result, giving the latter equal weight with one of these, we have a general mean of 35.236, $\pm .0335$.

For the ratio between silver chloride and chromic oxide, Siewert's two analyses of the anhydrochromate come out as follows. For 100 parts of AgCl we have of Cr₂O₈:

This figure, reduced to the standard of Berlin's work on the monochromate, becomes 26.525, $\pm .034$. Berlin's mean was 26.682, $\pm .0076$. The two means, combined, give a general mean of 26.676, $\pm .074$.

We may now consider the ratios before us, which are as follows:

- (1.) Percentage Cr.O. from Ag.CrO4, 23.014, ± .011
- (2.) Percentage Cr_2O_3 from $Ag_2Cr_2O_7$, 35.236, \pm .0335
- (3.) AgCl: Ag_2CrO_4 :: 100: 115.956, \pm .023
- (4.) AgCl: Ag, Cr_2O_7 :: 100: 150.816, \pm .074
- (5.) AgCl: Cr_2O_3 :: 100: 26.676, \pm .0074
- (6.) Percentage Cr₂O₃ in chromic sulphate, 39.1946, ± .0280
- (7.) Percentage Cr.O. in ammonia chrome alum, 16.143, ± .0125
- (8.) BaSO₄: BaCrO₄:: 100: 108.9815, ± .0369
- (9.) BaCrO₄: BaCl₂:: 100: 81.702, ± .014
- (10.) Molecular weight of $K_2Cr_2O_7$, 294.013, \pm .0697
- (11.) AgCl: $CrCl_2$:: 100: 36.842, \pm .0031

From these ratios we can at once deduce five values for the molecular weight of Cr.O., as follows:

For barium chromate we get two values:

From (8) ----- BaCrO₄ = 253.494,
$$\pm$$
 .094
" (9) ----- " = 253.976, \pm .067
General mean " = 253.816, \pm .054
From (3) we get Ag₂CrO₄ = 331.739, \pm .070
" (4) " Ag₂Cr₂O₇ = 431.470, \pm .215
" (11) " CrCl₃ = 158.102, \pm .018

Finally, from these intermediate data we derive six values for the atomic weight of chromium:

On account of the wide discrepancies between different data, and of the known constant errors vitiating some of the series of experiments, the foregoing general mean can have but little real value. In fact, a careful consideration of all the work represented in it will show that the most accurate estimate of the atomic weight of chromium must be deduced from the experiments of either Berlin, Kessler, or Siewert. Berlin's figures, taken by themselves, and combined, give, if the single analysis of silver anhydrochromate be assigned equal weight with a single analysis in the monochromate series, Cr = 52.389, \pm .019; or, if O = 16, Cr = 52.511.

Siewert's results, both for chromic chloride and the silver anhydrochromate, properly combined, give Cr = 52.009, $\pm .025$. If O = 16, this value becomes Cr = 52.129. In brief, the atomic weight of chromium may be nearly 52.5, or it may be 52. Only a revision of all the experiments could enable us to decide positively between these values. But as Siewert has pointed out probable sources of error in Berlin's work, I am inclined to give preference to the lower value.

MANGANESE.

Rejecting the early experiments of J. Davy and of Arfvedson, the first determinations of the atomic weight of manganese which we encounter are those of Turner* and of Berzelius.† Both of these chemists used the same method.

^{*} Trans. Roy. Soc. Edin., 11, 143. 1831.

[†] Lehrbuch, 5th Ed., 3, 1224.

The chloride of manganese was fused in a current of dry hydrochloric acid, and subsequently precipitated with a solution of silver. From the subjoined weighings I calculate the ratio given in the third column between MnCl, and 100 parts of AgCl:

```
4.20775 grm. MnCl<sub>2</sub> = 9.575 grm. AgCl. 43.945 } Berzelius. 3.063 " = 6.96912 " 43.950 } Berzelius. 12.47 grains MnCl<sub>2</sub> = 28.42 grains AgCl. 43.878—Turner. Mean, 43.924, ± .015
```

Hence the molecular weight of MnCl, is 125.662, \pm .045.

Many years later Dumas* also made the chloride of manganese the starting point of some atomic weight determinations. The salt was fused in a current of hydrochloric acid, and afterwards titrated with a standard solution of silver in the usual way. 100 parts of Ag are equivalent to the quantities of MnCl, given in the third column:

```
3.3672 grm. MnCl_2 = 5.774 grm. Ag. 58.317

3.0872 " 5.293 " 58.326

2.9671 " 5.0875 " 58.321

1.1244 " 1.928 " 58.320

1.3134 " 2.251 " 58.321

Mean, 58.321, \pm.001
```

Hence MnCl₂ = 125.594, \pm .011. This, combined with Berzelius and Turner's figures, gives MnCl₂ = 125.598, \pm .011. And Mn = 54.858, \pm .031.

An entirely different method of investigation was followed by v. Hauer,† who, as in the case of cadmium, ignited the sulphate in a stream of sulphuretted hydrogen, and determined the quantity of sulphide thus formed. I subjoin his weighings, and also the percentage of MnS in MnSO₄ as calculated from them:

^{*} Ann. Chem. Pharm., 113, 25. 1860. † Journ. für Prakt. Chem., 72, 360. 1857.

4.0626 gm	n. MnSO ₄ gave	2.3425	grm. MnS.	57.660 p	er cent.
4.9367	66	2.8442	66	57.613	66
5.2372	66	3.0192	66	57.649	46
7.0047	66	4.0347	44	57.600	44
4.9175	44	2.8297	66	57-543	66
4.8546	66	2.7955	66 [°]	57.585	66
4.9978	66	2.8799	44	57.625	66
4.6737	66	2.6934	66	57.629	66
4.7240	66	2.7197	46	57.572	66

Mean, 57.608, $\pm .008$

Hence Mn = 54.785, $\pm .031$.

This method of v. Hauer's, which seemed to give good results with cadmium, is, according to Schneider,* inapplicable to manganese; for the reason that the sulphide of the latter metal is liable to be contaminated with traces of oxysulphide. Such an impurity would bring the atomic weight out too high. The results of two different processes, one carried out by himself and the other in his laboratory by Rawack, are given by Schneider in this paper.

Rawack reduced manganoso-manganic oxide to manganous oxide by ignition in a stream of hydrogen, and weighed the water thus formed. From his weighings I get the values in the third column, which represent the Mn₃O₄ equivalent to one gramme of water:

```
4.149 grm. Mn<sub>2</sub>O<sub>4</sub> gave 0.330 grm. H<sub>2</sub>O.
                                                       12.5727
 4.649
                               .370
                                                       12.5643
 6.8865
                               .5485
                                                       12.5552
                               .5855
 7.356
                                                       12.5636
 8.9445
                               .7135
                                                       12.5361
11.584
                               .9225
                                                       12.5572
```

Mean, 12.5582, \pm .0034

Hence $Mn = 53.911, \pm .026$.

Here the most obvious source of error lies in the possible loss of water. Such a loss, however, would increase the apparent atomic weight of manganese; but we see that the value found is much lower than that obtained either by Dumas or v. Hauer.

^{*} Poggend. Annal., 107, 605.

Schneider himself effected the combustion of manganous oxalate with oxide of copper. The salt was not absolutely dry, so that it was necessary to collect both water and carbon dioxide. Then, upon deducting the weight of water from that of the original material, the weight of anhydrous oxalate was easily ascertained. Subtracting from this the CO₂, we get the weight of Mn. If we put CO₂ = 100, the quantities of manganese equivalent to it will be found in the last column:

```
1.5075 grm. oxalate gave .306 grm. H<sub>2</sub>O and .7445 grm. CO<sub>2</sub>. 61.3835
2.253 " .4555 " 1.1135 " 61.4291
3.1935 " .652 " 1.5745 " 61.4163
5.073 " 1.028 " 2.507 " 61.3482

Mean, 61.3943, ±.0122
```

Hence $Mn = 53.904, \pm .014$.

This result agrees beautifully with the value calculated from Rawack's experiments.

Now to combine the four independent values which we have thus far obtained:

The considerations already cited, however, go to show that this general mean must be slightly affected by some plus constant error. It is probable, therefore, that a more correct figure will result from rejecting the first and second values in the above combination, and taking the data furnished by Rawack and Schneider alone. Combining their figures, we get as follows. Mn = 53.906, $\pm .012$. Or, if O = 16, Mn = 54.029.

Since the foregoing calculations were made Dewar and Scott* have reported the following experiments. From the

^{*} Nature, Sept. 15, 1881, p. 470.

131 iron.

complete analysis of silver permanganate, putting Ag = 108 and O = 16, they find in three estimations Mn = 55.51, 54.04, and 54.45. From the analysis of pure MnO_2 , made from the nitrate, Mn = 53.3 to 53.6. Up to the date of writing a detailed account of the methods employed has not been published.

IRON.

The atomic weight of iron has been determined almost exclusively from the composition of ferric oxide. Beyond this there are only a few comparatively unimportant experiments by Dumas relative to ferrous and ferric chlorides.

Most of the earlier data relative to the percentage of metal and oxygen in ferric oxide we may reject at once, as set aside by later investigations. Among this no longer valuable material there is a series of experiments by Berzelius, another by Döbereiner, and a third by Capitaine. The work done by Stromeyer and by Wackenroder was probably good, but I am unable to find its details. The former found 30.15 per cent. of oxygen in the oxide under consideration, while Wackenroder obtained figures ranging from a minimum of 30.01 to a maximum of 30.38 per cent.*

In 1844 Berzelius† published two determinations of the ratio in question. He oxidized iron by means of nitric acid, and weighed the oxide thus formed. He thus found that when O = 100 Fe = 350.27 and 350.369.

Hence the following percentages of Fe in Fe₂O₃.

70.018 70.022 Mean, 70.020, ± .0013

About the same time Svanberg and Norlin ; published

^{*} For additional details concerning these earlier papers I must refer to Oudemans' monograph, pp. 140, 141.

[†] Ann. Chem. Pharm., 50, 432. Berz. Jahresb., 25, 43.

¹ Berzelius' Jahresbericht, 25, 42.

two elaborate series of experiments; one relating to the synthesis of ferric oxide, the other to its reduction. In the first set pure piano-forte wire was oxidized by nitric acid, and the amount of oxide thus formed was determined. The results were as follows:

```
1.5257 grm. Fe gave 2.1803 grm. Fe<sub>2</sub>O<sub>2</sub>.
                                                 69.977 per cent. Fe.
                                                 69.936
2.4051
                        3.4390
2.3212
                                                 69.928
                        3.3194
                        3.3183
                                                 69.968
2.32175
                                                 69.960
2.2772
                        3.2550
                                                              66
2.4782
                                     "
                                                 69.970
                        3.5418
               "
2.3582
                                                 69.935
                        3.3720
```

Mean, 69.9534, \pm .0050

In the second series ferric oxide was reduced by ignition in a current of hydrogen, yielding the subjoined percentages of metal:

```
2.98353 grm. Fe<sub>2</sub>O<sub>3</sub> gave 2.08915 grm. Fe.
                                                  70.025 per cent.
                            1.6910
                                                  70.015
2.41515
2.99175
                                                  70.014
                           2.09455
3.5783
                           2.505925
                                                  70.030
4.1922
                           2.9375
                                                  70.072
                                                  70.056
3.1015
                           2.17275
2.6886
                           1.88305
                                                  70.036
```

Mean, 70.0354, ± .0055

It is evident that one or both of these series must be vitiated by constant errors, and that these probably arise from impurities in the materials employed. Impurities in the wire taken for the oxidation series could hardly have been altogether avoided, and in the reduction series it is possible that weighable traces of hydrogen may have been retained by the iron. At all events it is probable that the errors of both series are in contrary directions, and, therefore, in some measure compensatory.

In 1844 there was also published an important paper by Erdmann and Marchand.* These chemists prepared ferric oxide by the ignition of pure ferrous oxalate, and submitted

^{*} Journ. für Prakt. Chem., 33, 1.

133

it to reduction in a stream of hydrogen. Two sets of results were obtained with two different samples of ferrous oxalate, prepared by two different methods. For present purposes, however, it is not necessary to discuss these sets separately. The percentages of iron in Fe₂O₃ come out as follows:

Mean, 70.0094, $\pm .0080$

In 1850 Maumené's* results appeared. He dissolved pure iron wire in aqua regia, precipitated with ammonia, filtered off the precipitate, washed thoroughly, ignited, and weighed, after the usual methods of quantitative analysis. The percentages of Fe in Fe₂O₃ are given in the third column:

```
1.482 grm. Fe gave 2.117 grm. Fe<sub>2</sub>O<sub>2</sub>.
                                                    70.005 per cent.
                                                    70.010
1.452
                        2.074
1.3585
                                                    69.990
                        1.941
                        2.0285
                                                    70.002
1.420
                                                    69.998
1.492
                        2.1315
                                                    70.000
                        2.220
1.554
```

Mean, 70.0008, \pm .0019

Two more results, obtained by Rivot† through the reduction of ferric oxide in hydrogen, remain to be noticed. The percentages are:

We have thus before us six series of results, which we may now combine.

^{*} Compt. Rend., Oct. 17, 1850.

[†] Ann. Chem. Pharm., 78, 214. 1851.

Berzelius	70.020,	±.	0013
Erdmann and Marchand	70.0094,	土。	.0080
Svanberg and Norlin, Oxyd	69.9534,	± .	.0050
" Reduc	70.0354,	士.	0055
Maumené	70.0008,	± .	0019
Rivot	69.33,	± .	.013
General mean	70.0075.	<u>+</u> .	.0010

From this we get Fe = 55.891, $\pm .012$; or, if O = 16, this becomes 56.0195.

Dumas'* results, obtained from the chlorides of iron, are of so little weight that they might safely be omitted from our present discussion. For the sake of completeness, however, we will include them.

Pure ferrous chloride, ignited in a stream of hydrochloric acid gas, was dissolved in water and titrated with a silver solution in the usual way. One hundred parts of silver are equivalent to the amounts of FeCl, given in the third column:

Ferric chloride, titrated in the same way, gave these results:

These give us two additional values for Fe, as follows:

From FeCl₂ Fe =
$$56.028$$
, $\pm .119$
" Fe₂Cl₂ = 56.189 , $\pm .062$

Combining these with the value deduced from the composition of Fe₂O₃, Fe = 55.891, \pm .012, we get this general mean, Fe = 55.913, \pm .012. If O = 16, this becomes Fe = 56.042.

^{*} Ann. Chem. Pharm., 113, 26. 1860.

COPPER.

The atomic weight of copper has been chiefly determined from the composition of the black oxide and the anhydrous sulphate. In dealing with the first named compound all experimenters have agreed in reducing it with a current of hydrogen, and weighing the metal thus set free.

The earliest experiments of any value were those of Berzelius,* whose results were as follows:

```
7.68075 grm. CuO lost 1.55 grm. O. 79.820 per cent. Cu in CuO. 9.6115 " 1.939 " 79.826 " " Mean, 79.823, ± .002
```

Erdmann and Marchand,† who come next in chronological order, corrected their results for weighing in air. Their weighings, thus corrected, give us the subjoined percentages of metal in CuO:

```
63.8962 grm. CuO gave 51.0391 grm. Cu. 79.878 per cent.
65.1590 " 52.0363 " 79.860 "
60.2878 " 48.1540 " 79.874 "
46.2700 " 36.9449 " 79.846 "

Mean, 79.8645, ± .0038
```

Still later we find a few analyses by Millon and Commaille.‡ These chemists not only reduced the oxide by hydrogen, but they also weighed, in addition to the metallic copper, the water formed in the experiments. In three determinations the results were as follows:

```
6.7145 grm. CuO gave 5.3565 grm. Cu and 1.5325 grm. H<sub>2</sub>O. 79.775 per cent.
3.3945 " 2.7085 " .7680 " 79.791 "
2.2240 grm. Cu. 79.770 "

Mean, 79.7787, ± .0043
```

For the third of these analyses the water estimation was not made, but for the other two it yielded results which, in

^{*} Poggend. Annal., 8, 177.

[†] Journ. für Prakt. Chem., 31, 389. 1844.

[‡] Fresenius' Zeitschrift, 2, 475. 1863.

the mean, would make the atomic weight of copper 63.087, \pm .222. This figure has so high a probable error that we need not consider it further.

The results obtained by Dumas* are wholly unavailable. Indeed, he does not even publish them in detail. He merely says that he reduced copper oxide, and also effected the synthesis of the subsulphide, but without getting figures which were wholly concordant. He puts Cu = 63.5.

Latest of all, and probably the best also, we have the determinations by Hampe.† First, he attempted to estimate the atomic weight of copper by the quantity of silver which the pure metal could precipitate from its solutions. This attempt failed to give satisfactory results, and he fell back upon the old method of reducing the oxide. From ten to twenty grammes of material were taken in each experiment, and the weights were reduced to a vacuum standard:

```
20.3260 grm. CuO gave · 16.2279 grm. Cu. 79.838 per cent.

20.68851 " 16.51669 " 79.835 "

10.10793 " 8.06926 " 79.831 "

Mean, 79.8347, ± .0013
```

Hampe also determined the quantity of copper in the anhydrous sulphate, CuSO₄. From 40 to 45 grammes of the salt were taken at a time, the metal was thrown down by electrolysis, and the weights were all corrected. I subjoin the results:

```
40.40300 grm. CuSO<sub>4</sub> gave 16.04958 grm. Cu. 39.724 per cent.

44.64280 " 17.73466 " 39.726 "

Mean, 39.725, ± .0007
```

We now have four series of experiments upon copper oxide, as follows:

```
      Berzelius
      79.823, ± .0020

      Erdmann and Marchand
      79.8645, ± .0038

      Millon and Commaille
      79.7787, ± .0043

      Hampe
      79.8347, ± .0013

      General mean
      79.830, ± .0010
```

^{*} Ann. d. Chim. et Phys., (3,) 55, 129.

[†] Fresenius' Zeitschrift, 13, 352.

For copper we have—

From composition of CuO....Cu = 63.181,
$$\pm$$
 .036
" CuSO₄, (Hampe) " = 63.171, \pm .012
General mean " = 63.173, \pm .011

If O = 16, then Cu becomes = 63.318.

The close agreement between the two independent values for Cu is certainly very striking. It will be seen that Hampe's two estimates upon the sulphate carry (perhaps accidentally) much greater weight than all the experiments upon the oxide. This might seem like giving them undue credit, were it not for the fact of the remarkable concordance of the results above referred to. Either estimate for Cu would be valid without the other.

MOLYBDENUM.

If we leave out of account the inaccurate determination made by Berzelius,* we shall find that the data for the atomic weight of molybdenum lead to two independent estimates of its value; one near 92, the other near 96. The earlier results found by Berlin and by Svanberg and Struve lead to the lower number; the more recent work of Debray, Dumas, and Lothar Meyer sustains the higher. The latter value is the more probable, although both may be vitiated by constant errors in opposite directions.

The earliest investigation which we need especially to consider is that of Svanberg and Struve.† These chemists tried a variety of different methods, but finally based their conclusions upon the two following: first, molybdenum trioxide was fused with potassium carbonate, and the carbon dioxide which was expelled was estimated; secondly, molybdenum disulphide was converted into the trioxide by

^{*} Poggend. Annal., 8, 1. 1826.

[†] Journ. für Prakt. Chem., 44, 301. 1848.

roasting, and the ratio between the weights of the two substances was determined.

By the first method it was found that 100 parts of MoO, will expel the following quantities of CO₂:

31.4954
31.3749
31.4705
—————
Mean, 31.4469, ± .0248

The carbon dioxide was determined simply from the loss of weight when the weighed quantities of trioxide and carbonate were fused together. It is plain that if, under these circumstances, a little of the trioxide should be volatilized, the total loss of weight would be slightly increased. A constant error of this kind would tend to bring out the atomic weight of molybdenum too low.

By the second method, the conversion by roasting of MoS, into MoO, Svanberg and Struve obtained these results. Two samples of artificial disulphide were taken, A and B, and yielded for each hundred parts the following of trioxide:

Mean, 89.7523, $\pm .0176$

Three other experiments in series B gave divergent results, and, although published, are rejected by the authors themselves. Hence it is not necessary to cite them in this discussion. We again encounter in these figures the same source of constant error which apparently vitiates the preceding series, namely, the possible volatilization of the trioxide. Here, also, such an error would tend to reduce the atomic weight of molybdenum.

Upon discussing the data given in the foregoing para-

graphs we get somewhat noticeable results. From the carbon dioxide series, Mo = 91.711, $\pm .113$, a figure having no unusual interest. From the other series, if S = 31.987 and O = 15.9633, we get Mo = 92.979, $\pm .354$; but if we take S = 32 and O = 16, then Mo becomes = 92.133. In this case the higher values for oxygen and sulphur lead to a lower number for molybdenum. In the carbonate series the assumption of 12 and 16 for C and O, respectively, makes Mo = 92.033. In other words, if we assume the ordinary even numbers for C, O, and S, S vanberg and S truve's two methods yield more nearly concordant results than when the revised values for these elements are taken.

Berlin,* a little later than Svanberg and Struve, determined the atomic weight of molybdenum by igniting a molybdate of ammonium and weighing the residual MoO₈. Here, again, a loss of the latter by volatilization may (and probably does) lead to too low a result. The salt used was $(NH_4)_4 Mo_8 O_{17}$. 3 H₂O, and in it these percentages of MoO₈ were found:

81.598 81.612 81.558 81.555 Mean, 81.581, ± .0095

Hence Mo = 91.9817, $\pm .0776$; a result agreeing quite well with those of Svanberg and Struve.

Until 1859 the value 92 was generally accepted on the basis of the foregoing researches, but in this year Dumas† published some figures tending to sustain a higher number. He prepared molybdenum trioxide by roasting the disulphide, and then reduced it to metal by ignition in hydrogen. At the beginning the hydrogen was allowed to act at a comparatively low temperature, in order to avoid volatilization of trioxide; but at the end of the operation the heat

^{*} Journ. für Prakt. Chem., 49, 444. 1850.

[†] Ann. Chem. Pharm., 105, 84, and 113, 23.

was raised sufficiently to insure a complete reduction. From the weighings I calculate the percentages of metal in MoO₂:

.448 gm	n. MoO ₃ gave	.299	grm.	Mo.		66.741 p	er cent.
.484	46	.323	46			66.736	66
.484	66	.322	46			66.529	66
.498	66	.332	46			66.667	44
·559	66	-373	46			66.726	66
.388	66	.258	46			66.495	a
					Mean,	66.649,	± .030

In 1868 the same method was employed by Debray.* His trioxide was purified by sublimation in a platinum tube. His percentages are as follows:

```
5.514 grm. MoO<sub>3</sub> gave 3.667 grm. Mo.

7.910

"
5.265

"
66.503 per cent.

66.561

"
9.031

"
66.604

Mean, 66.556, ± .020
```

This mean, combined with that of Dumas', gives a general mean of 66.585, $\pm .017$.

Hence Mo = 95.429, $\pm .057$.

Debray also made two experiments upon the precipitation of molybdenum trioxide in ammoniacal solution by nitrate of silver. In his results, as published, there is curious discrepancy, which, I have no doubt, is due to typographical error. These results I am, therefore, compelled to leave out of consideration. They could not, however, exert a very profound influence upon the final discussion.

The most recent investigation upon the atomic weight of molybdenum is the discussion by Lothar Meyer† of the experimental results obtained by Liechti and Kemp‡ in their analyses of the chlorides. Of these compounds there are four: MoCl₂, MoCl₂, MoCl₄, and MoCl₅. The chlorine in each was estimated as silver chloride, and the molybdenum as disulphide. From these analyses Meyer deduces three

^{*} Compt. Rend., 66, 734.

[†] Ann. Chem. Pharm., 169, 365. 1873.

[‡] Ann. Chem. Pharm., 169, 344.

sets of ratios, namely: between MoCl_n and n AgCl; between MoCl_n and MoS₂, and between MoS₂ and n AgCl. We will use only the first and last of these; the probable error of the atomic weight deduced from the second being relatively so high as to make the value connected with it comparatively unimportant. The analyses of the trichloride, being discordant, are here rejected.

By reducing the weighings published by Liechti and Kemp* to a common standard we get the following percentage results. In MoCl, the subjoined quantities of the original substance and of MoS, correspond to 100 parts of AgCl:

58.194 55.591 58.524 56.065		<u> </u>
--------------------------------	--	----------

Hence MoCl₂ = 166.902, $\pm .188$, and MoS₂ = 159.652, $\pm .268$.

With the tetrachloride similarly calculated we get these figures, corresponding to 100 parts AgCl:

$$MoCl_4$$
. MoS_2 .

41.492 27.957

41.319

————

Mean, 41.4055, \pm .0583

Hence $MoCl_4 = 236.914$, $\pm .358$, and MoS_2 , if given the weight of a single experiment in the dichloride series, = 159.964, $\pm .627$.

* These are as follows:

```
.2666 grm. MoCl<sub>2</sub> gave .2550 grm. MoS<sub>2</sub> and .4573 grm. AgCl.
.1811
                            .1730
                                                       .3112
.2530
                             .2422
                                                       .4320 ,
.4126 grm. MoCl<sub>4</sub> gave .2780
                                                       .9944
                                                       .4654
.1923
.5810 grm. MoCl<sub>s</sub> gave .3414
                                                      1.5222
.2466
                                                       .6465
                             .1441
```

For the pentachloride the following quantities balance 100 of AgCl:

$$MoCl_8$$
. MoS_2 .

38.168 22.428
38.057 22.289
...

Mean, 38.112, \pm .038 Mean, 22.3585, \pm .040

Hence MoCl₅ = 272.587, $\pm .271$, and MoS₂ = 159.914, $\pm .287$.

We have now the molecular weight of each chloride, and three values for that of the disulphide. Combining the latter we get a general mean, as follows:

With these data, in addition to those given by Dumas and by Debray, we get five estimates of the atomic weight of molybdenum:

```
Dumas and Debray's data _______ Mo = 95.429, \pm .057

From molecular weight of MoCl<sub>2</sub> ____ " = 96.262, \pm .190

" MoCl<sub>4</sub> ___ " = 95.434, \pm .363

" MoCl<sub>5</sub> ___ " = 95.737, \pm .280

" MoS<sub>2</sub> ___ " = 95.816, \pm .188

General mean ___ " = 95.527, \pm .051

Or, if O = 16, Mo = 95.747.
```

It will at once be seen that the most reliable results are those obtained by the reduction of molybdenum trioxide. Traces of oxychlorides may possibly have contaminated the chlorides and augmented their atomic weight. Our final figure, therefore, may be a trifle too high, but the early value, 92, is unquestionably very far too low.

Since the foregoing discussion was written a single experiment by Rammelsberg * has been brought to my netice.

^{*} Berlin Monatsbericht, 1877, 574.

Closely following Dumas' method, he reduced molybdenum trioxide to metal, finding in it 66.708 per cent. of the latter. This figure comes within the limits of variation of Dumas' experiments, and therefore gives them additional confirmation. Its introduction into the general mean, however, would exert too little influence upon the latter to justify the labor of recalculation.

TUNGSTEN.

The atomic weight of tungsten has been determined from analyses of the trioxide, the hexchloride, and the tungstates of iron, silver, and barium.

The composition of the trioxide has been the subject of many investigations. Malaguti* reduced this substance to the blue oxide, and from the difference between the weights of the two compounds obtained a result now known to be considerably too high. In general, however, the method of investigation has been to reduce WO₃ to W in a stream of hydrogen at a white heat, and afterwards to reoxidize the metal, thus getting from one sample of material two results for the percentage of tungsten. This method is unquestionably accurate, provided that the trioxide used be pure.

The first experiments which we need consider are, as usual, those of Berzelius. † 899 parts WO₃ gave, on reduction, 716 of metal. 676 of metal, reoxidized, gave 846 WO₃. Hence these percentages of W in WO₃:

79.644, by reduction.
79.905, by oxidation.
Mean, 79.7745, ± .0880

These figures are far too high, the error being undoubtedly due to the presence of alkaline impurity in the trioxide employed.

^{*} Journ. sür Prakt. Chem., 8, 179. 1836.

[†] Poggend. Annal., 8, 1. 1826.

Next in order of time comes the work of Schneider,* who, with characteristic carefulness, took every precaution to get pure material. His percentages of tungsten are as follows:

79.336
79.254
79.312
79.326
79.350

Mean, 79.3156, ± .0112

Oxidation Series.
79.329
79.324
79.328

Mean, 79.327, ± .0010

Closely agreeing with these figures are those of Marchand.† published in the following year:

Reduction Series.

79.307

79.302

Mean, 79.3045, .0017

Oxidation Series.

79.321

79.352

Mean, 79.3365, ± .0105

The figures obtained by v. Borch‡ agree in mean tolerably well with the foregoing. They are as follows:

Reduction Series.

79.310
79.212
79.289
79.313
79.225
79.290
79.302

Mean, 79.277, ± .0106

^{*} Journ. für Prakt. Chem., 50, 152. 1850. † Ann. Chem. Pharm., 77, 261. 1851.

[‡] Journ. für Prakt. Chem., 54, 254. 1851.

Oxidation Series.

79.359

79.339

Mean, 79.349, ± .0067

Dumas * gives only a reduction series, based upon trioxide obtained by the ignition of a pure ammonium tungsten. The reduction was effected in a porcelain boat, platinum being objectionable on account of the tendency of tungstate to alloy with it. Dumas publishes only weighings, from which I have calculated the percentages:

2.784	grm. WO ₃ gave	2.208	grm.	w.	79.310 p	er cent.
2.994	44	2.373	66		79.259	66
4.600	44	3.649	46		79.326	66
.985	"	.781	66		79.289	46
.917	66	.727	66		79.280	66
.917	c l	.728	66		79.389	66
1.717	44	1.362	66		79.324	66
2.988	46	2.370	"		79.317	"
				Mean,	79.312,	± .009

The data furnished by Bernoulli† differ widely from those just given. This chemist undoubtedly worked with impure material, the trioxide having a greenish tinge. Hence the results are too high. These are the percentages of W:

```
Reduction Series.

79.556
79.526
79.553
79.558
79.549
78.736
Mean, 79.413, ± .091

Oxidation Series.

79.558
79.656
79.555
```

79.554

Mean, 79.581, $\pm .017$

^{*} Ann. Chem. Pharm., 113, 23. 1860

[†] Poggend. Annal., 111, 573. 1860.

Two reduction experiments by Persoz* give the following results:

Finally, we have the work done by Roscoe.† This chemist used a porcelain boat and tube, and made six weighings, after successive reductions and oxidations, with the same sample of 7.884 grammes of trioxide. These weighings give me the following five percentages, which, for the sake of uniformity with foregoing series, I have classified under the usual, separate headings:

```
Reduction Series.

79.196

79.285

79.308

Mean, 79.263, ± .023

Oxidation Series.

79.230

79.299
```

Mean, 79.2645, ± .0233

There are still other experiments by Riche,‡ which I have not been able to get in detail. They cannot be of any value, however, for they give to tungsten an atomic weight of about ten units too low. We may therefore neglect this series, and go on to combine the others:

```
      Berzelius
      79.7745, ± .088

      Schneider, Reduction
      79.3156, ± .0112

      "Oxidation
      79.327, ± .0010

      Marchand, Reduction
      79.3045, ± .0017

      "Oxidation
      79.3365, ± .0105

      v. Borch, Reduction
      79.277, ± .0106

      "Oxidation
      79.349, ± .0067
```

^{. *} Zeit. Anal. Chem., 3, 260. 1864.

[†] Ann. Chem. Pharm., 162, 368. 1872.

¹ Journ. für Prakt. Chem., 69, 10. 1857.

```
Dumas ______ 79.312, ± .009

Bernouilli, Reduction _____ 79.413, ± .091

" Oxidation _____ 79.581, ± .017

Persoz _____ 79.314, ± .007

Roscoe, Reduction _____ 79.263, ± .023

" Oxidation _____ 79.2645, ± .0233

General mean _____ 79.3215, ± .00085
```

The rejection of the figures given by Berzelius and by Bernoulli exerts an unimportant influence upon the final result. There is, therefore, no practical objection to retaining them in the discussion.

In 1861 Scheibler* deduced the atomic weight of tungsten from analyses of barium metatungstate, BaO.4 WO₃.9 H₂O. In four experiments he estimated the barium as sulphate, getting closely concordant results, which were, however, very far too low. These, therefore, are rejected. But from the percentage of water in the salt a very good result was attained. The percentages of water are as follows:

13.053 13.054 13.045 13.010 13.022

Mean, 13.0368, \pm .0060

The work of Zettnow,† published in 1867, was somewhat more complicated than any of the foregoing researches. He prepared the pure tungstates of silver and of iron, and from their composition determined the atomic weight of tungsten.

In the case of the iron salt the method of working was this: The pure, artificial FeWO₄ was fused with sodium carbonate, the resulting sodium tungstate was extracted by water, and the thoroughly washed, residual ferric oxide was dissolved in hydrochloric acid. This solution was then reduced by zinc, and titrated for iron with potassium permanganate. Corrections were applied for the drop in excess of

^{*} Journ. für Prakt. Chem., 83, 324.

[†] Poggend. Annal.. 130, 30.

permanganate needed to produce distinct reddening, and for the iron contained in the zinc. 11.956 grammes of the latter metal contained iron corresponding to 0.6 cc. of the standard solution. The permanganate was standardized by comparison with pure ammonium-ferrous sulphate, Am, Fe(SO₄)₂. 6 H,O, so that, in point of fact, Zettnow establishes directly only the ratio between that salt and the ferrous tungstate. From Zettnow's four experiments in standardizing I find that 1 cc. of his solution corresponds to 0.0365457 grammes of the double sulphate, with a probable error of ± .0000012.

Three sets of titrations were made. In the first a quantity of ferrous tungstate was treated according to the process given above; the iron solution was diluted to 500 cc., and four titrations made upon 100 cc. at a time. The second set was like the first, except that three titrations were made with 100 cc. each, and a fourth upon 150 cc. In the third set the iron solution was diluted to 300 cc., and only two titrations upon 100 cc. each were made. In sets one and two thirty grammes of zinc were used for the reduction of each, while in number three but twenty grammes were taken. Zettnow's figures, as given by him, are quite complicated; therefore I have reduced them to a common standard. After applying all corrections the following quantities of tungstate, in grammes, correspond to 1 cc. of permanganate solution:

With the silver tungstate, Ag, WO4, Zettnow employed two methods. In two experiments the substance was de-

composed by nitric acid, and the silver thus taken into solution was titrated with standard sodium chloride. In three others the tungstate was treated directly with common salt, and the residual silver chloride collected and weighed. Here again, on account of some complexity in Zettnow's figures, I am compelled to reduce his data to a common standard. To 100 parts of AgCl the following quantities of Ag2WO4 correspond:

By First Method.

161.665

161.603

Mean, 161.634, ± .021

By Second Method.

161.687 161.651 161.613

Mean, 161.650, \pm .014

General mean from both series, 161.645, ± .012

Finally, we have two analyses by Roscoe of tungsten hexchloride, published in the same paper with his results upon the trioxide. In one experiment the chlorine was determined as AgCl; in the other the chloride was reduced by hydrogen, and the residual tungsten estimated. By bringing both results into one form of expression we have for the percentage of chlorine in WCl₆:*

We have now five ratios from which to calculate the atomic weight of tungsten:

- (1.) Percentage of W in WO₃, 79.3215, \pm .00085
- (2.) Percentage of H₂O in BaO.4WO₃.9H₂O, 13.0368, ± .0060

19.5700 grm. WCl₆ gave 42.4127 grm. AgCl. 10.4326 " 4.8374 " tungsten.

^{*} The actual figures are as follows:

- (3.) $Am_2Fe(SO_4)_2.6H_2O: FeWO_4::.0365457, \pm .0000012:.0283549, \pm .0000115$
- (4.) $AgCl: Ag_2WO_4:: 100: 161.645, \pm .012$
- (5.) Percentage of Cl in WCl₈, 53.610, ± .015

From these we get five values for tungsten, as follows:

URANIUM.

It is not the purpose of the present investigation to examine at all systematically such questions as are involved in the discussion whether the atomic weight of uranium is 120 or 240. For convenience we may use the formulæ based upon the smaller number, and, if eventually the larger value proves to be correct, it will be easy to double the figures which we obtain. Suffice it to say here, that the specific heat of the green oxide, according to Donath,* agrees best with the formula U_sO₄ and the lower atomic weight. On the other hand, the value 240 fits best into such schemes as that given by Mendelejeff in his paper on the periodic law. An accurate determination of the specific heat of the metal itself is much needed, for the material with which Regnault worked was of uncertain quality; furthermore, the vapor density of some volatile uranium compounds ought to be ascertained.† Until some such data have been rigidly

^{*}Ber. d. Deutsch. Chem. Gesell., 12, 742. 1879.

[†] The value of 240 for uranium is strongly sustained by the recent experiments of Zimmermann upon the vapor density of the tetrachlorid and tetrabromid. For UBr₄ the vapor density is 19.46, while theory (U = 240) requires 19.36. For UCl₄ the v. d. 13.33 was found. Theory, 13.21. (Ber. der Deutsch. Chem. Gesell., 14, s. 1934. 1881.)

established the controversy over the two rival values can hardly be satisfactorily settled.

The earlier attempts to determine the atomic weight of uranium were all vitiated by the erroneous supposition that the uranous oxide was really the metal. The supposition, of course, does not affect the weighings and analytical data which were obtained, although these, from their discordance with each other and with later and better results, have now only a historical value.

For present purposes the determinations made by Berzelius,* by Arfvedson,† and by Marchand,‡ may be left quite out of account. Berzelius employed various methods, while the others relied upon estimating the percentage of oxygen lost upon the reduction of U_sO₄ to UO. Rammelsberg's || results also, although very suggestive, need no full discussion. He analyzed the green chloride, UCl,; effected the synthesis of uranyl sulphate from uranous oxide; determined the amount of residue left upon the ignition of the sodio and bario-uranic acetates; estimated the quantity of magnesium uranate formed from a known weight of UO, and attempted also to fix the ratio between the green and the black oxides. His figures vary so widely that they could count for little in the establishing of any general mean; and, moreover, they lead to estimates of the atomic weight which are mostly below the true value. For instance, twelve lots of U,O, from several different sources were reduced to UO by heating in hydrogen. The percentages of loss varied from 3.83 to 4.67, the mean being 4.121. These figures give values for the atomic weight of uranium ranging from 92.66 to 117.65, or, in mean, 107.50. Such discordance is due partly to impurity in some of the material studied, and illustrates the difficulties inherent in the problem to be solved. Some of the uranoso-uranic oxide was prepared by

^{*}Schweigg. Journ., 22, 336. 1818. Poggend. Annal., 1, 359. 1825.

[†] Poggend. Annal., 1, 245. Berz. Jahr., 3, 120. 1822.

[‡] Journ. für Prakt. Chem., 23, 497. 1841.

[|] Poggend. Annal., 55, 318, 1842; 56, 125, 1842; 59, 9, 1843; 66, 91, 1845.

Journ. für Prakt. Chem., 29, 324.

calcining the oxalate, and retained an admixture of carbon. Many such points were worked up by Rammelsberg with much care, so that his papers should be scrupulously studied by any chemist who contemplates a redetermination of the atomic weight of uranium.

In 1841 and 1842 Peligot published certain papers* showing that the atomic weight of uranium must be somewhere near 120. A few years later the same chemist published fuller data concerning the constant in question, but in the time intervening between his earlier and his final researches other determinations were made by Ebelmen and by Wertheim. These investigations we may properly discuss in chronological order. For present purposes the early work of Peligot may be dismissed as only preliminary in character. It showed that what had been previously regarded as metallic uranium was in reality an oxide, but gave figures for the atomic weight of the metal which were merely approximations.

Ebelmen's † determinations of the atomic weight of uranium were based upon analyses of uranic oxalate. This salt was dried at 100°, and then, in weighed amount, ignited in hydrogen. The residual uranous oxide was weighed, and in some cases converted into U₈O₄ by heating in oxygen. The following weights are reduced to a vacuum standard:

```
10.1644 grm. oxalate gave 7.2939 grm. UO.
                                             Gain on oxidation, .3685
12.9985
                           9.3312
11.8007
                          8.4690
                                                                .3275
                           7.1731
                                                                .2812
9.9923
11.0887
                           7.9610
                                                                . 3105
                          7.2389
10.0830
                          4.8766
 6.7940
16.0594
                          11.5290
                                                                .4531
```

Reducing these figures to percentages, we may present the results in two columns. Column A gives the percentages of UO in the oxalate, while B represents the amount of U₃O₄ formed from 100 parts of UO:

^{*} Compt. Rend., 12, 735. 1841. Ann. Chim. Phys., (3,) 55. 1842.

[†] Journ. sür Prakt. Chem., 27, 385. 1842.

Α.

B.

Wertheim's * experiments were even simpler in character than those of Ebelmen. Sodio-uranic acetate, carefully dried at 200°, was ignited, leaving the following percentages of sodium uranate:

Hence the molecular weight of $Na_2U_4O_7 = 634.865$, $\pm .191$. And U = 119.282, $\pm .048$.

The final results of Peligot's† investigations appeared in 1846. Both the oxalate and the acetate of uranium were studied and subjected to combustion analysis. The oxalate was scrupulously purified by repeated crystallizations, and thirteen analyses, representing different fractions, were made. Seven of these gave imperfect results, due to incomplete purification of the material; six only, from the later crystallizations, need to be considered. In these the uranium

^{*} Journ. für Prakt. Chem., 29, 209. 1843.

[†] Compt. Rend., 22, 487.

was weighed as U₈O₄, and the carbon as CO₂. From the ratio between the CO₂ and U₈O₄ the atomic weight of uranium may be calculated without involving any error due to traces of moisture possibly present in the oxalate. I subjoin Peligot's weighings, and give, in the third column, the U₈O₄ proportional to 100 parts of CO₂:

CO ₃ .	U_3O_4 .	Ratio.
1.456 grm.	4.649 grm.	319.299
1.369 "	4.412 "	322.279
2.209 "	7.084 "	320.688
1.019 "	3.279 "	321.786
1.069 "	3.447 "	322.461
1.052 "	3.389 "	322.148

Mean, 321.443, $\pm .338$

Hence $U_3O_4 = 423.342, \pm .451$.

From the acetate, $C_2H_3(UO)O_2.H_2O$, the following percentages of U_3O_4 were obtained:

5.061	grm. acetate	gave 3.354	grm. U ₃ O ₄ .	66.2715 p	er cent.
4.601	66	3.057	66	66.4421	66
1.869	• 66	1.238	46	66.2386	64
3.817	46	2.541	44	66.5706	66
10.182	66	6.757	46	66.3622	64
4.393	46	2.920	46	66.4694	44
2.868	66	1.897	44	66.1437	66

Mean, 66.3569, $\pm .038$

The acetate also yielded the subjoined percentages of carbon and of water. Assuming that the figures for carbon were calculated from known weights of dioxide, with C=12 and O=16, I have added a third column, in which the carbon percentages are converted into percentages of CO_1 :

H_2O .	Ċ.	CO ₂ .
21.60	11.27	41.323
21.16	11.30	41.433
21.10	11.30	41.433
21.20	11.10	40.700
•		
Mean, 21.265, ± .187	11.24	41.222, ± .092

From all of these figures we may calculate the molecular weight of the uranic acetate as follows:

```
From percentage of U_3O_4——C_2H_3(UO)O_2.H_2O=212.629, \pm .242

" CO_2 — " =212.999, \pm .476

" H_2O—— " =211.184, \pm 1.863

General mean " =212.685, \pm .214
```

We have now before us the molecular weights of four uranium compounds, giving us four values for U:

- (1.) $UO = 134.652, \pm .097$ Ebelmen.
- (3.) $U_8O_4 = 423.342, \pm .451$ Peligot.
- (4.) $C_2H_2(UO)O_2.H_2O = 212.685, \pm .214$

The four values for uranium combine as follows:

Or, if O = 16, U = 119.515, or 239.030.

Considering Peligot's figures by themselves, and combining values 3 and 4, we have U = 119.849, $\pm .123$; or, if O = 16, U = 120.125, or 240.250.

It is plain that the atomic weight of uranium needs to be scrupulously revised. The foregoing figures are by no means satisfactory. Chemically considered, it is probable that Peligot's work is the best, and that his results should be given preference. His figures from the oxalate and the acetate tally well with each other, whereas Ebelmen's two sets of results vary widely. From the percentage of UO yielded by the oxalate, Ebelmen's figures give a low value for U. From his oxidation of UO to U₃O₄ we get a value nearly two units higher. Peligot, in his work with the oxalate, found it, even after three or four crystallizations, to be contaminated with oxalic acid, and rejected the figures obtained from impure material. Probably Ebelmen's low values are due to the same impurity.

ALUMINUM.

The atomic weight of aluminum has been determined by Berzelius, Mather, Tissier, Dumas, Isnard, Terreil, and Mallet. The early calculations of Davy and of Thomson we may properly disregard.

Berzelius'* determination rests upon a single experiment. He ignited 10 grammes of dry aluminum sulphate, $Al_{2}(SO_{4})_{3}$, and obtained 2.9934 grammes of $Al_{2}O_{3}$ as residue. Hence, if S = 31.987 and O = 15.9633, Al = 27.243.

In 1835† Mather published a single analysis of aluminum chloride, from which he sought to fix the atomic weight of the metal. 0.646 grm. of Al_2Cl_6 gave him 2.056 of AgCl and 0.2975 of Al_2O_3 . These figures give worthless values for Al, and are included here only for the sake of completeness. From the ratio between AgCl and Al_2Cl_6 , Al = 28.925.

Tissier's ‡ determination, also resting on a single experiment, appeared in 1858. Metallic aluminum, containing .135 per cent. of sodium, was dissolved in hydrochloric acid. The solution was evaporated with nitric acid to expel all chlorine, and the residue was strongly ignited until only alumina remained. 1.935 grm. of Al gave 3.645 grm of Al₂O₃. If we correct for the trace of sodium in the aluminum, we have Al = 27.073.

Essentially the same method of determination was adopted by Isnard, who, although not next in chronological order, may fittingly be mentioned here. He found that 9 grm. of aluminum gave 27 grm. of Al_2O_3 . Hence Al = 26.938.

In 1858 Dumas, in connection with his celebrated revision of the atomic weights, made seven experiments with aluminum chloride. The material was prepared in quantity,

^{*} Poggend. Annal., 8, 177.

[†] Silliman's Amer. Journ., 27, 241.

[‡] Compt. Rend., 46, 1105.

^{||} Compt. Rend., 66, 508. 1868.

[§] Ann. Chim. Phys., (3,) 55, 151. Ann. Chem. Pharm., 113, 26.

sublimed over iron filings, and finally resublimed from metallic aluminum. Each sample used was collected in a small glass tube, after sublimation from aluminum in a a stream of dry hydrogen, and hermetically enclosed. Having been weighed in the tube, it was dissolved in water, and the quantity of silver necessary for precipitating the chlorine was determined. Reducing to a common standard, his weighings give the quantities of Al₂Cl₆ stated in the third column, as proportional to 100 parts of silver:

1.8786 gr	m. Al ₂ Cl	= 4.543 g	rm. Ag.	41.352
3.021	66	7.292	64	41.459—Bad.
2.399	"	5.802	"	41.348
1.922	66	4.6525	44	41.311
1.697	46	4.1015	44	41.375
4.3165	44	10.448	44	41.314
6.728	46	16.265	44	41.365

In the second experiment the Al₂Cl₆ contained traces of iron. Rejecting this experiment the remaining six give a mean of 41.344, $\pm .007$. Hence Al = 27.441, $\pm .082$.

In consequence of these figures of Dumas, the atomic weight of aluminum has generally of late years been put at 27.5, and the lower results deduced from the work of other investigators have been disregarded.

In 1879 Terreil* published a new determination of the atomic weight under consideration, based upon a direct comparison of the metal with hydrogen. Metallic aluminum, contained in a tube of hard glass, was heated strongly in a current of dry hydrochloric acid. Hydrogen was set free, and was collected over a strong solution of caustic potash. 0.410 grm. of aluminum thus were found equivalent to 508.2 cc., or .0455 grm. of hydrogen. Hence Al = 27.033.

About a year after Terreil's determination appeared the lower value for aluminum was thoroughly confirmed by J. W. Mallet.† After giving a full resumé of the work done by others, exclusive of Isnard, the author describes his own experiments, which may be summarized as follows:

^{*} Bulletin de la Soc. Chimique, 31, 153.

[†] Phil. Trans., 1880, p. 1003.

Four methods of determination were employed, each one simple and direct, and at the same time independent of the First, pure ammonia alum was calcined, and the residue of aluminum oxide was estimated. Second, aluminum bromide was titrated with a standard solution of silver. Third, metallic aluminum was attacked by caustic soda, and the hydrogen evolved was measured. Fourth, hydrogen was set free by aluminum, and weighed as water. Every weight was carefully verified, the verification being based upon the direct comparison, by J. E. Hilgard, of a kilogramme weight with the standard kilogramme at Washington. The specific gravity of each piece was determined, and also of all materials and vessels used in the weighings. During each weighing both barometer and thermometer were observed, so that every result represents a real weight in vacuo.

The ammonium alum used in the first series of experiments was specially prepared, and was absolutely free from ascertainable impurities. The salt was found, however, to lose traces of water at ordinary temperatures; a circumstance which tended towards a slight elevation of the apparent atomic weight of aluminum as calculated from the weighings. Two sets of experiments were made with the alum; one upon a sample air-dried for two hours at 21°-25°, the other upon material dried for twenty-four hours at 19°-26°. These sets, marked A and B respectively, differ slightly; B being the less trustworthy of the two, judged from a chemical standpoint. Mathematically it is the better of the two. Calcination was effected with a great variety of precautions, concerning which the original memoir must be To Mallet's weighings I append the percentages of Al₂O₃ deduced from them:

			Series A.	1		
8.2144	grm. of	the alum	gave .9258 g	rm. Al ₂ O ₈ .	11.270 p	er cent.
14.0378		"	1.5825	66	11.273	44
5.6201		66	.6337	66	11.275	44
11.2227		44	1.2657	64	11.278	46
10.8435		44	1.2216	44	11.266	46

Mean, 11.2724, ± .0014

Series B.

12.1023	grm. of the alum gave	1.3660	grın. Al ₂ O ₃ .	11.287 pe	er cent.
10.4544	66	1.1796	46	11.283	46
6.7962	66	.7670	44	11.286	46
8.5601	46	.9654	46	11.278	66
4.8992	46	.5528	66	11.283	"
			Mean.	11.2834.	± .001

Combined, these series give a general mean of 11.2793, \pm .0008. Hence Al = 27.075, \pm .011.

The aluminum bromide used in the second series of experiments was prepared by the direct action of bromine upon the metal. The product was repeatedly distilled, the earlier portions of each distillate being rejected, until a constant boiling point of 263.°3 at 747 mm. pressure was noted. The last distillation was effected in an atmosphere of pure nitrogen, in order to avoid the possible formation of oxide or oxy-bromide of aluminum; and the distillate was collected in three portions, which proved to be sensibly identical. The individual samples of bromide were collected in thin glass tubes, which were hermetically sealed after nearly filling. For the titration pure silver was prepared, and after fusion upon charcoal it was heated in a Sprengel vacuum in order to eliminate occluded gases. This silver was dissolved in specially purified nitric acid, the latter but very slightly in excess. The altiminum bromide, weighed in the sealed tube, was dissolved in water, precautions being taken to avoid any loss by splashing or fuming which might result from the violence of the action. To the solution thus obtained the silver solution was added, the silver being something less than a decigramme in deficiency. The remaining amount of silver needed to complete the precipitation of the bromine was added from a burette, in the form of a standard solution containing one milligramme of metal to each cubic centimetre. The final results were as follows, the figures in the third column representing the quantities of bromide proportional to 100 parts of silver. Series A is from the first portion of the last distillate of Al, Br,; series

B from the second portion, and series C from the third portion:

		Series	A.	
6.0024	grm. Al ₂ Br ₆	= 7.2793	grm. Ag	g. 82.458
8.6492	64	10.4897	66	82.454
3.1808	66	3.8573	44	82.462
		Series	B .	
6.9617	46	8.4429	46	82.456
11.2041	44	13.5897	44	82.445
3.7621	64	4.5624	66	82.459
5.2842	66	6.4085	44	82.456
9.7338	46	11.8047	66	82.457
		Series	<i>C</i> .	
9.3515	46	11.3424	66	82.447
4,4426	46	5.3877	66	82.458
5.2750	44	6.3975	"	82.454
				Mean, 82.455, \pm .001

Hence A1 = 27.046, $\pm .061$.

The high probable error of this result is due to the high probable error of the atomic weight of bromine.

The experiments to determine the amount of hydrogen evolved by the action of caustic soda upon metallic aluminum were conducted with pure metal, specially prepared, and with caustic soda made from sodium. The soda solution was so strong as to scarcely lose a perceptible amount of water by the passage through it of a dry gas at ordinary temperature. As the details of the experiments are somewhat complex, the original memoir must be consulted for them. The following results were obtained, the weight of the hydrogen being calculated from the volume, by Regnault's data corrected for the latitude and elevation of the University of Virginia:

Weight of Al.	Vol. of H.	Wt. of H.	At. Wt.
.3697 grm.	458.8 c. c.	.04106 grm.	27.012
.3769 "	467.9 "	.04187 "	27.005
.3620 "	449. I "	.04019 "	27.022
·7579 "	941.5 "	.08425 "	26.998
.7314 "	907.9 "	.08125 "	27.006
.7541 "	936.4 "	.08380 "	26.996

Mean, 27.005, ± .0032

The closing series of experiments was made with larger quantities of aluminum than were used in the foregoing The hydrogen, evolved by the action of the caustic alkali, was dried by passing it through two drying tubes containing pumice stone and sulphuric acid, and two others containing asbestos and phosphorus pentoxide. Thence it passed through a combustion tube containing copper oxide heated to redness. A stream of dry nitrogen was employed to sweep the last traces of hydrogen into the combustion tube, and dry air was afterwards passed through the entire apparatus to reoxidize the surface of reduced copper, and to prevent the retention of occluded hydrogen. The water formed by the oxidation of the hydrogen was collected in three drying tubes. The results obtained were as follows. The third column gives the amount of water formed from 10 grammes of aluminum:

```
2.1704 grm. Al gave 2.1661 grm. H<sub>2</sub>O. 9.9802

2.9355 " 2.9292 " 9.9785

5.2632 " 5.2562 " 9.9867

Mean, 9.9818, \pm .0017

Hence Al = 26.998, \pm .007.
```

In combining the various determinations of the atomic weight of aluminum into one general mean, we must arbitrarily assign weight to the single experiments of Berzelius, Isnard, Tissier, and Terreil. This may fairly be done by giving to each the probable error, and therefore the weight, of a single observation in Dumas' series. Mather's work may be ignored altogether:

If O = 16, Al = 27.075. Taking Mallet's work alone, Al = 27.0089, $\pm .0028$.

Evidently all the data except Mallet's might be rejected without affecting sensibly the final result. Dumas' work is clearly vitiated by constant errors, but the determinations by Isnard, Tissier, and Terreil may be regarded as having some confirmative value.

GOLD.

The only determinations of the atomic weight of gold which are worthy of consideration are those of Berzelius and of Levol.

The earliest method adopted by Berzelius* was that of precipitating a solution of gold chloride by means of a weighed quantity of metallic mercury. The weight of gold thus thrown down gave the ratio between the atomic weights of the two metals. In the single experiment which Berzelius publishes, 142.9 parts of Hg precipitated 93.55 of Au. Hence, using the value for mercury given in a preceding chapter, 199.712, Au = 196.113.

In a later investigation[†] Berzelius resorted to the analysis of potassio-auric chloride, 2KCl.AuCl₃. Weighed quantities of this salt were ignited in hydrogen; the resulting gold and potassium chloride were separated by means of water, and both were collected and estimated. The loss of weight upon ignition was, of course, chlorine. As the salt could not be perfectly dried without loss of chlorine, the atomic weight under investigation must be determined by the ratio between the KCl and the Au. If we reduce to a common standard, and compare with 100 parts of KCl, the equivalent amounts of gold will be those which I give in the last of the subjoined columns:

^{*} Poggend. Annal., 8, 177.

[†] Lehrbuch, 5 Aufl., 3, 1212.

```
4.1445 grm. K<sub>2</sub>AuCl<sub>5</sub> gave .8185 grm. KCl and 2.159 grm. Au. 263.775
                                                 1.172
                                                                  263.815
2.2495
                          .44425
                                                2.67225
                                                                  263.600
5.1300
                         1.01375
                                                                  263.687
                           .674
                                                 1.77725
3.4130
                          .8295
                                                                  263.773
                                                 2.188
                                        "
4-19975
```

Mean, 263.730, $\pm .026$

Hence Au = 196.186, $\pm .101$.

Still a third series of experiments by Berzelius* may be included here. In order to establish the atomic weight of phosphorus he employed that substance to precipitate gold from a solution of gold chloride in excess. Between the weight of phosphorus taken and the weight of gold obtained it was easy to fix a ratio. Since the atomic weight of phosphorus has been better established by other methods, we may properly reverse this ratio and apply it to our discussion of gold. 100 parts of P precipitate the quantities of Au given in the third column:

Mean, 1051.44, ± .196

Hence Au = 195.303, $\pm .589$.

Levol's † estimation of the atomic weight under consideration can hardly have much value. A weighed quantity of gold was converted in a flask into AuCl₃. This was reduced by a stream of sulphur dioxide, and the resulting sulphuric acid was determined as BaSO₄. One gramme of gold gave 1.782 grm. BaSO₄. Hence Au = 195.794.

If we give this single experiment and Berzelius' single result with mercury each equal weight with one analysis in the potassio-auric chloride series, and include respectively the probable errors appertaining to Hg and to BaSO₄, we may combine all the data as follows:

^{*} Lehrbuch, 5 Aufl., 3, 1188.

[†] Ann. d. Chim. et d. Phys., (3,) 30, 355. 1850.

Or, if O = 16, Au = 196.606.

As gold is a metal which can be readily applied to the determination of the atomic weights of other elements, an experimental revision of its atomic weight is very desirable.

NICKEL AND COBALT.

On account of the close similarity of these metals to each other, their atomic weights, approximately if not actually identical, have received of late years much attention.

The first determinations, and the only ones up to 1852, were made by Rothhoff;* each with but a single experiment. For nickel 188 parts of the monoxide were dissolved in hydrochloric acid; the solution was evaporated to dryness, the residue was dissolved in water, and precipitated by silver nitrate. 718.2 parts of silver chloride were thus formed; whence Ni = 58.925. The same process was applied also to cobalt, 269.2 parts of the oxide being found equivalent to 1029.9 of AgCl. Hence Co = 58.817. These values are so nearly equal that their differences were naturally ascribable to experimental errors. They are, however, entitled to no special weight at present, since it cannot be certain from any evidence recorded that the oxide of either metal was absolutely free from traces of the other.

In 1852 Erdmann and Marchand† published some results, but without details, concerning the atomic weight of nickel. They reduced the oxide by heating in a current of

^{*} Cited by Berzelius. Poggend. Annal., 8, 184. 1826.

[†] Journ. für Prakt. Chem., 55, 202. 1852.

hydrogen, and obtained values ranging from 58.2 to 58.6, when O = 16. Their results were not very concordant, and the lowest was probably the best.

In 1856, incidentally to other work, Deville* found that 100 parts of pure metallic nickel yielded 262 of sulphate; whence Ni = 59.15.

To none of the foregoing estimations can any importance now be attached. The modern discussion of the atomic weights under consideration began with the researches of Schneider † in 1857. This chemist examined the oxalates of both metals, determining carbon by the combustion of the salts with copper oxide in a stream of dry air. The carbon dioxide thus formed was collected as usual in a potash bulb, which, in weighing, was counterpoised by a similar bulb, so as to eliminate errors due to the hygroscopic character of the glass. The metal in each oxalate was estimated, first by ignition in a stream of dry air, followed by intense heating in hydrogen. Pure nickel or cobalt was left behind in good condition for weighing. Four analyses of each oxalate were made, with the results given below. The nickel salt contained three molecules of water, and the cobalt salt two molecules:

```
NiC_{2}O_{4}.3H_{2}O.
1.1945 grm. gave .528 grm. CO<sub>2</sub>.
                                                  44.203 per cent.
                  1.12625
2.5555
                                                  44.072
                  1.408
                              66
3.199
                                                  44.014
          66
                  2.214
5.020
                                                  44.104
                                           Mean, 44.098, \pm .027
```

The following percentages of nickel were found in this salt:

```
29.107
       29.082
       29.066
       29.082
Mean, 29.084, \pm .006
```

^{*} Ann. Chim. Phys., (3,) 46, 182. 1856.

[†] Poggend. Annal., 101, 387. 1857.

$CoC_2O_4.2H_2O.$

1.6355	grin. gave	.781	grm. CO ₂ .	47·753 P	er cent.
1.107	66	.5295	46	47.832	44
2.309	66	1.101	"	47.683	4 4
3.007	66	1.435	44	47.722	66
					•

Mean, 47.7475, ± .0213

The following were the percentages found for cobalt:

32.552 32.619 32.528 32.523

Mean, 32.5555, ± .0149

In a later paper * Schneider also gives some results obtained with a nickel oxalate containing but two molecules of water. This gave him 47.605 per cent. of CO₂, and the following percentages of nickel:

The conclusion at which Schneider arrived was, that the atomic weights of cobalt and nickel are not identical, being about 60 and 58 respectively. The percentages given above will be discussed at the end of this chapter in connection with all the other data relative to the constants in question.

The next chemist to take up the discussion of these atomic weights was Marignac, in 1857.† His original paper is not accessible to me, and I am therefore obliged to give only such features of it as I can get from abstracts and reviews. He worked with the chlorides and sulphates of nickel and cobalt, using apparently common gravimetric methods. The sulphates, taken as anhydrous, were first ignited to expel SO, + O, after which the residues were heated with weighed amounts of lead silicate. The increase in weight

^{*} Poggend. Annal., 107, 616.

[†] Jahresbericht, 1857, 225. Bibl. Univ. de Genève, (nouv. s.,) 1, 373.

was CoO or NiO respectively. The anhydrous chlorides were prepared from the hydrated salts by ignition in dry chlorine or hydrochloric acid. With cobalt, the monohydrated chloride, dried at 100° , was also employed. For nickel he gives the following values, referred probably to 0 = 16, S = 32, Ag = 108, Cl = 35.5:

To cobalt these values are assigned:

That is, contrary to Schneider's view, the two atomic weights are approximately the same. The values for nickel, however, run a little lower than those for cobalt; a fact which is probably not without significance. Marignac criticizes Schneider's earlier paper, holding that the nickel oxalate may have contained some free oxalic acid, and that the cobalt salt was possibly contaminated with carbonate or with basic compounds. In his later papers Schneider rejects these suggestions as unfounded, and in turn criticizes Ma-The purity of anhydrous NiSO, is not easy to guarantee, and, according to Schneider, the anhydrous chlorides of cobalt and nickel are liable to be contaminated with This is the case even when the chlorides are heated oxides. in chlorine, unless the gas is carefully freed from all traces of air and moisture.

Dumas'* determinations of the two atomic weights were made with the chlorides of nickel and cobalt. The pure metals were dissolved in aqua regia, the solutions were repeatedly evaporated to dryness, and the residual chlorides were ignited in dry hydrochloric acid gas. The last two estimations in the nickel series were made upon NiCl, formed by heating the spongy metal in pure chlorine. In the third column I give the NiCl, or CoCl, equivalent to 100 parts of silver:

^{*} Ann. Chem. Pharm., 113, 25. 1860.

```
.9123 \text{ grm. NiCl}_2 = 1.515
                                                 60.218
                               grm. Ag.
                       3.8115
                                                  60.212
2.295
                       5.464
                                                 60.212
3.290
1.830
                       3.041
                                                 60.178
3.001
                       4.987
                                                  60.176
                                          Mean, 60.1992, \pm .0062
2.352 \text{ grm. } \text{CoCl}_2 = 3.9035 \text{ grm. Ag.}
                                                  60.254
4.210
                      6.990
                                                  60.229
                      5.960
                                                  60.268
3.592
                                                   60.186
                      4.1405
2.492
                                                   60.202
4.2295
                      7.0255
                                           Mean, 60.2278, ± .011
```

These results give values for Co and Ni differing by less than a tenth of a unit; here, as elsewhere, the figure for Ni being a trifle the lower.

In 1863* the idea that nickel and cobalt have equal atomic weights was strengthened by the researches of Russell. He found that the black oxide of cobalt, by intense heating in an atmosphere of carbon dioxide, became converted into a brown monoxide of constant composition. The ordinary oxide of nickel, on the other hand, was shown to be convertible into a definite monoxide by simple heating over the blast lamp. The pure oxides of the two metals, thus obtained, were reduced by ignition in hydrogen, and their exact composition thus ascertained. Several samples of each oxide were taken, yielding the following percentages of metal:

```
NiO.
              78.597 ]
                      } ist sample.
              78.584
              78.608
             78.581 )
2-80 } 2d sample.
            78.583
             78.616
                       3d sample.
             78.590
             78.588
             78.590
             78.594
                      } 4th sample.
              78.597
             78.588
Mean of all, 78.593, \pm .0018
```

^{*} Journ. Chem. Soc., (2,) 1, 51.

```
CoO.
78.591
78.588
78.550 } 1st sample.
78.598
78.614
78.603
78.591 } 2d sample.
78.591
78.588
78.592 } 3d sample.
78.597
        4th sample.
78.598 J
78.595
        5th sample.
78.596
```

Mean of all, 78.592, $\pm .0023$

These percentages are practically identical, and lead to essentially the same mean value for each atomic weight.

In a later paper Russell* confirmed the foregoing results by a different process. He dissolved metallic nickel and cobalt in hydrochloric acid and measured the hydrogen evolved. Thus the ratio between the metal and the ultimate standard was fixed without the intervention of any other element. About two-tenths of a gramme of metal, or less, was taken in each experiment. 100 parts by weight of Co or Ni give the following weights of H, calculated from the volume of the latter:

Ni.		Co.	
3.420]	3-395	} ist sample.
3.418		3.398	
3.416		3 ·39 7	
3.417	Ist sample.	3.398 .	
3.412		3.403)
3.415		3.401	2d sample.
3.416	İ	3.401)

^{*} Journ. Chem. Soc., (2,) 7, 494. 1869.

```
      Ni.
      Co.

      3.398
      3.404
      3d sample.

      3.404
      3.405
      3d sample.

      3.401
      3d sample.
      3d sample.

      3.412
      3d sample.
      3d sample.

      3.403
      3d sample.
      3d sample.

      3.412
      3d sample.
      3d sample.

      3.403
      3d sample.
      3d sample.

      3.403
      3d sample.
      3d sample.

      3.407
      3d sample.
      3d sample.

      3.408
      3d sample.
      3d sample.

      3.407
      4d sample.
      3d sample.

      3.408
      3d sample.
      3d sample.

      3.409
      4d sample.
```

Mean of all, 3.411, ± .001

A glance at the tabulated discussion which closes this chapter will show that these figures agree well with each other, and well with those found from the analyses of the oxides. The probable errors assigned in the hydrogen series may be a little too low, since they ought to be modified by the probable error of the weight of a unit volume of hydrogen. So insignificant a correction may, however, be neglected.

Some time after the publication of Russell's first paper, but before the appearance of his second, some other investigations were made known. Of these the first was by Sommaruga,* whose results, obtained by novel methods, closely confirmed those of Schneider and antagonized those of Dumas, Marignac, and Russell. The atomic weight of nickel Sommaruga deduced from analyses of the nickel potassium sulphate, K, Ni(SO₄), 6H, O, which, dried at 100°, has a perfectly definite composition. In this salt the sulphuric acid was determined in the usual way as barium sulphate, a process to which there are obvious objections. In the third column are given the quantities of the nickel salt proportional to 100 parts of BaSO₄:

0.9798	grm. gave	1.0462	grm. BaSO ₄ .	93.653	
1.0537	46	1.1251	44	93.654	
1.0802	66	1.1535	44	93.645	
1.1865	44	1.2669	46	93.654	
3.2100	44	3.4277	"	93.649	
3.2124	66	3.4303	44	93.648	
				Mean, 93.6505, ±	± .∞1

^{*} Sitzungsb. Wien Akad., 54, 2 Abth., 50. 1866.

For cobalt Sommaruga used the purpureo cobalt chloride of Gibbs and Genth. This salt, dried at 110°, is anhydrous and stable. Heated hotter, CoCl, remains. The latter, ignited in hydrogen, yields metallic cobalt. In every experiment the preliminary heating must be carried on cautiously until ammoniacal fumes no longer appear:

```
.6656 grm. gave .1588 grm. Co.
                                              23.858 per cent.
                 .2600
                                              23.814
1.0918
                 .2160
                                              23.846
 .9058
                                              23.813
1.5895
                 .3785
                                              23.847
2.9167
                 .6957
                                              23.806
                 .4378
1.8390
2.5010
                 .5968
                                              23.808
```

Mean, 23.827, \pm .006

Further along this series will be combined with a similar one by Lee. It may here be said that Sommaruga's paper was quickly followed by a critical essay from Schneider,* endorsing the former's work, and objecting to the results of Russell.

In 1867 still another new process for the estimation of these atomic weights was put forward by Winkler, † who determined the amount of gold which pure metallic nickel and cobalt could precipitate from a neutral solution of sodioauric chloride. Experimentally, the method seems to be quite accurate; practically, it involves a knowledge of the defectively ascertained atomic weight of gold. In order to obtain pure cobalt Winkler prepared purpureo-cobalt chloride, which, having been four or five times recrystallized, was ignited in hydrogen. His nickel was repeatedly purified by precipitation with sodium hypochlorite. From material thus obtained pure nickel chloride was prepared, which, after sublimation in dry chlorine, was also reduced by hydrogen. 100 parts of gold are precipitated by the quantities of nickel and cobalt given in the third columns respectively. In the cobalt series I include one experiment

^{*} Poggend. Annal., 130, 310.

[†] Zeit. Anal. Chem., 6, 18. 1867.

by Weselsky which was published by him in a paper presently to be cited:

```
.4360 grm. nickel precipitated .9648 grm. gold.
                                                   45.191
. .4367
                                .9666
                                                   45.179
 .5189
                               1.1457
                                                   45.291
  .6002
                               1.3286
                                                  45.175
                                           Mean, 45.209, \pm .019
 .5890 grm. cobalt precipitated 1.3045 grm. gold. 45.151
 .3147
                                .6981
                                                  45.080
 . 5829
                                1.2913
                                                  45.141
 .5111
                               1.1312
                                                  45.182
                               1.2848
 .5821
                                                  45.307
                  "
                                                  45.044—Weselsky.
                               1.241
 .559
                                           Mean, 45.151, \pm .025
```

Weselsky's paper,* already cited, relates only to cobalt. He ignited the cobalticyanides of ammonium and of phenylammonium in hydrogen, and from the determinations of cobalt thus made deduced its atomic weight. His results are as follows:

```
.7575 grm. (NH<sub>4</sub>)<sub>6</sub>Co<sub>2</sub>Cy<sub>12</sub> gave .166 grm. Co.
                                                                    21.914 per cent.
                                                                     21.972
                                              .113
  .5143
                                                           Mean, 21.943, ± .029
.8529 grm. (C<sub>6</sub>H<sub>8</sub>N)<sub>6</sub>Co<sub>2</sub>Cy<sub>12</sub> gave .1010 grm. Co. 11.842 per cent.
.6112
                                                                     11.829
                                             .0723
                                             .0850
                                                                     11.905
.7140
                                                                    11.890
                                                                                  44
                                             .1120
.9420
                                                           Mean, 11.8665, \pm .0124
```

Finally, we come to the work done by Lee† in the laboratory of Wolcott Gibbs. Like Weselsky, Lee ignited certain cobalticyanides and nickelocyanides in hydrogen and determined the residual metal. The double cyanides chosen were those of strychnia and brucia; salts of very high molecular weight, in which the percentages of metal are relatively low. A series of experiments with purpureo-cobalt

^{*} Ber. d. Deutsch. Chem. Gesell., 2, 592. 1868.

[†] Am. Journ. Sci. and Arts, (3,) 2, 44. 1871.

chloride was also carried out. In order to avoid admixture of carbon in the metallic residues, the salts were first ignited in air, and then in oxygen. Reduction by hydrogen followed. The salts were in each case covered by a porous septum of earthenware, through which the hydrogen diffused, and which served to prevent the mechanical carrying away of solid particles; furthermore, heat was applied from above. The results attained were very satisfactory, and assign to nickel and cobalt atomic weights varying from each other by about a unit; Ni being nearly 58, and Co about 59. The exact figures will appear later. The cobalt results agree remarkably well with those of Weselsky. The following are the percentages of metal found:

```
In brescia nickelocyanide, Ni_3Cy_{12}(C_{23}H_{26}N_2O_4)_6H_6. so H_3O.

5.724
5.729
5.733
5.712
5.729
```

In strychnia nickelocyanide, Ni₂Cy₁₂(C₂₁H₂₂N₂O₂)₆, H₆.8H₂O.

6.607 6.613 6.589 6.607 6.561

Mean, 5.7295, ± .0034

Mean, 6.595, $\pm .005$

6.595

In brucia cobalticyanide, $Co_2Cy_{12}(C_{23}H_{26}N_2O_4)_0$. $H_6.20H_2O$.

3.759 3.720 3.739 3.748 3.747 3.749

Mean, 3.7437, $\pm .0036$

```
In strychnia cobalticyanide, Co_2Cy_{12}(C_{21}H_{22}N_2O_2)_6. H_6. SH_2O.

4.583
4.596
4.554
4.564
4.577
4.549
Mean, 4.5705, \pm.005
```

In purpureo-cobalt chloride, Co2(NH2)10Cla.

23.575
23.587
23.586
23.579
23.569
23.581

Mean, 23.5795, ± .0019

The last series may be combined with Sommaruga's, thus:

In discussing the atomic weights of nickel and cobalt, we may ignore the work of Rothhoff, Erdmann and Marchand and Deville. That of Marignac must also be omitted, for want of sufficient data. For nickel we have the following ratios. The probable error assigned in No. 4, is that of a single experiment in No. 2:

```
(1.) Per cent. of Ni in
                                NiC_{\bullet}O_{\bullet}.3H_{\bullet}O_{\bullet}.29.084, \pm .006
                     CO<sub>2</sub> from
                                                44.098, \pm .027
 (2.)
                     Ni in NiC_2O_4.2H_2O_7, 31.4076, \pm .0026
 (3.)
                     CO<sub>2</sub> from
                                                47.605, \pm .053
 (4.)
                     Ni in NiO, 78.593, \pm .0018
 (5.)
                            brucia nickelocyanide, 5.7295, ± .0034
 (6.)
 (7.)
                            strychnia
                                                      6.595, \pm .005
 (8.) Ag: NiCl<sub>2</sub>:: 100: 60.1992, \pm .0062
 (9.) Ni : H :: 100 : 3.411, ± .001
(10.) Au : Ni :: 100 : 45.209, \pm .019
(11.) BaSO_4: K_2Ni(SO_4)_2.6H_2O::100:93.6505, \pm .001
```

Since the proportion of water in the oxalates is not an absolutely certain quantity, the data concerning such salts

are best handled by employing the ratios between the carbon dioxide and the metal. Accordingly ratios (1) and (2) give a single value for Ni, and ratios (3) and (4) another. In all, we have nine values for the atomic weight in question:

If O = 16, Ni = 58.682.

In the foregoing result it will be seen that the two sets of figures due to Russell receive very great weight. This is because the one set is referred directly to hydrogen, without the intervention of the probable error of any other element; while the second set involves only the atomic weight of oxygen, of which the probable error is small. As regards accuracy of methods, however, and certainty concerning the purity of material, Russell's work is no better than Schneider's, and probably inferior to Lee's. Now values one to five in the above table represent the tolerably concordant results of Schneider, Lee, and Sommaruga. They, combined by themselves, give a general mean of Ni = 57.928, $\pm .0215$; or, if O = 16, of Ni = 58.062. This value, taking everything into account, I cannot but regard as more likely to prove correct than the larger mean deduced from all the At all events, the atomic weight of nickel needs ratios. further careful investigation.

For cobalt these ratios are available:

```
(1.) Per cent. of Co in CoC<sub>2</sub>O<sub>4</sub>.2H<sub>2</sub>O, 32.5555, ± .0149

(2.) " CO<sub>2</sub> from " 47.7475, ± .0213

(3.) " Co in CoO, 78.592, ± .0023

" purpureo-cobalt chloride, 23.6045, ± .0018

" phenylammonium cobalticyanide, 11.8665, ± .0124

(6.) " ammonium " 21.943, ± .029
```

```
(7.) Per cent. of Co in brucia cobalticyanide, 3.7437, ± .0036
(8.) " " strychnia " 4.5705, ± .005
(9.) Ag: CoCl<sub>2</sub>:: 100: 60.2278, ± .011
(10.) Co: H:: 100: 3.4017, ± .0009
(11.) Au: Co:: 100: 45.151, ± .025
```

Hence we have ten values for Co, as follows:

If O = 16, $C_0 = 59.023$.

SELENIUM.

The atomic weight of this element was first determined by Berzelius,* who, saturating 100 parts of selenium with chlorine, found that 179 of chloride were produced. Further on these figures will be combined with similar results by Dumas.

We may omit, as unimportant for present purposes, the analyses of alkaline selenates made by Mitscherlich and Nitzsch,† and pass on to the experiments published by Sacc‡ in 1847. This chemist resorted to a variety of methods, some of which gave good results, while others were unsatisfactory. First, he sought to establish the exact composition of SeO₂, both by synthesis and by analysis. The former plan, according to which he oxidized pure selenium by

^{*} Poggend. Annal., 8, 1. 1826.

⁺ Poggend. Annal., 9, 623. 1827.

[†] Ann. d. Chim. et d. Phys., (3,) 21, 119.

nitric acid, gave poor results; better figures were obtained upon reducing SeO, with ammonium bisulphite and hydrochloric acid, and determining the percentage of selenium set free:

```
.6800 grm. SeO<sub>2</sub> gave .4828 grm. Se. 71.000 per cent. 3.5227 " 2.5047 " 71.102 " 71.161 " 71.161 " Mean, 71.088, ± .032
```

In a similar manner Sacc also reduced barium selenite, and weighed the resulting mixture of barium sulphate and free selenium. This process gave discordant results, and a better method was found in calcining BaSeO₃ with sulphuric acid, and estimating the resulting quantity of BaSO₄. In the third column I give the amounts of BaSO₄ equivalent to 100 of BaSeO₃:

```
.5573 grm. BaSeO<sub>4</sub> gave .4929 grm. BaSO<sub>4</sub>. 88.444

.9942 " .8797 " 88.383

.2351 " .2080 " 88.473

.9747 " .8621 " 88.448

Mean, 88.437, ± .013
```

Still other experiments were made with the selenites of silver and lead; but the figures were subject to such errors that they need no further discussion here.

A few years after Sacc's work was published, Erdmann and Marchand made with their usual care a series of experiments upon the atomic weight under consideration.* They alalyzed pure mercuric selenide, which had been repeatedly sublimed and was well crystallized. Their method of manipulation has already been described in the chapter upon mercury. These percentages of Hg in HgSe were found:

^{*} Journ. für Prakt. Chem., 55, 202. 1852.

The next determinations were made by Dumas,* who returned to the original method of Berzelius. Pure selenium was converted by dry chlorine into SeCl₄, and from the gain in weight the ratio between Se and Cl was easily deducible. I include Berzelius' single experiment, which I have already cited, and give in a third column the quantity of chlorine absorbed by 100 parts of selenium:

```
1.709 grm. Se absorb 3.049 grm. Cl.
                                              178.409
1.810
                                              177.845
                     3.219
1.679
                                              178.856
                     3.003
                     2.688
1.498
                                              179.439
                                             178.395
                     3.468
1.944
1.887
                     3.382
                                              179.226
                                              178.398
                     3.452
1.935
                                              179.000—Berzelius.
                                       Mean, 178.696, ± .125
```

The question may here be properly asked, whether it would be possible thus to form SeCl₄ and be certain of its absolute purity? A trace of oxychloride, if simultaneously formed, would increase the apparent atomic weight of selenium. In point of fact, this method gives a higher value for Se than any of the other processes which have been adopted, and that value has the largest probable error of any one in the entire series. A glance at the table which summarizes the discussion at the end of this chapter will render this point sufficiently clear.

Latest of all, we come to the determinations made by Ekman and Pettersson.* They tried various methods of investigation, and finally decided upon the two following:

First. Pure silver selenite, Ag, SeO, was ignited, leaving behind metallic silver in the subjoined percentages:

^{*} Ann. Chem. Pharm., 113, 32. 1860.

[†] Ber. d. Deutsch. Chem. Gesell., 9, 1210. 1876. Published in detail by the society at Upsala.

62.93 62.95 62.97 62.94 62.98 62.98

Mean, 62.957, $\pm .005$

Second. A warm aqueous solution of selenious acid was mixed with HCl, and reduced by a current of SO₂. The reduced Se was collected upon a glass filter, dried, and weighed. Percentages of Se in SeO₂:

This series, combined with that of Sacc, 71.088, \pm .032, gives a general mean of 71.1907, \pm .0016.

There are now five series of figures from which to deduce the atomic weight of selenium:

- (1.) Per cent. of Se in SeO₂, 71.1907, \pm .0016
- (2.) BaSeO₃: BaSO₄:: 100: 88.437, \pm .013
- (3.) Per cent. of Hg in HgSe, 71.7327, \pm .003
- (4.) Se: SeCl₄:: 100: 178.696, \pm .125
- (5.) Per cent. of Ag in Ag_2SeO_8 , 62.957, \pm .005

From these we get the following values for selenium:

If O = 16, Se = 78.978.

TELLURIUM.

Particular interest attaches to the atomic weight of tellurium, on account of the speculations of Mendelejeff. According to the "periodic law" of that chemist, tellurium should lie between antimony and iodine, having an atomic weight greater than 120, and less than 127. Theoretically, Mendelejeff assigns it a value of Te = 125; but all the published determinations lead to a mean number higher than would be admissible under the aforesaid "periodic law." Whether theory or experiment is at fault remains to be discovered.

The first, and for many years the only, determinations of the constant in question, were made by Berzelius.* By means of nitric acid he oxidized tellurium to the dioxide, and from the increase in weight deduced a value for the metal. He published only his final results; from which, if O = 100, Te = 802.121. The three separate experiments give Te = 801.74, 801.786, and 802.838; whence we can calculate the following percentages of metal in the dioxide:

80.057 80.036 80.034

Mean, 80.042, $\pm .005$

The next determinations were made by von Hauer,† who resorted to the analysis of the well crystallized double salt TeBr. 2KBr. In this compound the bromine was estimated as silver bromide, the values assumed for Ag and Br being respectively 108.1 and 80. Recalculating, with our newer atomic weights for the above named elements, we get from v. Hauer's analyses, for 100 parts of the salt, the quantities of AgBr which are put in the third column:

^{*} Poggend. Annal., 28, 395. 1833.

[†] Sitzungsb. Wien Akad., 25, 142.

```
2.000 grm. K<sub>2</sub>TeBr<sub>4</sub> gave 69.946 per cent. Br. 164.460
6.668 " 69.8443 " 164.221
2.934 " 69.9113 " 164.379
3.697 " 70.0163 " 164.626
1.000 " 69.901 " 164.355

Mean, 164.408, ± .045
```

From Berzelius' series we may calculate Te = 128.045, and from v. Hauer's Te = 127.419. Dumas,* by a method for which he gives absolutely no particulars, found Te = 129.

In 1879, with direct reference to Mendelejeff's speculations, the subject of the atomic weight of tellurium was taken up by Wills.† The methods of both Berzelius and von Hauer were employed, with various rigid precautions in the way of testing balance and weights, and to ensure purity of material. In the first series of experiments tellurium was oxidized by nitric acid to form TeO₂. The results gave figures ranging from Te = 126.31 to 129.34:

```
2.21613 grm. Te gave 2.77612 grm. TeO<sub>2</sub>. 79.828 per cent. Te. 1.45313 " 1.81542 " 80.044 " 2.67093 " 3.33838 " 80.007 " 4.77828 " 5.95748 " 80.207 " 2.65029 " 3.31331 " 79.989 " Mean, 80.015, \pm .041
```

In the second series tellurium was oxidized by aqua regia to TeO_2 ; with results varying from Te = 127.77 to 128.00:

```
2.85011 grm. Te gave 3.56158 grm. TeO<sub>3</sub>. 80.024 per cent. Te. 3.09673 " 3.86897 " 80.040 " 5.09365 " 6.36612 " 80.012 " 80.037 " 80.037 " 80.028, ± .004
```

Combining these series with that due to Berzelius, we have the following general mean:

^{*} Ann. d. Chim. et d. Phys., (3,) 55, 129. 1859.

[†] Journ. Chem. Society, Oct., 1879, p. 704.

```
Berzelius______ 80.042, ± .005
Wills, 1st series_____ 80.015, ± .041
" 2d " 80.028, ± .004
General mean_____ 80.035, ± .003
```

Hence Te = 127.986, $\pm .035$.

By von Hauer's process, the analysis of TeBr₄.2KBr, Will's figures give results ranging from Te = 126.07 to 127.61. Reduced to a common standard, 100 parts of the salt yield the quantities of AgBr given in the third column:

1.70673 gm	n. K ₂ TeBr _e gave	2.80499	grm. AgBr.	164.349
1.75225	66	2.88072	66	164.398
2.06938	66	3.40739	44	164.657
3.29794	"	5.43228	66	164.717
2.46545	66	4.05742	66	164.571
			Mean,	164.538, ± .048

Combined with von Hauer's mean, 164.408, \pm .045, this gives a general mean of 164.468, \pm .033. Hence Te = 127.170, \pm .173.

The two independent values for Te combine thus:

If O = 16, Te = 128.254.

A careful consideration of the foregoing figures, and of the experimental methods by which they were obtained, will show that they are not absolutely conclusive with regard to the place of tellurium under the periodic law. The atomic weight of iodine, calculated in a previous chapter, is 126.557. Wills' values for Te, rejecting his first series as relatively unimportant, range from 126.07 to 128.00; that is, some of them fall below the atomic weight of iodine, although none descend quite to the 125 assumed by Mendelejeff.

In considering the experimental methods, reference may properly be made to the controversy regarding the atomic weight of antimony. It will be seen that Dexter, estimating the latter constant by the conversion of the metal into Sb₂O₄, obtained a value approximately of Sb = 122. Dumas, working with SbCl₃, obtained a similar value. Schneider and Cooke, on the other hand, have established an atomic weight for antimony near 120, and Cooke in particular has traced out the constant errors which lurked unsuspected in the work of Dumas and Dexter. Now in some physical respects tellurium and antimony are quite similar. As constant errors vitiated the recently accepted values for Sb, so they may also effect our estimates for Te. The oxidation of Te by nitric acid resembles in minor particulars that of Sb. The analysis of K₂TeBr₆, gives a low value for Te, and yet the material may have contained traces of oxybromides, the presence of which would render even that lower value too high. A careful revision of the atomic weight of tellurium is still necessary.

VANADIUM.

Roscoe's determination of the atomic weight of vanadium is the only one having any present value. The results obtained by Berzelius* and by Czudnowicz † are unquestionably too high; the error being probably due to the presence of phosphoric acid in the vanadic acid employed. This particular impurity, as Roscoe has shown, prevents the complete reduction of V₂O₅ to V₂O₅ by means of hydrogen. All vanadium ores contain small quantities of phosphorus, which can only be detected with ammonium molybdate; a reaction unknown in Berzelius' time. Furthermore, the complete purification of vanadic acid from all traces of phosphoric acid is a matter of great difficulty, and probably never was accomplished until Roscoe undertook his researches.

In his determination of the atomic weight, Roscoe‡

^{*} Poggend. Annal., 22, 14. 1831.

[†] Poggend. Annal., 120, 17. 1863.

[‡] Journ. Chem. Soc., 6, pp. 330 and 344. 1868.

studied two compounds of vanadium; namely, the pentoxide, V₂O₅ and the oxychloride, VOCl₃. The pentoxide, absolutely pure, was reduced to V,O, by heating in hydrogen, with the following results:

```
17.533 per cent. of loss.
7.7397 grm. V<sub>2</sub>O<sub>5</sub> gave 6.3827 grm. V<sub>2</sub>O<sub>5</sub>.
                                                                            17.507
6.5819
                                 5.4296
                                                                            17.489
5.1895
                                  4.2819
                                 4.1614
                                                                            17.515
5.0450
5.4296 grm. V<sub>2</sub>O<sub>3</sub>, reoxidized, gave 6.5814 grm. V<sub>2</sub>O<sub>5</sub>. 17.501 per cent. difference.
```

Mean, 17.509, ± .005

Hence $V = 51.264, \pm .025$.

Upon the oxychloride, VOCl_a, two series of experiments were made, one volumetric, the other gravimetric. In the volumetric series the compound was titrated with solutions containing known weights of silver, which had been purified according to the methods recommended by Stas. publishes his weighings, and gives percentages deduced from them; his figures, reduced to a common standard, make the quantities of VOCl, given in the third column proportional to 100 parts of silver. He was assisted by two analysts:

Analyst A.

2.4322	grm. VOCl ₃ =	= 4.5525	grm. Ag.	53.425
4.6840	44	8.7505	44	53.528
4.2188	44	7.8807	44	53-533
3.9490	44	7-3799	66	53.510
.9243	46	1.7267	64	53.5 3 0
1.4330	46	2.6769	66	53.532
		Analyst	B .	
2.8530	46	5.2853	66	53.980
2.1252	46	3.9535	46	53.755
1.4248	46	2.6642	46	53-479

Mean, 53.586, \pm .039

The gravimetric series, of course, fixes the ratio between VOCl, and AgCl. If we put the latter at 100 parts, the proportion of VOCl_a comes out as given in the third column:

Analyst A.

1.8521	gnm. VOCl ₃ gave	4.5932	grm. A	AgCl.	40.323	
.7013	66	1.7303	46		40.531	
.7486	**	1.8467	66		40.537	
1.4408	**	3.5719	66		40.337	
·9 4 53	"	2.3399	66		40.399	
1.6183	44	4.0282	66		40.174	
		Analysi	<i>B</i> .			
2.1936	44	5.4039	66		40.391	
2.5054	66	6.2118	44		40.3 3 3	
				Mean,	40.378,	± .028

These two series give us two values for the molecular weight of VOCl₃:

From the volumetric series ___ VOCl₂ = 173.096,
$$\pm$$
 .126
" gravimetric " ___ " = 173.276, \pm .141
General mean____ " = 173.177, \pm .094

Hence $V = 51.104, \pm .104$.

Combining the two values for V we get the following result:

Or, if O = 16, V = 51.373.

ARSENIC.

For the determination of the atomic weight of arsenic two compounds have been studied; the chloride and the trioxide. The bromide may also be considered, since it was analyzed by Wallace in order to establish the atomic weight of bromine. His series, in the light of more recent knowledge, may properly be inverted, and applied to the determination of arsenic.

In 1826, Berzelius* heated arsenic trioxide with sulphur

^{*} Poggend. Annal., 8, 1.

in such a way that only SO, could escape. 2.203 grammes of As_2O_3 , thus treated, gave a loss of 1.069 of SO. Hence As = 74.840. This is a close estimation; but, being drawn from a single experiment, has so little weight that it need not be included in our final general mean.

In 1845 Pelouze* applied his method of titration with known quantities of pure silver to the analysis of the trichloride of arsenic, $AsCl_s$. Using the old Berzelian atomic weights, and putting Ag = 1349.01, and Cl = 443.2, he found in three experiments for As the values 937.9, 937.1, and 937.4. Hence 100 parts of silver balance the following quantities of $AsCl_s$:

Later, the same method was employed by Dumas,† whose weighings, reduced to the foregoing standard, give the following results:

```
4.298 grm. AsCl<sub>2</sub> = 7.673 grm. Ag. Ratio, 56.015

5.535 " 9.880 " 56.022

7.660 " 13.686 " 55.970

4.680 " 8.358 " 55.993

Mean, 56.000, ± .008
```

The two series of Pelouze and Dumas, combined, give a general mean of 56.014, $\pm .0035$, as the amount of AsCl, equivalent to 100 parts of silver. Hence As = 74.829, $\pm .048$, a value closely agreeing with that deduced from the single experiment of Berzelius.

The same process of titration with silver was applied by Wallace‡ to the analysis of arsenic tribromide, AsBr₃. This compound was repeatedly distilled to ensure purity, and was well crystallized. His weighings show that the quanti-

^{*} Compt. Rend., 20, 1047.

[†] Ann. Chim. Phys., (3,) 55, 174. 1859.

[†] Philosophical Magazine, (4,) 18, 279.

ties of bromide given in the third column are proportional to 100 parts of silver:

```
8.3246 grm. AsBr<sub>8</sub> = 8.58 grm. Ag. 97.023

4.4368 " 4.573 " 97.022

5.098 " 5.257 " 96.970

Mean, 97.005, \pm .012
```

Hence As = 74.046, $\pm .058$. Why this value should be so much lower than that from the chloride is unexplained.

The volumetric work done by Kessler,* for the purpose of establishing the atomic weights of chromium and of arsenic, has already been described in the chromium chapter. In that investigation the amount of potassium dichromate required to oxidize 100 parts of As₂O₃ to As₂O₅ was determined, and compared with the quantity of potassium chlorate necessary to produce the same effect. From the molecular weight of KClO₃, that of K₂Cr₂O₇ was then calculable.

From the same figures, the molecular weights of KClO₃ and of K₂Cr₂O₇ being both known, that of As₂O₃ may be easily determined. The quantities of the other compounds proportional to 100 parts of As₂O₃ are as follows:

$K_2Cr_2O_7$	KClO ₃ .
98.95	41.156
98.94	41.116
99.17	41.200
98.98	41.255
99.08	41.201
99.15	41.086
	·41.199
Mean, 99.045, ± .028	41.224
•	41.161
	41.193
	41.149
	41.126
	Mean, 41.172, ± .009

^{*} Poggend. Annal., 95, 204. 1855. Also 113, 134. 1861.

Another series with the bichromate gave the following figures:

99.08 99.06 99.10 98.97 98.97 98.97 Mean, 99.036, ± .019 Mean of previous series, 99.045, ± .028 General mean, 99.039, ± .016

Other defective series are given to illustrate the partial oxidation of the As₂O₃ by action of air. The foregoing figures give us two distinct values for the molecular weight of As₂O₃. In calculating from the bichromate results the value for chromium deduced from Siewert's determinations will be used, viz., Cr = 52.009, $\pm .025$.

Hence As = 75.002, $\pm .018$.

The general mean for As comes out as follows:

If O = 16, then As becomes = 75.090.

ANTIMONY.

After some earlier, unsatisfactory determinations, Berzelius,* in 1826, published his final estimation of the atomic weight of antimony. He oxidized the metal by means of nitric acid, and found that 100 parts of antimony gave 124.8 of Sb_2O_4 . Hence, if O=16, Sb=129.03. The

^{*} Poggend. Annal., 8, 1.

value 129 remained in general acceptance until 1855, when Kessler,* by special volumetric methods, showed that it was certainly much too high. Kessler's results will be considered more fully further along, in connection with a later paper; for present purposes a brief statement of his earlier conclusions will suffice. Antimony, and various compounds of antimony, were oxidized partly by potassium anhydrochromate and partly by potassium chlorate; and from the amounts of oxidizing agent required, the atomic weight in question was deduced:

The figures given are those calculated by Kessler himself. A recalculation with our newer atomic weights for O, K, Cl, Cr, S, and C, would yield slightly lower values. It will be seen that five of the estimates agree closely, while one diverges widely from the others. It will be shown hereafter that the concordant values are all vitiated by constant errors, and that the exceptional figure is after all the best.

Shortly after the appearance of Kessler's first paper, Schneider † published some results obtained by the reduction of antimony sulphide in hydrogen. The material chosen was a very pure stibnite from Arnsberg, of which the gangue was only quartz. This was corrected for, and corrections were also applied for traces of undecomposed sulphide carried off mechanically by the gas stream, and for traces of sulphur retained by the reduced antimony. The latter sulphur was estimated as barium sulphate. From 3.2 to 10.6 grammes of material were taken in each experiment. The final corrected percentages of S in Sb₂S₃ were as follows:

^{*} Poggend. Annal., 95, 215.

[†] Poggend. Annal., 98, 293. 1856. Preliminary note in Bd. 97.

28.559
28.557
28.501
28.554
28.532
28.485
28.492
28.481
———
Mean, 28.520, ± .008

Hence, if S = 32, Sb = 120.3.

Immediately after the appearance of Schneider's memoir, Rose * published the result of a single analysis of antimony trichloride, previously made under his supervision by Weber. This analysis, if Cl = 35.5, makes Sb = 120.7, a value of no great weight, but in a measure confirmatory of that obtained by Schneider.

The next research upon the atomic weight of antimony was that of Dexter,† published in 1857. This chemist, having tried to determine the amount of gold precipitable by a known weight of antimony, and having obtained discordant results, finally resorted to the original method of Berzelius. Antimony, purified with extreme care, was oxidized by nitric acid, and the gain in weight was determined. From 1.5 to 3.3 grammes of metal were used in each experiment. The reduction of the weights to a vacuum standard was neglected as being superfluous. From the data obtained, we get the following percentages of Sb in Sb₂O₄:

79.268 79.272 79.255 79.266 79.253 79.271 79.264 79.260 79.286

^{*} Poggend. Annal., 98, 455. 1856.

[†] Poggend. Annal., 100, 563.

79.274 79.232 79.395 79.379 Mean, 79.283, ± .009

Hence, if O = 16, Sb = 122.46.

The determinations of Dumas* were published in 1859. This chemist sought to fix the ratio between silver and antimonicus chloride, and obtained results for the atomic weight of antimony quite near to those of Dexter. SbCl₃ was prepared by the action of dry chlorine upon pure antimony; it was distilled several times over antimony powder, and it seemed to be perfectly pure. Known weights of this preparation were added to solutions of tartaric acid in water, and the silver chloride was precipitated without previous removal of the antimony. Here, as Cooke has since shown, is a possible source of error, for under such circumstances the crystalline argento-antimonious tartrate may also be thrown down and contaminate the chloride of silver. But be that as it may; Dumas' weighings, reduced to a common standard, give as proportional to 100 parts of silver, the quantities of SbCl_s which are stated in the third of the subjoined columns:

1.876 gr	m. SbCl ₈	= 2.660 g	rm. Ag.	70.526
4.336	46	6. 148	46	70.527
5.065	44	7.175	66	70.592
3-475	•6	4.930	46	70.487
3.767	66	5.350	66	70.411
5.910	"	8.393	44	70.416
4.828	66	6.836	44	70.626

Mean, 70.512, ± .021

Hence, if Ag = 108, and Cl = 355, Sb = 122.

In 1861 Kessler's second paper † relative to the atomic weight of antimony appeared. Kessler's methods were somewhat complicated, and for full details the original memoirs must be consulted. A standard solution of potassium anhydrochromate was prepared, containing 6.1466

^{*} Ann. Chim. Phys., (3,) 55, 175.

[†] Poggend. Annal., 113, 145.

grammes to the litre. With this, solutions containing known quantities of antimony or of antimony compounds were titrated, the end reaction being adjusted with a standard solution of ferrous chloride. In some cases the titration was preceded by the addition of a definite weight of potassium chlorate, insufficient for complete oxidation; the anhydrochromate then served to finish the reaction. The object in view was to determine the amount of oxidizing agent, and therefore of oxygen, necessary for the conversion of known quantities of antimonious into antimonic compounds.

In the later paper Kessler refers to his earlier work, and shows that the values then found for antimony were all too high, except in the case of the series made with tartar emetic. That series he merely states, and subsequently ignores, evidently believing it to be unworthy of further consideration. For the remaining series he points out the sources of error. These need not be rediscussed here, as the discussion would have no value for present purposes; suffice it to say that in the series representing the oxidation of Sb_*O_* with anhydrochromate and chlorate, the material used was found to be impure. Upon estimating the impurity and correcting for it, the earlier value of Sb = 123.80 becomes Sb = 122.36, according to Kessler's calculations.

In the paper now under consideration four series of results are given. The first represents experiments made upon a pure antimony trioxide which had been sublimed, and which consisted of shining colorless needles. This was dissolved, together with some potassium chlorate, in hydrochloric acid, and titrated with anhydrochromate solution. Six experiments were made, but Kessler rejects the first and second as untrustworthy. The data for the others are as follows:

Sb_2O_3 .	KClO3.	K ₂ Cr ₂ O ₇ sol. in cc.
1.7888 grm.	.4527 grm.	19.2 cc.
1.6523 "	.4506 "	3.9 "
3.2998 "	.8806 "	16.5 "
1.3438 "	.3492 "	10.2 "

From these figures Kessler deduces Sb = 122.16.

These data, reduced to a common standard, give the following quantities of oxygen needed to oxidize 100 parts of Sb_2O_3 to Sb_2O_5 . Each cubic centimetre of the $K_2Cr_2O_7$ solution corresponds to one milligramme of O:

In the second series of experiments pure antimony was dissolved in hydrochloric acid with the aid of an unweighed quantity of potassium chlorate. The solution, containing both antimonious and antimonic compounds, was then reduced entirely to the antimonious condition by means of stannous chloride. The excess of the latter was corrected with a strong hydrochloric acid solution of mercuric chloride, then, after diluting and filtering, a weighed quantity of potassium chlorate was added, and the titration with anhydrochromate was performed as usual. Calculated as above, the percentages of oxygen given in the last column correspond to 100 parts of antimony:

Sb.	KClO ₃ .	K2Cr2O7 sol. cc.	Per cent. O.
1.636 grm.	0.5000 grm.	18.3	13.088
3.0825• "	0.9500 "	30.2	13.050
4.5652 "	1.4106 "	45-5	13.098
-			
		**	

Mean, 13.079, ± .0096

This series gave Kessler Sb = 122.34.

The third and fourth series of experiments were made with pure antimony trichloride, SbCl₃, prepared by the action of mercuric chloride upon metallic antimony. This preparation, in the third series, was dissolved in hydrochloric acid, and titrated. In one experiment solid K₂Cr₂O₇ in weighed amount was added before titration: in the other two estimations KClO₃ was taken as usual. If, according to Siewert's work, we take Cr = 52.009, the percentages of oxygen in the last column correspond to 100 parts of SbCl₃:

							F	Per cent. O.
1.8576 gm	n. SbCl ₃	needed . 5967	grm.	K ₂ Cr ₂ O ₇	and	33-4	cc. sol.	7.0338
1.9118	66	.3019	"	KClO ₃	"	16.2	44	7.0321
4.1235	46	.6801	64	44	44	23.2	44	7.0222
			•	_			Mean,	7.0294, ± .0024

The fourth set of experiments was gravimetric. The solution of SbCl₃, mixed with tartaric acid, was first precipitated by hydrogen sulphide, in order to remove the antimony. The excess of H₂S was corrected by copper sulphate, and then the chlorine was estimated as silver chloride in the ordinary manner. 100 parts of AgCl correspond to the amounts of SbCl₃ given in the third column.

1.8662 gm	m. SbCl ₃ gave	3.483 gr	m. AgCl.	53.580
1.6832	46	3.141	66	53.588
2.7437	4.6	5.1115	66	53.677
2.6798	44	5.0025	44	53.569
5.047	46	9.411	46	53.629
3.8975	44	7.2585	••	53.696
			3	Mean, 53.623 , $\pm .015$

The volumetric series with SbCl₃ gave Kessler values for Sb ranging from 121.16 to 121.47. The gravimetric series, on the other hand, yielded results from Sb = 124.12 to 124.67. This discrepancy Kessler rightly attributes to the presence of oxygen in the chloride; and, ingeniously correcting for this error, he deduces from both sets combined, the value of Sb = 122.37.

The several mean results for antimony agree so fairly with each other, and with the estimates obtained by Dexter and Dumas, that we cannot wonder that Kessler felt satisfied of their general correctness, and of the inaccuracy of the figures published by Schneider. Still, the old series of data obtained by the titration of tartar emetic with anhydrochromate contained no evident errors, and was not accounted for. This series,* if we reduce all of Kessler's figures to a single common standard, give a ratio between K₂Cr₂O₃ and C₄H₄KSbO₇. H₂O. 100 parts of the former will oxidize of the latter:

^{*} Poggend. Annal., 95, 217.

336.64 338.01 336.83 337.93 338.59 335.79

Mean, 337.30, ± .29

From this, if $K_2Cr_2O_7 = 294.64$, Sb = 119.8.

The newer atomic weights found in the previous chapters of this work will be applied to the discussion of all these series further along. It may, however, be properly noted at this point, that the probable errors assigned to the percentages of oxygen in three of Kessler's series are too low. These percentages are calculated from the quantities of KClO₃ involved in the several reactions, and their probable errors should be increased with reference to the probable error of the molecular weight of that salt. The necessary calculations would be more laborious than the importance of the figures would warrant, and, accordingly, in computing the final general mean for antimony, Kessler's figures will receive somewhat higher weight than they are legitimately entitled to.

Naturally, the concordant results of Dexter, Kessler, and Dumas led to the general acceptance of the value of 122 for antimony as against the lower figure 120 of Schneider. Still, in 1871, Unger* published the results of a single analysis of Schlippe's salt, Na₈SbS₄.9H₂O. This analysis gave Sb = 119.76, if S = 32 and Na = 23, but no great weight could be attached to the determination. It served, nevertheless, to show that the controversy over the atomic weight of antimony was not finally settled.

More than ten years after the appearance of Kessler's second paper the subject of the atomic weight of antimony was again taken up, this time by Professor Cooke. His results appeared in the autumn of 1877,† and were conclusive in favor of the lower value, approximately 120. For full

^{*} Archiv. der Pharmacie, 197, 194. Quoted by Cooke.

[†] Proceedings American Academy, v. 13.

details the original memoir must be consulted; only a few of the leading points can be cited here.

Schneider analyzed a sulphide of antimony which was already formed. Cooke, reversing the method, effected the synthesis of this compound. Known weights of pure antimony were dissolved in hydrochloric acid containing a little nitric acid. In this solution weighed balls of antimony were boiled until the liquid became colorless; subsequently the weight of metal lost by the balls was ascertained. To the solution, which now contained only antimonious compounds, tartaric acid was added, and then, with a supersaturated aqueous sulphhydric acid, antimony trisulphide was precipitated. The precipitate was collected by an ingenious process of reverse filtration, converted into the black modification by drying at 210°, and weighed. After weighing, the Sb, S, was dissolved in hydrochloric acid, leaving a carbonaceous residue unacted upon. This was carefully estimated and corrected for. About two grammes of antimony were taken in each experiment and thirteen syntheses were In two of these, however, the antimony triperformed. sulphide was weighed only in the red modification, and the results were uncorrected by conversion into the black variety and estimation of the carbonaceous residue. In fact, every such conversion and correction was preceded by a weighing of the red modification of the Sb,S. The mean result of these weighings, if S = 32, gave Sb = 119.994. The mean result of the corrected syntheses gave Sb = 120.295. In these eleven experiments the following percentages of S in Sb₂S₃ were established:

```
28.57

28.60

28.57

28.43

28.42

28.53

28.50

28.49

28.58

28.50

28.50

28.51

Mean, 28.5182, ± .0120
```

These results, confirmatory of the work of Schneider, were presented to the American Academy in 1876. Still, before publication, Cooke thought it best to repeat the work of Dumas, in order to detect the cause of the old discrepancy between the values Sb = 120 and Sb = 122. Accordingly, various samples of antimony trichloride were taken, and purified by repeated distillations. The final distillate was further subjected to several recrystallizations from the fused state; or, in one case, from a saturated solution in bisulphide of carbon. The portions analyzed were dissolved in concentrated aqueous tartaric acid, and precipitated by silver nitrate, many precautions being observed. The silver chloride was collected by reverse filtration, and dried at temperatures from 110° to 120°. In one experiment the antimony was first removed by H₂S. Seventeen experiments were made, giving, if Ag = 108 and Cl = 35.5, a mean value of Sb = 121.94. If we reduce to a common standard, Cooke's analyses give, as proportional to 100 parts of AgCl, the quantities of SbCl_n stated in the third column:

```
1.5974 grm. SbCl, gave 3.0124 grm. AgCl.
                                               53.028
                         2.3620
1.2533
                                               53.061
 .8876
                         1.6754
                                               52.978
 .8336
                         1.5674
                                               53.184
 .5326
                         1.0021
                                               53.148
 .7270
                         1.3691
                                               53.101
1.2679
                         2.3883
                                               53.088
                         3.6646
1.9422
                                               52.999
                         3.3384
1.7702
                                               53.025
2.5030
                         4.7184
                                               53.048
2.1450
                         4.0410
                                               53.081
1.7697
                         3.3281
                                     46
                                               53.175
2.3435
                                               53.072
                         4.4157
1.3686
                         2.5813
                                    46
                                               53.020
1.8638
                         3.5146
                                               53.030
2.0300
                         3.8282
                                               53.028
                         4.6086
2.4450
                                               53.053
                                        Mean, 53.066, \pm .0096
```

This mean may be combined with that of Kessler's series, as follows:

General mean	53.2311.	** .008
Cooke	53.066,	± .0096
Kessler	53.623,	± .015

The results thus obtained with SbCl, confirmed Dumas' determination of the atomic weight of antimony as remarkably as the syntheses of Sb,S, had sustained the work of Schneider. Evidently, in one or the other series a constant error must be hidden, and much time was spent by Cooke in searching for it. It was eventually found that the chloride of antimony invariably contained traces of oxychloride, an impurity which tended to increase the apparent atomic weight of the metal under consideration. If was also found, in the course of the investigation, that hydrochloric acid solutions of antimonious compounds oxidize in the air during boiling as rapidly as ferrous compounds; a fact which explains the high values for antimony found by Kessler.

In order to render "assurance doubly sure," Professor Cooke also undertook the analysis of the bromide and the iodide of antimony. The bromide, SbBr₃, was prepared by adding the finely powdered metal to a solution of bromine in carbon disulphide. It was purified by repeated distillation over pulverized antimony, and by several recrystallizations from bisulphide of carbon. The bromine determinations resembled those of chlorine, and gave, if Ag = 108 and Br = 80, a mean value for antimony of Sb = 120. Reduced to a common standard, the fifteen analyses give the subjoined quantities of SbBr₃ proportional to 100 parts of silver bromide:

1.8621	grm. SbBr, gave	2.9216	grm. AgBr.	63.736
.9856	66	1.5422	46	63.909
1.8650	66	2.9268	66	63.721
1.5330	"	2.4030	66	63.795
1.3689	44	2.1445	66	63.833
1.2124	66	1.8991	46	63.841
.9417	66	1.4749	**	63.848
2.5404	44	3.9755	66	63.901
1.5269	66	2.3905	46	63.874
1.8604	66	2.9180	64	63.756
1.7298	44	2.7083	66	63.870

```
3.2838 grm. SbBr<sub>8</sub> gave 5.1398 grm. AgBr. 63.890
2.3589 " 3.6959 " 63.825
1.3323 " 2.0863 " 63.859
2.6974 " 4.2285 " 63.791

Mean, 63.830, ± .008
```

The iodide of antimony was prepared like the bromide, and analyzed in the same way. At first, discordant results were obtained, due to the presence of oxyiodide in the iodide studied. The impurity, however, was removed by subliming the iodide in an atmosphere of dry carbon dioxide. With this purer material, seven estimations of iodine were made, giving, if Ag = 108 and I = 127, a value for antimony of Sb = 120. Reduced to a uniform standard, Cooke's weighings give the following quantities of SbI_3 proportional to 100 parts of silver iodide:

1.1877	grm. SbI ₃ gave	1.6727	grm. AgI.	71.005
.4610	44	.6497	46	70.956
3.2527	44	4.5716	44	71.150
1.8068	64	2.5389	44	71.165
1.5970	66	2.2456	66	71.117
2.3201	66	3.2645	44	71.071
.3496	"	.4927	• •	7 0.956

Mean, 71.060, \pm .023

Although Cooke's work was practically conclusive, as between the rival values for antimony, his results were severely criticized by Kessler,* who, evidently, had read Cooke's paper in a very careless way. On the other hand, Schneider published in Poggendorff's Annalen a friendly review of the new determinations, which so splendidly vindicated his own accuracy. In reply to Kessler, Cooke undertook still another series of experiments with antimony bromide,† and obtained absolute confirmation of his previous results. To a solution of antimony bromide was added a solution containing a known weight of silver not quite sufficient to precipitate all the bromine. The excess

^{*} Berichte d. Deutsch. Chem. Gesell., 12, 1044. 1879.

[†] Amer. Journ. Sci. and Arts, May, 1880. Berichte, 13, 951.

The r
determi
ably as
Schneid
error m
in searc
ride of;
an imp
weight
in the
solution
ing boi
explair
In o
Cooke;

iodide.

adding

in cark

tion ov

tions fi

tions r

and B

Reduc

the sul

of silv

I

I

I

I

of the latter was estimated by titration with a normal silver solution. Five analyses gave values for antimony ranging from 119.98 to 120.02, when Ag = 108 and Br = 80. Reduced to a common standard, the weights obtained gave the amounts of SbBr, stated in the third column as proportional to 100 parts of silver:

2.5032	grm. SbBr ₃ =	2.2528	grm. Ag.		111.115
2.0567	и	1.8509	46		111.119
2.6512	44	2.3860	44		111.115
3.3053	44	2.9749	44		111.106
2.7495	46	2.4745	**		111.113
				Mean,	111.114, ± .0014

Schneider,* also, in order to more fully answer Kessler's objections, repeated his work upon the Arnsberg stibnite. This he reduced in hydrogen as before, correcting scrupulously for impurities. The following percentages of sulphur were found:

These figures confirm his old results, and may be fairly combined with them and with the percentages found by Cooke, as follows:

```
Schneider, early series 28.520, ± .008

" late " 28.541, ± .0024

Cooke 28.5182, ± .0120

General mean 28.5385, ± .0023
```

We have now before us the following ratios, good and bad, from which to calculate the atomic weight of antimony. The single results obtained by Weber and by Unger, being unimportant, are not included:

	" §	Sb in Sb.O.	28.5385, ± .0023 4, 79.283, ± .009
O nec	ded to oxi	idize 100 l	parts SbCl ₂ , 7.0294, ± .0024
0	66	46	Sb_2O_2 , 10.953, \pm .0075
0	44	"	Sb, 13.079, ± .0096

* Journ. für Prakt. Chem., (2,) 22, 131.

```
(6.) K<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub>: tartar emetic:: 100: 337.30, ± .29
(7.) Ag: SbCl<sub>3</sub>:: 100: 70.512, ± .021
(8.) AgCl: SbCl<sub>3</sub>:: 100: 53.2311, ± .008
(9.) Ag: SbBr<sub>3</sub>:: 100: 111.114, ± .0014
(10.) AgBr: SbBr<sub>3</sub>:: 100: 63.830, ± .008
. (11.) AgI: SbI<sub>3</sub>:: 100: 71.060, ± .023
```

Three of these ratios give estimates for the molecular veight of antimony trichloride, and two give corresponding values for the bromide. These values may be combined, as follows: First, for the chloride we have—

From (3)-----SbCl₃ = 227.094,
$$\pm$$
 .115
" (7)----- " = 227.771, \pm .091
" (8)---- " = 228.433, \pm .039
General mean " = 228.225, \pm .034

Hence Sb = 122.115, $\pm .055$.

For the bromide we get:

: 13

Hence Sb = 119.625, $\pm .063$.

From all the data eight values for Sb may be deduced. These fall into two groups; the one near the number 120, the other not far from 122. In making the calculation the atomic weights found in previous chapters are applied; the value selected for chromium being that deduced from Siewert's experiments:

```
1. From Sb_2S_3, ratio (1)_____Sb = 120.145, \pm .045
      SbBr_8 = 119.625, \pm .063
      SbI_3, ratio (11)----- " = 119.665, \pm .179
      tartar emetic, ratio (6) = 118.690, \pm .278
      Sb_2O_4, ratio (2) ---- " = 122.181, \pm .061
5.
      6.
                                            High.
      ratio (4) = 121.798, \pm .105
7.
       " (5) = 122.053, \pm .094
8.
    General mean ____ " = 121.027, ± .025
    General mean of values 1 to 4_- " = 119.935, ± .036
                    5 " 8 ... " = 122.092, ± .035
```

Although the means of the four lower values and of the four higher values are thus shown to be approximately

equal in weight, we know from Cooke's experiments that the larger mean is vitiated by serious constant errors. Only in value 5, the result calculated from Dexter's experiments, has the constant error not been pointed out. Cooke considers it probable, however, that the Sb₂O₄ involved in this work contained traces of some lower oxide, which, if present, would render the atomic weight of antimony apparently too high. Chemically considered, the preponderance of evidence is strongly in favor of values 1 to 3, deduced from the experiments of Schneider and of Cooke. These give a general mean of Sb = 119.955, $\pm .036$; or, if O = 16, this becomes Sb = 120.231.

This we may accept as most nearly the true result, and reject the data of Dexter, Dumas, and Kessler altogether.

Since this chapter was written, Pfeifer has compared the amount of antimony thrown down electrolytically, with the quantity of silver deposited by the same current in the same time. From rather meagre data he concludes that the atomic weight of antimony, thus determined, may be 121. Additional investigation is promised. The figures thus far published would weigh little as against Cooke's experiments. (Ann. Chem. Pharm., 209, 161. 1881.)

BISMUTH.

Early in the century the combining weight of bismuth was approximately fixed through the experiments of Lagerhjelm.* Effecting the direct union of bismuth and sulphur, he found that ten parts of the metal yield the following quantities of trisulphide:

^{*} Annals of Philosophy, 4, 358. 1814. Results adopted by Berzelius.

Hence B=215 in round numbers, a value now known to be much too high. Lagerhjelm also oxidized bismuth with nitric acid, and, after ignition, weighed the trioxide thus formed. Ten parts of metal gave the following quantities of Bi_2O_3 :

11.1382 11.1275 ———— Mean, 11.13285

Hence, if O = 16, Bi = 211.85, a figure still too high.

In 1851 the subject of the atomic weight of bismuth was taken up by Schneider,* who, like Lagerhjelm, studied the oxidation of the metal with nitric acid. The work was executed with a variety of experimental refinements, by means of which every error due to possible loss of material was carefully avoided. For full details the original paper must be consulted; there is only room in these pages for the actual results, as follows. The figures represent the percentages of Bi in Bi, O₃:

89.652 89.682 89.644 89.634 89.656 89.666 89.653

Mean, 89.6552, ± .0034

Hence Bi = 207.523, $\pm .082$; or, if O = 16, Bi = 208.001.

Finally, we come to the results obtained by Dumas.† Bismuth trichloride was prepared by the action of dry chlorine upon bismuth, and repeatedly rectified by distillation over bismuth powder. The product was weighed in a closed tube, dissolved in water, and precipitated with sodium carbonate. In the filtrate, after strongly acidulating.

^{*} Poggend. Annal., 82, 303. 1851.

[†] Ann. de Chim. et de Phys., (3,) 55, 176. 1859.

with nitric acid, the chlorine was precipitated by a known amount of silver. The figures in the third column show the quantities of BiCl, proportional to 100 parts of silver:

3.506 grm	a. BiCl _s =	= 3.545 g	rm. Ag.	98.900
1.149	"	1.168	"	98.373
1.5965	"	1.629	66	98.005
2. 176 7	"	2.225	"	9 7.829
3.081	"	3.144	66	9 7.99 6
2.4158	44	2.470	66	97.806
1.7107	"	1.752	66	97.643
3.523	46	3.6055	46	97.712
5.241	66	5.361	66	97.762
				Mean, 08.002, + .000

Hence Bi = 210.464, $\pm .294$.

The first three of the foregoing series of experiments were made with slightly discolored material, and may therefore be rejected. The remaining six percentages give a mean of 97.791; whence Bi = 209.78; or, if O = 16, Bi = 210.26.

As between the unaccordant results of Schneider and of Dumas, those of the former chemist are probably nearest correct. His method of determination was the more reliable, and the details which he gives concerning his manipulations afford strong presumptions of accuracy. Doubtless the bismuth trichloride used by Dumas, contained, like the corresponding antimony compounds, traces of oxychloride. We may fairly assume, for all practical purposes, that the atomic weight of bismuth cannot be far from 208.

TIN

Stannic oxide and stannic chloride are the compounds which have been studied in estimating the atomic weight of tin.

The composition of stannic oxide has been fixed in two ways; by synthesis from the metal, and by reduction in hydrogen. For the first method we may consider the work of Berzelius, Mulder and Vlaanderen, and Dumas.

TIN. 205

Berzelius* oxidized 100 parts of tin by nitric acid, and found that 127.2 parts of SnO₂ were formed.

The work done by Mulder and Vlaanderen † was done in connection with a long investigation into the composition of Banca tin, which was found to be almost absolutely pure. For the atomic weight determinations, however, really pure tin was taken, prepared from pure tin oxide. This metal was oxidized by nitric acid, with the following results. 100 parts of tin gave of SnO₂:

Dumas ‡ oxidized pure tin by nitric acid in a flask of glass. The resulting SnO₂ was strongly ignited, first in the flask, and afterwards in platinum. His weighings, reduced to the foregoing standard, give for dioxide from 100 parts of tin the amounts stated in the third column:

In an investigation later than that previously cited, Vlaanderen || found that when tin was oxidized in glass or porcelain vessels, and the resulting oxide ignited in them, traces of nitric acid were retained. When, on the other hand, the oxide was strongly heated in platinum, the latter was perceptibly attacked, so much so as to render the results uncertain. He therefore, in order to fix the atomic weight of tin, reduced the oxide by heating it in a porcelain boat in a stream of hydrogen. Two experiments gave Sn = 118.08, and Sn = 118.24. These, when O = 16, become, if reduced to the above common standard,

^{*} Poggend. Annal., 8, 177.

[†] Journ. für Prakt. Chem., 49, 35. 1849.

¹ Ann. Chem. Pharm., 113, 26.

^{||} Jahresbericht, 1858, 183.

We have now four series of results showing the quantity of SnO, formed from 100 parts of tin. To Berzelius' single value may be assigned the probable error of a single experiment in Mulder and Vlaanderen's series:

General mean	127.143, ± .0098
Vlaanderen	127.082, ± .012—Reduction.
Dumas	127.105, ± .024— "
Mulder and Vlaanderen	127.517, ± .029— "
Berzelius	127.200, \pm .041—Oxidation.

Dumas, in the paper previously quoted, also gives the results of some experiments with stannic chloride, SnCl₄. This was titrated with a solution containing a known weight of silver. From the weighings given, 100 parts of silver correspond to the quantities of SnCl₄ named in the third column:

All these data properly combined give us the following values for the atomic weight of tin:

If O = 16, this becomes Sn = 117.968.

TITANIUM.

The earliest determinations of the atomic weight of titanium are due to Heinrich Rose.* In his first investigation he studied the conversion of titanium sulphide into titanic acid, and obtained erroneous results; later, in 1829, he published his analyses of the chloride.† This compound was purified by repeated rectifications over mercury and over potassium, and was weighed in bulbs of thin glass. These were broken under water in tightly stoppered flasks; the titanic acid was precipitated by ammonia, and the chlorine was estimated as silver chloride. The following results were obtained. In a fourth column I give the TiO, in percentages referred to TiCl, as 100; and in a fifth column the quantity of TiCl, proportional to 100 parts of AgCl:

TiCl4.	TiO ₂ .	AgCl.	Per cent. TiO ₂ .	AgCl Ratio.
.885 grm.	.379 grm.	2.661 grm.	42.825	33.258
2.6365 "	1.120 "	7.954 "	42.481	33.147
1.7157 "	.732 "	5.172 "	42.665	33.173
3.0455 "	1.322 "	9.198"	43.423	33.100
2.4403 "	1.056 "	7.372 "	43.273	33.102
		M	ean, 42.933, ± .12	1 33.156, ± .019

If we directly compare the AgCl with the TiO, we shall find 100 parts of the former proportional to the following quantities of the latter:

From all these figures we can get three values for Ti, thus:

[#] Gilbert's Annalen, 1823, 67 and 129.

⁺ Poggend. Annal., 15, 145. Berz. Lehrbuch, 3, 1210.

These results will be discussed further along in connection with others.

Shortly after the appearance of Rose's paper, Mosander* published some figures giving the percentages of oxygen in titanium dioxide, from which a value for the atomic weight of titanium was deduced. Although no details are furnished as to experimental methods, and no actual weighings are given, I cite his percentages for whatever they may be worth:

These figures give values for Ti ranging from 46.277 to 48.231; or, in mean, Ti = 47.045. They are not, however, sufficiently explicit to deserve any further consideration. It will be noticed that the highest value nearly coincides with Rose's lowest.

In 1847 Isidor Pierre made public a series of important determinations.† Titanium chloride, free from silicon and from iron, was prepared by the action of chlorine upon a mixture of carbon with pure, artificial, titanic acid. This chloride was weighed in sealed tubes, these were broken under water, and the resulting hydrochloric acid was titrated with a standard solution of silver after the method

^{*} Berz. Jahresbericht, 10, 108. 1831.

[†] Ann. de Chim. et Phys., (3,) 20, 257.

of Pelouze. I subjoin Pierre's weighings, and add, in a third column, the ratio of TiCl₄ to 100 parts of silver:

TiCl ₄ .	Ag.	Ratio.
.8215 grm.	1.84523 grm.	44.520
.7740 "	1.73909 "	44.506
·7775 "	1.74613 "	44.527
.7160 "	1.61219 "	44.412
.8085 "	1.82344 "	44.339
.6325 "	1.42230 "	44.470
.8155 "	1.83705 "	44.392
.8165 "	1.83899 "	44-399
.8065 "	1.81965 "	44.322

Mean, 44.432, \pm .0173

It will be seen that the first three of these results agree well with each other and are much higher than the remaining six. The last four experiments were made purposely with tubes which had been previously opened, in order to determine the cause of the discrepancy. According to Pierre, the opening of a tube of titanium chloride admits a trace of atmospheric moisture. This causes a deposit of titanic acid near the mouth of the tube, and liberates hydrochloric acid. The latter gas being heavy, a part of it falls back into the tube, so that the remaining chloride is richer in chlorine and poorer in titanium than it should be. Hence, upon titration, too low figures for the atomic weight of titanium are obtained. Pierre accordingly rejects all but the first three of the above estimations:

From all of Pierre's......Ti = 49.889,
$$\pm$$
 .096 " the first three......" = 50.259, \pm .063

The memoir of Pierre upon the atomic weight of titanium was soon followed by a paper from Demoly,* who obtained much higher results. He also started out from titanic chloride, which was prepared from rutile. The latter substance was found to contain 1.8 per cent. of silica; whence Demoly inferred that the TiCl₄ investigated by Rose and by Pierre

^{*} Ann. Chem. Pharm., 72, 214. 1849. Berz. Jahresb., 30, 58.

might have been contaminated with SiCl, an impurity which would lower the value deduced for the atomic weight under consideration. Accordingly, in order to eliminate all such possible impurities, this process was resorted to: the chloride, after rectification over mercury and potassium, was acted upon by dry ammonia, whereupon the compound TiCl₄.4NH₃ was deposited as a white powder. This was ignited in dry ammonia gas, and the residue, by means of chlorine, was reconverted into titanic chloride, which was again repeatedly rectified over mercury, potassium, and potassium amalgam. The product boiled steadily at 135°. This chloride, after weighing in a glass bulb, was decomposed by water, the titanic acid was precipitated by ammonia, and the chlorine was estimated in the filtrate as silver chloride. Three analyses were performed, yielding the following results. I give the actual weighings:

```
1.470 grm. TiCl<sub>4</sub> gave 4.241 grm. AgCl and .565 grm. TiO<sub>2</sub>.
2.330 " 6.752 " .801 "
2.880 " 8.330 " 1.088 "
```

The ".801" in the last column is certainly a misprint for .901. Assuming this correction, the results may be given in three ratios, thus:

Per cent. TiO2 from TiCl4.	TiCl4: 100 AgCl.	TiO ₂ : 100 AgCl.
38.435	34.662	13.322
38.669	34.508	13.344
37.778	34-574	13.061
Mean, 38.294 , $\pm .180$	34.581, ± .030	13.242, ± .061

These three ratios give three widely divergent values for the atomic weight of titaniun;

The value assumed by Demoly is 56; who employs but one ratio and ignores practically the others.

Upon comparing Demoly's figures with those obtained by Rose, certain points of similarity are plainly to be noted. Both sets of results were reached by essentially the same method; and in both the discordance between the percentages of titanic acid and of silver chloride is glaring. discordance can rationally be accounted for by assuming that the titanic chloride was in neither case absolutely what it purported to be; that, in brief, it must have contained impurities; such for example as hydrochloric acid, as shown in the experiments of Pierre, or possibly traces of oxychlorides. Considerations of this kind also throw doubt upon the results attained by Pierre, for he neglected the direct estimation of the titanic acid altogether, thus leaving us without means for correctly judging as to the character of his material. In fact, not one of the determinations of the atomic weight of titanium can be regarded as trustworthy. All depend upon the chloride, and the volatile chlorides of metals are as a class especially liable to contaminations of a kind most difficult to recognize. Possibly a series of good determinations might be based upon analyses of some of the titanofluorides. I subjoin a combination of the foregoing mean values, feeling that such a general average is a little better than any one set of determinations taken singly:

```
From Rose's analyses...... Ti = 48.710, \pm .105

"Pierre's " = 49.889, ± .096

"Demoly's " = 52.191, ± .153

General mean..... = 49.846, ± .064

Or, if O = 16, Ti = 49.961.
```

This mean agrees with the average of all of Pierre's experiments.

ZIRCONIUM.

The atomic weight of zirconium has been determined by Berzelius, by Hermann, and by Marignac. Berzelius* ignited the neutral sulphate, and thus ascertained the ratio in it between the ZrO₂ and the SO₃. Putting SO₃ at 100, he gives the following proportional quantities of ZrO₂:

```
75.84

75.92

75.80

75.74

75.97

75.85

Mean, 75.853, ± .023
```

Hence Zr = 89.255, $\pm .039$; or, if O = 16, then Zr = 89.461.

Hermann's † estimate of the atomic weight of zirconium was based upon analyses of the chloride, concerning which he gives no details or weighings. From sublimed zirconium chloride he finds Zr = 831.8, when O = 100; and from two lots of the basic chloride $2ZrOCl_{.9}H_{.0}$, Zr = 835.65 and 851.40 respectively. The mean of all three is 839.62; whence, with modern formulæ and O = 15.9633, Zr becomes = 89.354.

Marignac's results ‡ were obtained by analyzing the double fluoride of zirconium and potassium. His weights are as follows:

```
1.000 grm. gave .431 grm. ZrO<sub>2</sub> and .613 grm. K<sub>2</sub>SO<sub>4</sub>.
2.000 " .864 " 1.232 "
.654 " .282 " .399 "
5.000 " 2.169 " 3.078 "
```

These figures give us three ratios. A, the ZrO, from 100

^{*} Poggend. Annal., 4, 126. 1825.

[†] Journ. für Prakt. Chem., 31, 77. Berz. Jahresb., 25, 147.

[‡] Ann. Chim. Phys., (3,) 60, 270. 1860.

parts of salt; B, the K₂SO₄ from 100 parts of salt; and C, the ZrO₂ proportional to 100 parts of K₂SO₄:

Or, if O = 16, Zr = 90.536.

Combining with Berzelius' work we get this result:

Or, if O = 16, Zr = 89.573.

These figures need little criticism. They show conclusively that the atomic weight of zirconium ought to be redetermined. Probably the method employed by Berzelius was the best with respect to manipulation, while on the other hand it is likely that Marignac worked with purer material. Hermann's experiments could hardly have yielded certain results, since the zirconium chloride might so easily become contaminated with traces of moisture and thence of oxygen.

THORIUM.

The atomic weight of thorium has been determined from analyses of the sulphate, oxalate, formate, and acetate, with widely varying results. The earliest figures are due to Berzelius,* who worked with the sulphate, and with the double sulphate of potassium and thorium. The thoria was precipitated by ammonia, and the sulphuric acid was estimated as BaSO₄. The sulphate gave the following ratios in two experiments. The third column represents the weight of ThO₂ proportional to 100 parts of BaSO₄:

The double potassium sulphate gave .265 grm. ThO₂, .156 grm. SO₃, and .3435 K₂SO₄. The SO₃, with the Berzelian atomic weights, represents .4537 grm. BaSO₄. Hence 100 BaSO₄ is equivalent to 58.408 ThO₂. This figure, combined with the two previous values for the same ratio, give a mean of 58.026, \pm .214. Hence ThO₂ = 269.940, \pm .997.

From the ratio between the K_2SO_4 and the ThO₂ in the double sulphate, ThO₂ = 268.284.

In 1861 new determinations were published by Chydenius,† whose memoir is accessible to me only in an abstract; which gives results without details. Thoria is regarded as a monoxide, ThO, and the old equivalents (O = 8) are used. The following values are assigned for the molecular weight of ThO, as found from analyses of several salts:

From Sulphate.	From K. Th. Sulphate.
66.33	67.02
67.13	
67.75	
68. 03	
Mean, 67	$7.252, \pm .201$

^{*} Poggend. Annal., 16, 398. 1829. Lehrbuch, 3, 1224.

[†] Kemisk undersökning af Thorjord och Thorsalter. Helsingfors, 1861. An academic dissertation.

[†] Poggend. Annal., 119, 55. 1863.

From Acetate.	From Formate.	From Oxalate.
67.31	68.06	65.87 \ Two results
66.59	67.89	65.95 by Berlin.
67.27	68.94	65.75
67.06	•	65.13
68.40	Mean, 68.297, ± .219	66.54
· ·		65.85
Mean, 67.326 , $\pm .201$		
	N	fean, 65.85 , $\pm .123$

We may fairly assume that these figures were calculated with O = 8, C = 6, and S = 16. Correcting by the values for these elements which have been found in previous chapters, ThO, becomes as follows:

The single result from the double potassium sulphate is included with the column from the ordinary sulphate, and the influence of the atomic weight of potassium is ignored.

Chydenius was soon followed by Marc Delafontaine, whose researches appeared in 1863.* This chemist especially studied thorium sulphate; partly in its most hydrous form, partly as thrown down by boiling. In Th(SO₄)₂.9H₂O, the following percentages of ThO₂ were found:

Hence ThO₂ = 263.637, $\pm .256$.

The lower hydrate, $2\text{Th}(SO_4)_2.9\text{H}_2\text{O}$, was more thoroughly investigated. The thoria was estimated in two ways; first, (A,) by precipitation as oxalate and subsequent ignition; second, (B,) by direct calcination. These percentages of ThO, were found:

^{*} Arch. des Sci. Phys. et Nat., (2,) 18, 343.

Mean, 52.511, ± .04

Hence ThO₂ = 266.025, $\pm .363$.

In three experiments with this lower hydrate the sulphuric acid was also estimated, being thrown down as barium sulphate after removal of the thoria:

```
1.2425 grm. gave .400 SO<sub>3</sub>. (1.1656 grm. BaSO<sub>4</sub>.)
1.138 " .366 " (1.0665 " )
.734 " .2306 " (.6720 " )
```

The figures in parenthesis are reproduced by myself from Delafontaine's results, he having calculated his analyses with O = 100, S = 200, and Ba = 857. These data may be reduced to a common standard, so as to represent the quantity of $2\text{Th}(SO_4)_2.9\text{H}_2\text{O}$ equivalent to 100 parts of $BaSO_4$. We then have the following results:

Hence ThO, = 259.555, ± 2.725 .

Delafontaine seems himself to have calculated from the ratio between the percentages of SO, and ThO₂; whence, with our revised values for S, Ba, and O, ThO₂ = 262.643. Delafontaine's work was soon confirmed by Hermann,*

^{*} Journ. für Prakt. Chem., 93, 114.

who published a single analysis of the lower hydrated sulphate, as follows:

Hence, from the ratio between SO_s and ThO₂, ThO₂ = 263.030. Probably the SO_s percentage was loss upon calcination.

The latest, and probably also the best determinations, are those of Cleve,* whose results, obtained from both the sulphate and the oxalate of thorium, agree admirably. The anhydrous sulphate, calcined, gave the subjoined percentages of thoria:

Hence ThO₂ = 265.380, $\pm .123$.

The oxalate was subjected to a combustion analysis, whereby both thoria and carbonic acid could be estimated. From the direct percentages of these constituents no accurate value can be deduced, there having undoubtedly been moisture in the material studied. From the ratio between CO₂ and ThO₂, however, good results are attainable. This ratio I put in a fourth column, making the thoria proportional to 100 parts of carbon dioxide:

Oxalate.	ThO_2 .	CO_2 .	Ratio.
1.7135 grm.	1.0189 grm.	.6736 grm.	151.262
1.3800 "	.8210 "	·5433 "	151.114
1.1850 "	.7030 "	.4650 "	151.183
1.0755 "	.6398 "	.4240 "	150.896
Hence ThO.	005 957		, 151.114, ± .053

^{*} K. Svenska Vet. Akad. Handlinger. Bd. 2, No. 6. 1874.

There are now before us twelve estimates for the molecular weight of thoria. Two of these represent single experiments, and have no probable error attached to them; namely, the one due to Hermann, and the one deduced from Berzelius' K₂SO₄: ThO₂ ratio. A third value, from Delafontaine's sulphuric acid estimations, has so high a probable error that it could be rejected without influencing the general mean. These three values might all be excluded without serious objection; but it is perhaps better to arbitrarily assign them equal weight, average them together, and give their mean the same probable error as that which attaches to Berzelius' BaSO₄: ThO₂ series. This mean is indicated as "A" in the following combination:

```
= 269.940, \pm .997
Berzelius _____
Chydenius—Sulphate ....
                           = 268.584, \pm .803
                           = 268.735, \pm .805
       Acetate _____
                           = 272.586, \pm .877
       Formate
                           = 262.804, \pm .493
       Oxalate _____
Delafontaine—Higher hydrate
                           = 263.637, \pm .256
        Lower "
                           = 266.025, \pm .363
                           = 265.380, \pm .123
Cleve—Sulphate
                           = 265.357, \pm .104
    Oxalate _____
    General mean ____ " = 265.341, ± .072
```

Hence Th = 233.414, \pm .0725; or, if O = 16, Th = 233.951.

These values vary from those derived from Cleve's experiments alone only in the second decimal.

GALLIUM.

Gallium has been so recently discovered, and obtained in such small quantities, that its atomic weight has not as yet been determined with much precision. The following data were fixed by the discoverer, Lecoq de Boisbaudran:*

^{*} Journ. Chem. Soc., 1878, p. 646.

3.1044 grammes gallium ammonium alum, upon ignition, left .5885 grm. Ga, O₃.

Hence Ga = 68.071. If O = 16, Ga = 68.233.

.4481 grammes gallium, converted into nitrate and ignited, gave .6024 grm. Ga₂O₃.

• Hence Ga = 69.538. If O = 16, Ga = 69.693.

These values, assigned equal weight, give these means:

If O = 15.9633, Ga = 68.854. If O = 16, Ga = 68.963.

In brief, for all practical purposes, 69 may be assumed as the atomic weight of gallium.

INDIUM.

Reich and Richter, the discoverers of indium, were also the first to determine its atomic weight.* They dissolved weighed quantities of the metal in nitric acid, precipitated the solution with ammonia, ignited the precipitate, and ascertained its weight. Two experiments were made, as follows:

Hence, in mean, In = 110.61, if O = 16; a value known now to be too low.

An unweighed quantity of fresh, moist indium sulphide was also dissolved in nitric acid, yielding, on precipitation,

.2105 grm. In₂O₈ and .542 grm. BaSO₄.

Hence, with $BaSO_4 = 233$, In = 111.544; also too low.

Soon after the publication of Reich and Richter's paper the subject was taken up by Winkler.† He dissolved indium in nitric acid, evaporated to dryness, ignited the residue, and weighed the oxide thus obtained.

^{*} Journ. für Prakt. Chem., 92, 484.

[†] Journ. für Prakt. Chem., 94, 8.

```
.5574 grm. In gave .6817 grm. In<sub>2</sub>O<sub>3</sub>.
.6661 " .8144 "
.5011 " .6126 "
```

Hence, in mean, if O = 16, In = 107.76; a result even lower than the values already cited.

In a later paper by Winkler* better results were obtained. Two methods were employed. First, metallic indium was placed in a solution of pure, neutral, sodio-auric chloride, and the amount of gold precipitated was weighed. I give the weighings and, in a third column, the amount of indium proportional to 100 parts of gold:

In.	Au.	Ratio.
.4471 grm.	.8205 grm.	57.782
.8445 "	1.4596 "	57.858
		Mean, 57.820 , $\pm .026$

Hence, if Au = 196.155, $\pm .095$, In = 113.417, $\pm .074$.

Winkler also repeated his earlier process, converting indium into oxide by solution in nitric acid and ignition of the residue. An additional experiment, the third as given below, was made after the method of Reich and Richter. The third column gives the percentage of In in In,O,:

1.124 grm.	In gave	1.3616	grm. In ₂ O ₃ .	Per cent.,	82.550
1.015	46	1.2291	66	46	82.581
.6376	46	.7725	44	4.6	82.537

These figures were confirmed by a single experiment of Bunsen's,† published simultaneously with the specific heat determinations which showed that the oxide of indium was In₂O₃, and not InO as had been previously supposed:

```
1.0592 grm. In gave 1.2825 grm. In<sub>1</sub>O<sub>3</sub>. Per cent. In, 82.589
```

For convenience we may add this figure in with Winkler's series, which gives us a mean percentage of In in In $_2O_3$ of 82.564, \pm .0082. Hence, if O=15.9633, \pm .0035, In = 113.385, \pm .060.

^{*} Journ. für Prakt. Chem., 102, 282.

[†] Poggend. Annal., 141, 28.

Combining results, we have the following general mean:

From gold series In = 113.417,
$$\pm$$
 .074

" oxide " = 113.385, \pm .060

General mean " = 113.398, \pm .047

Or, if O = 16, In = 113.659.

CERIUM.

Although cerium was discovered almost at the beginning of the present century, its atomic weight was not properly determined until after the discovery of lanthanum and didymium by Mosander. In 1842 the investigation was undertaken by Beringer,* who employed several methods. His cerium salts, however, were all rose-colored, and therefore were not wholly free from didymium; and his results are further affected by a negligence on his part to fully describe his analytical processes.

First, a neutral solution of cerium chloride was prepared by dissolving the carbonate in hydrochloric acid. This gave weights of ceroso-ceric oxide and silver chloride as follows. The third column shows the amount of CeO₂ proportional to 100 parts of AgCl:

CeO ₂ .	AgCl.	Ratio.
.5755 grm.	1.419 grm.	40.557
.6715 "	1.6595 "	40.464
1.1300 "	2.786 "	40.560
.5366 "	1.3316 "	40.297
	N	lean, 40.469, ± .0415

The analysis of the dry cerium sulphate gave results as follows. In a fourth column I show the amount of CeO, proportional to 100 parts of BaSO₄:

^{*} Ann. Chem. Pharm., 42, 134.

Sulphate.	CeO ₂ .	$BaSO_4$.	Ratio.
1.379 grm.	.8495 grm.	1.711 grm.	49.649
1.276 "	.7875 "	1.580 "	49.836
1.246 "	.7690 "	1.543 "	49.838
1.553 "	·959 5 "	1.921 "	49.948
		Mean	${49.819, \pm .042}$

Beringer also gives a single analysis of the formate and the results of one conversion of the sulphide into oxide. The figures are, however, not valuable enough to cite.

The foregoing data involve one variation from Beringer's paper. Where I put CeO, as found he puts Ce,O₃. The latter is plainly inadmissible, although the atomic weights calculated from it agree curiously well with some other determinations. For instance, in the chloride series, the assumption of Ce,O₃ as the formula of the oxide obtained, gives Ce = 137.749, while CeO, makes Ce = 141.636. The former agrees with the results of Wolf, Wing, and others quite fairly; the latter is near the value obtained by Bührig. Obviously, the presence of didymium in the salts analyzed should tend to raise rather than to lower the apparent atomic weight of cerium.

Shortly after Beringer, Hermann * published the results of one experiment. 23.532 grm. of anhydrous cerium sulphate gave 29.160 grm. of BaSO₄. Hence 100 parts of the sulphate correspond to 123.926 of BaSO₄.

In 1848 similar figures were published by Marignac,† who found the following amounts of BaSO₄ proportional to 100 of dry cerium sulphate:

If we give Hermann's single result the weight of one experiment in this series, and combine, we get a mean value of 123.019, $\pm .113$.

^{*} Journ. für Prakt. Chem., 30, 185. 1843.

[†] Arch. des Sciences Phys. et Nat., (1,) 8, 273. 1848.

Still another method was employed by Marignac. A definite mixture was made of solutions of cerium sulphate and barium chloride. To this were added, volumetrically, solutions of each salt successively, until equilibrium was attained. The figures published give maxima and minima for the BaCl, proportional to each lot of Ce₂(SO₄)₈. In another column, using the mean value for BaCl, in each case, I put the ratio between 100 parts of this salt and the equivalent quantity of sulphate. The latter compound was several times recrystallized:

	$Ce_2(SO_4)$	8-	$BaCl_2$.		Ratio.
First crys	tallization	. 11.011 grm.	11.990 — 12.050 g	rm.	91.606
44		. 13.194 "	14.365 — 14.425	66	91.657
Second	"	. 13.961 "	15.225 — 15.285	44	91.518
44		. 12.627 "	13.761 — 13.821	4.6	91.559
66	••	11.915 "	12.970 — 13.030	"	91.654
Third	46	. 14.888 "	16.223 — 16.283	66	91.602
44	44	. 14.113 "	15.383 — 15.423	6.	91.755
Fourth		13.111 "	14.270 — 14.330	66	91.685
44		13.970 "	15.223 — 15.283	"	91.588
				Mean,	91.625, ± .016

Omitting the valueless experiments of Kjerulf,* we come next to the figures published by Bunsen and Jegel† in 1858. From the air dried sulphate of cerium the metal was precipitated as oxalate, which, ignited, gave CeO₂. In the filtrate from the oxalate the sulphuric acid was estimated as BaSO₄:

```
1.5726 grm. sulphate gave .7899 grm. CeO<sub>2</sub> and 1.6185 grm. BaSO<sub>4</sub>.
1.6967 " .8504 " 1.7500 "
```

Hence, for 100 parts BaSO₄, the CeO₂ is as follows:

One experiment was also made upon the oxalate:
.3530 grm. oxalate gave .1913 CeO, and .0506 H,O.

Hence, in the dry salt, we have 63.261 per cent. of CeO₂.

^{*} Ann. Chem. Pharm., 87, 12.

[†] Ann. Chem. Pharm., 105, 45.

In each sample of CeO, the excess of oxygen over true Ce₂O₃ was estimated by an iodometric titration; but the data thus obtained need not be further considered.

In two papers by Rammelsberg * data are given for the atomic weight of cerium, as follows. In the earlier paper cerium sulphate is analyzed, the cerium being thrown down by caustic potash, and the acid precipitated from the filtrate as barium sulphate:

.413 grm. Ce₂(SO₄)₃ gave .244 grm. CeO₂ and .513 grm BaSO₄.

Hence 100 BaSO₄ = 47.563 CeO₂, a value which may be combined with others, thus; this figure being assigned a weight equal to one experiment in Bunsen's series:

Beringer	• • • • • • • • • • • • • • • • • • • •
Rammelsberg	
General mean	49.360, ± .035

It should be noted here that this mean is somewhat arbitrary, since Bunsen and Rammelsberg's cerium salts were undoubtedly freer from didymium than the material studied by Beringer.

In his later paper Rammelsberg gives these figures concerning cerium oxalate. 100 parts gave 10.43 of carbon and 21.73 of water. Hence the dry salt should yield 48.862 per cent. of CO_2 , whence Ce = 137.83.

In all of the foregoing experiments the ceroso-ceric oxide was somewhat colored, the tint ranging from one shade to another of light brown according to the amount of didymium present. Still, at the best, a faint color remained, which was supposed to be characteristic of the oxide itself. In 1868, however, some experiments of Dr. C. Wolf† were post-humously made public, which went to show that pure ceroso-ceric oxide is white, and that all samples previously studied were contaminated with some other earth, not necessarily didymium but possibly a new substance, the removal of

^{*} Poggend. Annal., 55, 65; 108, 44.

[†] Amer. Journ. Science and Arts, (2,) 46, 53.

which tended to lower the apparent atomic weight of cerium very perceptibly.

Cerium sulphate was recrystallized at least ten times. Even after twenty recrystallizations it still showed spectroscopic traces of didymium. The water contained in each sample of the salt was cautiously estimated, and the cerium was thrown down by boiling concentrated solutions of oxalic acid. The resulting oxalate was ignited with great care. I deduce from the weighings the percentage of CeO₂ given by the anhydrous sulphate:

Sulphate.	Water.	CeO2. Per	r cent. CeO ₂ .
1.4542 grm. 1.4104 " 1.35027 "	.19419 grm. .1898 " .1820 "	.76305 grm7377 " .70665 "	60.559 60.437 60.487
	•	Mean,	 60.494, ± .024

After the foregoing experiments the sulphate was further purified by solution in nitric acid and pouring into a large quantity of boiling water. The precipitate was converted into sulphate and analyzed as before:

Sulphate.	Water.	C&O2.	Per cent. CeO ₂ .
1.4327 grm.	.2733 grm.	.69925 grm.	60.311
1.5056 "	.2775 "	.7405 "	60.296
1.44045 "	.2710 "	.7052 "	60.300
		Me	ean, 60.302, ± .004

From another purification the following weights were obtained:

1.4684 grm. .1880 grm. .7717 grm. 60.270 per cent.

A last purification gave a still lower percentage:

1.3756 grm. .1832 grm. .7186 grm. 60.265 per cent.

The last oxide was perfectly white, and was spectroscopically free from didymium. In each case the CeO, was titrated iodometrically for its excess of oxygen. It will be noticed that in the successive series of determinations the percentage of CeO, steadily and strikingly diminishes, to an extent for which no ordinary impurity of didymium can

account. The death of Dr. Wolf interrupted the investigation, the results of which were edited and published by Professor F. A. Genth.

The experiments of Wolf seem to have hitherto escaped general notice, except from Wing, who has partially verified them.* This chemist, incidentally to other researches, purified some eerium sulphate after the method of Wolf, and made two similar analyses of it, as follows:

Sulphale.	Water.	CeO2.	Per cent. CeO2.
1.2885 grm.	.1707 grm.	.6732 grm.	60.225
1.4090 "	.1857 "	·7372 "	60.263
		Me	an, 60.244, ± .012

The ceroso-ceric oxide in this case was perfectly white. The cerium oxalate which yielded it was precipitated boiling by a boiling concentrated solution of oxalic acid. The precipitate stood twenty-four hours before filtering.

We may now combine the results of Wolf and of Wing, as follows. The two concordant experiments of Wolf's series three and four may be united, giving a mean of 60.267, $\pm .001$:

```
Wolf, 1st series 60.494, ± .024

" 2d " 60.302, ± .004

" 3d and 4th series 60.267, ± .001

Wing 60.244, ± .012

General mean 60.271, ± .001
```

This mean, the percentage of CeO_2 in the anhydrous sulphate, gives Ce = 137.724; or, if O = 16, Ce = 138.039. This varies widely from the ordinarily accepted value as determined by Buehrig.

In 1875 Buehrig's † paper upon the atomic weight of cerium was issued. He first studied the sulphate, which, after eight crystallizations, still retained traces of free sulphuric acid. He found furthermore that the salt obstinately retained traces of water, which could not be wholly expelled by heat without partial decomposition of the material.

^{*} Amer. Journ. Sci. and Arts, (2,) 49, 358. 1870.

⁺ Journ. für Prakt. Chem., 120, 222.

227

These sources of error probably affect all the previously cited series of experiments; although, in the case of Wolf's work, it is doubtful whether they could have influenced the atomic weight of cerium by more than one or two tenths of Buehrig also found, as Marignac had earlier shown, that upon precipitation of cerium sulphate with barium chloride the barium sulphate invariably carried down traces of cerium. Furthermore, the ceroso-ceric oxide from the filtrate always contained barium. For these reasons the sulphate was abandoned, and the atomic weight determinations of Buehrig were made with air-dried oxalate. This salt was placed in a series of platinum boats in a combustion tube behind copper oxide. It was then burned in a stream of pure, dry oxygen, and the carbonic acid and water were collected after the usual method. Ten experiments were made; in all of them the above named products were estimated, and in five analyses the resulting cerosoceric oxide was also weighed. By deducting the water found from the weight of the air-dried oxalate, the weight of the anhydrous oxalate is obtained, and the percentages of its constituents are easily determined. In weighing, the articles weighed were always counterpoised with similar materials. The following weights were found:

Oxala	te.	Water.		CO ₂ .		CeO ₂	
9.8541	grm.	2.1987 g	rm.	3.6942	grm.		
9.5368	66	2.1269	"	3.5752	44		
9.2956	44	2.0735	66	3.4845	44	****	
10.0495	46	2.2364	66	3.7704	46		
10.8249	66	2.4145	44	4.0586	46		
9.3679	66	2.0907	•6	3.5118	44	4.6150	grm.
9.7646	66	2.1769	44	3.6616	46	4.8133	"
9.9026	66	2.2073	"	3.7139	66	4.8824	"
9.9376	66	2.2170	48	3.7251	66	4.8971	"
9.5324	46	2.1267	66	3.5735	46	4.6974	66

These figures give us the following percentages for CO, and CeO, in the anhydrous oxalate:

CO ₂ .	CeO_2 .
48.256	*
48.249	
48.248	20

$$CeO_2$$
.
 CO_3 .

 48.257

 48.258
 63.417
 48.257
 63.436
 48.262
 63.446
 48.249
 63.429
 48.253
 63.430

 Mean, 48.2546 , \pm .001
 63.4316 , \pm .0032

From percentage
$$CO_2$$
 ---- $Ce = 141.228$, $\pm .025$
" CeO_2 ---- " = 141.141, $\pm .020$

Obviously the single oxalate experiments of Jegel and of Rammelsberg would exert no appreciable influence upon these mean results. They may therefore be ignored.

In combining all of these data in one general mean, we may begin as usual by tabulating our ratios:

```
(1.) BaSO<sub>4</sub>: Ce<sub>2</sub>(SO<sub>4</sub>)<sub>3</sub>:: 100: 123.019, \pm .113

(2.) BaSO<sub>4</sub>: CeO<sub>2</sub>:: 100: 49.360, \pm .035

(3.) BaCl<sub>2</sub>: Ce<sub>2</sub>(SO<sub>4</sub>)<sub>3</sub>:: 100: 91.625, \pm .016

(4.) AgCl: CeO<sub>2</sub>:: 100: 40.469, \pm .0415

(5.) Percentage CeO<sub>2</sub> from anhydrous sulphate, 60.271, \pm .001

(6.) " " oxalate, 63.4316, \pm .0032

(7.) " CO<sub>2</sub> " 48.2546, \pm .001
```

These ratios give us four values for the molecular weight of CeO, and two values for Ce₂(SO₄)₃:

Hence we have three independent values for the atomic weight of cerium, as follows:

Or, if O = 16, Ce = 140.747.

Buehrig's results alone, both sets combined, give Ce = 141.198, $\pm .020$; or, if O = 16, Ce = 141.523.

Wolf and Wing's figures alone make Ce = 137.724; or, if O = 16, Ce = 138.039.

The latter result is subject to the errors pointed out by Buehrig as involved in the use of cerium sulphate; but the ceroso-ceric oxide obtained in the analyses was pure white. Buehrig's ceroso-ceric oxide, on the other hand, was yellow. In neither case was didymium present. All things considered, therefore, it is probable that the lower result is too low and the higher result too high. How near the general mean of all may be to the truth we have no evidence to show. It is clear that new determinations are needed, made with material yielding white ceroso-ceric oxide, and with avoidance of the sources of error which Buehrig pointed out.

LANTHANUM.

Leaving out of account the work of Mosander, and the valueless experiments of Choubine, we may consider the estimates of the atomic weight of lanthanum which are due to Hermann, Rammelsberg, Marignac, Czudnowicz, Holzmann, Zschiesche, Erk, and Cleve.

From Rammelsberg* we have but one analysis. .700 grm. of lanthanum sulphate gave .883 grm. of barium sulphate. Hence 100 parts of BaSO₄ are equivalent to 79.276 of La₂(SO₄)₃.

^{*} Poggend. Annal., 55, 65.

Marignac,* working also with the sulphate of lanthanum, employed two methods. First, the salt in solution was mixed with a slight excess of barium chloride. The resulting barium sulphate was filtered off and weighed; but, as it contained some occluded lanthanum compounds, its weight was too high. In the filtrate the excess of barium was estimated, also as sulphate. This last weight of sulphate, deducted from the total sulphate which the whole amount of barium chloride could form, gave the sulphate actually proportional to the lanthanum compound. The following weights are given:

$La_2(SO_4)_3.$	$BaCl_2$.	st BaSO4.	2d BaSO4.
4.346 grm.	4.758 grm.	5.364 grm.	.115 grm.
4.733 "	5.178 "	5.848 "	.147 "

Hence we have the following quantities of La₂(SO₄)₃ proportional to 100 parts of BaSO₄. Column A is deduced from the first BaSO₄ and column B from the second, after the manner above described:

	A.	В.
	81.022	83.281
	80.934	83.662
Mean,	80.978, ± .030	Mean, 83.471, ± .128
From	A	La = 138.776
46	B	" = 147.474

A agrees best with other determinations, although, theoretically, it is not so good as B.

Marignac's second method, described in the same paper with the foregoing experiments, consisted in mixing solutions of La₂(SO₄)₃ with solutions of BaCl₂, titrating one with the other until equilibrium was established. The method has already been described under cerium. The weighings give maxima and minima for BaCl₂. In another column I give La₂(SO₄)₃ proportional to 100 parts of BaCl₂, mean weights being taken for the latter:

^{*} Archives des Sci Phys. et Naturelles, (1,) 11, 29. 1849.

$La_2(SO_4)_3$.	$BaCl_2$.	Ratio.
II.644 grm.	12.765 — 12.825 grm.	91.004
12.035 "	13.195 — 13.265 "	90.968
10.690 "	11.669 — 11.749 "	91.297
12.750 "	13.920 — 14.000 "	91.332
10.757 "	11.734 11.814 "	91.362
12.672 "	13.813 — 13.893 "	91.475
9.246 "	10.080 — 10.160 "	91.364
10.292 "	11.264 — 11.264 "	91.615
10.192 "	11.111 — 11.171 "	91.482

Mean, 91.322, \pm .048

Hence La = 140.484.

Although not next in chronological order, some still more recent work of Marignac's * may properly be considered here. The salt studied was the sulphate of lanthanum, purified by repeated crystallizations. In two experiments the salt was calcined, and the residual oxide weighed; in two others the lanthanum was precipitated as oxalate, and converted into oxide by ignition. The following percentages are given for La₂O₃:

$$57.56$$
 By calcination.
 57.58 Ppt. as oxalate.
 57.55 Ppt. as oxalate.
Mean, 57.5475 , \pm .0115

The atomic weight determinations of Holzmann† were made by analyses of the sulphate and iodate of lanthanum, and the double nitrate of magnesium and lanthanum. In the sulphate experiments the lanthanum was first thrown down as oxalate, which, on ignition, yielded oxide. The sulphuric acid was precipitated as BaSO₄ in the filtrate.

$La_2(SO_4)_8$.	La_2O_3 .	$BaSO_{\bullet}$.	
.9663 grm.	.5157 grm.	1.1093 grm.	
.6226 "	·33 ² 3 "	.7123 "	
.8669 "	.4626 "	.9869 "	

^{*} Ann. de Chim. et de Phys., (4,) 30, 68. 1873.

[†] Journ. für Prakt. Chem., 75, 321. 1858.

These results are best used by taking the ratio between the BaSO₄, put at 100, and the La₂O₃. The figures are then as follows:

In the analyses of the iodate the lanthanum was thrown down as oxalate, as before. The iodic acid was also estimated volumetrically, but the figures are hardly available for present discussion. The following percentages of La₂O₃ were found:

The formula of this salt is La₂(IO₈)₆.3H₂O.

The double nitrate, La₂(NO₃)₆.3Mg(NO₃)₂.24H₂O, gave the following analytical data:

Salt.	H_2O .	MgO.	La_2O_3 .
.5327 grm.	.1569 grm.	.0417 grm.	.1131 grm.
.5931 "	.1734 "	.0467 "	.1262 "
.5662 "	.1647 "	.0442 "	.1197 "
·3757 "		.0297 "	.0813
.3263 "	*	.0256 "	.0693 "

These weighings give the subjoined percentages of La,O,:

These data of Holzmann give values for the molecular weight of La₂O₃ as follows:

From sulphate _____ La₂O₃ = 325.674,
$$\pm$$
 .522

" iodate ____ " = 322.419, \pm .113

" magnesian nitrate _ " = 324.355, \pm .923

Czudnowicz* based his determination of the atomic weight of lanthanum upon one analysis of the air-dried sulphate. The salt contained 22.741 per cent. of water.

.598 grm. gave .272 grm. La₂O₂ and .586 grm. BaSO₄.

The La₂O₃ was found by precipitation as oxalate and ignition. The BaSO₄ was thrown down from the filtrate. Reduced to the standards already adopted these data give for the percentage of La₂O₃ in the anhydrous sulphate the figure 58.668. 79.117 parts of the salt are proportional to 100 parts of BaSO₄.

Hermann† studied both the sulphate and the carbonate of lanthanum. From the anhydrous sulphate, by precipitation as oxalate and ignition, the following percentages of La,O, were obtained:

The carbonate, dried at 100°, gave the following percentages:

Reckoning from the ratio between CO, and La₂O₃ the molecular weight of the latter becomes 325.896.

Zschiesche's ‡ experiments consist of six analyses of lanthanum sulphate, which salt was dehydrated at 230°, and afterwards calcined. I subjoin his percentages, and in a fourth column deduce from them the percentage of La₂O₃ in the anhydrous salt:

H_2O .	SO ₃ .	La_2O_3 .	La ₂ O ₃ in anhydrous salt.
22.629	33.470	43.909	56.745
22.562	33.306	44.132	56.9 64
22.730	33.200	44.070	57.034

^{*} Journ. für Prakt. Chem., 80, 33. 1860.

[†] Journ. sür Prakt. Chem., 82, 396. 1861.

[†] Journ. für Prakt. Chem., 104, 174.

H_2O .	SO ₃ .	La_2O_3 .	La202 in anhydrous salt.
22.570	33-333	44.090	56.947
22.610	33.160	44.240	57.150
22.630	33.051	44.310	57.277
			Mean, 57.021, ± .051

Erk * found that .474 grm. of La₂(SO₄)₃, by precipitation as oxalate and ignition, gave .2705 grm. of La₂O₃, or 57.068 per cent. .7045 grm. of the sulphate also gave .8815 grm. of BaSO₄. Hence 100 parts of BaSO₄ are equivalent to 79.921 of La₂(SO₄)₃.

Last of all, and probably best of all, we come to the determinations of Cleve.† Strongly calcined La,O,, spectroscopically pure, was dissolved in nitric acid, and then, by evaporation with sulphuric acid, converted into sulphate:

From the last column, which indicates the percentage of La₂O₃ in La(SO₄)₃, we get, if SO₃ = 80, La = 139.15.

We may now combine the similar means into general means, and deduce a value for the atomic weight of lanthanum. For the percentage of oxide in sulphate we have six estimates, as follows. The single experiments of Czudnowicz and of Erk are assigned the probable error and weight of a single experiment in Hermann's series:

Czudnowicz	58.668,	士 .027
Erk	57.068,	土 .027
Hermann	57.654,	± .016
Zschiesche	57.021,	± .051
Marignac		
Cleve	57.619,	± .0085
General mean	57.620,	±.0059

^{*} Jenaisches' Zeitschrift, 6, 306. 1871.

[†] K. Svenska Vet. Akad. Handlingar, Bd. 2, No. 7. 1874.

For the quantity of La₂(SO₄), proportional to 100 parts of BaSO₄, we have five experiments, which may be given equal weight and averaged together:

Marignac	81.022
***************************************	80.934
Rammelsberg	79.276
Czudnowicz	79.117
Erk	79.921
Mean,	80.054, ± .270

In all, there are seven ratios from which to calculate:

- (1.) Percentage of La_2O_3 in $La_2(SO_4)_3$, 57.620, \pm .0059
- (2.) BaCl₂: La₂(SO₄)₃:: 100: 91.322, \pm .048—Marignac.
- (3.) BaSO₄: La₂(SO₄)₃:: 100: 80.054, \pm .270
- (4.) BaSO₄: La₂O₃:: 100: 46.671, \pm .075—Holzmann.
- (5.) Percentage of La₂O₃ in iodate, 23.447, ± .0216—Holzmann.
- (6.) " magnesian nitrate, 21.3056, ± .058—Holzmann.
- (7.) " carbonate, 68.47—Hermann.

These ratios give five values for the molecular weight of lanthanum oxide, and two for that of the sulphate:

From (2).....La₂(SO₄)₈ = 568.488,
$$\pm$$
 .320
" (3)..... " = 558.624, \pm 1.888
General mean, " = 568.212, \pm .316

Hence La = 140.346, $\pm .160$.

From (1) ______ La₂O₃ = 325.791,
$$\pm$$
 .074

" (4) _____ " = 325.674, \pm .522

" (5) ____ " = 322.419, \pm .113

" (6) ___ " = 324.355, \pm .923

" (7) ___ " = 325.896, \pm .488

General mean ___ " = 324.810, \pm .061

Here the value derived from ratio (7) is given the weight of a single experiment in ratio (1.) Hence La = 138.460, $\pm .031$.

Combining the two values for La, we get this final result:

From La₂O₃La = 138.460,
$$\pm$$
 .031

" La₂(SO₄)₃ " = 140.346, \pm .160

General mean " = 138.526, \pm .030

Or, if O = 16, La = 138.844.

Since this value is a little under and Cleve's a little over 139, the latter figure may fairly be used in all calculations involving a knowledge of the atomic weight of lanthanum.

DIDYMIUM.

The atomic weight of didymium has been determined by Marignac, Hermann, Zschiesche, Erk, and Cleve. Mosander's early experiments we may leave out of account.

Marignac* mixed a solution of the sulphate with a slight excess of barium chloride, filtered, weighed the precipitate, and estimated the excess of barium in the filtrate by the ordinary method. The first precipitate always contained didymium, and therefore weighed too much. By deducting the weight of the second precipitate, representing the excess of the barium chloride, from the weight of barium sulphate theoretically formable, the weight of the latter proportional to the quantity of didymium salt taken was found:

$Di_2(SO_4)_3$.	$BaCl_2$.	rst BaSO ₄ .	ed BaSO4.
3.633 grm.	3.902 grm.	4.412 grm.	.084 grm.
3.862 "	4.227 "	4.679 "	.075 "
3.330 "	3.552 "	4.027 "	.088 "
1.386 "	1.477 "	1.681 "	.014 "

These figures give us a ratio between the sulphates of didymium and barium which we may express as follows. Column A gives the Di₂(SO₄)₃ proportional to 100 parts of BaSO₄, as calculated from the first precipitate of the latter. Column B gives a similar ratio calculated with the second BaSO₄ precipitate, this being deduced from the total BaSO₄ which the chloride used could form:

A.	В.
82.344	84.685
82.539	82.626
82.692	85.545
82.451	84.425
82.247—Erk.	
·	Mean, 84.320, ± .414
Mean, 82.455, ± .052	

^{*} Arch. des Sci. Phys. et Naturelles, (1,) 11, 29. 1849.

To A I have added a single result of Erk's, to be described further along. It will be seen that although A is theoretically defective, its figures are much more concordant than those in B. In fact, the latter would almost vanish for the final general mean for the atomic weight of didymium:

In a later paper* Marignac adopts two other methods for establishing the atomic weight of didymium. The carefully dehydrated sulphate was taken, the didymium was precipitated as oxalate, and the latter, ignited, yielded oxide. The following percentages of oxide were found:

58.22 58.24 58.29 58.31 58.29 Mean, 58.27, ± .0115

The chloride of didymium was also studied. As the anhydrous salt could not be obtained in an absolutely definite state, Marignae prepared neutral solutions of it and determined the ratio between didymium oxide and silver chloride. The latter compound was first precipitated in the usual way, and filtered off; the excess of silver in the filtrate was removed by hydrochloric acid, and after that the didymium was thrown down as oxalate and weighed as oxide. The subjoined weights of AgCl and Di,O, were found. In a third column I give the ratio between the two compounds, putting AgCl at 100:

AgCl.	Di_2O_3 .	Ratio.
10.058 grm.	3.946 grm.	39.232
5.029 "	1.960 "	38.974
5.844 "	2.276 "	38.946

Mean, 39.051, ± .061

Hence Di = 143.637, $\pm .263$.

^{*} Ann. d. Chim. et d. Phys., (3,) 38, 148. 1853

Hermann's* determination of the atomic weight of didymium rests on a single experiment with the sulphate. By precipitation as oxalate and subsequent ignition, he found that this salt yielded 58.14 per cent. of Di,O₃.

Zschiesche † also analyzed didymium sulphate, which he dehydrated at 230°, and afterwards converted into oxide by calcination. I give his percentages, and also, in a fourth column, the percentage of oxide from the anhydrous sulphate as deduced from his figures:

H_2O .	SO ₃ .	Di_2O_3 .	Di ₂ O ₃ in anhyd. salt.
23.19	32.97	43.83	57.070
23.03	32.39	44.58	57.919
23.00	32.56	44.95	58.006
23.547	31.938	44.515	58. 225
22.550	32.870	44.570	57.554

The salt used in the first experiment probably contained lanthanum. Rejecting this, the mean of the figures remaining in the fourth column is 57.926, $\pm .094$. Hence Di = 141.007.

Erk,‡ to whom reference has already been made, estimated didymium in the sulphate by precipitation as oxalate and calcination to oxide:

$Di_2(SO_4)_8$	Di_2O_3 .	Per cent. Di ₂ O ₃ .	
.556 grm.	.323 grm.	58.094	
.674 "	.3915 "	58.087	

Hermann's single result for this percentage, 58.14, agrees more nearly with Erk's series than with any other. It may therefore be averaged in with Erk's two experiments, giving a mean of 58.107, \pm .0112. Erk also obtained from .7065 grm. of sulphate .859 grm. BaSO₄. This experiment has already been averaged with Marignac's earlier results.

The latest determinations of the atomic weight of didymium were published by Cleve || in 1874. Strongly calcined

^{*} Journ. für Prakt. Chem., 82, 367. 1861.

⁺ Journ. für Prakt. Chem., 107, 74.

[†] Jenaisches' Zeitschrift, 6, 306. 1871.

^{||} K. Svenska Vet. Akad. Handlingar, Bd. 2, No. 8. These figures were kindly transcribed for me by Professor Delafontaine of Chicago, as I had not access to a copy of the original memoir.

didymium oxide was dissolved in nitric acid, the solution was evaporated with sulphuric acid, and the weight of the resulting sulphate was ascertained. I subjoin the weighings and the percentage of Di₂O₃ in Di₂(SO₄)₃:

Di_2O_3 .	$Di_2(SO_4)_8$.	Per cent. Di ₂ O ₃ .
2.257 grm:	3.844 grm.	58.715
1.086 "	1.8485 "	58.750
1.1525 "	1.9615 "	58.756
1.3635 " .	2.319 "	58.797
1.9655 "	3-3435 "	58.786
1.528 "	2.599 "	58.792

Mean, 58.766, $\pm .0087$

Hence Di = 146.804. If $SO_3 = 80$, Di = 147.021.

This determination is undoubtedly the best of all, and might properly be accepted to the exclusion of the others. Still, it is worth while to combine all the figures into one general mean. For the percentage of Di₂O₃ in Di₂(SO₄)₃ we have the following data:

Marignac	58.270, ± .0115
Erk and Hermann	$58.107, \pm .0112$
Zschiesche	57.926, ± .094
Cleve	$58.766, \pm .0087$
General mean	58.451, ± .0059

For the atomic weight of didymium we have now three independent values:

```
From per cent. Di_2O_3 in Di_2(SO_4)_3......Di = 144.604, \pm .031

"Marignac's chloride analyses......" = 143.637, \pm .263

"Marignac's and Erk's BaSO<sub>4</sub> ratio..." = 143.929, \pm .189

General mean....." = 144.573, \pm .0306

If O = 16, Di = 144.906.
```

THE YTTRIUM GROUP.

The atomic weights of the metals in this group can only be said to have been determined approximately. Not only do great difficulties attend the purification of the material used for study and the separation of the earths from each other, but there have been and still are grave doubts as to the actual nature of some of the latter. The figures for scandium, yttrium, and ytterbium seem to be tolerably good; those for decipium, philippium, thulium, erbium, and terbium are little more than estimates; for samarium we have no data whatever. All the atomic weights in this group are based upon analyses or syntheses of sulphates; and from analogy to the cerium metals all of these elements are regarded as forming sesquioxides.

SCANDIUM.

Cleve,* who was the first to make accurate experiments on the atomic weight of this metal, obtained the following data. 1.451 grm. of sulphate, ignited, gave .5293 grm. of Sc.O., .4479 grm. of Sc.O., converted into sulphate, yielded 1.2255 grm. of the latter, which, upon ignition, gave .4479 grm. of Sc.O., Hence, for the percentage of Sc.O., in Sc.(SO.), we have:

36.478 36.556 36.556 Mean, 36.530

Hence, if $SO_{2} = 80$, Sc = 45.044.

Later and better results are those of Nilson,† who converted scandium oxide into the sulphate. I give in a third column the percentage of oxide in sulphate:

^{*} Compt. Rend., 89, 419.

[†] Compt. Rend., 91, 118.

```
.3379 grm. Sc_2O_3 gave .9343 grm. Sc_2(SO_4)_3. 36.166 per cent.

.3015 " .8330 " 36.194 "

.2998 " .8257 " 36.187 "

.3192 " .8823 " 36.178 "

Mean, 36.181, \pm .004
```

Hence Sc = 43.980, $\pm .015$; or, if O = 16, then Sc = 44.081. If $SO_3 = 80$, then Sc = 44.032. These values are doubtless very nearly correct.

YTTRIUM.

For yttrium we need consider only the determinations of Popp, Delafontaine, Bahr and Bunsen, and Cleve.

Popp* evidently worked with material not wholly free from earths of higher molecular weight than yttria. The yttrium sulphate was dehydrated at 200°; the sulphuric acid was then estimated as barium sulphate; and after the excess of barium in the filtrate had been removed, the yttrium was thrown down as oxalate, and ignited to yield oxide. The following are the weights given by Popp:

Sulphate.	$BaSO_4$.	Yt_2O_3 .	H_2O .
1.1805 grm.	1.3145 grm.	.4742 grm.	.255 grm.
1.4295 "	1.593 "	·5745 "	.308 "
.8455 "	.9407 "	.3392 "	.1825 "
1.045 "	1.1635 "	.4195 "	.2258 "

Eliminating water, these figures give us for the percentages of Yt₂O₃ in Yt₂(SO₄)₃ the values in column A. In column B I put the quantities of Yt₂O₃ proportional to 100 parts of BaSO₄:

A.	В
51.237	36.075
51.226	36.064
51.161	36. 0 5 8
51.209	36.055
Mean, 51.208, ± .011	Mean, 36.063, ± .003

From B, Yt = 101.880. The values in A will be combined with similar data from other experimenters.

^{*} Ann. Chem. Pharm., 131, 179.

In 1865 Delafontaine* published some results obtained from yttrium sulphate, the yttrium being thrown down as oxalate and weighed as oxide. In the fourth column I give the percentages of Yt,O, reckoned from the anhydrous sulphate:

Sulphate.	Yt_2O_3 .	H_2O .	Per cent. Yt, O3.
.9545 grm.	.371 grm.	.216 grm.	50.237
2.485 "	.9585 "	.565 "	49.922
2.153 "	.827 "	· 49 35 "	49.834
		M	ean, 49.998, ± .081

In another paper † Delafontaine gives the following percentages of Yt₂O₃ in dry sulphate. The mode of estimation was the same as before:

Bahr and Bunsen,‡ and likewise Cleve, adopted the method of converting dry yttrium oxide into anhydrous sulphate, and noting the gain in weight. Bahr and Bunsen give us the two following results. I add the usual percentage column:

Yt_2O_3 .	$Yt_2(SO_4)_3$.	Per cent. Yt2O3.
.7266 grm.	1.4737 grm.	49.304
.7856 "	1.5956 "	49.235
•		
		Mean, 49.2695 , $\pm .0233$

Cleve's || results are published in a joint memoir by Cleve and Hoeglund, and are as follows:

^{*} Ann. Chem. Pharm., 134, 108.

⁺ Arch. des Sci. Phys. et Nat., (2,) 25, 119. 1866.

[†] Ann. Chem. Pharm., 137, 21. 1866.

^{||} K. Svenska Vet. Akad. Handlingar, Bd. 1, No. 8.

Yt_2O_3 .	$Yt_2(SO_4)_3$.	Per cent. Yt, O.
1.4060 grm.	2.8925 grm.	48.608
1.0930 "	2.2515 "	48.545
1.4540 "	2.9895 "	48.637
1.3285 "	2.7320 "	48.627
2.3500 "	4.8330 "	48.624
2.5780 "	5.3055 "	48.591
	•	

Mean, 48.605, $\pm .0096$

This series is unquestionably the best of all. From it, if $SO_3 = 80$, Yt = 89.485.

Combining all these data we have the subjoined general mean for the percentage of Yt,O₃:

Popp	51.208,	110. 土
Delafontaine, 1st	49.998,	180. 土
" 2d	48.230,	±.055
Bahr and Bunsen	49.2695,	± .0233
Cleve	48.605,	± .0096
General mean	49.637,	± .0069
Rejecting Popp	48.705,	± .0087

From the general mean of all, Yt = 97.616. From the mean after excluding Popp's work, Yt = 89.816, \pm .067; or, if O = 16, Yt = 90.023.

YTTERBIUM.

For ytterbium we have one very good set of determinations by Nilson.* The oxide was converted into the sulphate after the usual manner:

Yb ₂ O ₃ .	$Yb_{2}(SO_{4})_{8}$.	Per cent. Yb2O3.
1.0063 gm.	1.6186 grm.	62.171
1.0139 "	1.6314 "	62.149
.8509 "	1.3690 "	62.155
·737 ¹ "	1.1861 "	62.145
1.0005 "	1.6099 "	62.147
.8090 "	1.3022 "	62.126
1.0059 "	1.6189 "	62.134
		Mean, 62.147, ± .0036

^{*} Compt. Rend., 91, 56. 1880.

Hence Yb = 172.761, $\pm .038$. If O = 16, then Yb = 173.158. If SO₈ = 80, Yb = 173.016. The true number cannot be far from 173.

ERBIUM.

Since the earth which was formerly regarded as the oxide of this metal is now known to be a mixture of two or three different oxides, the older determinations of its molecular weight have little more than historical interest. Nevertle-less the work done by several investigators may properly be cited, since it sheds some light upon certain important problems.

First, Delafontaine's * early investigations may be considered. A sulphate, regarded as erbium sulphate, gave the following data. An oxalate was thrown down from it, which, upon ignition, gave oxide. The percentages in the fourth column refer to the anhydrous sulphate. In the last experiment water was not estimated, and I assume for its water the mean percentage of the four preceding experiments:

Sulphate.	Er_2O_8 .	H_2O .	Per cent, Er ₂ O ₃ .
.827 grm.	.353 grm.	.177 grm.	54.308
1.0485 "	·4475 "	.226 "	54.407
.8 03 "	.3415 "	.171 "	54.035
1.232 "	.523 "	.264 "	54.028
1.1505 "	·495 "		54.760
		Me	an, 54.308, ± .0915

Bahr and Bunsen† give a series of results, representing successive purifications of the earth which was studied. The final result, obtained by the conversion of oxide into sulphate, was as follows:

.7870 grm. oxide gave 1.2765 grm. sulphate. 61.653 per cent. oxide.

Hoeglund,‡ following the method of Bahr and Bunsen, secured these results:

^{*} Ann. Chem. Pharm., 134, 108. 1865.

[†] Ann. Chem. Pharm., 137, 21. 1866.

[†] K. Svenska Vet. Akad. Handlingar, Bd. 1, No. 6.

Er_2O_3 .	$Er_2(SO_4)_3$.	Per cent. Er2O3.
1.8760 grm.	3.0360 grm.	61.792
1.7990 "	2.9100 "	61.821
2.8410 "	4.5935 "	61.848
1.2850 "	2.0775 "	61.853
1.1300 "	1.827 "	61.850
.8475 "	1.370 "	61.861
		-

Mean, 61.8375, $\pm .0063$

Humpidge and Burney* give data as follows:

```
1.9596 grm. Er_2(SO_4)_8 gave 1.2147 grm. Er_2O_3. 61.987 per cent.

1.9011 " 1.1781 " 61.965 " Mean, 61.976, \pm .0074
```

Combining all four series we get the subjoined general mean for the percentage of oxide in sulphate. Bahr and Bunsen's single experiment is given the probable error of one experiment in Hoeglund's series:

```
Delafontaine 54.308, ± .0915
Bahr and Bunsen 61.653, ± .0178
Hoeglund 61.8375, ± .0063
Humpidge and Burney 61.976, ± .0074

General mean 61.860, ± .0046
Rejecting the first 61.880, ± .0046
```

From the mean of all, Er = 170.379, $\pm .082$; or, if O = 16, Er = 170.770. From Bahr and Bunsen's determination, Er = 168.683; and from Humpidge and Burney's highest, Er = 171.428.

The foregoing data were all published before the composite nature of the supposed erbia was fully recognized. It will be seen, however, that three sets of results were fairly comparable, while Delafontaine evidently studied an earth widely different from that investigated by the others. Since the discovery of ytterbium, some light has been thrown on the matter. The old erbia is a mixture of at least three earths, to one of which, a rose-colored body, the name erbia is now restricted. For the atomic weight of the true erbium

^{*} Journ. Chem. Society, Feb., 1879, p. 116.

Cleve* gives three values, but without data concerning weighings or methods. Doubtless the oxide was converted into sulphate, and the calculations were made with SO₃ = 80:

166.00 166.21 166.25 ——— Mean, 166.153

With SO₈ = 79.874, this becomes 165.891, and if only 0 = 16, 166.273. These figures are undoubtedly the nearest yet reached to the true value. According to Thalén,† who reasons from spectroscopic evidence, the erbium of Hoeglund was largely ytterbium.

TERBIUM, SAMARIUM, PHILIPPIUM, DECIPIUM, THULIUM, HOLMIUM, AND SORET'S EARTH X.

Concerning these substances, real or alleged, the data are exceedingly vague. For phillippium Delafontaine‡ gives an atomic weight approximating to 123 or 125, and in the same memoir decipium is put at 171. It seems probable that philippium may be identical with Cleve's holmium and the metal of Soret's earth X, while decipium comes near Cleve's thulium, for which the discoverer gives a value of about 170.7.|| If decipium and thulium are identical, or if either proves to be erbium or ytterbium contaminated with the other, then we shall have a triad of metals with atomic weights ranging from Er = 166 to Yb = 173, strikingly parallel with lanthanum, cerium, and didymium. take the natural arrangement of the elements as tabulated after Mendelejeff's plan, somewhat modified in Roscoe and Schorlemmer's "Treatise on Chemistry, we find that such a triad should exist, and, furthermore, that another similar

^{*} Compt. Rend., 91,-382.

[†] Poggend. Beiblätter, 5, 122. 1881.

[†] Arch. des Sci. Phys. et Nat., Mars, 1880.

^{||} Compt. Rend., 91, 329. 1880.

[&]amp; Vol. 2, Part 2, p. 507.

group ought to lie between indium and tin. The latter triad should have atomic weights ranging from 114 to 117; and here possibly, or else forming a triad with yttrium, the other metals of this group may lie.

COLUMBIUM.*

The atomic weight of this metal has been determined by Rose, Hermann, Blomstrand, and Marignac. Rose† analyzed a compound which he supposed to be chloride, but which, according to Rammelsberg,‡ must have been nearly pure oxychloride. If it was chloride, then the widely varying results give approximately Cb = 122; if it was oxychloride, the value becomes nearly 94. If it was chloride, it was doubtless contaminated with tantalum compounds.

Hermann's || results seem to have no present value, and as for Blomstrand's, I am not able to get at a copy of his original memoir. The results of the latter chemist are thus summed up in Becker's "Digest." Three chlorine estimations in the pentachloride give, in mean, Cb = 96.67. Eleven weighings of columbic acid from the same compound make Cb = 96.16. Other experiments on sodium columbate lead Blomstrand to regard 95 as the most probable value.

Marignac¶ made about twenty analyses of the potassium fluoxycolumbate, CbOF₃.2KF.H₂O. 100 parts of this salt give the following percentages:

```
Cb<sub>2</sub>O<sub>5</sub> ----- Extremes 44.15 to 44.60 Mean, 44.36

K<sub>2</sub>SO<sub>4</sub>----- " 57.60 " 58.05

H<sub>2</sub>O ----- " 5.75 " 5.98

F<sub>----</sub> " 30.62 " 32.22
```

^{*}This name has priority over the more generally accepted "niobium," and therefore deserves preference.

[†] Poggend. Annal., 104, 439. 1858.

[†] Poggend. Annal., 136, 353. 1869.

^{||} Journ. filr Prakt. Chem., 68, 73. 1856.

[&]amp; Acta Univ. Lund, 1864.

[¶] Archives des Sci. Phys. et Nat., (2,) 23, 258. 1865.

From the mean percentage of Cb_2O_5 , Cb = 93.217. If O = 16, this becomes 93.431.

From the mean between the extremes given for $K_{\bullet}SO_{\bullet}$, Cb = 93.812. If O = 16, this becomes 94.027.

As Deville and Troost's * results for the vapor density of the chloride and oxychloride agree fairly well with Cb = 94, we may adopt this value as approximately correct.

TANTALUM.

The results obtained for the atomic weight of this metal by Berzelius,† Rose,‡ and Hermann || may be fairly left out of account as valueless. These chemists could not have worked with pure preparations, and their data are sufficiently summed up in Becker's "Digest."

Marignac § made four analyses of a pure potassium fluotantalate, and four more experiments upon the ammonium salt. The potassium compound, K_2 Ta F_7 was treated with sulphuric acid, and the mixture was then evaporated to dryness. The potassium sulphate was then dissolved out by water, while the residue was ignited and weighed as Ta $_2$ O $_3$. 100 parts of the salt gave the following quantities of Ta $_2$ O $_3$ and K_2 SO $_4$:

Ta_2O_5 .	K_2SO_4 .
56.50	44-37
56.75	44.35
56.55	44.22
56.56	44.24
	
Mean, 56.59, ± .037	Mean, 44.295, ± .026

^{*} Comptes Rend., 56, 891. 1863.

[†] Poggend. Annal., 4, 14. 1825. Lehrbuch, 3, 1209.

[†] Poggend. Annal., 99, 80. 1856.

^{||} Journ. für Prakt. Chem., 70, 193. 1857.

³ Archives des Sci. Phys. et Nat., 26, 89, serie 2. 1866.

From these figures, 100 parts of K,SO₄ correspond to the subjoined quantities of Ta,O₅:

The ammonium salt, (NH₄), TaF₇, ignited with sulphuric acid, gave these percentages of Ta₂O₅. The figures are corrected for a trace of K₂SO₄ which was always present:

Hence we have four values for Ta:

Or, if O = 16, Ta = 182.562.

If we assume K = 39, O = 16, F = 19, S = 32, and N = 14; the percentage of K_2SO_4 from K_2TaF_7 gives Ta = 181.912; and the analyses of the ammonium salt make Ta = 182.020. Evidently, 182 is not far from the true value.

PLATINUM.

For this metal we have to consider only experiments by Berzelius, by Andrews, and by Seubert. In an early paper Berzelius* reduced platinous chloride, and found it to contain 73.3 per cent. of platinum. Hence, Pt = 194.204, a

^{*} Poggend. Annal., 8, 177. 1826.

value very near that obtained most recently by Seubert. In his later investigations, Berzelius* studied the potassium chloroplatinate, K₂PtCl₆. 6.981 parts of this salt, ignited in hydrogen, lost 2.024 of chlorine. The residue consisted of 2.822 platinum, and 2.135 potassium chloride. From these data we may calculate the atomic weight of platinum in four ways:

```
1st. From loss of Cl upon ignition ______ Pt = 197.722
2d. "weight of Pt in residue _____ " = 196.942
3d. " KCl " = 196.215
4th. "ratio between KCl and Pt _____ " = 196.652
```

The last of these values is undoubtedly the most reliable, since it involves no errors due to the possible presence of moisture in the salt analyzed. If O = 16, the value becomes Pt = 197.104.

The work done by Andrews † is even less satisfactory than the foregoing, for the reason that its full details seem never to have been published. Andrews dried potassium chloroplatinate at 105°, and then decomposed it by means of zinc and water. The excess of zinc having been dissolved by treatment with acetic and nitric acids, the platinum was collected upon a filter and weighed, while the chlorine in the filtrate was estimated by Pelouze's method. Three determinations gave as follows for the atomic weight of platinum:

If we assume that these values were calculated with K = 39 and Cl = 35.5, the mean, corrected by our later figures for these elements, becomes Pt = 197.382. If O = 16, this becomes Pt = 197.836. Unfortunately, Andrews does not, in his brief note upon the subject, indicate the manner by which his calculations were made.

^{*} Poggend. Annal., 13, 468. 1828.

[†] British Association Report, 1852. Chem. Gazette, 10, 380.

Latest of all we have to consider the experiments of Seubert.* This chemist prepared very pure chloroplatinates of ammonium and potassium, and from their composition deduced the atomic weight of the metal under consideration. The ammonium salt, (NH₄), PtCl₆ was analyzed by heating in a stream of hydrogen, expelling the excess of that gas by a current of carbon dioxide, and weighing the residual metal. In three experiments the hydrochloric acid formed during such a reduction was collected in an absorption apparatus, and estimated by precipitation as silver chloride. Three series of results are given for the percentage of platinum in this salt, together with another single result which may be considered alone. Here are the figures:

Series I.	Series II.	Series III.
43.957	43.871	43.990
43.948	43.876	43.986
43.960	43.872	44.001
43.946	43.881	44.020
43.963	43.875	43.994
43.961	43.879	43.996
		44.004
Mean, 43.956 , $\pm .002$	Mean, 43.876, \pm .001	44.026
		43.998
		Mean, 44.001, ± .003

These series represent three preparations. The additional single experiment above referred to was made with material belonging to series II, but recrystallized from water. This salt gave 43.955 per cent. of platinum, a figure to which we may assign the probable error of one experiment in the first series. Combining, we get the subjoined general mean percentage of Pt in (NH₄)₂PtCl₆:

Series	I	$43.956, \pm .002$
66	II	$43.876, \pm .001$
44	III	$44.001, \pm .003$
Extra	experiment	$43.955, \pm .004$
	General mean	$43.907, \pm .0009$

^{*}Ber. der Deutsch. Chem. Gesell., 14, 865. 1881.

Hence Pt = 194.314, $\pm .078$. If N = 14, and Cl = 35.5, then Pt = 194.906. Calculating with Stas' values for N and Cl, Seubert gets from the four results combined above, the following figures for Pt, respectively: 194.685, 194.039, 195.034, 194.665.

For the chlorine estimations in the ammonium salt the subjoined weighings are given:

Salt.	Pt.	AgCl.
2.7054 grm.	1.1871 grm.	5.2226 grm.
2.2748 "	.9958 "	4.3758 "
3.0822 "	1.3561 "	5.9496 "

Hence 100 parts of AgCl correspond to the following quantities of salt:

Hence, calculating directly from the ratio between 6AgCl and $(NH_4)_2$ PtCl₆, Pt = 196.871, \pm .363.

Seubert himself reckons the percentage of chlorine from the weight of silver chloride, and then calculates the ratio between Cl₅ and Pt. He thus finds, with Stas' value for Cl₇, Pt = 195.330.

The potassium salt, K, PtCl, was also analyzed by ignition in hydrogen, treatment with water, and weighing both the platinum and the potassium chloride. These percentages were found:

Pt.	KCI.
40.119	30.706
40.120	30.728
40.076	30.698
40.070	30.66 6
40.107	30.700
40.120	30.627
40.114	30.710
40.130	30.621
Mean 40.107, ± .005	Mean, 30.682, ± .009

From the first column———Pt = 194.370,
$$\pm$$
 .068 "second" ——— "= 194.645, \pm .213

If K = 39, and Cl = 35.5, the first column gives Pt = 194.933. Seubert, from the percentage of platinum, gets Pt = 194.392; and from the ratio 2KCl: Pt he finds Pt = 194.494.

As with the ammonium salt, three experiments were made upon the potassium compound to determine the amount of chlorine lost upon reduction in hydrogen. I cite the weighings, and add in a fourth column the quantity of K₂PtCl₆ proportional to 100 parts of AgCl. This AgCl represents but four atoms of the chlorine:

Salt.	Pt.	AgCl.	Ratio.
6.7771 grm.	2.7158 grm.	7.9725 grm.	85.006
3-5834 "	1.4372 "	4.2270 "	84.774
4-4139 "	1.7713 "	5.2144 "	84.648
		Mean,	84.809, ± .071

Hence Pt = 195.002, $\pm .415$. If K = 39, Ag = 108, and Cl = 35.5, then Pt = 194.955. Seubert, calculating the percentage of chlorine and thence the ratio Cl₄: Pt, gets Pt = 194.631.

Combining all the values we have the following result for the atomic weight of platinum:

```
1. From per cent. Pt in (NH_4)_2PtCl<sub>6</sub> ------Pt = 194.314, \pm .078

2. " 6AgCl: (NH_4)_2PtCl<sub>6</sub> ratio ----- " = 196.871, \pm .363

3. " per cent. Pt in K_2PtCl<sub>6</sub> ----- " = 194.370, \pm .068

4. " KCl " = 194.645, \pm .213

5. " 4AgCl: K_2PtCl<sub>6</sub> ratio ---- " = 195.002, \pm .415

General mean ---- " = 194.415, \pm .049
```

Or, if O = 16, Pt = 194.867.

Seubert, taking the arithmetical mean of his eight values, gets Pt = 194.620. He regards, however, those results as best which are dependent upon the percentage of platinum in the ammonium salt, and upon the complete analysis of the potassium compound. These give him a mean of Pt = 194.461, which, if corrected by reduction to a vacuum standard, becomes Pt = 194.34.

In will be noticed that three of the ratios, calculated with

K = 39, N = 14, Ag = 108, and Cl = 35.5, give nearly Pt = 195, namely:

194.906 194.933 194.955

The general mean of all, if O = 16, gives Pt = 194.867. Hence, for all practical calculations, the value 195 may be safely employed.

OSMIUM.

The atomic weight of this metal has been determined by Berzelius and by Fremy.

Berzelius* analyzed potassium osmichloride, igniting it in hydrogen like the corresponding platinum salt. 1.3165 grammes lost .3805 of chlorine, and the residue consisted of .401 grm. of potassium chloride, with .535 grm. of osmium. Calculating only from the ratio between the Os and the KCl, we have, Os = 198.494; or, if O = 16, Os = 198.951.

Fremy's determination \uparrow is based upon the composition of osmium tetroxide. No details as to weighings or methods are given; barely the final result is stated. This, if O = 15.9633, is Os = 199.190. If O = 16, Os = 199.648.

Berzelius' work is evidently entitled to preference, although neither determination is in any sense equal to the present requirements of chemical science. The values given are doubtless several units too high.

IRIDIUM.

The only early determination of the atomic weight of iridium was made by Berzelius,‡ who analyzed potassium iridichloride by the same method employed with the platinum and the osmium salts. The result found from a single

^{*} Poggend. Annal., 13, 530. 1828.

[†] Compt. Rend., 19, 468. Journ. für Prakt. Chem., 33, 410. 1844.

[†] Poggend. Annal., 13, 435. 1828.

IRIDIUM. 255

analysis was not far from Ir = 196.7. This is now known to be too high. I have not, therefore, thought it worth while to recalculate Berzelius' figures, but give his estimation as it is stated in Roscoe and Schorlemmer's "Treatise on Chemistry."

In 1878 the matter was taken up by Seubert,* who had at his disposal 150 grammes of pure iridium. From this he prepared the iridichlorides of ammonium and potassium, $(NH_4)_2$ IrCl₆ and K₂IrCl₆, which salts were made the basis of his determinations. The potassium salt was dried by gentle heating in a stream of dry chlorine.

Upon ignition of the ammonium salt in hydrogen, metallic iridium was left behind in white coherent laminæ. The percentages of metal found in seven estimations were as follows:

$$43.742$$
 43.725
 43.745
 43.739
 43.726
 43.739
 43.75

Mean, 43.732 , $\pm .0035$

The potassium salt was also analyzed by decomposition in hydrogen with special precautions. In the residue the iridium and the potassium chloride were separated after the usual method, and both were estimated. Eight analyses gave the following results, expressed in percentages:

Ir.	aKCl.	Cl ₄ .
39.881	30.829	29.290
39.890	30.842	29.277
39.868	30.813	29.300
39.876	30.835	29.289
39.877	30.825	29.287
39.879	30.811	29.310
39:882	30.814	29.285
39.883	30.792	29.288
Mean, 39.880, ± .0015	${30.820, \pm .0037}$	29.291, ± .0024
-		

^{*}Ber. d. Deutsch. Chem. Gesell., 11, 1767.

From these data several values for the atomic weight of iridium may be calculated:

If O = 16, this becomes Ir = 193.145.

In the potassium salt, instead of calculating from the percentages directly, we may reckon upon the ratios between Ir and Cl₄, and between Ir and 2KCl:

Again, we may combine this mean with the value derived from the ammonium iridichloride, and so estimate the relative importance of the latter:

If O = 16, this becomes Ir = 193.094.

We may assume, then, from all the facts before us, that if O = 16, the atomic weight of iridium varies from the even number 193 only within the limits of experimental error.

PALLADIUM.

The atomic weight of palladium has been studied by Berzelius and by Quintus Icilius. In an early paper Berzelius* found that 100 parts of the metal united with 28.15 of sulphur. Hence Pd = 113.63, a result which is unquestionably far too high.

^{*} Poggend. Annal., 8, 177. 1826.

In a later paper* Berzelius published two analyses of potassium palladiochloride, K, PdCl₄. The salt was decomposed by ignition in hydrogen, as was the case with the double chlorides of potassium with platinum, osmium, and iridium. Reducing his results to percentages, we get the following composition for the substance in question:

Pd.	2KCl.	Cl ₂ .
32.726	46.044	21.229
32.655	45.741	21.604
	-	
Mean, 32.690	45.892	21.416

From these percentages, calculating directly, very discordant results are obtained:

Obviously, the only way to get satisfactory figures is to calculate from the ratio between the Pd and 2KCl. Doing this, we get, Pd = 105.737; or, if O = 16, Pd = 105.981.

This last value varies so slightly from the even number 106 that the latter may be safely used for all purposes of chemical calculation.

The determination made by Quintus Icilius* need be given only for the sake of completeness. He ignited potassium palladichloride in hydrogen, and found the following amounts of residue. His weights are here recalculated into percentages:

64.708 64.965 64.781 Mean, 64.818

From this mean, Pd = 111.879. Upon looking at the values deduced from Berzelius' figures, it will be seen that

^{*} Poggend. Annal., 13, 454. 1828.

[†] Die Atomgewichte vom Pd, K, Cl, Ag, C, und H, nach der Methode der kleinsten Quadrate berechnet." Inaug. Diss. Göttingen, 1847. Contains no other original analyses.

the highest, 110.796, is calculated from the chlorine lost upon igniting the palladiochloride. The same kind of error which vitiates that result probably affects also these data drawn from the palladiochloride.

RHODIUM.

Berzelius* determined the atomic weight of this metal by the analysis of sodium and potassium rhodiochlorides, Na₈RhCl₆, and K₂RhCl₅. The latter salt was dried by heating in chlorine. The compounds were analyzed by reduction in hydrogen, after the usual manner. Reduced to percentages the analyses come out as follows:

•	In Na ₈ RhCl ₆ .	
Rh.	3NaCl.	CI ₃ .
26.959	45.853	27.189
27.229	45.301	27.470
		27.616
Mean, 27.094	45-577	
		27.425
	In K ₂ RhCl ₅ .	
Rh.	2KCl.	Cl ₃ .
28.989	41.450	29.561

From the analyses of the sodium salt we get the following values for Rh:

These are discordant figures, and indicate some doubt as to purity of material. The last value is fairly good, however, and is confirmed by results from the potassium compound:

^{*} Poggend. Annal., 13, 435. 1828.

If O = 16, this becomes Rh = 104.285.

RUTHENIUM.


The atomic weight of this metal has been determined only by Claus.* Although he employed several methods, the only results worthy of present notice come from the analysis of potassium rutheniochloride, K,RuCl₅. The salt was dried by heating to 200° in chlorine gas, but even then retained a trace of water. The percentage results of analysis are as follows:

Ru.	2KCl.	Cl ₈ .
28.96	40.80	30.24
28.48	41.39	30.22
28.91	41.08	30.04
Mean, 28.78	41.09	, 30.17

Reckoning directly from the percentages we get the following discordant values for Ru:

Obviously, the best result is to be obtained from the ratio between Ru and 2KCl. This gives Ru = 104.217; or, if O = 16, Ru = 104.457. But little weight can be attached to this determination.

^{*} Journ. für Prakt. Chem., 34, 435. 1845.



APPENDIX.

ON DUMAS' CORRECTION AND PROUT'S HYPOTHESIS.

In the year 1815 Prout put forth his famous hypothesis that the atomic weights of all the elements were multiples of that of hydrogen. His views were adopted by many chemists, but opposed by others; among them Berzelius and Turner; and down to the present day "Prout's Law" has been the subject of earnest controversy. Of course the fact was early recognized that in its original form the hypothesis could not stand, and accordingly it was modified by Dumas in such manner that half and quarter multiples of the atomic weight of hydrogen were considered as well as the whole numbers.

But of late years Prout's hypothesis, even with its elastic modification, has been in disfavor. Only a few chemists still clung to it as the representative of a veritable law. The researches of Stas were especially directed towards ascertaining its truth or falsity; and his results, as well as those obtained by Marignac, were such as to lead most chemists to the belief that it had been forever overthrown. The atomic weights determined by Stas agreed neither with whole, half, nor quarter multiples of that of hydrogen, and the variations seemed to be wholly outside the range of recognizable experimental errors.

In 1878, however, a probable source of error in some of Stas' researches was pointed out by Dumas.* Many of Stas' ratios had involved the use of pure metallic silver, which had been fused under a cover of borax containing a little

nitre. Such silver Dumas heated to redness in a Sprengel vacuum, and found that it gave up weighable quantities of oxygen, which had been absorbed by the metal when in the melted state. In one experiment a kilogramme of silver gave 82 milligrammes of occluded gas, and in three other cases 226,140, and 249 milligrammes respectively were found. In other words, the silver which had been considered pure by Stas and others, was really not pure, and a correction became necessary in nearly all series of atomic weight determinations.

The amount of this correction, which I think may hereafter be appropriately designated as "Dumas' correction," will naturally vary in different cases, and in no particular case can we tell, without actual examination of the silver employed, exactly how great it should be. We may, however, assume that all the metallic silver heretofore used in establishing atomic weight ratios was subject to it; and, reckoning from the largest error indicated in the experiments of Dumas, namely, 249 milligrammes of oxygen in the kilogramme of metal, we may ascertain its tendency with reference to Prout's law.

In the chapter upon the atomic weights of silver, chlorine, bromine, iodine, potassium, sodium, and sulphur, twenty ratios are given, of which nine are subject to Dumas' correction. Applying it as suggested above, we get the following results. The values previously found and given in the chapter just quoted, we may designate as uncorrected. For convenience in future reference I assume that O = 16:

	Uncorrected.	Corrected.	Difference.
Silver	107.923	107.896	027
Chlorine	35.451	35.478	+ .027
Bromine	79.951	79.978	+ .027
Iodine	126.848	126.875	+ .027
Potassium	39.109	39.083	026
Sodium	23.051	23.024	027
Sulphur	32.058	32.058	***

The result of the correction, it will be seen, is generally favorable to Prout's hypothesis. Of the seven elements

under consideration, one has its atomic weight unaffected, one is rendered less in accord with the hypothesis, and five approximate more closely than before to even multiples or multiples half of hydrogen.

In the later chapters of this work the effect of Dumas' correction is generally less striking. One general statement, however, may be made concerning it. Whenever the atomic weight of a metal is calculated from the ratio between its haloid salts and metallic silver, the total effect of Dumas' correction, including the above corrections for the halogens themselves, will be to lower the final result. This point will be further considered presently. Only chlorine, bromine, and iodine have their atomic weights raised by the correction.

In view of Dumas' correction the question naturally arises as to how far other metals, used in atomic weight researches, may occlude gaseous impurities. For example, when the atomic weight of oxygen is fixed by the synthesis of water over copper oxide, may not the copper occlude appreciable quantities of the hydrogen in which it cools? If it does, then the apparent weight of metallic copper would be too high, and the atomic weight of oxygen would come out too low. Such an error might possibly account for the difference between 16 and 15.9633 in the atomic weight of oxygen, and it would also increase the atomic weight of copper as determined by the same process. At all events, every metal of which the atomic weight has been determined by the reduction of its compounds in hydrogen, ought to be scrupulously investigated with reference to the possible occlusion of gaseous impurities. With all of these metals the effect of such impurities would be to render the apparent atomic weights decidedly too high.

Although every series of atomic weight determinations must be considered by itself, and weighed on its own merits, it may not be out of place for me just here to point out two general sources of error in addition to the one we have been considering. First, every value after oxygen, with one or two partial exceptions, involves whatever error may attach

to the atomic weight of oxygen. If the latter be 16, instead of 15.9633, this error in some instances becomes multiplied to a large fraction of a unit, as the subjoined example will show.

Other similar errors are repeated continually. The value assigned to any element is necessarily affected by whatever errors may attach to the atomic weights of those other elements through whose medium it is compared with the standard, hydrogen. Thus, the atomic weight of carbon depends upon that of oxygen; calcium depends upon both carbon and oxygen; and fluorine, as determined from calcium fluoride, involves the foregoing elements, together with sulphur, silver, and chlorine. Since, however, some atomic weights are affected by plus errors and others by minus errors, there is a fortunate tendency to compensation of errors in cases like that of fluorine, and, in reality, better results are obtained than considerations such as these would lead us to look for.

Another general source of error is to be found in the fact that some of the weighings involved in our discussions had been reduced to absolute standards, while others were merely uncorrected weighings in air. The errors thus introduced into the work are doubtless small, but still they ought not to be absolutely ignored.

Now, having considered the larger classes of errors, we may properly pass on to a comparison of our atomic weights with reference to Prout's hypothesis. In order to facilitate work, I have tabulated the figures in two columns, one giving atomic weights referred to hydrogen as unity, the other based upon the standard of oxygen as exactly sixteen. Such imperfectly known elements as decipium, philippium, samarium, terbium, and thulium are not included.

TABLE OF ATOMIC WEIGHTS.

			,
	H = 1.	O = 16.	Remarks.
A1i	an an a	07.07.5	
Aluminum	27.009, ± .003	27.075	Cooke's and Schneider's data.
Antimony		120.231	Cooke sand Schneider's data.
Arsenic Barium		75.090	
Bismuth		137.007	From Schneider's data.
Boron	, , ,		From Schneider's data.
		10.966	
Bromine	79.768, ± .019	79.951	
CadmiumCæsium		112.092	
Calcium	J J J - 1	132.918 40.082	
Carbon	39.990, ± .010	•	
Cerium	$11.9736, \pm .0028$		Buehrig's data give 141.523.
Octifutti	$140.424, \pm .017$	140.747	(0 = 16.)
Chlorine	$35.370, \pm .014$	25 451	(0 = 10.)
Chromium	33.31 · ·	35.45I	From Siewert's data.
Cobalt	$52.009, \pm .025$ $58.887, \pm .008$	52.129	From Siewere's data.
Columbium		59.023	From one ratio only.
Copper		94.027 63.318	Trom one rado only.
	0		Cleve's data give 147.021.
Didymium	$144.573, \pm .031$	144.906	$(SO_8 = 80.)$
Erbium	165.891	166.273	From Cleve's data only.
Fluorine	$18.984, \pm .0065$		Trom Cieve's data omy.
Gallium'	68.854	19. 027 68.963	Imperfectly determined.
Glucinum	<u> </u>	9.106	Nilson and Pettersson's data.
Gold	$9.085, \pm .0055$	196.606	14115011 and 1 ettersson 5 data.
Hydrogen	196.155, ± .095 1.0000	1.0023	
Indium	_	113.659	
Iodine	113.398, \pm .047	126.848	
Iridium		193.094	Seubert's data.
Iron .		56.042	Scubert's data.
Lanthanum		138.844	
Lead	$138.526, \pm .030$ $206.471, \pm .021$	206.946	
Lithium	-	7.0235	
Magnesium	$7.0073, \pm .007$ 23.959, $\pm .005$	24.014	Marchand and Scheerer's data.
Manganese	$23.959, \pm .005$ $53.906, \pm .012$	54.029	Schneider and Rawack's data.
Mercury		200.171	Definition and Nawack 5 data.
Molybdenum	199.712, \pm .042		
Nickel	$95.527, \pm .051$ 57.928, $\pm .022$	95.747 58.062	Schneider, Sommaruga, and
***************************************	37.920, ± .022	30.002	Lee.
Nitrogen	14.0210, ± .0035	14.029	Dec.
Osmium	198.494	198.951	Very doubtful.
Oxygen	$15.9633, \pm .0035$	16.000	i
Palladium	19.9033, 12.0033	105.981	Badly determined.
Phosphorus	$30.958, \pm .007$	31.029	Dudiy determined.
Platinum	$194.415, \pm .049$	194.867	Seubert's data.
Potassium	$39.019, \pm .012$	39.109	Cabor b dam.
Rhodium	104.055	104.285	Badly determined.
Rubidium	$85.251, \pm .018$	85.529	2001
Ruthenium	104.217	104.457	Badly determined.
Scandium		44.081	
Selenium	$78.797, \pm .011$	78.978	
	. 11717 14-4	12.212	,

TABLE OF ATOMIC WEIGHTS-CONTINUED.

	H == 1.	0 = 16.	Remarks.			
Silicon Silver Sodium Strontium Sulphur Tantalum Tellurium Thallium Thorium Tin Titanium Tungsten Uranium Vanadium Ytterbium Yttrium	日 = 1. 28.195, ±.066 107.675, ±.0096 22.998, ±.011 87.374, ±.032 31.984, ±.012 182.144, ±.166 127.960, ±.034 203.715, ±.0365 233.414, ±.073 117.698, ±.040 49.846, ±.064 183.610, ±.032 238.482, ±.082 51.256, ±.024 172.761, ±.038 89.816, ±.067	28.260 107.923 23.051 87.575 32.058 182.562 128.254 204.183 233.951 117.968 49.961 184.032 239.030 51.373 173.158 90.023	Remarks. Very badly determined. Imperfectly determined. Crookes' data. Imperfectly determined. If SO ₃ = 80, Yb = 173.016. Doubtful.			
ZincZirconium	$64.9045, \pm .019$ $89.367, \pm .037$	65.054 89.573	Axel Erdmann's data. Doubtful.			

At the close of his admirable paper on the atomic weight of aluminum Mallet makes substantially the following argument in favor of Prout's hypothesis. Citing the atomic weights of eighteen elements which he considers well determined, he shows that ten of them have values falling within one-tenth of a unit of whole numbers. Now, what is the mathematical probability that this close approximation to conformity, with Prout's law, in ten cases out of eighteen, is purely accidental, as those chemists who reject the hypothesis seem to hold? Working this problem out, Mallet finds the probability in favor of mere coincidence to be in the ratio of 1:1097.8, and hence he concludes that Prout's views are still worthy of respectful consideration.

Applying Mallet's reasoning to the table of atomic weights now before us, we find that in the first column, when H=1, twenty-five elements out of sixty-six have values falling within the limits of one-tenth of a unit variation from whole numbers. But many of the figures which fall without this limit involve the variation of oxygen multiplied many times over. We must therefore study the second column, which assumes that the atomic weight of oxygen is exactly six-

teen. Here we have forty elements falling within the limit of variation assigned by Mallet, and twenty-six falling without. The variations we may properly study in some detail.

Taking first the elements whose atomic weights vary from even multiples of unity by less than a tenth of a unit, we have to consider the following: aluminum, arsenic, barium, bismuth, boron, bromine, cadmium, cæsium, calcium, carbon, cobalt, columbium, didymium, fluorine, gallium, hydrogen, iridium, iron, lead, lithium, magnesium, manganese, nickel nitrogen, osmium, oxygen, palladium, phosphorus, scandium, selenium, silver, sodium, sulphur, thorium, tin, titanium, tungsten, uranium, yttrium, and zinc. Of these, aluminum, arsenic, barium, bismuth, cadmium, calcium, carbon, cobalt, columbium, fluorine, hydrogen, iridium, iron, lithium, magnesium, manganese, nickel, nitrogen, phosphorus, scandium, sodium, sulphur, tungsten, uranium, yttrium, and zinc have plus variations, while boron, bromine, cæsium, didymium, gallium, lead, osmium, palladium, selenium, silver, thorium, tin, and titanium fall slightly under the units to which they approximate. gen, as the standard of comparison, of course shows here no variation, its possible error having been transferred to hydrogen.

Of the foregoing elements it will be seen that twenty-six have plus variations from whole numbers, while thirteen Among the latter, boron, gallium, osmium, palare minus. ladium, thorium, and titanium have been but roughly determined. Bromine, by Dumas' correction, has its variation diminished. In the cases of lead, cæsium, selenium, and tin, the cause of variation, supposing one to exist, remains to be determined. The value for osmium is undoubtedly several units too high, so that its agreement with Prout's law may be considered purely accidental. As for didymium, the figure assigned is the mean of all determinations; whereas Cleve's data, calculated with SO₃ = 80, make Di = 147.021, a variation which, like most of the others, is far within the limits of ordinary experimental error. In the case of silver it has already been shown that Dumas' correction is unfavorable to it considered in its bearings upon Prout's law. Silver is the only element among those having minus variations which could carry very much weight against the hypothesis.

Among the elements whose variations are plus, columbium, uranium, and yttrium have been poorly determined. Yttrium especially may be considered doubtful. The atomic weights of aluminum, arsenic, barium, cadmium, lithium, phosphorus, and sodium involve Dumas' correction to a greater or less extent, and will be lowered by its application, that is, brought nearer to whole numbers. For aluminum, certain other causes for variation were pointed out in the chapter upon that metal; and it may be noted that the direct ratio between it and hydrogen gives Al = 27.998 \pm .007. Here the variation is less than the probable error. For calcium, and consequently for fluorine also, sources of plus error were indicated in the discussion of their respective atomic weights, and reiteration here is unnecessary. iridium, iron, nickel, and tungsten all involve such errors as may arise from the possible occlusion of hydrogen by the metals after reduction from their compounds. For scandium, the atomic weight, calculated with SO₃ = 80, becomes 44.032, a variation much within the limits of experimental For carbon and bismuth the variations are insignifierror. In short, in the majority of instances the errors may be diminished by corrections which are in all probability needed, and which can be easily pointed out. The more carefully we scrutinize the data the more probable Prout's hypothesis appears.

Among the twenty-six elements whose atomic weights are removed by more than a tenth of a unit from whole numbers, chlorine, rubidium, and strontium have values nearly half multiples of that of hydrogen, and in each case Dumas' correction will make the approximation still closer. Erbium, gold, indium, lanthanum, rhodium, ruthenium, silicon, and zirconium may be dismissed from consideration as too imperfectly determined to carry much weight in the present

discussion. For chromium, copper, molybdenum, and vanadium I have no criticisms to offer; but the remaining elements may be considered individually.

The value assigned to antimony, 120.231, is the general mean of Cooke's and Schneider's work upon the bromide, iodide, and sulphide. If Ag = 108, Br = 80, and I = 127, Cooke's data for the bromide and iodide give the following values for Sb, all of which fall within a tenth of a unit of the whole number 120:

Early bromid	le serie	S	Sb	=	119.901
Late	44		"	==	120.009
Iodide series			"	=	119.973

In the case of cerium, the value assigned in the table is the general mean of all reputable determinations. But it is subject to doubt on account of the facts observed by Wolf and by Wing, whose ceroso-ceric oxide was white, while that of all other observers was yellowish. Wolf's and Wing's data, calculated with O = 16, give Ce = 138.039. Cerium, then, is not an established exception to Prout's law.

Glucinum and ytterbium have their atomic weights calculated from analyses of the sulphates. But if Prout's law is true, SO₃ = 80. Calculated with this figure, we have Gl = 9.096 and Yb = 173.016. Both elements thus fall within reasonable limits of variation from the hypothetical values.

Iodine is one of the most important seeming exceptions. If we assume Ag = 108, and calculate the atomic weight of iodine only from the direct ratio between iodine and silver, we have, with Dumas' correction applied, I = 126.966; that is, it comes within one-tenth of a unit of the whole number 127.

The atomic weight of mercury depends upon analyses of the chloride, oxide, and sulphide. Of these three compounds the purity of the chloride is most easily assured. Calculated from its composition, with Cl = 35.5, Hg = 199.971. With so high an atomic weight small errors are easily multiplied.

For the atomic weight of platinum Seubert's data give five values, ranging both above and below the round number 195. Calculated with integer values for the other elements, three of these figures fall very close to 195, as follows:

From per cent. Pt in
$$(NH_4)_2$$
PtCl₆----Pt = 194.906
" K_2 PtCl₆------ " = 194.933
From chlorine estimation in K_2 PtCl₆-- " = 194.955

Potassium is the most serious exception of all. But if O = 16 and Dumas' correction be applied, the general mean from all the available data becomes K = 39.083. That is, potassium falls within the limit of 0.1 variation.


The atomic weight assigned to tantalum is the mean of four values. Two of these, recalculated with integers, come out as follows:

For tellurium I need only call attention to the discrepancies between the several sets of determinations made by Wills. A reference to the chapter on tellurium will show that his figures give results ranging from Te = 126.07 to Te = 129.34. The mean value is therefore too much subject to doubt to carry weight as an exception.

As for thallium, the last case to be considered, I have already shown that Crookes' data, recalculated with integer values for N and O, give Tl = 204.008. That is, instead of an exception, we have here an admirable instance in support of Prout's hypothesis.

Enough has been said in this brief resumé to show that none of the seeming exceptions to Prout's law are inexplicable. Some of them, indeed, carefully investigated, support it strongly. In short, admitting half multiples as legitimate, it is more probable that the few apparent exceptions are due to undetected constant errors, than that the great number of close agreements should be merely accidental. I began this recalculation of the atomic weights

with a strong prejudice against Prout's hypothesis, but the facts as they came before me have forced me to give it a very respectful consideration. All chemists must at least admit that the strife over it is not yet ended, and that its opponents cannot thus far claim a perfect victory.



INDEX.

A .	Barium chloride and lanthanum sul-
	phate 230
Allen and Johnson 91	and silver 57
Allen and Pepys6	and silver chloride 60
Aluminum156	chromate 121, 122
bromide 159	metatungstate 147
chloride156	nitrate 61
oxide 156	selenite 177
sulphate156	silicofluoride 85
Ammonia alum 158	sulphate and barium chlo-
Ammonia chrome alum 120	ride 61
Ammonium chloride 40	and barium chromate_ 121
chloroplatinate 251	and barium fluoride 80
cobalticyanide 172	and barium nitrate 61
fluotantalate 249	and barium selenite 177
iridichloride 255	and cerium dioxide,222,224
molybdate 139	and didymium sulphate, 236
Anderson 76	and glucinum sulphate, 97
Andrews 57, 250	and gold 163 and indium oxide 219
Antimony 188, 269	
bromide 198, 200	and lanthanum sul-
chloride, 190, 191, 193, 194, 197	phate 229
iodide 199	and magnesium sul-
oxide 188, 190	phate 100
sulphide 189, 196, 200	and nickel potassium
Antimony compounds, oxidation	sulphate 170
of 189, 191, 192, 193, 194	and thallium sulphate, 93
Antimony and potassium tartrate_ 194	and thorium sulphate_ 214
Arago	and yttrium oxide 241
Arfvedson 87, 127, 151	uranate 151
Arsenic 185	and uranyl acetate 151
bromide 186	Beringer 221
chloride 186	Berlin 118, 139
trioxide 123, 187	Bernoulli 145
Arsenious oxide and potassium an-	Beryllium. See Glucinum 96
hydrochromate 123, 187	Berzelius 1, 6, 10, 14, 19, 21, 27, 31,
Arsenious oxide and potassium	39, 60, 61, 68, 70, 72, 78, 82, 84,
chlorate 187	86, 87, 100, 108, 117, 127, 131, 135,
Atomic weights, table of 265, 266	137, 143, 151, 156, 162, 176, 180,
Awdejew 97	183, 185, 188, 205, 212, 214, 248,
_	249, 254, 256, 258
B.	Biot and Arago 6, 39
	Bismuth 202
Bahr 102, 242, 244	chloride 203
Bahr and Bunsen 242, 244	oxide 203
Balard 21	sulphide202
Barium 57	Blomstrand 247
chloride and barium chro-	Boisbaudran 218
mate 122	Borax 84
and barium sulphate 70	Borch 144
and cerium sulphate 223	Boron 84
18	273
	• •

Boron bromide 84	Cobalticyanide of phenylammo-
chloride 84	nium 172
Brauner 96	of strychnia 174
Bromine 9, 21	Columbic acid 247
Brucia cobalticyanide 173	Columbium 247
nickelocyanide 173	oxychloride 247
Buehrig 226	pentachloride 247
Buff6	Cooke 29, 112, 195, 198, 199
Bunsen 90, 91, 92, 220, 242, 244	Copper 135
Bunsen and Jegel 223	oxide 1, 8, 56, 135
Burney 245	sulphate136
_	subsulphide136
C.	Crookes 95
	Czudnowicz 183, 233
Cadmium III	
bromide II2	_
chloride 112	D .
oxalate112	
sulphate	Davy 6, 78, 127
sulphide 111	Debray 98, 140
Cæsium 91	Decipium 246
bitartrate 91	Delafontaine 215, 242, 244
chloride 91	De Luca 79
Calcium 67	Demoly 209
carbonate 67	De Saussure
and calcium sulphate 69	Deville 84, 248
chloride 70	Deville and Troost248
fluoride 78	Dewar and Scott 130
oxide and calcium sulphate, 70	Dexter 190
sulphate 69, 70	Diamond54
Capitaine 131	Didymium 236
Carbon 50	chloride 237
dioxide 54, 56	oxalate 237
monoxide 55 Cavendish 6	oxide 237
Cerium 221, 269	sulphate236
chloride21	Diehl 87
formate 222	Dœbereiner 72, 131
oxalate 223, 224, 227	Dulong and Berzelius 1, 6, 39
oxide 221, 224	Dumas, 1, 2, 16, 27, 28, 30, 58, 64, 67
sulphate, 221, 222, 223, 224, 225	70, 75, 79, 80, 83, 84, 85, 106, 112.
sulphide223	128, 134, 136, 139, 145, 156, 167.
Chlorine 9	178, 181, 186, 191, 203, 205, 206,
Choubine 229	261
Chrome alum	Dumas and Boussingault 6, 39
Chromium 117	and Stas 54
acetate 118	Dumas' correction 261
chloride 124	
sulphate 120	E.
Chydenius 214	E.
Clark2	
Claus 259	Ebelmen 152
Cleve 217, 234, 238, 240, 242, 246	Ekman and Petterson 178
Cleve and Hoeglund 242	Erbium 244
Cobalt 164	oxide 244
chloride 166, 167	sulphate244
oxalate 166, 168	Erdmann (Axel) 108
sulphate 166 Cobalticyanide of ammonium 172	Erdmann and Marchand 1, 4, 28, 54,
Cobalticyanide of ammonium 172	68, 69, 115, 132, 135, 164, 177
of brucia 173	Erk 234, 23 [§]

r.	Indium sulphide 219
377	Iodine9, 269
Faget 12	and silver 2
Favre 109	Iridium 254
Ferric chloride	Iron 131
oxide	chlorides 134
Ferrous chloride134	oxide 131
tungstate 147	tungstate 147
Fluorine 78	Isnard 150
Fluor spar 78	154444 222222222222222222222222222222222
	T
Fourcroy 6	J.
Fremy 254	V
~	Jacquelain 82, 101, 108, 118
G.	Jegel 223
~	Johnson and Allen 91
Gallium 218	
ammonium alum 219	K.
nitrate 219	
oxide 219	Kemp 140
Gay-Lussac 108	Kessler122, 187, 189, 191, 199
Genth 226	Kirwan
Gerhardt 11	Kjerulf 223
	l
Glucinum 96, 269	Klatzo 98
and ammonium oxalate 98	Kralovanzky 87
sulphate 97	_
Gmelin 87	L.
Godeffroy 90, 91	
Gold162	Lagerhjelm 202
and cobalt 172	Lamy 93
indium 220	Lanthanum 229
mercury 162	carbonate 233
nickel 172	iodate232
	memorium mituate
phosphorus 163	magnesium nitrate 232
Graphite 54	oxalate231
	oxide 232
H.	sulphate 229
•	Laurent 9, 84
Hagen 87	Lavoisier 6, 39
Hampe 136	Lead 72
Hauer 111, 128, 180	chloride 74
Hebberling 93	chromate I18
Hermann, 87, 212, 216, 222, 233, 238,	fluoride 80
247, 248	
Hilgard 158	nitrate 75 and lead oxide 76
Hoeglund 242, 244	l
Holmium 246	oxide 72
Holzmann 231	and lead sulphate 74
Humpidge 96, 245	sulphate 72
Humpidge and Burney 245	and lead fluoride 80
Huntington 112	and lead nitrate 74
Hydrogen 1, 3, 6, 7	and lead oxide 74
and aluminum 157, 160	Le Conte 7, 40
cobalt 169	Lee 172
nickel 169	Lefort 121
109	Lenssen 112
T.	
	Liebig 21, 51
٠. ١	Liebig and Redtenbacher 51
Iceland spar67	Liechti and Kemp 140
7ndium 219	Lithium 87
oride 210	carbonate 87

Lithium chloride 87, 89	Nickel oxalate 165, 166
nitrate 89	oxide 164, 168
	potassium sulphate 170
•	
Loewig 2I	sulphate 165, 166
Longchamp 72	Nilson 96, 99, 240, 244
Louyet 78, 79	and Pettersson 96, 99
30	Niobium. See Columbium 247
M.	Nitrogen 39
	Nitzsch 176
Macdonnell 100	Nordenfeldt 102
Magnesite 103	Norlin 131
Magnesium 100	_
carbonate 103	0.
chloride 106	
lanthanum nitrate 232	Osmium 254
oxalate IO2	OxygenI
oxide 101	
sulphate 100	P.
uranate 151	
Malaguti 143	Palladium 256
Mallet 87, 157, 267	sulphide 256
Manganese 127	Peligot 118, 151, 153
chloride 128	Pelouze, 11, 32, 40, 57, 64, 82, 85, 186
dioxide 131	Penny 10, 15, 32, 41, 45, 47
oxalate 130	Pepys6
sulphate 128	Persoz146
sulphide 128	Pettersson 96, 99, 178
Manganoso-manganic oxide 129	Pfeifer 202
Marchand, 1, 4, 28, 54, 68, 69, 103,	Phenylammonium cobalticyanide 172
	Philippium 246
115, 132, 135, 144, 151, 164, 177 Marchand and Scherer 103	
Marchand and Scheerer 103	Phosphorus 82
Marignac, 9, 11, 12, 14, 15, 16, 18, 19,	and gold 163
22, 23, 25, 40, 42, 44, 57, 59, 60,	pentoxide 82
62, 65, 74, 166, 212, 222, 230, 231,	trichloride 83
236, 237, 247, 248	Piccard 90
Mather 156 Maumené 9, 12, 19, 52, 133	Pletinum 208
Mandalaine 1 9, 12, 19, 52, 133	Platinum 249, 270
Mendelejeff's law 96, 150, 180, 246	dichloride 249
Mercury 114, 269	Popp 241
and gold 162	Potassium 9, 270
chloride116, 269	anhydrochromate 123, 187
oxide II5	and tartar emetic 194
selenide 177	aurichloride162
sulphide 116	bromate 2I
Meyer, Lothar 96, 140	bromide 23
Millon 24, 116, 135	chlorate10
and Commaille 135	and arsenious oxide 187
Mitscherlich and Nitzsch 176	and potassium nitrate_ 47
Moberg120	chloride and gold 162
Molybdenum 137	iridium 256
chlorides 140	osmium 254
sulphide 137	palladium 257
trioxide 137	platinum 250
Mosander 208, 229, 236	potassium nitrate_ 46
Mulder and Vlaanderen 205	rhodium 259
·	ruthenium 259
N.	
	silver 18
	silver 18 silver chloride 19
Nickel 164 chloride 166, 167	silver 18


INDEX.

Potassium fluoxycolumbate 247	Scheibler 147
iodate 24	Schiel
iridichloride 254, 255	Schlippe's salt195
nitrate and potassium chlo-	Schneider, 129, 144, 165, 166, 171, 189
rate 47	199, 200, 203
nitrate and potassium chlo-	Schrötter S2
ride 46	Sefström 114
osmichloride 254	Seguin (
palladichloride 257	Selenium 176
palladiochloride 257	chloride 176, 178
perchlorate II	dioxide 176, 179
platinochloride 250, 252	Seubert 251, 255
rhodiochloride 258	Siewert 124
rutheniochloride 259	Silicon 85
sulphate and potassium fluo-	chloride 85
ride 70	dioxide 85
and potassium tantalo-	Silver
fluoride 248	and aluminum bromide 160
and thoria 214	chloride 157
and zirconia 213	ammonium chloride 40
tantalofluoride248	antimony 202
tartrantimonite 194	bromide 200
zircofluoride 213	chloride 191
Prout's hypothesis 261	arsenious bromide 187
Purpureo-cobalt chloride 171, 174	chloride 186
- mparco coom canonacte 1/1, 1/4	barium chloride 57
•	bismuth chloride 204
Q.	cadmium bromide 113
Ovietve Teiline	chloride 112
Quintus Icilius 257	calcium chloride 70
_	cobalt chloride 168
R.	ferric chloride 134
	ferrous chloride 134
Rammelsberg, 142, 151, 224, 229, 247	lead chloride 74
Rawack 129	lithium chloride
Rawack 129 Redtenbacher 51	lithium chloride 89
Rawack 129 Redtenbacher 51 Regnault 6, 39, 150	lithium chloride 89 magnesium chloride_ 106
Rawack 129 Redtenbacher 51 Regnault 6, 39, 150 Reich and Richter 219	lithium chloride 89 magnesium chloride 106 manganese chloride 128
Rawack 129 Redtenbacher 51 Regnault 6, 39, 150 Reich and Richter 219 Reynolds 96	lithium chloride 80 magnesium chloride 106 manganese chloride 128 nickel chloride 168
Rawack 129 Redtenbacher 51 Regnault 6, 39, 150 Reich and Richter 219 Reynolds 96 Rhodium 258	lithium chloride 89 magnesium chloride 106 manganese chloride 128 nickel chloride 168 phosphorus trichloride, 83
Rawack 129 Redtenbacher 51 Regnault 6, 39, 150 Reich and Richter 219 Reynolds 96 Rhodium 258 Riche 146	lithium chloride 89 magnesium chloride 106 manganese chloride 128 nickel chloride 168 phosphorus trichloride, 83
Rawack 129 Redtenbacher 51 Regnault 6, 39, 150 Reich and Richter 219 Reynolds 96 Rhodium 258 Riche 146 Richter 219	lithium chloride 80 magnesium chloride 106 manganese chloride 128 nickel chloride 168 phosphorus trichloride, 83 potassium bromide 23 chloride 18
Rawack 129 Redtenbacher 51 Regnault 6, 39, 150 Reich and Richter 219 Reynolds 96 Rhodium 258 Riche 146	lithium chloride 80 magnesium chloride 106 manganese chloride 128 nickel chloride 168 phosphorus trichloride, 83 potassium bromide 23 chloride 18
Rawack 129 Redtenbacher 51 Regnault 6, 39, 150 Reich and Richter 219 Reynolds 96 Rhodium 258 Riche 146 Richter 219	lithium chloride 80 magnesium chloride 106 manganese chloride 128 nickel chloride 168 phosphorus trichloride, 83 potassium bromide 23 chloride 18 iodide 25 silicon chloride 85
Rawack 129 Redtenbacher 51 Regnault 6, 39, 150 Reich and Richter 219 Reynolds 96 Rhodium 258 Riche 146 Richter 219 Rivot 133 Roscoe 146, 183 Rose 190, 207, 247, 248	lithium chloride 80 magnesium chloride 106 manganese chloride 128 nickel chloride 168 phosphorus trichloride, 83 potassium bromide 23 chloride 18 iodide 25 silicon chloride 85 silver nitrate 41
Rawack 129 Redtenbacher 51 Regnault 6, 39, 150 Reich and Richter 219 Reynolds 96 Rhodium 258 Riche 146 Richter 219 Rivot 133 Roscoe 146, 183 Rose 190, 207, 247, 248 Rothhoft 164	lithium chloride 89 magnesium chloride 106 manganese chloride 128 nickel chloride 168 phosphorus trichloride, 83 potassium bromide 23 chloride 18 iodide 25 silicon chloride 85 silver nitrate 41 sodium chloride 32
Rawack 129 Redtenbacher 51 Regnault 6, 39, 150 Reich and Richter 219 Reynolds 96 Rhodium 258 Riche 146 Richter 219 Rivot 133 Roscoe 146, 183 Rose 190, 207, 247, 248	lithium chloride 89 magnesium chloride 106 manganese chloride 128 nickel chloride 168 phosphorus trichloride, 83 potassium bromide 23 chloride 18 iodide 25 silicon chloride 85 silver nitrate 41 sodium chloride 32 strontium chloride 64
Rawack 129 Redtenbacher 51 Regnault 6, 39, 150 Reich and Richter 219 Reynolds 96 Rhodium 258 Riche 146 Richter 219 Rivot 133 Roscoe 146, 183 Rose 190, 207, 247, 248 Rothhoft 164 Rubidium 90 chloride 90	lithium chloride 80 magnesium chloride 106 manganese chloride 128 nickel chloride 168 phosphorus trichloride, 83 potassium bromide 23 chloride 18 iodide 25 silicon chloride 85 silver nitrate 41 sodium chloride 32 strontium chloride 32 tin tetrachloride 206
Rawack 129 Redtenbacher 51 Regnault 6, 39, 150 Reich and Richter 219 Reynolds 96 Rhodium 258 Riche 146 Richter 219 Rivot 133 Roscoe 146, 183 Rose 190, 207, 247, 248 Rothhoft 164 Rubidium 90 chloride 90	lithium chloride 89 magnesium chloride 106 manganese chloride 168 nickel chloride 168 phosphorus trichloride, 83 potassium bromide 23 chloride 18 iodide 25 silicon chloride 85 silver nitrate 41 sodium chloride 32 strontium chloride 64 tin tetrachloride 208
Rawack 129 Redtenbacher 51 Regnault 6, 39, 150 Reich and Richter 219 Reynolds 96 Rhodium 258 Riche 146 Richter 219 Rivot 133 Roscoe 146, 183 Rose 190, 207, 247, 248 Rothhoft 164 Rubidium 90 chloride 90 Russell 168, 169	lithium chloride 80 magnesium chloride 106 manganese chloride 168 nickel chloride 168 phosphorus trichloride, 83 potassium bromide 23 chloride 18 iodide 25 silicon chloride 85 silver nitrate 41 sodium chloride 32 strontium chloride 206 titanium chloride 206 vanadium oxychloride, 184
Rawack 129 Redtenbacher 51 Regnault 6, 39, 150 Reich and Richter 219 Reynolds 96 Rhodium 258 Riche 146 Richter 219 Rivot 133 Roscoe 146, 183 Rose 190, 207, 247, 248 Rothhoft 164 Rubidium 90 chloride 90	lithium chloride 80 magnesium chloride 106 manganese chloride 128 nickel chloride 168 phosphorus trichloride, 83 potassium bromide 23 chloride 18 iodide 25 silicon chloride 85 silver nitrate 41 sodium chloride 32 strontium chloride 364 tin tetrachloride 208 titanium chloride 208 vanadium oxychloride, 184 occlusion of oxygen by 262
Rawack 129 Redtenbacher 51 Regnault 6, 39, 150 Reich and Richter 219 Reynolds 96 Rhodium 258 Riche 146 Richter 219 Rivot 133 Roscoe 146, 183 Rose 190, 207, 247, 248 Rothhoff 164 Rubidium 90 chloride 90 Russell 168, 169 Ruthenium 259	lithium chloride 80 magnesium chloride 106 manganese chloride 168 nickel chloride 168 phosphorus trichloride, 83 potassium bromide 23 chloride 18 iodide 18 silicon chloride 85 silver nitrate 41 sodium chloride 32 strontium chloride 32 strontium chloride 206 titanium chloride 206 vanadium oxychloride, 184 occlusion of oxygen by 262 acetate 51, 53
Rawack 129 Redtenbacher 51 Regnault 6, 39, 150 Reich and Richter 219 Reynolds 96 Rhodium 258 Riche 146 Richter 219 Rivot 133 Roscoe 146, 183 Rose 190, 207, 247, 248 Rothhoft 164 Rubidium 90 chloride 90 Russell 168, 169	lithium chloride 89 magnesium chloride 106 manganese chloride 128 nickel chloride 168 phosphorus trichloride, 83 potassium bromide 23 chloride 18 iodide 25 silicon chloride 85 silver nitrate 41 sodium chloride 32 strontium chloride 208 tin tetrachloride 208 vanadium oxychloride, 184 occlusion of oxygen by 262 acetate 51, 53 anhydrochromate 119, 125
Rawack 129 Redtenbacher 51 Regnault 6, 39, 150 Reich and Richter 219 Reynolds 96 Rhodium 258 Riche 146 Richter 219 Rivot 133 Roscoe 146, 183 Rose 190, 207, 247, 248 Rothhoft 164 Rubidium 90 chloride 90 Russell 168, 169 Ruthenium 259	lithium chloride 89 magnesium chloride 106 manganese chloride 168 nickel chloride 168 phosphorus trichloride, 83 potassium bromide 23 chloride 18 iodide 25 silicon chloride 85 silver nitrate 41 sodium chloride 32 strontium chloride 206 tin tetrachloride 206 titanium chloride 206 vanadium oxychloride, 184 occlusion of oxygen by 262 acetate 51, 53 anhydrochromate 119, 125 bromate 21
Rawack 129 Redtenbacher 51 Regnault 6, 39, 150 Reich and Richter 219 Reynolds 96 Rhodium 258 Riche 146 Richter 219 Rivot 133 Roscoe 146, 183 Rose 190, 207, 247, 248 Rothhoft 164 Rubidium 90 chloride 90 Russell 168, 169 Ruthenium 259	lithium chloride 106 magnesium chloride 106 manganese chloride 168 nickel chloride 168 phosphorus trichloride, 83 potassium bromide 23 chloride 18 iodide 25 silicon chloride 85 silver nitrate 41 sodium chloride 32 strontium chloride 206 tin tetrachloride 206 titanium chloride 206 vanadium oxychloride, 184 occlusion of oxygen by 262 acetate 51, 53 anhydrochromate 119, 125 bromate 21 bromide 22
Rawack 129 Redtenbacher 51 Regnault 6, 39, 150 Reich and Richter 219 Reynolds 96 Rhodium 258 Riche 146 Richter 219 Rivot 133 Roscoe 146, 183 Rose 190, 207, 247, 248 Rothhoff 164 Rubidium 90 chloride 90 Russell 168, 169 Ruthenium 259 Salvetat 57, 64, 67	lithium chloride 106 magnesium chloride 106 manganese chloride 168 nickel chloride 168 phosphorus trichloride, 83 potassium bromide 23 chloride 18 iodide 25 silicon chloride 85 silver nitrate 41 sodium chloride 32 strontium chloride 206 titanium chloride 206 titanium chloride 208 vanadium oxychloride, 184 occlusion of oxygen by 262 acetate 51, 53 anhydrochromate 119, 125 bromate 21 bromide 22 and antimony bromide 198
Rawack 129 Redtenbacher 51 Regnault 6, 39, 150 Reich and Richter 219 Reynolds 96 Rhodium 258 Riche 146 Richter 219 Rivot 133 Roscoe 146, 183 Rose 190, 207, 247, 248 Rothhoff 164 Rubidium 90 chloride 90 Russell 168, 169 Ruthenium 259 Salvetat 57, 64, 67 Samarium 246	lithium chloride
Rawack 129 Redtenbacher 51 Regnault 6, 39, 150 Reich and Richter 219 Reynolds 96 Rhodium 258 Riche 146 Richter 219 Rivot 133 Roscoe 146, 183 Rose 190, 207, 247, 248 Rothhoff 164 Rubidium 90 chloride 90 Russell 168, 169 Ruthenium 259 Salvetat 57, 64, 67 Samarium 246 Scandium 240	lithium chloride
Rawack 129 Redtenbacher 51 Regnault 6, 39, 150 Reich and Richter 219 Reynolds 96 Rhodium 258 Riche 146 Richter 219 Rivot 133 Roscoe 146, 183 Rose 190, 207, 247, 248 Rothhoft 164 Rubidium 90 chloride 90 Russell 168, 169 Ruthenium 259 Sacc 57, 64, 67 Samarium 246 Scandium 240 oxide 240	lithium chloride
Rawack 129 Redtenbacher 51 Regnault 6, 39, 150 Reich and Richter 219 Reynolds 96 Rhodium 258 Riche 146 Richter 219 Rivot 133 Roscoe 146, 183 Rose 190, 207, 247, 248 Rothhoff 164 Rubidium 90 chloride 90 Russell 168, 169 Ruthenium 259 Sacc 176 Salvetat 57, 64, 67 Samarium 246 Scandium 240 oxide 240 sulphate 240	lithium chloride 80 magnesium chloride 106 manganese chloride 168 nickel chloride 168 phosphorus trichloride, 83 potassium bromide 23 chloride 18 iodide 25 silicon chloride 85 silver nitrate 41 sodium chloride 32 strontium chloride 206 titanium chloride 206 titanium chloride 206 vanadium oxychloride, 184 occlusion of oxygen by 262 acetate 51, 53 anhydrochromate 119, 125 bromate 21 bromide 22 and antimony bromide 198 and boron bromide 84 and cadmium bromide, 112 and silver chloride 21 and tellurium potassi-
Rawack 129 Redtenbacher 51 Regnault 6, 39, 150 Reich and Richter 219 Reynolds 96 Rhodium 258 Riche 146 Richter 219 Rivot 133 Roscoe 146, 183 Rose 190, 207, 247, 248 Rothhoft 164 Rubidium 90 chloride 90 Russell 168, 169 Ruthenium 259 Sacc 176 Salvetat 57, 64, 67 Samarium 246 Scandium 240 oxide 240	lithium chloride

Silver chlorate	14 !	Sodium susulphantimonite 195
	14	uranate 153
and ammonium platin-		Sommaruga 170
chloride2	252	Soret's earth X 246
and antimony chloride,		Stas, 13, 14, 16, 18, 21, 22, 23, 24, 25,
194, 197	ľ	26, 28, 30, 32, 38, 40, 41, 42, 43,
and barium chloride	60	44, 46, 48, 54, 55, 73, 75, 89
and boron chloride	84	Stibnite 189, 200 Stromeyer 64, 87, 111, 131
and ceroso-ceric oxide,		Stromeyer 64, 87, 111, 131
and chromic chloride_ 1	- 1	Strontium 64
and didymium oxide_ 2	— -	chloride and silver 64
and lithium chloride	87	and strontium sulphate, 66
and manganese chlo- ride	128	Struve 30, 31, 61, 137 Strychnia cobalticyanide 174
and molybdenum chlo-	120	nickelocyanide 173
rides1	141	Sulphur 9, 27
and potassium chlo-		Svanberg,31, 76, 102, 116, 131, 137,
ride	19	and Nordenfeldt 102
and potassium platin-		and Norlin 131
chloride 2	253	and Struve 31, 137
and rubidium chloride,	90	
and silicon chloride	86	T.
and silver bromide	21	
chromate 1	-	Tantalum 248, 270
iodide	27	oxide 248
nitrate	43	Tartar emetic 194
sulphide	31	Tellurium 180, 270 dioxide 180, 181
tungstate i and thallium chloride,	149	potassium bromide 180, 181
93, 94		Terbium 246
and titanium chloride, a	207	Terreil
dioxide_ 207, 2		Thalén246
and vanadium oxychlo-	ļ	Thallium 93
ride 1	85	chloride 93
	119	iodide 94
iodate	24	nitrate 95
iodide	25	sulphate 93 Thomsen 8
and antimony iodide I		
and silver chloride	27	Thomson 6, 39
and thallium iodide malate	94	Thorium 214
•	52 41	formate 215
	44	oxalate 215, 217
••••••	43	oxide 214
oxalate	53	sulphate 214, 215, 217
	31	Thulium 246
racemate	51	Tin 204
	78	dioxide 205
	30	tetrachloride 206
	28	Tissier 156
	31	Titanium 207
	51	
tungstate I Sodio-uranic acetate I5I, I	1	chloride207, 209, 210 dioxide 207, 208
Sodium9,		sulphide 207
chlorate	32	Troost 88, 248
	32	Tungsten
	47	oxides 143
fluoride	70	Turner_15, 60, 61, 73, 114, 115, 116,
rhodiochloride 2	258	127

INDEX.

U.	Winklef 171, 219
	Wolf 224
Unger 195	Wrede6
Uranium 150	
acetate 153	X.
oxalate152	
oxides	X, Soret's earth 246
sulphate 151	
tetrabromide 150	Y.
tetrachloride 150, 151	
	Ytterbium 243, 269
▼.	oxide 243
•	sulphate 243
Vanadium 183	Yttrium 241
oxides 183	oxide 241
oxychloride 184	sulphate 241
Vauquelin6	•
Vlaanderen 205	Z.
W.	Zettnow 147
***	Zimmermann 150
Wackenroder 131	Zinc 108
Wallace 21, 186	oxalate109
Weber 190	oxide 108
Weeren 97	Zirconium 212
Wertheim 153	chloride 212
Weselsky 172	dioxide 212
Wildenstein122	potassium fluoride 212
Wills 181	sulphate212
Wing 226	Zschiesche 233, 238
· · · · · · · · · · · · · · · · · · ·	



SMITHSONIAN MISCELLANEOUS COLLECTIONS.

437

CHECK LIST

04

PUBLICATIONS

OF THE

SMITHSONIAN INSTITUTION,

DECEMBER, 1881.



WASHINGTON, D. C.: 1881.



•

~

•

•

•

·

•

CHECK LIST

OF

PUBLICATIONS OF THE SMITHSONIAN INSTITUTION,

To December, 1881.

Where no price is affixed the work cannot be furnished.

Publications marked * do not appear in the Contributions, Col ections, or Reports.

No.	А отнов.	TITLE.	Size or Sentes.	PAGES	DATE.	PRICE.
A	•	Journal of Regents,	8 v o.#	32	1846	
В		Report of Organization Committee	8 v o.≉	32	1847	
C		Digest of Act of Congress,	8 ▼ 0. *	8	1847	
D	Dallas, G. M.	Address at Laying Corner Stone,	8 v o #	8	1847	
E	Henry, Jos.	Exposition of Bequest,	8 v o.#	8	1847	
F		First Report of Secretary,	8 v o.#	48	1848	
G		First Report of the Institution,	8vo.#	3 8	1847	
H		Second Report of Institution,	8vo.#	208	1848	
I		Third Report of Institution,	8vo.#	64	1849	
J	1	Programme of Organization,	4to.#	4	1847	l
K		Correspondence, Squier & Davis,	8 v o.#	8	1847	
L	! !	First Report of Organization Com- mittee,	8 v o.#	8	1846	ļ
N		Reports of Institution up to Jan. 1849,	8 v o.#	72	1849	
N		Officers, Regents, Act, &c.,	8 v o.#	14	1846	
0		Act to establish Smithsonian Institution,	8 v o.#	8	1846	
P	Owen, R. D.	Hints on Public Architecture,	4to.#	140	1849	10.00
Q		Check List of Periodicals,	4to.#	28	1853	
1	Squier, E. G. and Davis, E. H.	Ancient Monuments of Mississippi Valley,	8. C. 1,	346	1848	
2		Smithsonian Contributions to Knowledge	B. C. 1,	360	1848	

No.	Author.	Title.	Size On Series.	PAGES.	DATE.	PRICE.
3	Walker, S. C.	Researches, Planet Neptune	8. C. 11,	60	1849	
4	Walker, S. C.	Ephemeris of Neptune for 1848,	8. C. 11,	8	1848	
5	Walker, S. C.	Ephemeris of Neptune for 1849,	8. C. 11,	32	1849	•
6	Walker, S. C.	Ephemeris of Neptune for 1850,	8. C. 11,	10	1850	
7	Walker, S. C.	Ephemeris of Neptune for 1851,	S. C. 11,	10	1850	ļ
8	Downes, John	Occultations in 1848,	4to.#	12	1848	
9	Downes, John	Occultations in 1849,	4to.#	24	1848	
10	Downes, John	Occultations in 1850,	4to.#	26	1849	
11	Downes, John	Occultations in 1851,	8. C. 11,	26	1850	
12	Lieber, Francis	Vocal Sounds of L. Bridgeman,	S. C. 11,	32	1850	
13	Ellet, Charles	Physical Geography of U. S.	S. C. 11,	64	1850	•
14	Gibbes, R. W.	Memoir on Mosasaurus,	S. C. 11,	14	1850	
15	Squier, E. G.	Aboriginal Monuments of N. Y.	S. C. 11,	188	1850	
16	Agassiz, Louis	Classification of Insects,	8. C. 11,	28	1850	
17	Hare, Robert	Explosiveness of Nitre,	8. C. 11,	20	1850	
18	Gould, Jr., B. A.	Discovery of Neptune,	8 v o.#	565	1850	
19	Guyot, A.	Directions for Meteorological Observations,	8 v o.#	40	1850	
2 0	Bailey, J. W.	Microscopic Examination of Soundings,	S. C. 11,	16	1851	
21		Annual Report of Smithsonian Institution for 1849	8 v o.	272	1850	
22	Gray, Asa	Plants Wrightians. Part I,	8. C. 111,	146	1852	
23	Bailey, J. W.	Microscopic Observations in S. Carolina, Georgia, and Florida,	S. C. 11,	48	1851	
24	Walker, S. C	Ephemeris of Neptune, 1852. Appendix I,	S. C. 111,	10	1853	
2 5	Jewett, Chas. C.	Public Libraries of United States,	8vo.	21 0	1851	.5
26		Smithsonian Contributions to Knowledge,	S. C. 11,	56-4	1851	
27	Booth, J. C. and Morfit, C.	Improvements in Chemical Arts,	M. C. 11,	216	1852	.50
2 8		Annual Report of Smithsonian Institution for 1850,	8 v o.	326	1851	
29	Downes, John	Occultations in 1852,	S. C. 111,	84	851	

No.	AUTHOB.	Title.	Size or · Series.	PAGES.	DATE.	PRICE.
30	Girard, Charles	Fresh-Water Fishes of N. America	8. C. 111,	80	1851	
31	Guyot, A.	Meteorological Tables,		212	1852	
32	Harvey, Wm. H.	Marine Algo of North America. Part I,	8. C. 111,	152	1852	
33	Davis, Chas. H.	Law of Deposit of Flood Tide,	8. C. 111,	14	185 2	.75
34		Directions for Collecting Specimens,	M. C. 11,	40	18 59	free
3 5	Locke, John	Observations on Terrestrial Mag- netism,	8. C. 111,	30	18 52	
36	Secchi, A.	Researches on Electrical Rheo- metry,	8. C. 111,	60	185 2	
37	Whittlesey, Ch.	Ancient Works in Ohio,	8. C. 111,	20	1851	
38		Smithsonian Contributions to Knowledge,	8. C. 111,	564	1852	
39	•	Smithsonian Contributions to Knowledge,	8. C. 1v,	426	1852	
40	Riggs, S. R.	Dakota Grammar and Dictionary,	S. C. 1▼,	414	1852	
41	Leidy, Joseph	Extinct American Ox,	8. C. v,	20	1852	
42	Gray, Asa	Plants Wrightians. Part II,	S. C. v,	120	1853	
43	Harvey, Wm. H.	Marine Alges of North America. Part II,	8. C. v,	262	1853	
44	Leidy, Joseph	Flora and Fauna within Living Animals,	8. C. v,	68	1858	
45	Wyman, Jeffries	Anatomy of Rana Pipiens,	S. C. ▼,	52	1853	
46	Torrey, John	Plants Fremontians,	8. C. vi,	24	1853	-
47	Jewett, Chas. C.	Construction of Catalogues of Libraries,	8vo.#	108	1853	-50
48	Girard, Charles	Bibliographia Americana Ilis- torico Naturalis,	8 vo.#	64	1852	
49	Baird, S. F. and Girard C.	Catalogue of Serpents,	M. C. 11,	188	1853	1.00
50	Stimpson, Wm.	Marine Invertebrata of Gr. Manan	S. C. v1,	68	1853	1.50
51		Annual Report of Smithsonian Institution for 1851,	8 v o.	104	1852	
52	Coffin, Jas. H.	Winds of the Northern Hemisphere,	S. C. vi,	200	1853	
53	Stanley, J. M.	Catalogue of Portraits of Indians,	M. C. 11,	76	1852	
54	Downes, John	Occultations in 1853,	S. C. VI,	86	185 3	

No.	Author.	Title.	SIZE OR Series	PAGES.	DATE.	PRICE.
55	i	Smithsonian Contributions to Knowledge,	S. C. v,	5 3 8	1853	
56	† 1	Smithsonian Contributions to Knowledge,	S. C. v1,	484	1854	1
57	 - 	Annual Report of Smithsonian Institution for 1852,	8 ▼ 0.	96	1853	¦
58	Leidy, Joseph	Ancient Fauna of Nebraska,	8. C. vi,	126	185 3	i
59	Chappelsmith, J.	Tornado in Indiana,	S. C. v11,	! ! 32	! 1855	.
60	Torrey, John	Batis Maritima,	S. C. VI,	! . 8	1853)
61	Torrey, John	Darlingtonia Californica,	S. C. v1,	8	1853	
62	Melsheimer, F.E.	Catalogue of Coleoptera,	8vo.*	190	1853	2.0
63	Bailey, J. W.	New Species of Microscopic Organisms,	S. C. VII,	16	1854	
64		List of Foreign Correspondents of Smithsonian Institution,	8 ∀ 0. *	16	1 1856	i i
65		Registry of Period. Phenomena,	folio,#	4	1854)
66		Annular Eclipse, May 26, 1854	8 vo. •	14	1854	
67		Annual Report of Smithsonian Institution for 1853,	8vo.	310	1854	
68	Mitchell, B. R. & Turner, W. W.	Vocabulary of Jargon of Oregon,	8vo.#	22	1853	
69		List of American Correspondents of Smithsonian Institution,	8 v o.#	16	1853	ı
70	Lapham, I. A.	Antiquities of Wisconsin,	S. C. v 11,	108	1855	
71	Haven, S. F.	Archmology of the United States,	S. C. v111,	172	1856	l
72	Leidy, Joseph	Extinct Sloth Tribe of N. America,	S. C. V11,	70	1855	1
73		Publications of Societies in Smith- sonian Library,	S. C. VII,	40	1855	
74		Catalogue of Smithsonian Publications,	M. C. v,	52	1862	
75		Annual Report of Smithsonian Institution for 1854,	8 v o.	464	18 5 5	1
76		Smithsonian Contributions to Knowledge,	S. C. v 11,	262	1855	; ;
77		Annual Report of Smithsonian Institution for 1855,	8 v o.	440	1856	!
78		Smithsonian Contributions to Knowledge,	S. C. v 111,	566	1856	

No.	Author.	Title.	SIZE OR SERIES.	PAGES.	DATE.	PRICE.
79	Runkle, John D.	Tables for Planetary Motion,	8. C. 1x,	64	1856	1.00
80	Alvord, Benj.	Tangencies of Circles and Spheres,	S. C. VIII,	16	1856	1.00
81	Olmsted, D.	Secular Period of Aurora Borealis	S. C. VIII,	52	1856	
82	Jones, Joseph	Investigation on A. Vertebrata,	S. C. viii,	150	1856	1.50
83	Meech, L. W.	Relative Intensity of Heat and Light of the Sun,	S. C. 1x,	58	1856	
84	Force, Peter	Auroral Phenomena in North Latitudes,	S. C. VIII,	122	1856	
85	; ,	Publications of Societies in Smith- onian Library. Part II,	S. C. VIII,	3 8	1856	
86	Mayer, Brants	Mexican History and Archæology	8. C. 1x,	36	1856	
87	Coffin, Jas. H.	Psychrometrical Tables,	M. C. 1,	20	1856	.25
88	Gibbs, W. and Genth, F. A.	Ammonia Cobalt Bases,	8. C. 1x,	72	185 6	1.00
89	Brewer, Th. M.	North American Oology. Part I,	8. C. x1,	140	1857	5.00
90	Hitchcock, E.	Illustrations of Surface Geology,	S. C. 1x,	164	1857	
91		Annual Report of Smithsonian Institution for 1856,	8 v o.	468	1857	
92		Smithsonian Contributions to Knowledge,	S. C. 1x,	482	1857	
93		Meteorological Observations for 1855,	8vo.#	118	1857	
94	Runkle, John D.	Asteroid Supplement to New Tables for $b = \frac{(i)}{s}$,	S. C. 1x,	72	1857	.50
95	Harvey, Wm. H.	Marine Algee of North America. Part III,	S. C. x,	142	1858	
96	Harvey, Wm. H.	Marine Algæ of North America. 3 parts complete,	4to.	568	1858	
97	Kane, E. K.	Magnetic Observations in the Arctic Seus,	S. C. x,	72	1859	1.00
98	Bowen, T. J.	Yoruba Grammar and Dictionary,	S. C. x,	232	1858	
99		Smithsonian Contributions to Knowledge,	S. C. x,	462	1858	
00	Gillis, J. M.	Eclipse of the Sun, Sept. 7, 1858,	S. C. x1,	22	1859	.50
01	Hill Thos.	Map of Solar Eclipse, Mar. 15, '58,	8 v o.#	8 I	1858	free
.02	Osten Sacken, R.	Catalogue of Diptera of North America,	M. C. 111,	112	1858	

No.	Author.	TITLE.	SIZE OR SERIES.	PAGES.	DATE.	PRICE.
103	Caswell, A.	Meteorological Observations, Providence, R. I.,	8. C. x11,	188	1860	2.00
104	Kane, E. K.	Meteorological Observations in Arctic Seas,	8. C. x1,	120	18 59	
105	Baird, S. F.	Catalogue of North American Mammals,	4to.#	! 2 2	1857	.50
106	Baird, S. F.	Catalogue of North American Birds,	4to.*	42	1858	.50
107		Annual Report of Smithsonian Institution for 1857,	8 v o.	438	1858	1
108	Baird, S. F.	Catalogue of N. American Birds,	M. C. 11,	24	 1859 	,25
109		Annual Report of Smithsonian Institution for 1858,	8 ₹0 .	44	 - 1859	I
110	,	Annual Report of Smithsonian Institution for 1859,	`8 ▼ 0.	4.5	1860	<u> </u>
111	•.	Smithsonian Contributions to Knowledge,	S. C. x1,	50	1859)
112		Smithsonian Contributions to Knowledge,	8. C. x 11,	54) 0 1860))
113	Bache, A. D.	Magnetic and Meteorological Observations at Girard Coll. Pt. I,		2	, 2 1859	.25
114	Sonntag, A.	Terrestrial Magnetism in Mexico,	S. C. x1,	9	2 185	1.25
115		Report on Invention of Electro- Magnetic Telegraph,	М. С. п,	4	186	free
116	Rhees, Wm. J.	List of Public Libraries, &c.	8 v o.#	84	1859	
117		Catalogue of Publications, &c., in Smithsonian Library,	M. C. 111,	264	1859	2.00
118	Morris, John G.	Catalogue of Lepidoptera of North America,	M. C. 111,	76	1860	1.00
119	Whittlesey, Ch.	Fluctuations of Level in N. A. Lakes,	S. C. x11,	; 28	1860	1.00
120	Hildreth, S. P. and Wood, J.	Meteorological Observations at Marietta, O.,	S. C. xvi,	5 2	1867	1.00
121	Bache, A. D.	Magnetic and Meteorological Observations at Girard Coll. Pt. II,		28	18 62	.25
122	1	Smithsonian Miscellaneous Collections,	M. C. 1,	738	1862	
123		Smithsonian Miscellaneous Collections,	M. C. 11,	715	1862	
134		Smithsonian Miscellaneous Collections,	M. C. 111,	772	1862	

	1		Sizz		i	•
No.	AUTHOB.	Title.	or Series.	PAGES	DATE.	PRICE.
125		Smithsonian Miscellaneous Collections,	M. C. 1V,	762	1862	•
126	Le Conte, John L.	Coleoptera of Kansas and New Mexico,	S. C. x1,	64	1859	1.25
127	Loomis, E.	Storms in Europe and America, Dec. 1836,	S. C. x1,	2 8	1860	1.25
128	Lea, Carpenter,	Check List of Shells in N.America	M. C. 11,	52	1860	.25
129	Kane, E. K.	Astronomical Observations in the Arctic Seas,	S. C. x11,	56	1860	1.00
130	Kane, E. K.	Tidal Observations in the Arctic Seas,	S. C. x 1111,	90	1860	1.50
131	Smith, N. D.	Meteorological Observations in Arkansas from 1840 to 1859,	S. C. x11,	96	1860	1.25
132	Bache, A. D.	Magnetic and Meteorological Observations at Girard Coll. Pt.III		16	1862	.25
133	Morris, John G.	Synopsis of Lepidoptera of North America. Part I,	M. C. 1V,	386	1862	2.00
134	Hagen, H.	Synopsis of Neuropters of North America,	M. C. 1 v ,	368	1861	
135	Mitchell, S. W.	Venom of the Rattlesnake,	S. C. X11,	156	1860	
136	Le Conte, John L.	Classification of Colcoptera of North America. Part I,	M. C. 111,	312	1862	
137		Circular to Officers of Hudson's Bay Co.,	M.C. VIII,	6	1860	
138	Morgan, L. H.	Circular as to Degrees of Relationship,	M. C. 11,	34	1860	
139		Collecting Nests and Eggs of North American Birds,	M. C. 11,	22	1860	free
140	Le Conte, John L.	List of Coleoptera of North America. Part I,	M. C. v1,	82	1866	.75
141	Loew, H. and Osten Sacken	Monographs of Diptera. Part I,	M. C. vi,	246	1862	1.50
142	Binney, W. G.	Bibliography of North American Conchology. Part I,	M. C. v,	658	1863	3.00
143	Binney, W. G.	Land and Fresh-Water Shells of North America. Part II,	M. C. v11,	172	1865	1.25
144	Binney, W. G.	Land and Fresh-Water Shells of North America. Part III,	M. C. v11,	128	1865	1.00
145	Prime, Temple	Monograph of American Corbiculade,	M. C. vii,	92	 2 1865	.75

No.	Author.	Title.	or or Series.	PAGES.	DATE.	PRICE.
146	M'Clintock, Sir F. L.	Meteorological Observations in the Arctic Seas,	S. C. XIII,	164	1862	1.50
147		Annual Report of Smithsonian Institution for 1860,	8 v o.	448	1861	
148	•	Directions for Meteorological Observations,	M. C. 1,	72	1860	
149	 - -	Annual Report of Smithsonian Institution for 1861,	8 v o.	464	1862	
150	•	Annual Report of Smithsonian Institution for 1862,	8 v o.	446	1863	
151	i	Smithsonian Contributions to Knowledge,	S. C. x 111,	55 8	1863	
152	Carpenter, P. P.	Lectures on Mollusca,	8 v o.#	140	1861	
153	Guyot, A.	Tables, Meteorological and Physical,	M. C. 1,	63 8	1859	
154		List of Foreign Correspondents of Smithsonian Institution,	M. C. v,	56	1862	
155	Whittlesey, Ch.	Ancient Mining on Lake Superior	S. C. xiii,	34	1863	1
156	Egleston, T.	Catalogue of Minerals,	M. C. VII,	56	1863	.50
157	Henry, Jos. and Coffin, J. H.	Results of Meteorological Observations from 1854 to 1859,	4to.#	1270	1661	
158	, 1	Smithsonian Miscellaneous Collections,	M. C. ▼,	774	1864	
159	Mitchell, S. W. & Morehouse, G. R.	Anatomy and Physiology of Respiration in Chelonia,	S. C. XIII,	50	1863	1.00
160	Gibbs, G.	Instructions for Ethnology and Philology,	M. C. v11,	56	1863	I
161	Gibbs, G.	Dictionary of the Chinook Jargon	M. C. v11,	60	1863	.50
162	Bache, A. D.	Magnetic and Meteorological Obs. at Girard Coll. Pt. IV, V,&VI,	1	78	1862	1.00
163	i	Circular on History of Grass- hoppers,	M. C. 11,	4	1860	free
164	!	Smithsonian Museum Miscellanea	M.C. v111,	88	1862	. 50
165	Allen, H.	Monograph of the Bats of North America,	M. C. v11,	•	1864	ł
166	Bache, A. D.	Magnetic Survey of Pennsylvania	S. C. x 111,	88	1863	' 1.00
167	Le Conte, Jno. L.	New Species of North America Coleoptera. Part I,	M. C. vi,	180	1866	; ;,00
168		Circular Relative to Birds from Middle and South America,	 M.C. viii,	2	1863	ltee.

No.	Author.	Title.	SIZE OR SERIES.	PAGES	DATE.	PRICE.
69		Smithsonian Miscellaneous Collections,	M. C. v1,	888	1867	
170		Comparative Vocabulary,	4to.*	2 0	1863	free
71	Loew, H.	Monograph of the Diptera of North America. Part II,	M. C. vi,	372	1864	
72	Meek, F. B. and Hayden, F. V.	Palmontology of the Upper Missouri. Part I,	S. C. xiv,	168	1865	
73	Dean, John	Gray Substance of the Medulla Oblongata,	S. C. xvi,	80	1864	2.50
74	Binney, W. G.	Bibliography of North American Conchology. Part II,	M. C. 1x,	302	1864	2.00
75	Bache, A. D.	Mag. and Met. Observ. at Girard Coll. Parts VII, VIII, & IX,		72	1864	1.00
76		Circular, Collecting North American Shells,	M. C. 11,	4	1860	free
77	Meek, F. B.	Check List of Invertebrate Fossils of North America,	M. C. v11,	42	1864	.25
78		Circular to Entomologists,	M.C. v111,	2	1860	free
'9	•	Catalogue of Publications of Societies,	M. C. 1x,	596	1866	
30	Draper, H.	Construction of a Silvered Glass Telescope,	S. C. xiv,	60	1864	1.00
81	Baird, S. F.	Review of American Birds in Smithsonian Museum. Part I,		484	1866	2.00
8 2		Results of Meteorological Observations from 1854-1859. Vol. II,		546	1864	2.50
83	Meek, F. B.	Check List of Invertebrate Fos- sils of North America,	M. C. VII,	34	1864	.25
84		Smithsonian Contributions to Knowledge,	S. C. xiv,	490	1865	
85		List of Birds in Mexico, &c.,	8vo.*	8	1863	free
86	Bache, A. D.	Mag. and Met. Observ. at Girard College. Parts X, XI, & XII,		42	1865	.50
87		Annual Report of Smithsonian Institution for 1863,	8 v o.	420	1864	
.88		Annual Report of Smithsonian Institution for 1864,	8vo.	450	1865	
89	Scudder, S. H.	Catalogue of Orthoptera of North America,	M C. viii,	110	1868	1.00
190	Henry, Joseph	Queries Relative to Tornadoes,	M. C. x,	4	1865	free

N T -	A		Siz E o R	AGES.		ICE.
No.	AUTHOR.	TITLE.	SERIES.	PA	DATE	<u>R</u>
191		Smithsonian Miscellaneous Collections,	M. C. v11,	878	1867	
192	Leidy, Joseph	Cretaceous Reptiles of the U.S.,	S. C. xiv,	142	1865	
193		Duplicate Shells from Expedition of Capt. Wilkes,	8 v o.#	4	1865	free
194	Binney, W. G & Bland, T.	Land and Fresh-Water Shells of North America. Part I,	M.C. VIII,	328	1869	
195	Bache, A. D.	Girard College Observations. Complete. Parts I to XII,	s. c.	262	1865	
196	Hayes, I. I.	Physical Observations in the Arctic Seas,	S. C. xv,	286	1867	
197	Whittlesey, Ch.	Glacial Drift of Northwestern States,	S. C. xv,	38	1866	<u>!</u>
198	Kane, E. K.	Physical Observations in the Arctic Seas. Complete,	8. C .	340	1860	
199	Newcomb, 8.	Orbit of Neptune,	S. C. xv,	116	1866	
200	Conrad, T. A.	Check List of the Invertebrate Fossils of North America,	M. C. vii,	46	1866	.1
201	Stimpson, Wm.	Hydrobiins and Allied Forms,	M. C. v11,	64	1865	
202	Pumpelly, B.	Geological Researches in China, Mongolia, &c.	S. C. xv,	173	1866	3.5
203		List of Works published by Smithsonian Institution,	M. C. v11.	12	1866	
204	Cleaveland, P.	Meteorological Observations, Brunswick, Me., 1807-1859,	S. C. xvi,	- 60	1867	1.0
205		Circular for Archæology and Eth- nology,	M.C. v111,	2	1867	(Ice
206		Smithsonian Contributions to Knowledge,	8. C. xv,	620	1867	
207		Relative to Scientific Investiga- tions in Russian America,	M.C. v 111,	10	1867	
208	Pickering, Chas.	Gliddon Mummy Case in Smith- sonian Institution,	S. C. xv1,	6	1867	.\ .\
209		Annual Report of the Smithsonian Institution for 1865,	8 v o.	496	1866	
210			M.C. v111,		1860	
211		Smithsonian Contributions to Knowledge,	S. C. xv i,	408	1870	þ

No.	AUTHOR.	Title.	Size OR Series.	PAGES.	DATE.	PRICE.
212		Smithsonian Miscellaneous Collections,	M.C. v111,	921	1869	
213		Smithsonian Miscellaneous Collections,	M. C. 1x,	918	18 6 9	
214		Annual Report of Smithsonian Institution for 1866,	8 v o.	470	1867	
215		Annual Report of Smithsonian Institution for 1867,	8 v o.	506	1868	
216		List of Photograph Portraits of North American Indians,	M. C. xıv,	42	1867	
217	Hoek, M.	Meteoric Shower, 1867, Nov. 13,	8 v o.#	4	1867	
218	Morgan, L. H.	Systems of Consanguinity and Affinity,	S. C. xv 11,	616	1869	
219	Osten Sacken, R.	Monograph of Diptera of North America. Part IV,	M .C. v111,	3 58	1869	2.00
2 20	Swan, Jas. G.	Indians of Cape Flattery,	S. C. xv1,	118	1870	
221	Coffin, James H.	Orbit, &c., of Meteoric Fire Ball, July 20, 1860,	S. C. xv1,	56	1869	1.00
222	Schott, Chas. A.	Tables of Rain and Snow in United States,	S.C.xviii,	178	1872	3 .00
223	Gould, B. A.	On the Transatlantic Longitude,	S. C. xvi,	110	1869	1.00
224		Annual Report of Smithsonian Institution for 1868,	8 v o.#	473	1869	
225		List of Foreign Correspondents of Smithsonian Institution,	8 v o.#	56	1870	
226		List of Publications of Smithson- ian Institution,	8 v o.	34	1869	
227	Gill, Theod.	Families of Mollusks,	M. C. x,	65	1871	.25
228		Annual Report of Smithsonian Institution for 1869,	8vo.	430	1871	
229		Smithsonian Contributions to Knowledge,	S. C. xv 11,	616	1871	į
230	Gill, Theod.	List of Families of Mammals,	M. C. x1,	104	1872	.25
231	Baird, S. F.	Memoranda Food Fishes,	M. C. x,	8	1871	
232	Stockwell, J. N.	Secular Variations of Orbits of Planets,	S. C. xv iii,	220	1872	2.00
233	Ferrel, Wm.	Converging Series, Ratio of Diameter, and Circum. of Circles,		6	1871	.25
234	Baird, S. F.	Circular Relative to Food Fishes,	M. C. x,	14	1871	free

	 -		í	1.	1 1	
No.	Autror.	Title.	Size or	AGES.	ATR.	R:OK.
			SERIES.	ď	a	A .
2 35	Henry, Joseph	Circular Relative to Thunder- storms,	M. C. x,	2	1871	free
236	Henry, Joseph	Circular Relative to Heights,	M. C. x,	2	1871	free
237	Henry, Joseph	Circular Relative to Lightning-rods,	M. C. x,	4	1871	free
238	Rhees, Wm. J.	List of American Libraries, and Public Institutions,	M. C. x,	256	1872	free
239	Harkness, Wm.	Magnetic Observations on the Monadnock,	s. C. xvni,	226	1872	2.00
240	Barnard, J. G.	Problems of Rotary Motion,	8. C. xix,	56	1872	
241	Wood, H. C.	Fresh-Water Algæ of N. America,	S. C. xix,	272	1872	7.50
242	Clark, H. J.	Lucernariæ.	8. C. xxu,	138	1878	7.50
243		List of Foreign Correspondents of Smithsonian Institution,	M. C. x,	68	1872	
244		Annual Report of Smithsonian Institution for 1870,	810.	494	1871	i
245		Check List of Smithsonian Publications to July, 1872,	M. C. x,	21	1872	
246		Smithsonian Contributions to Knowledge,	8. C. xvm,	646	1872	
247	Gill, Theod.	List of Families of Fishes,	M. C. x1,	96	1872	
248	Hilgard, E. W.	Geology of Lower Louisiana,	S. C. xxiii,	38	1872	
249		Annual Report of Smithsonian Institution for 1871,	8 v o.	473	1873	
25 0		Smithsonian Miscellaneous Col- lections,	M. C. x,	1	1873	
2 51	Rau, Charles	Memoir of Von Martius,	R. 1869,		1871	,
252	Carpenter, P. P.	American Mollusca,	M. C. x,	ł	1872	
253	Tryon, G. W.	Mon. of Strepomatidæ. Land and Fresh Water Shells. Part IV,	M. C. xvi,	1	1873	
254	De Saussure, H.	Monograph of Hymenoptera,	M. C. xi.;		ľ	2.00
2 55	Clarke, F. W.	Specific Gravity Tables. Part I,	M. C. x11,	1	1873	
256	Loew, H.	Monograph Diptera. Part III,	M. C. x1,	Ĺ	I	200
257	Baird, S. F.	Systematic List of For. Corresp.	M. C. x,		1872	
258	Watson, S.	Botanical Index,	M. C. xv,	484	1878	2.00
259	Jones, Jos.	Antiquities of Tennessee,	S. C. xx11,	181	1876	3.94
260		Regulations of Smith. Inst'n,	*	42	1872	

No.	Auteor.	Title.	Size or Series.	PAGES.	DATE.	PRICE.
261	Packard, A. S.	Directions for Collecting and Preserving Insects,	M. C. x1,	60	1873	free
262	Newcomb, S.	Orbit of Uranus,	8. C. x1x,	296	1873	3.00
2 63		Astronomical Telegram Circular,	M. C. x11,	4	 18 73	ı
264	LeConte, J. L.	New Species Coleoptera. Part II,	M. C. x1,	74	1873	
26 5	LeConte, J. L.	Classification Coleopters. Part II,	M. C. x1,	72	1873	
266	Woodward, J. J.	Toner Lecture I. Cancerous Tumors,	M. C. xv,	44	1873	.25
267	Swan, J. G.	Haidah Indians,	8. C. xx1,	22	1874	1.00
26 8	Coffin, J. H.	Winds of the Globe,	8. C. xx,	781	1875	ì
	Rabel, Simeon, Osten Sacken,	Sculptures of Santa Lucia Cosumal- whusps in Gustemuls.	S. C. xx11,	94	1878	
	C. B.	Catalogue of Diptera of North Am.,	M. C. xvi,	324	1878	2.00
2 71		Annual Report of Smithsonian Institution for 1872,	8 v o.	456	1873	
272	 - 	Smithsonian Contributions,	S. C. xix,	660	1874	
273		Miscellaneous Collections,	M. C. x1,	796	1874	İ
274		Miscellaneous Collections,	M. C. x11,	767	1874	l
2 75	<u> </u>	Annual Report of Smithsonian Institution for 1873,	8 v o.	452	 1874	ı
276	Clarke, F. W.	Specific Heat Tables. Part II,	M. C. xiv,	58	1876	.50
277	Schott, C. A.	Temperature Tables,	S. O. xxi,	360	1876	3.00
2 78		Check List of Smithsonian Publications,	M. C.	24	1874	
279	DaCosta, J. M.	Toner Lecture III. The Heart,	M. C. xv,	32	1874	. 25
2 80	Alexander, S.	Harmonies of Solar System,	S. C. xxi,	104	1875	1.00
281	Newcomb, S.	Planetary Motion,	8. C. xxi,	40	1874	
282	Wood, H. C.	Toner Lecture IV. Study of Fever,	M. O. xv,	50	1875	.25
2 83	Gill, Theod.	Catalogue of Fishes,	M. C. XIV,	56	1875	.25
284		Smithsonian Contributions,	S. C. xx,	794	1876	l
2 85		Smithsonian Contributions,	S. C. xxi,	543	1876	ı
286		Annual Report of Smithsonian Institution for 1874,	8 v o.	416	1875	
2 87	Rau, Charles,	Archæological Collection, Nat. Museum,	8. C. xx11,	118	1876	2.00

No.	Author.	TITLE.	SIZE OR SERIES.	PAGES.	DATH.	PRIOR.
288	Clarke, F. W.	Specific Gravity Tables. Supp. I,	M. C. x17,	62	1876	,50
289	Clarke, F. W.	Tables, Expan. by Heat. Pt. III.	M. C. XIV,	59	1876	.50
290		List of Smithsonian Publications,	8 v o. X 111,	12	1876	
291	Brown-Séquard,	Toner Lecture II. The Brain,	M. C. xv,	26	1877	.25
292	Cope, E. D.	Batrachia. Bulletin National Museum, No. 1,	M. C. x111,	106	! 1875	
29 3	Kidder, J. L., Coues, E.	Birds Kerguelen Island. Bulle- tin National Museum, No. 2,	M. C. x111,	61	1875	
294	Kidder, J. L., and others,	Nat. Hist. Kerguelen Island. Bulletin Nat. Museum, No. 3,	 M . C. x1111,	122	 1876 	ı
295	Lawrence, G. N.	Birds of Mexico. Bulletin Nat. Museum, No. 4,	M. C. xiti,	56	1875	
296	Goode, G. B.	Fishes of Bermuda. Bull. Nat. Museum, No. 5,	M. C. x111,	82	 1876 	
297	Goode, G. B.	Classification of Animal Resources, etc. Bulletin Nat. Museum, No. 6,	M. C. xiii,	139	1876	
298		Annual Report of Smithsonian Institution for 1875,		422	1876	l
299		Annual Report of Smithsonian Institution for 1876,	8 v o.	488	1877 	1
300	Keen, W. W.	Toner Lecture V. Continued Fevers,	M. C. xv,	72	 1877 	.25
301		Check List of Smithsonian Publications to July, 1877,	M. C. xiv,	72	1877	I
802	Adams, Wm.	9	M. C. xv.	20	 1877 	.25
303	Streets, Thos. H.	Natural History of Hawaiian and Fanning Islands and Lower California. Bulletin National Museum, No. 7,	M. C. xiii,	172	1877	
304	Dall, Wm. H.	Index to names applied to Bra- chiopoda. Bulletin Nat. Mus., No. 8,	M. C. xiii,	88	1877	
305	Jordan, David S.	North Amer. Ichthyology, No. 1. Review of Rafinesque's N. Am. Fishes. Bulletin of Nat. Mus., No. 9,	M. C. xnī,	53	1877	
306	Jordan, David S.	North Amer. Ichthyology, No. 2. Notes on Cottidae, &c. Bull. Nat. Mus., No. 10.	M. C. xnı,	120	1877	
807	Baird, S. F.	Report on Centennial Exhibition of 1876,	R. 1876,	22	1877	

No.	Author.	Title.	SIZE OR SERIES.	PAGES.	DATE.	PRICE.
308	Jordan, D. S., and Brayton, A. W.	North Amer. Ichthyology, No. 3 Distribution of Fishes of S. C Ga., and Tenn. Bulletin Nat Museum, No. 12,	м. с.	237	1878	
309		List of Foreign Correspondents of the Smithsonian Institution to Jan. 1878,	M. C. xv,	120	1878	
310	Barnard, J. G.	Internal Structure of the Earth,	S. C. xxiii,	19	1877	.25
311	Holden, Edw. S.	Index Catalogue of Books Relat- ing to Nebulæ,	M. C. xiv,	126	1877	.50
312	,	Smithsonian Miscellaneous Coll. (Bull. N. M., 1-10.)	M. C. x111,	982	1878	
3 13	Eggers, Baron,	Flora of St. Croix and Virgin Islands. Bull. Nat. Mus., No. 13,		136	1879	
314		Smithsonian Miscellaneous Collections.	M. C. xıv,	911	1878	
315		Smithsonian Miscellaneous Collections.	M. C. xv,	880	1878	
316		Circular in Reference to American Archæology,	M. C. xv,	15	1878	free
317	Elliot, D. G.	Classification and Synopsis of Trochilids,	S. C. xxiii,	289	1879	3. 0 0
318	Dall, Wm. H.	Remains of Man from Caves in Aleutian Islands.	S. C. xx11,	44	1878	2.00
8 19	Baird, S. F.	Circular. Inquiries Relative to Crawfish and Crustaces,	M. C. xv,	8	1878	free
320	Baird, S. F.	Circular Relating to Collections of Living Reptiles,	M. C. xv,	2	1878	free
3 21	Shakespeare, E. O.	Toner Lecture, VIL Inflamma- tion in Arteries atter Ligature, etc.		74	1879	.25
322)	Smithsonian Miscellaneous Collections.	M. C. xvi,	950	1880	
323		Annual Report of the Smithsonian Institution for 1877,	8 vo.	50 0	1878	
324		Circular relative to Scientific and Literary Exchanges,	M. C. MVI,	2	1879	free
325		Circular. Business Arrangements of the Smithsonian Institution,		7	1879	free
3 26	Goode, G. Brown,	Catalogue of Collection of Animal Resources and Fisheries of U.S. Bulletin Nat. Museum, No. 14,		367	1879	,

No.	Author.	Title.	SIZE OR SERIES.	PAGER.	DATE.	Paica.
327	Smithson, James,	Scientific writings of,	M. C. xxi,	166	1879	.75
32 8	Rhees, Wm. J.	Smithsonian Institution. Docu- ments Relative to its Origin and History,	ı	1027	1879	2.50
329	Rhees, Wm. J.	Smithsonian Institution. Jour- nals of Board of Regents and Reports of Committees,	M.C.xvIII,	851	1879	2 00
330	Rhees, Wm. J.	Smithson and his Bequest,	M. C. xxi,	76	1879	.75
3 31	Rau, Charles,	The Palenque Tablet,	S. C. xxII,	90	1879	2.00
332	1	Proceedings of the Nat. Museum for 1878. Vol I,	M. C. xix,	524	1879	1.00
333		Proceedings of the Nat. Museum for 1879. Vol. II,	M. C. xix,	504	1680	1.00
334	Elliot, D. G.	List of Described Species of Hum- ming-Birds,	M. C. xvi,	22	1879	ن2.
335	Rhees, Wm. J.	List of Principal American Libra- ries, Museums, Societies, etc.,		6	1879	free
336		Smithsonian Miscellaneous Collections,	M C. xvii,	1034	1880	
337	<u>.</u>	Smithsonian Miscellaneous Collections,	M.C. xviii,	851	1880	! !
338	Welling, J. C.	Life and Character of Joseph Henry,	M. C. xxi,	30	1880	
339	Taylor, Wm. B.	The Scientific Work of Joseph Henry,	M. C. xxi,	225	1880	
340	<u> </u> - 	Smithsonian Contributions to Knowledge,	8. C. xx11,	544	1880	·
34 1		Annual Report of Smithsonian Institution for 1878,	.8 ∀o .	575	1879	
342	Kumlien, L.	Contributions to Natural History of Arctic America. Bulletin of the National Museum, No. 15,		179	1879	
343	Henry, Joseph,	Annual Reports of Sec. of S. I., 1865-77,	 R.	5 4 8	1880	free
344		Check List of Smithsonian Publications,	M. C. xvi,	16	1879	free
345		Annual Report of Smithsonian Institution for 1879,	¦ 8 ∀0.	631	1850	'
346		Smithsonian Contributions to Knowledge,	8. C. xx111,	766	1881	
347	;	Cloud Charts,	•	2	1851	[160

No.	Author.	Title.	Size or Series.	PAGES.	DATE.	PRICE.
348	Baird, S. F.	Fishes of New Jersey Coast,	R. 1854,	40	1855	
349	Waring, G. E.	Toner Lecture VIII. Sanitary Drainage of Washington.	M. C.	26	1880	.25
350		Map of Stars. (For Aurora obs.)	*	1	1856	free
351	Cooper, J. G.	Forests of North America,	R. 1858,	36	1859 	I
352	Whitney, W. D.	Lectures on Linguistics,	R. 1863,	22	1864	free
353	Schott, C. A.	Tables of Rain Fall, (2d edition).	s. c.	269	1881	3.00
354	Delaunay, M.	Essay on Velocity of Light,	R. 1864,	31	1864	free
355	Wetherill, C. M.	Ozone and Antozone,	R. 1864,	12	1864	
356		Memorial of Joseph Henry,	M. C. xxi,	532	1880	free
357	Wood, H. C.	Researches on Fever,	8. C. xx1113	263	1878	2.50
358	Becker, G. J.	Atomic Weights. Constants of Nature. Part IV,	M. C.	152	1880	1.00
359		Planisphere of the Heavens,	*	1	1864	free
360	Desor, E.	Palafittes, or Lacustrian Constructions,	R. 1865,	53	1865	
361	Baegert, J.	Aborigines of California,	R. 1863-4,	41	1865	
362	Rau, Charles,	Artificial Shell Deposits in New Jersey,	R. 1864,	6	1865	
363	Lewis, James,	Instructions for collecting Land and Fresh Water Shells,	R. 1866,	8	18 6 0	free
364	Lilljeborg, W.	Classification of Birds,	R 1865,	16	1866	free
36 5	Ross, Hardisty, Jones, Gibbs,	Tinneh or Chepewyan Indians,	R. 1866,	25	1856	
366	Edwards, A. M.	Directions for collecting Diatomacea,	. *	7	1867	free
367	Rothrock, J. T.	Flora of Alaska,	R. 1867,	33	1867	free
368	Rau, Charles,	Indian Pottery,	R. 1866,	11	1867	<u> </u>
369	Abbe, C.	Dorpat and Poulkova,	R. 1867,	23	1867	
370	Rau, Charles,	Flint Implements in Illinois,	R. 1858,	9	1868	
371	Newton, H. A.	Metric Tables,	R. 1865,	23	1868	free
372	Rau, Charles,	Drilling in Stone without Metal,	R. 1868,	11	1869	
373	1	Meteorological Stations and Observers,	R. 1868,	42	1869	free
374	Schott, C. A.	Rain Charts for Summer, Winter, and Year,	S. C. XVIII,	3	1870	free

	·	R. 1870,		1870	free
	Chemistry of the Earth,	R. 1869.	98		
Babinet, M.	1	;	20	1871	free
	Diamond and Precious Stones,	R. 1870,	3 3	1872	free
Roehrig, F. L. O.	Dakota Language,	R. 1871,	19	1872	free
Henry, Joseph,	Eulogy on Alex. D. Bache,	R. 1870,	28	1872	free
Peabody, A. P.	Scientific Education of Mechan- ics,	R. 18 72,	13	1873	free
Schott, C. A.	Temperature Chart of U. S. for Year, etc.,	•	1	1873	free
Rau, Charles,	North American Stone Implements,	R. 1872,	16	1873	
Bransford, J. F.	Archmological Researches in Nicaragua,	s. c.	100	1881	2.5
Baird, S. F.	Circular for Shipping Fresh Fish,	M. C.	4	188	free
Rau, Charles,	Ancient Aboriginal Trade in North America,	R. 1872,	49	1873	
Brezina, A.	Crystallography,	B. 1872,	36	1874	free
Schott, C. A.	Temperature Charts, Summer Winter, Year,	8. C. xxi,	3	1874	free
Schott, C. A.	Temperature Chart—Year,	*	1	1874	free
Henry, Joseph,	Investigation of Illuminants,	R. 1880,	25	1881	free
Hilgard, J. E.	Tides and Tidal Action in Harbors,	R. 1874,	22	1875	free
	Act to Establish Smithsonian Institution,	M.C.xviii,	10	1875	free
Romer, F. F.			9	1877	
Gillman, Henry,	Mound-builders. Ancient Man in Michigan,			ı	
Abbott, C. C.	Stone-age in New Jersey,	R. 1875,			
Taylor, Wm. B.	Kinetic Theories of Gravitation,	R. 1876,	80	1877	fre
McParlin, T. A.	History and Climate of New Mexico,	R. 1877,	30	1877	frei
Mason, Otis T.	Latimer Collection of Antiqui- ties,	R. 1876,	23	1877	fre
Abbe C.	Meteorological Memoirs (Trans- lations),	R. 1877,	104	1878 1878	fret
	Schott, C. A. Rau, Charles, Bransford, J. F. Baird, S. F. Rau, Charles, Bresina, A. Schott, C. A. Henry, Joseph, Hilgard, J. E. Romer, F. F. Gillman, Henry, Abbott, C. C. Taylor, Wm. B. McParlin, T. A. Mason, Otis T.	ics, Schott, C. A. Temperature Chart of U. S. for Year, etc., Rau, Charles, Bransford, J. F. Archæological Researches in Nicaragua, Baird, S. F. Circular for Shipping Fresh Fish, Ancient Aboriginal Trade in North America, Bresina, A. Crystallography, Schott, C. A. Temperature Charts, Summer Winter, Year, Schott, C. A. Henry, Joseph, Investigation of Illuminants, Hilgard, J. E. Tides and Tidal Action in Harbors, Act to Establish Smithsonian Institution, Romer, F. F. Prehistoric Antiquities of Hungary, Gillman, Henry, Mound-builders. Ancient Man in Michigan, Abbott, C. C. Stone-age in New Jersey, Kinetic Theories of Gravitation, MoParlin, T. A. Mistory and Climate of New Mexico, Mason, Otis T. Latimer Collection of Antiquities, Meteorological Memoirs (Trans-	ics, Schott, C. A. Bau, Charles, Bransford, J. F. Bransford, J. F. Circular for Shipping Fresh Fish, M. C. Ancient Aboriginal Trade in North America, Bresina, A. Crystallography, Schott, C. A. Temperature Charts, Summer Winter, Year, Schott, G. A. Henry, Joseph, Hilgard, J. E. Tides and Tidal Action in Harbors, Act to Establish Smithsonian Institution, Romer, F. F. Gillman, Henry, Abbott, C. C. Taylor, Wm. B. Kinetic Theories of Gravitation, McParlin, T. A. History and Climate of New Mexico, Masson, Otis T. Latimer Collection of Antiquities, R. 1876, Meteorological Memoirs (Trans-	Schott, C. A. Temperature Chart of U. S. for Year, etc., Rau, Charles, Bransford, J. F. Archæological Researches in Nicaragua, Baird, S. F. Circular for Shipping Fresh Fish, M. C. Rau, Charles, Brexina, A. Crystallography, Brexina, A. Crystallography, Bchott, C. A. Temperature Charts, Summer Winter, Year, Schott, C. A. Temperature Chart—Year, Investigation of Illuminants, R. 1872, Act to Establish Smithsonian Institution, Romer, F. F. Prehistoric Antiquities of Hungary, Mound-builders. Ancient Man in Michigan, Abbott, C. C. Stone-age in New Jersey, R. 1875, Mason, Otis T. Mason, Otis T. Meteorological Memoirs (Trans-	Schott, G. A. Temperature Chart of U. S. for Year, etc., Rau, Charles, Bransford, J. F. Archæological Researches in Nicaragua, Baird, S. F. Circular for Shipping Fresh Fish, M. C. Ancient Aboriginal Trade in North American, Bresina, A. Crystallography, Schott, C. A. Temperature Charts, Summer Winter, Year, Schott, G. A. Temperature Chart—Year, R. 1872, Schott, G. A. Temperature Chart—Year, Kenry, Joseph, Hilgard, J. E. Tides and Tidal Action in Harbors, Act to Establish Smithsonian Institution, Prehistoric Antiquities of Hungary, Gillman, Henry, Mound-builders. Ancient Man in Michigan, Abbott, G. C. Stone-age in New Jersey, R. 1876, McParlin, T. A. History and Climate of New Mexico, Masson, Otis T. Latimer Collection of Antiquities. Meteorological Memoirs (Trans- Meteorological Memoirs (Trans-

No.	AUTHOR.	TITLE.	Size or Series.	PAGES.	DATE.	PRICE.
399	Holmgren, F.	Color-blindness,	R. 1877,	72	1878	free
400	Jones, Chas. C.	Aboriginal Structures in Ga.,	R. 1877,	13	1878	free
401	Weismann, A.	Change of Mexican Axolotl to Amblystoma,	R. 1877,	29	1878	free
403	Rau, Charles,	Stock in - trade of Aboriginal Lapidary,	R. 1877,	9	1878	free
403	Rau, Charles,	Gold Ornament from Florida,	R. 1877,	6	1878	free
404	Haldeman, S. S.	Polychrome Bead from Fiorida,	R. 1877,	6	1878	free
405	Taylor, Wm. B.	Henry and the Telegraph,	R. 1878,	103	1879	free
406	Henry, Joseph,	Researches in Sound,	R. 1878,	106	1879	free
407	Gray, Asa,	Memoir of Joseph Henry,	R. 1878,	35	1879	free
408	Baird, S. F.	Report of Secretary of Smith- sonian Institution for 1878,	R. 1878,	60	1879	free
409	Sherman, W. T. Parker, P. and Baird, S. F.	Report of Museum Building Commission and of the Archi tects for 1879,	R. 1879,	18	1880	free
410	Holden, E. S.	Reports of Astronomical Observatories for 1879,	R. 1879,	60	1880	free
411	Lautenbach, B. F.	Irritation of Polarised Nerve,	R. 1878,	59	1880	free
412	Cope, Edw. D	Zoological Position of Texas. Bull. Nat. Mus., No. 17.	M C.	51	1880	
413	Goode, G. B.	Berlin Fishery Exhibit. Bull. Nat. Mus., No. 18.	M. C.	278	1880	
4i4	Schott, C. A.	Base-Chart of the U.S.	*	1	1880	.15
+15	Knight, Edw. H.	Savage Weapons,	R. 1879,	90	1880	free
4!8		Smithsonian Miscellaneous Collections,	M. C. xix,	1034	1880	
417	Mayer, Alfred M.	Henry as a Discoverer,	M. C. xxi,	36	1880	free
418	Baird, S. F.	Report of Secretary of Smith- sonian Institution for 1879,	R. 1879,	76	1880	free
419	Baird, S. F.	Report of Secretary of Smith- sonian Institution for 1880,	R. 1880,	88	1881	free
420	Mason, Otis T.	Anthropological Investigations in 1879,	R. 1879,	30	1881	free
421	Boehmer, G. H.	Index to Smithsonian Anthropological Articles,	R. 1879,	10	1881	free
422	Ridgway, Robt.	Nomenclature of N. American Birds Bull. Nat. Mus., No. 21,	M. C.	94	1881	.25

No.	Author.	TITLE.	Size or Series.	PAGES.	DATR.	Paics.
42 3		Smithsonian Miscellaneous Collections,	M. C. xx,	846	1881	
424		Smithsonian Miscellaneous Collections,	M. C. xx:,	773	1881	
42 5		Proceedings of National Museum for 1880. Vol. III,	M. C.	594	1881	
4 26	Holden, E. S. and Hastings, C. S.	Synopsis of Herschel's Writings,	R. 1880,	118	1881	fre
427	Holden, E. S.	Recent Progress in Astronomy,	R. 1880,	39	1881	fre
42 8	Hawes, G. W.	Recent Progress in Geology and Mineralogy,	R. 188 0 ,	30	1881	fre
129	Barker, G. F.	Recent Progress in Physics and Chemistry,	R. 1880,	65	1881	ſre
13 0	Farlow, W. G.	Recent Progress in Botany,	R. 1880,	19	1881	fre
431	Gill, Theodore,	Recent Progress in Zoology,	R. 18 £0,	62	1881	fre
132	Mason, Otis T.	Recent Progress in Anthro-	R. 1880,	51	1881	fre
133	Mason, Otis T.	Visit to Luray Cavern, Va.	R. 1880,	12	1881	fre
434	Sherman, W. T. Parker, P. and Baird, S. F.	Report of National Museum Building Commission and of the Architects for 1880,		12	1881	fre
435	Loud, F. H.	Discussion of Snell's Barometric Observations,	R. 1880,	23	1881	fre
436		List of Periodicals received by Smithsonian Institution,	R. 1880,	9	1881	íre
437		Check List of Publications of Smithsonian Institution,	M. C.	22	1881	fre
43 8	Holden E. S. and Boehmer, G. H.	Reports of Astronomical Observatories for 1880,	R. 1880,	1 2 8	1881	fre

WORKS IN PREPARATION.

Monograph of Chitonidæ. By P. P. CARPENTER.

Bibliography of the Fishes of Western North America. By Theodore Gill. Bulletin of the National Museum, No. 11.

Synopsis of the Fishes of the United States. By DAVID S. JORDAN. Bulletin of the National Museum, No. 16.

An Index of Names used for Zoological Genera, comprising 70,000 titles. By Samuel H. Scudder. 8vo. Bulletin of the National Museum, No. 19.

The Writings of American Zoologists. Index Bibliography, No. I. Publications of Spencer Fullerton Baird. By G. Brown Goode. 8vo. Bulletin of the National Museum, No. 20.

Flora of the District of Columbia. By LESTER F. WARD. Bulletin of the National Museum, No. 22.

Collector's Manual of Marine Zoology. By RICHARD RATHBUN. Bulletin of the National Museum, No. 23.

Tables showing the amount of Precipitation of Rain and Snow for each Month and Year at upwards of 2000 stations in the United States. By Charles A. Schott and E. H. Courtenay.

Regulations for the Telegraphic Announcement of Astronomical Discoveries. By S. F. Baird.

The Constants of Nature. Part V. A Revision of the Atomic Weights. By Frank W. Clarke.

List and Index of Publications of the Smithsonian Institution from 1846 to 1881. By Wm. J. Rhees.

List of Foreign Correspondents. By GEO. H. BOEHMER.

			•	•
•			•	
		•		
	•		•	
		•		
		•	•	
	•			
	•	•		
·				
				-





SMITHSONIAN MISCELLANEOUS COLLECTIONS.

CATALOGUE OF PUBLICATIONS

OF THE

SMITHSONIAN INSTITUTION,

(1846 - 1882,)

WITH AN

ALPHABETICAL INDEX OF ARTICLES

IN THE

SMITHSONIAN CONTRIBUTIONS TO KNOWLEDGE, MISCELLANEOUS COL-LECTIONS, ANNUAL REPORTS, BULLETINS AND PROCEEDINGS OF THE U. S. NATIONAL MUSEUM, AND REPORT OF THE BUREAU OF ETHNOLOGY.

BY WILLIAM J. RHEES,

CHIEF CLERK OF THE INSTITUTION.

WASHINGTON: SMITHSONIAN INSTITUTION. 1882.

JUDD & DETWEILER, PRINTERS, WASHINGTON, D. C.

CONTENTS.

									Page.
1.	Smithsonian Contributions	•	•	•	•	•	•	•	V
2.	Miscellaneous Collections	. (•	•	•	•	•	VI
3.	Annual Reports	•	•	•	•	•	•	•	VII
4.	Bulletins of the U.S. National	Mu	seum		•	•	•	•	VIII
5.	Proceedings of the U.S. Nation	nal l	Muser	ım	•	•	•	•	IX
6.	Report of Bureau of Ethnology	7	•	•	•	•	•	•	IX
7.	Copyright	•	•	•	•	•	•	•	IX
8.	Use of illustrations						•		x
9.	Size of editions	•	•	•	•	•	•		x
10.			•	_	•	•	_	•	xı
11.	Rules of distribution .	•			•	•	•	•	XII
	Form of application for publica	Ition	a.	•	•	•	_	•	XIII
13.	Price list of publications .			•	•	•	•	•	XIV
	-	•	•	•	•	•	•	•	1
	of Publications, A to Q	•	1 47	•	•	•	•	• •	9
	of Publications, Regular Ser	-			•	• •	•	•	3
JLA	SIFIED LIST OF SEPARATE PU				•	•	•	•	91 92
	Anatomy, Physiology, Medicin Anthropology	це, г	mu s	arker	y	•	•	•	92 92
	Architecture	•	•	•	•	•	•	•	94
	Astronomy	•	•	•	•	•	•	•	94
	Bibliography	•	•	•				•	95
	Biography	•	•	•	•	•	•		95
	Biology	•	•	•	•	•		•	96
	Botany	•	•	•	• .	•	•	•	96
	Chemistry and Technology	•	•	•	•	•	•	•	97
	Electricity and Magnetism	•	•	•	•	•	•	•	97
	Geology	•	•	•	•	•	•	•	98
	Mathematics	•	•	•	•	•	•	•	98
	Meteorology	•	•	•	•	•	•	•	99
	Microscopy	•	•	•	•	•	•	•	100
	Mineralogy	•	•	•	•	•		•	100
	Miscellaneous	•	•	•	•	•	•	• •	100
	Palæontology	•	•	•	•	•	•	•	102
	Philology	•	•	•	•	•	•	•	102
							(1	II)	

CLASSIFIED	LIST OF SEPAR	ATE P	TIRT.	TC A TT	\V a	-Conti	inuad			Page.
	ical Geography		UDL	LUZII		COM	mucu.			103
· Phys		•	•	•	•	•	•	•	•	103
•	estrial Physics	•	•	•	•	•	•	•	•	104
	ogy—Birds .	•	•	•	•	•	•	•	•	104
250010	Fishes .	•	•	• -	•	•	•	•	•	105
	Insects .	•	•	•	•	•	•	•	•	105
	Mammals	•	•	•	•	•	•	•	•	106
	Mollusks	•	•	•	•	•	•	•	•	106
	Radiates	•	•	•	•	•	-	•	•	107
	Reptiles		· •	•	•	•	•	•	•	107
SMITHBONIA	N CONTRIBUTION	or a	Kno	WLEI	GE—	-				
	e showing date, n						s, &c.	•	•	108
	EOUS COLLECTION		of r	ages.	illust	ration	s. &c.	•		109
	EPORTS OF THE		_				5, 65 00	•	•	
	e showing date, no				illust	ration	s, &c.	•	•	110
	ONS OF THE U. S									
Table	e showing titles, n	umber	in 8	Smiths	sonia	n serie	s, &c.	•	•	111
Bulletin	NS OF THE U.S.	NATI	ONA	L Mu	SEUM	—				
Table	e showing date, ni	ımber	of p	ages,	illust	ration	s, &c.	•	•	112
Bulletin	NS OF THE U.S. N	ATION	NAL	Muse	um, c	ollecte	ed in v	olun	nes-	_
Table	e showing date, ni	ımber	of p	ages,	illust	ration	s, &c.	•	•	112
PROCEED	INGS OF THE U.	S. NA	- OITA	NAL I	Muse	UM				
	e showing date, nu							_		113
	NS OF THE BUR		-	•			,	•	•	
	e showing date, nu						s. &c.			113
	having separate		_	•				ion l	•	***
	ded in the Annua							·	ut	114
Publications	not included in	regula	ır sei	ries of	" Co	ntribu	tions,"		ol-	448
	ons," or "Reports				•		•		•	116
	ers published in t						rledge,	givi	ng	
autho	ors, titles, and con	missio	ons o	f refe	rence	•	•	•	•	117
ALPHARETT	CAL INDEX of al	1 artic	les i	n Smi	theon	ian m	hlicet	ione.		191

PREFACE.

The present catalogue embraces all the articles published by the Smithsonian Institution from its organization in 1846 to the first of July, 1882, a period of thirty-six years.

At the beginning nothing was issued but pamphlets explanatory of the plan of the Institution and brief annual reports of the proceedings of the Board of Regents, indicated in the catalogue by the letters A, B, C, D, E, F, G, H, I, J, K, L, M, N, O, Q. An elaborate work, (P in the catalogue), by ROBERT DALE OWEN, on public architecture, with special reference to the plans of the Smithsonian Institution, prepared on behalf of the Building Committee, was printed at the expense of the Institution in 1849, but did not form part of the regular series organized by the Secretary of the Institution, Prof. Henry.

1. SMITHSONIAN CONTRIBUTIONS TO KNOWLEDGE.

The series entitled "Smithsonian Contributions to Knowledge," in quarto form, was commenced in 1848 by the publication of Squier and Davis' Ancient Monuments of the Mississippi Valley. The following "Advertisement" of the first volume, prepared by Prof. Henry, has been inserted in every succeeding volume to indicate the character and design of the series:

"This volume is intended to form the first of a series of volumes, consisting of original memoirs on different branches of knowledge published at the expense and under the direction of the Smithsonian Institution. The publication of this series forms part of a general plan adopted for carrying into effect the benevolent intentions of James Smithson, Esq., of England. This gentleman left his property in trust to the United States of America to found at Washington an institution which should bear his own name, and have for its objects 'the increase and diffusion of knowledge among men.' This trust was accepted by the Government of the United States, and an act of Congress was passed August 10, 1846, constituting the President and the other principal executive officers of the General Government, the Chief Justice of the Supreme Court, the Mayor of Washington, and such other persons as they might elect honorary members, an establishment under the name of the 'Smithsonian Institution, for the increase and diffusion of knowledge among

men.' The members and honorary members of this establishment are to hold stated and special meetings for the supervision of the affairs of the Institution and for the advice and instruction of a Board of Regents, to whom the financial and other affairs are entrusted.

"The Board of Regents consists of three members ex-officio of the establishment, namely, the Vice-President of the United States, the Chief Justice of the Supreme Court, and the Mayor of Washington, together with twelve other members, three of whom are appointed by the Senate from its own

body, three by the House of Representatives from its members, and six citizens appointed by a joint resolution of both houses. To this Board is given the power of electing a Secretary and other officers, for conducting the active

operations of the Institution.

"To carry into effect the purposes of the testator, the plan of organization should evidently embrace two objects, one, the increase of knowledge by the addition of new truths to the existing stock; the other, the diffusion of knowledge thus increased among men. No restriction is made in favor of any kind of knowledge, and hence each branch is entitled to and should receive a share of attention.

"The act of Congress, establishing the Institution, directs, as part of the plan of organization, the formation of a Library, a Museum, and a Gallery of Art, together with provisions for physical research and popular lectures, while it leaves to the Regents the power of adopting such other parts of an organization as they may deem best suited to promote the objects of the

bequest.

"After much deliberation, the Regents resolved to divide the annual income, thirty thousand nine hundred and fifty dollars, into two equal parts, one part to be devoted to the increase and diffusion of knowledge by means of original research and publications, the other half of the income to be applied in accordance with the requirements of the act of Congress to the gradual formation of a Library, a Museum, and a Gallery of Art."

(The Programme of Organization, adopted December 8, 1847, follows.)

"In accordance with the rules adopted in the Programme of Organization, each memoir in this volume has been favorably reported on by a Commission appointed for its examination. It is however, impossible, in most cases, to verify the statements of an author; and, therefore, neither the Commission nor the Institution can be responsible for more than the general character of a memoir."

The total number of papers published in the 23 volumes of "Contributions" is 119, with an aggregate of 12,456 pages, 1,567 wood cuts, 523 plates, and 16 maps, each volume averaging 541½ pages.

2. MISCELLANEOUS COLLECTIONS.

In the year 1862, another series was instituted, entitled "Smithsonian Miscellaneous Collections" each volume of which has the following preface:

"The present series, entitled "Smithsonian Miscellaneous Collections," is intended to embrace all the publications issued directly by the Smithsonian

PREFACE. VII

Institution in octavo form; those in quarto constituting the "Smithsonian Contributions to Knowledge." The quarto series includes memoirs, embracing the records of extended original investigations and researches, resulting in what are believed to be new truths, and constituting positive additions to the sum of human knowledge. The octavo series is designed to contain reports on the present state of our knowledge of particular branches of science; instructions for collecting and digesting facts and materials for research; lists and synopses of species of the organic and inorganic world; museum catalogues; reports of explorations; aids to bibliographical investigations, etc.; generally prepared at the express request of the Institution and at its expense.

"The position of a work in one or the other of the two series will sometimes depend upon whether the required illustrations can be presented more

conveniently in the quarto or the octavo form.

"In the Smithsonian Contributions to Knowledge, as well as in the present series, each article is separately paged and indexed, and the actual date of its publication is that given on its special title page, and not that of the volume in which it is placed. In many cases works have been published and largely distributed years before their combination into volumes.

"While due care is taken on the part of the Smithsonian Institution to insure a proper standard of excellence in its publications, it will be readily understood that it cannot hold itself responsible for the facts and conclusions of the authors, as it is impossible in most cases to verify their statements."

The total number of papers published in the 23 volumes of "Miscellaneous Collections" is 122, each volume averaging 882½ pages, with an aggregate of 20,299 pages, 2,868 wood cuts, and 43 plates.

3. Annual Reports.

By the act of Congress organizing the Institution it was made the duty of the "Board of Regents to submit at each session a report of the operations, expenditures, and condition of the Institution." These Annual Reports form a third series of Smithsonian publications. They consist of the reports of the Secretary to the Board of Regents of the operations and condition of the Institution; the reports of committees of the Board; reports of lectures; extracts from correspondence; original or translated articles relating to the history and progress of science, etc.

The first report was submitted by the Board to the second session of the 29th Congress, 1847, and formed an octavo pamphlet of 38 pages. A similar report was presented annually thereafter, varying in size from 64 pages to 326, printed in pamphlet form with paper covers up to 1853, when Congress ordered the report to be bound in cloth. In the volume for that year the essential portion of the contents of the preceding seven reports was reprinted,

VIII PREFACE.

and this is now considered as the first of a set of Smithsonian Reports. The number of pages was limited between 1854 and 1876 to 400. In the latter year this restriction was removed, and since then the average number of pages has been 600.

The number of copies of these reports for general distribution ordered by Congress has been very variable, the largest being 7,500 in 1874 and 1875, and the smallest 150 in 1847. The number of copies granted the Institution each year is shown in the following table:

Number of extra copies furnished the Institution by Congress for distribution.

FOR THE YEAR.	No. of Copies.	FOR THE YEAR.	No. of Copies.	FOR THE YEAR.	No. of Copies.

FOR THE YEAR.	No. of Copies.	FOR THE YEAR.	No. of Copies.	FOR THE YEAR.	No. of Copies.
1847	150	1859	2,000	1871	5,000
1848	f,000	1860	2,000	1872	5,000
1849	500	1861	2,000	1873	6,00 0
1850	1,000	1862	2,000	1874	7,500
1851	2,000	1863	2,000	1875	7,500
1852	2,000	1864	2,000	1876	6,500
1853	3,000	1865	2,000	1877	6,500
1854	2,500	1866	2,000	1878	6,500
1855	2,500	1867	2,000	1879	7,000
1856	2,500	1868	2,000	1880	7,000
1857	5,000	1869	3,000]	
1858	5,000	1870	5,000	1	

The total number of pages in the 35 volumes of Annual Reports is 14,419, average 412 pages; total number of woodcuts, 1,898.

4. Bulletins of the U.S. National Museum.

In the year 1875 a fourth series of publications (octavo) was commenced. entitled "Bulletins of the National Museum," intended to illustrate the collections of natural history and ethnology belonging to the United States. constituting the National Museum, of which the Smithsonian Institution is the custodian.

Twenty of these Bulletins have been published, with an aggregate of 3,103 pages, 45 plates, and 1 map.

5. Proceedings of the U.S. National Museum.

In imitation of the practice of those learned societies which publish periodically descriptions of new species, &c., in the form of proceedings of weekly or monthly meetings, and thus present to the world the discoveries connected with the establishment at the earliest practicable moment, it appeared to be very desirable that the National Museum should have some medium of prompt publication for announcing descriptions of specimens received, (many of which are new species,) as well as other interesting facts relative to natural history furnished by correspondents of the Institution. To meet this want a fifth series of publications, (octavo,) entitled "Proceedings of the National Museum," was commenced in 1880. They are printed in successive signatures as fast as material sufficient for 16 pages is prepared, and distributed at once to scientific societies and leading active working naturalists in this country and in Europe,* each signature having printed at the bottom of its first page the date of actual issue, for settling any questions as to priority of publication. Of this series four volumes have been published, comprising 2,221 pages, with 28 cuts and 19 plates.

6. Reports of the Bureau of Ethnology.

The sixth series of publications is the annual report (in Imperial octavo) of the Bureau of Ethnology, placed by Congress in charge of the Smithsonian Institution. The first volume of this series was issued in 1881, and consists of 638 pages, with 343 cuts, 54 plates, and 1 map.

The distribution of this volume to individuals is wholly by Members of Congress and the Director of the Bureau, Major J. W. Powell—the Institution having copies at its disposal only for the libraries on its regular list of distribution for its own full series.

7. COPYRIGHT.

No copyright has ever been secured on the publications of the Institution. They are left free to be used by compilers of books without any restrictions, except that full credit shall be given to the name of Smithson for any extracts which may be made from them.

^{*} Prof. Baird's report for 1880.

X

8. Use of Illustrations.

Copies of the wood cuts used by the Institution are granted to authors or publishers on payment of the actual cost of production of electrotypes, and promise to give proper reference to the article in which they originally appeared.

9. Size of Editions.

In the first experiments of the Smithsonian system of publication, the proper magnitude of the editions necessary to meet the immediate and future demand could not be accurately ascertained. The number of copies of the Contributions then fixed upon, has since been found inadequate, although it was larger than that usually issued by other institutions. The edition has, therefore, been augmented, until at the present time 1,000 copies of each article are set aside to be combined into volumes, and an extra number, varying with the probable demand, struck off for separate distribution, and for sale.

Each article is complete in itself, with separate paging, title, and index, and without any necessary relationship to others combined with it in the same volume.

Of the early volumes of Smithsonian Contributions, the edition, for reasons already explained, was less than of the succeeding ones, so that complete sets cannot now be furnished.

In the year 1862, the plan of stereotyping every article printed by the Institution was adopted, the plates being carefully preserved, thus making it practicable at any time to issue new editions except where expensive lithographic plates were used, a limited number, only, of impressions from these having been taken.

A number of the earlier articles in octavo were out of print before the commencement of the series of "Miscellaneous Collections," and consequently are not included in them.

The printing of the "Bulletins" and "Proceedings" is authorized by the Department of the Interior and paid for out of its fund. An edition of 1,000 copies is published, of which one-half is distributed by the Department of the Interior and one-half by the Institution. As the pages are stereotyped, the cost of additional copies is alight; and for the purpose of making sure that a sufficient number of sets will be accessible forever to

students in all parts of the world, it has been considered expedient to print 1,500 additional copies of each for incorporation in the Miscellaneous Collections.*

10. DISTRIBUTION OF PUBLICATIONS.

The distribution of the publications of the Institution is a matter which requires much care and judicious selection, the great object being to make known to the world the truths which may result from the expenditure of the Smithson fund. For this purpose the Contributions are so distributed as to be accessible to the greatest number of readers; that is, to large central libraries.

The volumes of Contributions are presented on the express condition that, while they are carefully preserved, they shall be accessible at all times to students and others who may desire to consult them, and be returned to the Institution in case the establishments to which they are presented at any time cease to exist.

These works, it must be recollected, are not of a popular character, but require profound study to fully understand them; they are, however, of importance to the professional teacher and the popular expounder of science. They contain the materials from which general treatises on special subjects may be elaborated.†

Full sets of the publications cannot be given to all who apply for them, since this is impossible with the limited income of the Institution, and, indeed, if care be not exercised in the distribution, so large a portion of the income will be annually expended on the production of copies for distribution of what has already been published that nothing further can be done in the way of new publications. It must be recollected that every addition to the list of distribution not only involves the giving of the publications which have already been made, but also of those which are to be made hereafter.‡

The rules governing the distribution of the Smithsonian publications are appended. To enable institutions not coming within their provisos, as well as individuals, to procure copies of such as may be desired, a small number is set aside and sold by the Institution at a price which is intended merely to cover the actual cost of their publication.

^{*} Prof. Baird's report for 1880.

[†] Prof. Henry's report for 1876.

[‡] Prof. Henry's report for 1873.

XII PREFACE.

11. Rules for Distribution of the Publications of the Smithsonian Institution.

To Institutions.

The publications of the Smithsonian Institution are furnished:

- 1st. To learned societies of the first class, which present complete series of their publications to the Institution.
- 2d. To libraries of the first class, which give in exchange their catalogues and other publications; or an equivalent, from their duplicate volumes.
- 3d. To colleges of the first class, which furnish catalogues of their libraries and of their students, and all publications relative to their organization and history.
 - 4th. To public libraries containing 25,000 volumes.
- 5th. To smaller public libraries, where a large district would be otherwise unsupplied.
- 6th. Institutions devoted exclusively to the promotion of particular branches of knowledge may receive such Smithsonian publications as relate to their respective objects.

To Individuals.

The gratuitous distribution to individuals, of the publications of the Institution, is restricted:

- 1st. To those who are engaged in original research in the branch of science to which the book asked for pertains.
 - 2d. To those who require it in the business of instruction.
 - 3d. To donors to the museum or library of the Institution.

18

12. Form of Application for Publications.

Date,

To the	. Smithsonian	Institution.	Washington.	D. C.
	~::::::::::::::::::::::::::::::::::::::		, ,, ,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	_, _,

In behalf of the	, we respectfully apply			
for the publications of the Smithsonian In	stitution, on condition that all vol-			
umes received shall be carefully preserved	l, be accessible to any person who			
may wish to consult them, and be returned	to the Smithsonian Institution in			
case the establishment at any time ceases t	o exist.			
1. Name of Establishment	· · · ·			
2. Location—Town				
State				
3. When established				
4. Character				
5. Buildings and property				
6. Permanent fund				
7. Annual income				
8. Volumes in library				
9. Number of persons having use of books				
to. Date of last catalogue of library				
11. What publications made				
(Send printed list if possible.)				
12. Names of Officers: President				
Secretary				
Librarian				
13. Addresses of principal scientific men connecte	d with the Establishment and subjects in			
which specially interested				
I recommend the above application.				
	Member of Congress,			
Distr	ictState.			

13. PRICE LIST OF SMITHSONIAN PUBLICATIONS.

Where no price is given the work is out of print, and cannot be furnished. Of those marked "free" the edition is limited, and copies are only given to those specially interested in the subjects to which they pertain, who are collaborators of the Institution or contributors to its library or museum.

41 0 00	80	61 00	100	es 00	100	81 00	940		900	en os	960	fraa	490	free.
♣TA M								47 80		3 0 23		11.66		free.
										95		free		.\$) 35
														6 91
				0 00										6 99
				6 00							365		425	2 (0
	66		126	1 25	186	50	246	12 00	306	50	366	free.	426	fre.
	67		127	1 25	187	50	247	50	307		367	free.	427	free
					188	50	248			_				غمنا
												_		free
		6 00				free.		A						free.
4 00	71	9 00				F 00		Iree.		60				free.
										60				free
		20		Z 00		iree.		2 00						free
								200						free
4 00		12 00		ı				2 60		_				free
50						1 00								fre
		12 00		free.		_ 00		2 00						free.
						2 00		3 00					439	3
1 00	80	1 00	140	75	200	25	260		320	free.	380	free.	440	ž
1 00	81		141	1 50	201	50	261	free.	321	25	381	free.	441	1 56
						3 50		3 00		6 00				80
2 00														1 (0
														30
50		25												ine
80		05		1 90										free. free.
				•				1 00						free.
1 00				Hee.		80				5 00				free
						95		2 00						free
		0 00		12 00				2 00		2 00				free.
		12 00						12 00						free
75										2 00			453	free
free.	94	50	154		214	1 00	274	6 00	334	25	394	free.	454	f.ee
	95	6 00			215		275		335	free.	395	free.		free
1 00				50		25								free
										6 00				free
				7 60										free
				1 00		2 00				30.00				free free
1 00				R/\	220	1 00								free
1 00		1100.						96						free.
		2 00				1 00				_				X
		2 00				1 00								free
		50												free.
				1 00	226	free.	286						466	tee
50													467	3 (1)
	108	25	168	free.	228	50	288	50	348		408	free.		G IF
1 00				8 00		12 00	289							free
1 50				fr e e.		25				free.				3d 2
					231								471	free.
				0.50	232									free free
					233									free
50										1166				6 00
										RA.				•
1 m		200												free
														73
25	119	î 👸	179	3 00	239	2 00	299	50	350	free.	419	free.	I	
	1 00 2 00 50 1 00	61 62 63 64 65 66 67 68 69 70 71 4 00 72 2 00 73 1 00 75 76 50 77 78 79 1 00 80 1 00 81 82 2 00 83 84 50 85 86 50 87 1 00 88 89 90 91 92 75 93 free. 94 1 00 96 97 98 99 100 1 00 101 102 103 104 105 106 107 108 109 110 101 102 103 104 105 106 107 108 109 110 101 101 102 103 104 105 106 107 108 108 109 110 101 101 102 103 104 105 106 107 108 108 108 108 108 108 108 108 108 108	61 1 00 62 2 00 63 1 00 64 free. 65 66 67 68 69 70 6 00 71 2 3 00 2 00 73 25 1 00 74 4 4 00 75 76 12 00 76 12 00 79 1 00 1 00 80 1 00 1 00 81 1 50 82 1 50 83 1 25 84 1 50 85 85 25 86 87 25 86 87 25 1 00 88 1 00 89 5 00 90 5 00 91 12 00 91 12 00 92 12 00 93 12 00 94 50 95 6 00 96 20 00 97 1 00 98 4 00 99 12 00 101 free. 102 103 2 00 104 105 50 106 50 107 108 25 100 109 110 111 12 00 50 106 50 107 108 25 108 25 109 12 12 00	61 1 00 121 62 2 00 123 64 free. 65 126 66 127 68 69 129 70 6 00 130 71 131 4 00 72 3 00 132 2 00 73 25 133 1 00 74 134 4 00 75 135 76 12 00 136 77 1 00 139 1 00 80 1 00 140 1 00 81 125 86 146 50 87 25 145 86 146 50 87 25 145 86 146 50 87 25 145 86 146 50 87 25 145 100 88 1 00 149 90 5 00 150 91 90 5 00 150 91 100 151 92 12 00 152 75 93 153 free. 94 50 154 95 6 00 155 1 00 96 20 00 156 97 1 00 157 98 4 00 158 99 12 00 156 1 00 101 free. 102 103 2 00 163 104 105 50 160 1 00 100 50 160 1 00 101 free. 1 00 101 free. 1 00 101 free. 1 00 102 103 1 00 104 1 00 105 106 1 00 107 108 1 109 1 151 1 12 00 170 1 100 170 1 111 12 00 170 1 100 170 1 111 12 00 170 1 170 1 111 12 00 170 1 170 1 111 12 00 170 1 170 1 111 12 00 170 1 170 1 111 12 00 170 1 170 1 111 12 00 170 1 170 1 171 170 1	61 1 00 121 25 6 00 62 2 00 63 1 00 64 free. 124 65 125 6 00 126 1 25 6 00 127 6 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	61 1 00 121 25 181 62 2 00 122 6 00 182 63 1 00 123 6 00 182 64 free. 124 184 65 66 126 127 1 25 186 67 127 1 25 187 68 128 1 00 189 70 6 00 130 1 50 190 71 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	61 1 00 121 625 181 2 00 62 2 00 63 1 00 123 6 00 183 25 66 66 126 125 186 50 67 127 1 25 187 50 68 128 25 188 50 69 129 1 00 189 1 00 70 6 00 130 1 50 100 free. 71 131 1 25 191 4 00 72 3 00 132 25 192 5 00 20 73 25 133 2 00 193 free. 194 4 00 75 135 195 195 195 195 195 195 195 195 195 19	61 1 00 121 25 181 2 00 241 63 100 123 6 00 183 25 243 100 123 6 00 183 25 243 184 65 125 6 00 185 free. 246 66 126 1 25 186 60 246 67 127 1 25 187 50 247 68 128 25 188 60 246 67 127 1 25 187 50 247 68 128 25 188 60 246 67 71 128 129 100 189 1 00 246 70 6 00 130 1 50 190 free. 250 71 1 125 191 10 251 125 191 10 125 19	61 1 00 121 25 600 182 2 50 241 \$7 50 63 1 00 123 6 00 183 25 243 free. 64 free. 124 6 125 186 50 244 1 00 125 6 00 185 free. 246 free. 128 1 25 186 50 247 80 247 80 127 1 25 187 50 247 80 248 12 00 127 1 25 187 50 247 80 248 12 00 127 1 25 187 50 247 80 248 12 00 127 1 25 187 50 247 80 248 12 00 127 1 25 187 50 247 80 248 12 00 129 1 00 189 1 00 249 250 70 6 00 130 1 50 100 free. 125 1 100 100 100 100 100 100 100 100 100	61 1 00 121 25 181 2 00 241 \$7 50 301 62 2 00 122 6 00 182 2 50 242 7 50 302 63 1 00 123 6 00 183 25 243 free. 303 64 free. 124 184 12 00 244 1 00 304 64 free. 124 184 12 00 244 1 00 306 125 125 186 50 246 120 306 67 127 1 25 187 50 247 50 307 66 122 1 100 189 1 00 249 309 309 70 6 00 129 1 00 189 1 00 249 309 309 70 6 00 130 1 50 100 free. 250 310 171 131 1 25 191 251 free. 311 131 1 25 191 251 free. 311 131 1 25 191 251 free. 311 132 25 192 5 00 252 313 20 103 free. 253 313 1 00 74 134 194 254 2 00 316 135 100 76 120 138 free. 198 2 556 2 60 316 135 777 137 78 12 00 138 free. 198 2 556 2 00 318 100 78 120 138 free. 198 2 00 257 3 00 138 free. 198 2 00 257 3 00 319 100 180 100 140 75 200 25 260 310 320 100 80 1 00 140 75 200 25 260 3 00 319 100 80 1 00 140 75 200 25 260 3 00 320 100 81 141 1 50 201 50 261 free. 321 324 50 84 1 50 144 1 00 204 1 00 264 324 50 324 5	61 1 00 121 650 181 2 00 241 87 50 301 2 6 68 1 00 123 6 00 183 2 55 243 free. 303 50 60 66 126 125 185 60 186 free. 245 free. 305 50 60 67 126 125 187 50 244 1 00 304 50 60 67 127 125 187 50 247 50 307 50 60 67 127 125 187 50 247 50 307 50 60 67 127 125 187 50 247 50 307 50 60 67 127 125 187 50 247 50 307 50 60 60 120 120 180 100 244 1 00 306 50 100 127 127 125 187 50 247 50 307 50 60 120 120 120 120 120 120 120 120 120 12	61 1 00 121 25 181 2 00 241 \$7 50 301 362 25 362 260 363 1 00 123 6 00 182 25 02 242 7 50 302 25 360 363 66	61 1 00 121 20 50 182 250 241 \$7 50 301 561 662 664 free. 64 free. 65 126 600 183 25 243 free. 305 50 363 free. 66 66 126 125 186 50 246 free. 246 free. 65 126 125 187 50 247 50 307 366 free. 66 127 1 25 187 50 247 50 307 366 free. 68 128 120 189 100 189 100 170 130 1 50 170 189 100 170 131 1 25 187 50 249 309 free. 366 free. 246 120 308 50 366 free. 246 120 308 50 366 free. 360 371 free. 371 free. 371 free. 372 free. 373 free.	61

LIST OF PUBLICATIONS

OF THE

SMITHSONIAN INSTITUTION.

Note. —A to Q indicate early publications not embraced in the regular series.

- A. Journal of Proceedings of the Regents of the Smithsonian Institution, at the city of Washington, beginning on the first Monday of September, 1846. 1846. 8vo., pp. 32.
- B. Report of the Organization Committee of the Smithsonian Institution, with the resolutions accompanying the same and adopted by the Board of Regents; also, the Will of the testator, the Act accepting the bequest, and the Act organizing the Institution. 1847. 8vo., pp. 32.
- C. Digest of the Act of Congress establishing the Smithsonian Institution. August 10, 1846. 8vo., pp. 8.
- D. Address delivered on occasion of laying the Corner Stone of the Smithsonian Institution, May 1, 1847. By George M. Dallas, Chancellor of the Institution. 1847. 8vo., pp. 8.
- E. Smithson's Bequest. Professor Henry's exposition before the New Jersey Historical Society, at its meeting in Princeton, on Wednesday, September 27. 1847. 8vo., pp. 8.
- F. First Report of the Secretary of the Smithsonian Institution to the Board of Regents; giving a Programme of Organization, and an account of the operations during the year. Presented December 8, 1847. 1848. 8vo., pp. 48.
- G. [First] Report from the Board of Regents, submitted to Congress, of the operations, expenditures, and condition of the Smithsonian Institution. Senate Doc. 211; 29th Congress, 2d Session. 1847. 8vo., pp. 38.
 - I. Second Report of the Board of Regents of the Smithsonian Institution, to the Senate and House of Representatives, showing the operations, expenditures, and condition of the Institution during the year 1847. 30th Congress, 1st Session. Senate Miscellaneous No. 23. 1848. 8vo., pp. 208.

(1)

CLASSIFIED LIS	T OF SEPARA	Ame 1	Piiri.	TC A TTC	Ng_	-Conti	hand		Page.
	Geography		. 011				aucu.	_	. 103
· Physics	· ·	•	•	•	•	•	_	_	. 103
	al Physics	•	•	•	•	•	_	_	. 104
Zoölogy-	•	•	•	•	•	•	•		. 104
	Fishes .	•	•	•	•	•	•	•	. 105
	Insects .	•	•	•	•	•	•	•	. 105
	Mammals	•	•	•	•	•	•	•	. 106
	Mollusks	•	•		•	•		•	. 106
	Radiates	•	•	•	•	•	•	•	. 107
	Reptiles	•		•	•	•	•	•	. 107
Smithsonian C	ONTRIBUTION	s то	KNO	WLED	GE-	-			
Table sho	owing date, nu	ımber	of p	ages,	illust	ration	s, &c.	•	. 108
MISCELLANEOUS	COLLECTION	vs—							
Table sho	owing date, nu	ımbeı	r of p	ages,	illust	rations	s, &c.	•	. 109
ANNUAL REPOR	RTS OF THE	[neti	TUTIC)N					
Table sho	wing date, nu	ımbeı	r of p	ages,	illust	rations	s, &c.		. 110
Publications of	OF THE U. S.	. Na	TION	L M	USEU	M —			
Table sho	wing titles, n	umbe	r in 8	Smiths	oniai	n serie	s, &c.	•	. 111
Bulletins o	•								
	owing date, nu				_		8, &c.	•	. 112
Bulletins of	•		-	•			-		
	owing date, nu								. 112
Proceedings	•		_	•			,	•	•
	wing date, nu						. Ac		. 113
			_	•			, u .	•	. 110
Publications (e _		110
	owing date, nu		_	_					
Publications hav included	ing separate : in the Annua							es bu	ut . 114
Publications not	included in	regul	ar ser	ies of	" Co	ntribu	tions,"	" C o	
•	or "Reports				-	•		•	. 116
List of papers p							ledge,	givin	_
authors, t	itles, and com	missi	ons o	f refe	rence	•	•	•	. 117
ALPHABETICAL	INDEX of all	l arti	cles i	n Smi	theon	ian nu	blicati	വ	121

PREFACE.

The present catalogue embraces all the articles published by the Smithsonian Institution from its organization in 1846 to the first of July, 1882, a period of thirty-six years.

At the beginning nothing was issued but pamphlets explanatory of the plan of the Institution and brief annual reports of the proceedings of the Board of Regents, indicated in the catalogue by the letters A, B, C, D, E, F, G, H, I, J, K, L, M, N, O, Q. An elaborate work, (P in the catalogue), by ROBERT DALE OWEN, on public architecture, with special reference to the plans of the Smithsonian Institution, prepared on behalf of the Building Committee, was printed at the expense of the Institution in 1849, but did not form part of the regular series organized by the Secretary of the Institution, Prof. Henry.

1. SMITHSONIAN CONTRIBUTIONS TO KNOWLEDGE.

The series entitled "Smithsonian Contributions to Knowledge," in quarto form, was commenced in 1848 by the publication of Squier and Davis' Ancient Monuments of the Mississippi Valley. The following "Advertisement" of the first volume, prepared by Prof. Henry, has been inserted in every succeeding volume to indicate the character and design of the series:

"This volume is intended to form the first of a series of volumes, consisting of original memoirs on different branches of knowledge published at the expense and under the direction of the Smithsonian Institution. The publication of this series forms part of a general plan adopted for carrying into effect the benevolent intentions of James Smithson, Esq., of England. This gentleman left his property in trust to the United States of America to found at Washington an institution which should bear his own name, and have for its objects 'the increase and diffusion of knowledge among men.' This trust was accepted by the Government of the United States, and an act of Congress was passed August 10, 1846, constituting the President and the other principal executive officers of the General Government, the Chief Justice of the Supreme Court, the Mayor of Washington, and such other persons as they might elect honorary members, an establishment under the name of the Smithsonian Institution, for the increase and diffusion of knowledge among

- other rivers. By Charles Ellet, Jr. 1850. 4to., pp. 64, 2 woodcuts, 1 plate. (S. C. 11.)
- 14. A Memoir on Mosasaurus, and the three Allied New Genera, Holcodus, Conosaurus, and Amphorosteus. By Robert W. Gibbes. November, 1850. 4to., pp. 14, 3 plates of 28 figures. (S. C. 11.)
- 15. Aboriginal Monuments of the State of New York. Comprising the results of original surveys and explorations; with an illustrative appendix. By E. G. SQUIER. 1850. 4to., pp. 188, 79 woodcuts, 14 plates of 33 figures. (S. C. 11.)
- 16. The Classification of Insects from Embryological Data. By Louis Agassiz, 1850. 4to., pp. 28, 8 woodcuts, one plate of 23 figures. (S. C. II.)
- 17. Memoir on the Explosiveness of Nitre, with a view to elucidate its agency in the tremendous explosion of July, 1845, in New York. By Robert Hare. 1850. 4to., pp. 20. (S. C. II.)
- 18. Report on the History of the Discovery of Neptune. By BENJAMIN APTHORP GOULD, Jr. 1850. 8vo., pp. 56.
- 19. Directions for Meteorological Observations, intended for the first class of observers. By Arnold Guyot. 1850. 8vo., pp. 40, 9 woodcuts.
- 20. Microscopical Examination of Soundings, made by the United States Coast Survey off the Atlantic coast of the United States. By J. W. Bailey. January, 1851. 4to., pp. 16 and 1 plate of 68 figures. (S. C. II.)
- 21. Fourth Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1849. 31st Congress, 1st Session. Senate Miscellaneous No. 120, 8vo., pp. 64, with appendix of 207 pp. House of Representatives Miscellaneous No. 50. 1850. 8vo., pp. 272.

Report of Prof. J. HENRY, and Proceedings of the Board.

GRAY, ASA. Account of Lindheimer's, Fendler's and Wright's botanical explorations in New Mexico and California.

Agassiz, Louis. On the formation of a museum.

List of meteorological observers.

JEWETT, G. C. Report on library and catalogue system.

JEWETT, C. C. Report on public libraries of the United States.

22. Plantæ Wrightianæ Texano-Neo-Mexicanæ. By Asa Gray. Part I. March, 1852. 4to., pp. 146, 10 plates of 127 figures. (S. C. III.

An account of a collection of plants made by Charles Wright in Western Texas, New Mexico, and Sonora, in the years 1851 and 1852.

23. Microscopical Observations made in South Carolina, Georgia, and Florida. By J. W. Bailey. 1851. 4to., pp. 48, 3 plates of 83 figures. (S. C. II.)

- 24. Ephemeris of the Planet Neptune for the year 1852. By Sears C. Walker. 1853. 4to. pp. 10. (S. C. III.)
- 25. Notices of Public Libraries in the United States of America. By Chas. C. Jewett. Printed by order of Congress as an appendix to the Fourth Annual Report of the Board of Regents of the Smithsonian Institution. 1851. 8vo., pp. 210.
- 26. Smithsonian Contributions to Knowledge. Vol. II. 1851. 4to., pp. 572, 89 woodcuts, 24 plates.

WALKER, S. C. Researches relative to Neptune. No. 3.

LIEBER, F. Vocal sounds of Laura Bridgman. No. 12.

BAILEY, J. W. Microscopical soundings off Atlantic Coast. No. 20.

ELLET, C. Physical geography of the Mississippi Valley. No. 13.

GIBBES, R. W. Mosasaurus and three allied genera. No. 14.

AGASSIZ, L. Classification of insects from embryological data. No. 16.

HARE, R. Explosiveness of nitre. No. 17.

BAILEY, J. W. Microscopical observations in S. C., Ga., Fla. No. 23.

SQUIER, E. G. Aboriginal monuments of State of New York. No. 15.

WALKER, S. C. Ephemeris of Neptune for 1848. No. 4.

WALKER, S. C. Ephemeris of Neptune for 1846, '47, '48, '49. No. 5.

WALKER, S. C. Ephemeris of Neptune for 1850. No. 6.

WALKER, S. C. Ephemeris of Neptune for 1851. No. 7.

Downes, J. Occultations visible in the United States in 1851. No. 11.

- 27. On Recent Improvements in the Chemical Arts. By James C. Booth and Campbell Morfit. 1852. 8vo., pp. 216. (M. C. 11.)
- 28. Fifth Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1850. Special session, March, 1851. Senate Miscellaneous No. 1. 1851. 8vo., pp. 145. (Extra edition of 326 pp.)

CONTENTS.

Report of Prof. J. HENRY, and Proceedings of the Board.

JEWETT, C. C. General catalogue system for libraries.

BAIRD, S. F. Report on museum, and statistics of British Museum.

Memorial of the Regents to Congress, relative to the Smithson Fund.

SQUIER, E. G. Antiquities of Nicaragua.

Report of Commission on General Stereotype Catalogue of Pub. Libraries.

Culbertson, T. A. Expedition to the Mauvaises Terres and Upper Missouri.

PORTER, T. C. List of plants of Upper Missouri.

HARRIS, E. List of birds and mammalia of Missouri river.

CULBERTSON, T. A. Indian tribes of the Upper Missouri.

JEWETT, C. C. Copyright books from 1846-1849.

29. Occultations visible in the United States during the year 1852. By John Downes. 1851. 4to., pp. 34. (S. C. III.)

- 30. Contributions to the Natural History of the Fresh Water Fishes of North America. By Charles Girard. Part I.—A monograph of the Cottoids. December, 1851. 4to., pp. 80, 3 plates of 48 figures. (S. C. III.)
- 31. A Collection of Meteorological Tables, with other tables useful in Practical Meteorology. By Arnold Guyot. 1852. 8vo., pp. 212.
- 32. Nereis Boreali-Americana: or, Contributions to a History of the Marine Algæ of North America. By William Henry Harvey. Part I.—Melanospermeæ. January, 1852. 4to., pp. 152, 12 colored plates of 29 figures. (S. C. III.)
- 33. The Law of Deposit of the Flood Tide: its Dynamical Action and Office. By Charles Henry Davis. 1852. 4to., pp. 14. (S. C. III.)
- 34. Directions for Collecting, Preserving, and Transporting Specimens of Natural History. March, 1859. 8vo., pp. 40, 6 woodcuts. (M. C. 11.)
- 35. Observations on Terrestrial Magnetism. By John Locke. April. 1852. 4to., pp. 30. (S. C. III.)
- 36. Researches on Electrical Rheometry. By A. SECCHI. May, 1852. 4to., pp. 60, 3 plates of 15 figures. (S. C. III.)
- 37. Descriptions of Ancient Works in Ohio. By Chas. WHITTLESEY. 1851. 4to., pp. 20, 7 plates of 18 figures. (S. C. III.)
- 38. Smithsonian Contributions to Knowledge. Vol. III. 1852. 4to., pp. 562, and 35 plates.

LOCKE, J. Terrestrial magnetism. No. 85.

SECCHI, A. Electrical rheometry. No. 36.

GIRARD, C. Monograph of the cottoids. No. 80.

HARVEY, W. H. Marine algo of North America. Part I. No. 32.

GRAY, A. Plantæ Wrightianæ Texano-Neo-Mexicanæ. Part I. No. 22.

DAVIS, C. H. Law of deposit of the flood tide. No. 33.

WHITTLESEY, C. Descriptions of ancient works in Ohio. No. 37.

WALKER, S. C. Ephemeris of the planet Neptune for 1852. No. 24.

Downes, J. Occultations visible in United States during 1852. No. 22.

39. Smithsonian Contributions to Knowledge. Vol. IV. 1852. 4to., pp. 426.

CONTENTS.

Riggs, S. R. Dakota Grammar and Dictionary. No. 40.

40. Grammar and Dictionary of the Dakota Language. Collected by the members of the Dakota mission. Edited by S. R. RIGGS. 1852. 4to., pp. 414. (S. C. IV.)

- 41. Memoir on the Extinct Species of American Ox. By Joseph Leidy. December, 1852. 4to., pp. 20, 4 plates of 15 figures. (S. C. v.)
- 42. Plantæ Wrightianæ Texano-Neo-Mexicanæ. By Asa Gray. Part II. February, 1853. 4to., pp. 120, 4 plates of 39 figures. (S. C. v.)
- 43. Nereis Boreali-Americana; or, Contributions to a History of the Marine Algæ of North America. By W. H. HARVEY. Part II.—Rhodospermeæ. March, 1853. 4to., pp. 262, 24 plates, colored, of 64 figures. (S. C. v.)
- 44. A Flora and Fauna within Living Animals. By JOSEPH LEIDY. April, 1853. 4to., pp. 68, 10 plates of 140 figures. (S. C. v.)
- 45. Anatomy of the Nervous System of Rana pipiens. By JEFFRIES WY-MAN. March, 1853. 4to., pp. 52, 4 woodcuts, 2 plates of 29 figures. (S. C. v.)
- 46. Plantæ Frémontianæ: or, Descriptions of Plants collected by J. C. Frémont in California. By John Torrey. 1853. 4to., pp. 24, 10 plates of 89 figures. (S. C. vi.)
- 47. On the Construction of Catalogues of Libraries, and their publication by means of separate stereotyped titles. With rules and examples. By Charles C. Jewett. 1852. 8vo., pp. 78. 1853. 8vo., pp. 108.
- 48. Bibliographia Americana Historico-Naturalis; or Bibliography of American Natural History for the year 1851. 1851. By Charles Girard. December, 1852. 8vo., pp. 64.
- 49. Catalogue of North American Reptiles in the Museum of the Smithsonian Institution. By S. F. BAIRD and C. GIRARD. Part I.—Serpents. January, 1853. 8vo., pp. 188. (M. C. II.)
 - 60. Synopsis of the Marine Invertebrata of Grand Manan: or the region about the mouth of the Bay of Fundy, New Brunswick. By Wm. STIMPSON. March, 1853. 4to., pp. 68, 3 plates of 37 figures. (S. C. vi.)
- 51. Sixth Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1851. 32d Congress, 1st session, Senate Mis. No. 108. 1852. 8vo., pp. 104.

Report of Prof. J. HENRY, and Proceedings of the Board.

JEWETT, C. C. Smithsonian library and copyright system.

BAIRD, S. F. Natural history explorations in the United States in 1851.

FOREMAN, E. Meteorological system and correspondence.

LEIDY, J. Report on fossils from Nebraska.

TURNER, W. W. Indian philology.

51. Report for 1851—Continued.

Report of Committee of American Association for Promotion of Science, on a system of combined meteorological observations for North America.

- 52. Winds of the Northern Hemisphere. By James H. Copfin. November, 1853. 4to., pp. 200, 6 woodcuts, 13 plates of 238 figures. S. C. vi.)
- 53. Catalogue of Portraits of North American Indians, and Sketches of Scenery, etc., painted by J. M. Stanley. Deposited with the Smithsonian Institution. December, 1852. 8vo., pp. 76. (M. C. II.)
- 54. Occultations of Planets and Stars by the Moon, during the year 1853. Ву Јони Downes. 1853. 4to., pp. 36. (S. C. vi.)
- 55. Smithsonian Contributions to Knowledge. Vol. V. 1853. 4to., pp. 538, 4 woodcuts, 45 plates.

CONTENTS.

LEIDY, J. Flora and fauna within living animals. No. 44.

LEIDY, J. Extinct species of American ox. No. 41.

WYMAN, J. Anatomy of the nervous system of Rana pipiens. No. 45. HARVEY, W. H. Marine algo of North America. Part II. No. 43.

GRAY, A. Plantæ Wrightianæ. Part II. No. 42.

56. Smithsonian Contributions to Knowledge. Vol. VI. 1854. 4to., pp. 484, 9 woodcuts, 53 plates.

CONTENTS.

Torrey, J. Plantæ Frémontianæ. No. 46.

TORREY, J. Batis maritima. No. 60.

TORREY, J. Darlingtonia californica. No. 61.

STIMPSON, W. Marine invertebrata of Grand Manan. No. 50.

COFFIN, J. H. Winds of the Northern Hemisphere. No. 52.

LEIDY, J. Ancient fauna of Nebraska. No. 58.

Downes, J. Occultations during the year 1853. No. 54.

57. Seventh Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1852. 32d Congress, 2d Session, Senate Mis. No. 53. 1853. 8vo., pp. 96.

CONTENTS.

Report of Prof. J. HENRY, and Proceedings of the Board.

JEWETT, C. C. Report on library and the Halliwell manuscripts.

BAIRD, S. F. Scientific explorations in America in 1852.

FOREMAN, E. Report on meteorological system.

- 58. The Ancient Fauna of Nebraska: or, a Description of Remains of Extinct Mammalia and Chelonia, from the Mauvaises Terres of Nebraska. By Joseph Leidy. June, 1853. 4to., pp. 126, 3 woodcuts. 25 plates of 126 figures. (S. C. vi.)
- 59. Account of a Tornado near New Harmony, Indiana, April 30, 1852.

with a map of the track, etc. By John Chappelsmith. April, 1855. 4to., pp. 12, 2 woodcuts, 1 map, 1 plate. (S. C. VII.)

- 60. Observations on the Batis maritima of Linnæus. By John Torrey. April, 1853. 4to., pp. 8, 1 plate of 21 figures. (S. C. vi.)
- 61. On the Darlingtonia Californica; a new pitcher-plant from Northern California. By John Torrey. April, 1858. 4to., pp. 8, 1 plate of 9 figures. (S. C. vi.)
- 62. Catalogue of the Described Coleoptera of the United States. By F. E. Melsheimer. July, 1853. 8vo., pp. 190.
- 63. Notes on New Species and Localities of Microscopical Organisms. By J. W. Bailey, February, 1854. 4to., pp. 16, 7 woodcuts, 1 plate of 39 figures. (S. C. VII.)
- 64. List of Foreign Institutions in Correspondence with the Smithsonian Institution. 1856. 8vo., pp. 16.
- 65. Registry of Periodical Phenomena. Folio, pp. 4.
- 66. The Annular Eclipse of May 26, 1854. 8vo., pp. 14, 1 map.
- 67. Eighth Annual Report of the Board of Regents of the Smithsonian Institution for the year 1853. 33d Congress, 1st Session, Senate Doc. No. 73, pp. 269. House of Representatives, Mis. Doc. No. 97. 1854. 8vo., pp. 310.*

CONTENTS.

Report of Prof. J. HENRY, and Proceedings of the Board.

JEWETT, C. C. Report on library.

BAIRD, S. F. Report on publications, exchanges, museum, and explorations.

BLODGET, L. List of meteorological observers.

PEARCE, J. A. Report of Committee of Regents on distribution of Smithsonian income.

MEACHAM, J. Minority report of Committee on income.

SMITHSON, J. Will of

RUSH, R. Letter from, relative to James Smithson.

GILBERT, D., President of the Royal Society. Notice of Smithson.

SMITHSON, J. List of papers presented by, to the Royal Society.

SMITHSON, J. Contributions to the Annals of Philosophy.

Act of Congress accepting Bequest, July 1, 1836.

Act of Congress to establish the Smithsonian Institution, August 10, 1846.

BERRIEN, J. McP. Construction of the Act establishing the Smithsonian Institution.

HENRY, J. Address on the Smithsonian Institution.

^{*}This is the first of the series of annual reports published by Congress as a bound volume.

- 67. Report for 1853—Continued.
 - EVERETT, E.; SPARKS, J.; PIERCE, B.; LONGFELLOW, H. W.; GRAY, A. Report of American Academy of Arts and Sciences on Organization of Smithsonian Institution.
 - HENRY, J. First Report of the Secretary, Dec. 8, 1847. (Reprint.)

 "Second Report of the Secretary for 1848.

 "Third Report of the Secretary for 1849.

 "Fourth Report of the Secretary for 1850.
 - " Fifth Report of the Secretary for 1851. "
 Sixth Report of the Secretary for 1852. "
- 68. Vocabulary of the Jargon or Trade Language of Oregon. By B. Rush Mitchell, with additions by W. W. Turner. April, 1853. 8vo., pp. 22.
- 69. List of Domestic Institutions in correspondence with the Smithsonian Institution. 1853. 8vo., pp. 16.
- 70. The Antiquities of Wisconsin, as Surveyed and Described, by I. A. LAPHAM. May, 1855. 4to., pp. 108, 65 woodcuts, one map, 54 plates of 112 figures. (S. I. VII.)
- 71. Archæology of the United States; or Sketches, Historical and Bibliographical, of the progress of information and opinion respecting Vettiges of Antiquity in the United States. By SAMUEL F. HAVEN. July, 1856. 4to., pp. 172. (S. C. VIII.)
- 72. A Memoir on the Extinct Sloth tribe of North America. By Joseph Leidy. June, 1855. 4to., pp. 70, 16 plates of 139 figures. S. C. VII.)
- 73. Publications of Learned Societies and Periodicals in the Library of the Smithsonian Institution. December 31, 1854. Part 1. 1855. 4to., pp. 40. (S. C. VII.)
- 74. Catalogue of Publications of the Smithsonian Institution. Corrected to June, 1862. 8vo., pp. 52. (M. C. v.)
- 75. Ninth Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1854. Senate Mis. Doc. No. 24, 33d Congress, 2d session. House of Representatives, Mis. Doc. No. 35. 1855. 8vo., pp. 464, 4 woodcuts.

Report of Prof. J. HENRY, and Proceedings of the Board.

BAIRD, S. F. Report on publications, exchanges, museum, and explorations in the years 1858 and 1854.

ALEXANDER, B. S. Report of architect.

List of meteorological stations and observers.

MARSH, G. P. Lecture on the camel.

Brainard, D. Lecture on nature and cure of bites of serpents and the wounds of poisoned arrows.

Loomis, E. Lecture on the zone of small planets between him and Jupiter.

75. Report for 1854—Continued.

CHANNING, W. F. Lecture on the American fire alarm telegraph.

REED, H. Lectures on the Union.

RUSSELL, R.; HENRY, J. Lectures and notes on meteorology.

HARE, R. On John Wise's observation of a thunderstorm.

GIBBONS, H. Climate of San Francisco.

LOGAN, T. M. Meteorological observations at Sacramento, California.

HATCH, F. W. Meteorological observations at Sacramento, California.

FROEBEL, J. Remarks contributing to the physical geography of the North American Continent.

STRANG, J. J. Natural history of Beaver Island, Michigan.

EOFF, J. Habits of the black bass of the Ohio.

HEAD, J. E. Natural history of the country about Fort Ripley, Minn.

PARVIN, J. B. Habits of the gopher of Illinois.

Mann, C. Habits of a species of salamander.

Hoy, P. R. On the amblystoma luridum, a salamander inhabiting Wis.

CARLETON, J. H. Diary of an excursion in New Mexico.

BAIRD, S. F. Fishes on the coast of New Jersey and Long Island.

Jackson, C. T. Catalogue of rocks, minerals, and ores collected on geological survey in Michigan.

LOCKE, J. Catalogue of rocks, minerals, ores, and fossils.

FOSTER, J. W. Catalogue of rocks, minerals, etc.

WHITNEY, J. D. Catalogue of rocks, minerals, etc.

OWEN, D. D. Catalogue of geological specimens.

BERLANDIER, L. Catalogue of collection of historical and geographical manuscripts, maps, etc.

HENRY, J. Circular respecting new report on libraries.

HENRY, J. Circular respecting copyrights.

76. Smithsonian Contributions to Knowledge. Vol. VII. 1855. 4to., pp. 260, 74 woodcuts, 72 plates, two maps.

CONTENTS.

CHAPPELSMITH, J. Tornado near New Harmony, Indiana. No. 59.

BAILEY, J. W. New species and localities of microscopic organisms. No. 63.

LAPHAM, I. A. Antiquities of Wisconsin. No. 70.

LEIDY, J. Extinct sloth tribe of North America. No. 72.

Publications of societies and periodicals in Smithsonian Library. Part I. No. 78.

77. Tenth Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1855. 34th Congress, 1st session, Senate Mis. Doc. 73. House of Representatives Mis. Doc. 113. 1856. 8vo., pp. 440, 79 woodcuts.

CONTENTS.

Report of Prof. J. HENRY, and Proceedings of the Board.

BAIRD, S. F. Report on publications, exchanges, museum, and explorations.

List of meteorological observers.

Correspondence:

Hamilton College, Clinton, N. Y. Examination of Spencer's Telescope.

77. Report for 1855—Continued.

AMERICAN ACADEMY OF ARTS AND SCIENCES, Boston, Mass. Thanks to Smithsonian Institution for Exchanges.

STONE, WM. J. On plaster casts of antique and modern statues, etc. ILLINOIS STATE BOARD OF EDUCATION. Meteorological system for every State.

BUTLER, A. P. Report of Senate Judiciary Committee on the Management of the Smithsonian Institution, Feb. 6, 1855.

HARVEY, W. H. Lecture on marine algæ.

Morris, J. G. Lecture on natural history as applied to farming and gardening.

Morris, J. G. Lecture on insect instincts and transformations.

CHACE, G. I. Lecture on oxygen and its combinations.

SMITH, J. L. Lecture on meteoric stones.

SNELL, E. S. Lecture on planetary disturbances.

LOGAN, T. M. On the climate of California.

Morris, O. W.; Henry, J. Quantity of rain at different heights.

GUYOT, A.; HENRY J. Directions for meteorological observations.

HENRY, J. Earthquake directions.

HENRY, J. Aurora directions.

GREEN, J. Account of a new barometer.

HENRY, J. Registration of periodical phenomena.

MASTERMAN, S. Observations on thunder and lightning.

LETTERMAN, J. Sketch of the Navajo Indians.

CLINGMAN, T. L. Topography of Black Mountain, North Carolina.

ROBINSON, E.; LUDEWIG, H. E.; SQUIER, E. G.; MURPHEY, H. C.; HODGSON, W. B.; IRVING, W.; PRESCOTT, W. H.; SPARKS, J.; BANCROFT, G.; HAWKS, F. L. Communications relative to publication of Spanish works on New Mexico, by Buckingham Smith. Muller, J. Report on recent progress in physics—Galvanism.

78. Smithsonian Contributions to Knowledge. Vol. VIII. 1856. 4to., pp. 564, 52 woodcuts, 9 plates.

CONTENTS.

HAVEN, S. F. Archæology of the United States. No. 71.

OLMSTED, D. Recent secular period of aurora borealis. No. 81.

ALVORD, B. Tangencies of circles and of spheres. No. 80.

Jones, J. Chemical and physiological investigations relative to vertebrata. No. 82.

Force, P. Auroral phenomena in higher northern latitudes. No. 84. Publications of societies and periodicals in Smithsonian Library. Part II. No. 85.

- 79. New Tables for determining the Values of the Coefficients in the Perturbative Function of Planetary Motion, which depend upon the ratio of the mean distances. By John D. Runkle. November, 1856. 4to., pp. 64. (S. C. 1x.)
- 80. The Tangencies of Circles and of Spheres. By Benjamin Alvord. January, 1856. 4to., pp. 16, 25 woodcuts, 9 plates of 20 figures. (S. C. VIII.)

- 81. On the recent Secular Period of the Aurora Borealis. By Denison Olmsted. May, 1856. 4to., pp. 52. (S. C. VIII.)
- 82. Investigations, Chemical and Physiological, relative to certain American Vertebrata. By JOSEPH JONES. July, 1856. 4to., pp. 150, 27 woodcuts. (S. C. VIII.)
- 83. On the Relative Intensity of the Heat and Light of the Sun upon Different Latitudes of the Earth. By L. W. MEECH. November, 1856. 4to., pp. 58, 5 woodcuts, 6 plates of 9 figures. (S. C. IX.)
- 84. Record of Auroral Phenomena observed in the higher Northern Latitudes. By Peter Force. July, 1856. 4to., pp. 122. (S. C. VIII.)
- 85. Publications of Learned Societies and Periodicals in the Library of the Smithsonian Institution, Part II. May, 1856. 4to., pp. 38. (S. C. VIII.)
- 86. Observations on Mexican History and Archæology, with a special notice of Zapotec Remains, as delineated in Mr. J. G. Sawkins's drawings of Mitla, etc. By Brantz Mayer. November, 1856. 4to., pp. 36, 17 woodcuts, 4 plates of 6 figures. (S. C. IX.)
- 87. Psychrometrical Table for Determining the Elastic Force of Aqueous Vapor, and the Relative Humidity of the Atmosphere from indications of the Wet and the Dry Bulb Thermometer, Fahrenheit. By James H. Coffin. 1856. 8vo., pp. 20. (M. C. 1.)
- 88. Researches on the Ammonia-cobalt Bases. By Wolcott Gibbs and Frederick Aug. Genth. December, 1856. 4to., pp. 72, 21 woodcuts. (S. C. IX.)
- 89. North American Oölogy. By Thomas M. Brewer. Part I. Raptores and Fissirostres. 1857. 4to., pp. 140, 5 plates of 193 figures. (S. C. x1.)
- 90. Illustrations of Surface Geology. By Edward Hitchcock. April, 1857. 4to., pp. 164, 2 woodcuts, 12 plates of 89 figures. (S. C. 1x.)
- 91. Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1856. 34th Congress, 3d session, Senate, No. 54, House of Representatives, Mis. Doc. No. 55. 1857. 8vo., pp. 468, 69 woodcuts.

Report of Prof. J. HENRY, and Proceedings of the Board.

BAIRD, S. F. Report on publications, exchanges, museum, and explorations.

List of meteorological stations and observers.

Kohl, J. G. On a collection of the charts and maps of America.

REID, D. B. Architecture in relation to ventilation, warming, lighting, fire-proofing, acoustics, and the general preservation of health.

91. Report for 1856—Continued.

HENRY, J. Syllabus of a course of lectures on physics.

HENRY, J. Acoustics applied to public buildings.

BAIRD, S. F. Directions for collecting, preserving, and transporting specimens of natural history.

GILL, T. Fishes of New York.

Guest, W. E. Ancient Indian remains near Prescott, Canada West.

SHARPLESS, T., and PATTERSON, R. Phonography.

Institutions in which phonography is taught.

WALL, G. P., and SAWKINS, J. G. Report on the survey of the economic geology of Trinidad.

BABBAGE, C. On tables of the constants of nature and art.

HENRY, J. On the mode of testing building materials, and an account of the marble used in the extension of the United States Capitol.

SMALLWOOD, C. Description of the observatory at St. Martin, Isle Jess. Canada East.

MEECH, L. W. Relative intensity of the heat and light of the sun.

MULLER, J. Report on recent progress in physics—electricity.

92. Smithsonian Contributions to Knowledge. Vol. IX. 1857. 4to., pp. 480, 45 woodcuts, 22 plates.

CONTENTS.

MEECH, L. W. Intensity of heat and light of sun upon different latted. No. 88.

HITCHCOCK, E. Illustrations of surface geology. No. 90.

MAYER, B. Mexican history and archæology, and Zapotec remains. No. 86.

GIBBS, W. and GENTH, F. A. Researches on ammonia-cobalt bases. No. 88.

RUNKLE, J. D. New tables, planetary motion. No. 79.

RUNKLE, J. D. Asteroid supplement to new tables. No. 94.

- 93. Smithsonian Meteorological Observations for the year 1855. (Printed for examination by the observers.) 1857. 8vo., pp. 118.
- 94. Asteroid Supplement to New Tables for determining the Values of $b^{(i)}$ and its derivatives. By John D. Runkle. May, 1857. 40., pp. 72. (S. C. 1x.)
- 95. Nereis Boreali-Americana: or, Contributions to the History of the Marine Algæ of North America. By William Henry Harvey. Part III.—Chlorospermeæ. March, 1858. 4to., pp. 142, 14 plates of 44 figures. (S. C. x.)
- 96. Nereis Boreali-Americana: or, Contributions to a History of the Marine Algæ of North America. By William Henry Harvey. Three parts in one vol., with 50 plates. May, 1858. 4to., pp. 568. [Nos. 32, 43, 95.]
- 97. Magnetical Observations in the Arctic Seas. By ELISHA KENT KANE.

 Made during the Second Grinnell Expedition in search of Sir John

- Franklin, in 1853-'54-'55, at Van Rensselaer Harbor, and other points on the west coast of Greenland. Reduced and discussed by Charles A. Schott. 1859. 4to., pp. 72, 1 woodcut, 2 plates. (S. C. x.)
- 98. Grammar and Dictionary of the Yoruba Language. With an introductory description of the country and people of Yoruba. By T. J. Bowen. June, 1858. 4to., pp. 232, 1 map. (S. C. x.)
- 99. Smithsonian Contributions to Knowledge. Vol. X. 1858. 4to., pp. 462, 1 woodcut, 16 plates, 1 map.

HARVEY, W. H. Marine algæ. Part III. Chlorospermeæ. No. 95. KANE, E. K. Magnetical observations in the Arctic seas. No. 97. Bowen, T. J. Grammar and dictionary of the Yoruba language. No. 98.

- 100. An Account of the Total Eclipse of the Sun on September 7, 1858, as observed near Olmos, Peru. By J. M. Gilliss. April, 1859.

 4to., pp. 22, 1 woodcut, 1 plate. (S. C. xi.)
- 101. Map of the Solar Eclipse of March 15, 1858. By Thomas Hill. January, 1858. 8vo., pp. 8, 1 plate.
- 102. Catalogue of the described Diptera of North America. By R. Osten Sacken. January, 1858. 8vo., pp. 116. October, 1859. (M. C. III.)
- 103. Meteorological Observations made at Providence, Rhode Island, from December, 1831, to May, 1860. By A. CASWELL. October, 1860. 4to., pp. 188. (S. C. XII.)
- 104. Meteorological Observations in the Arctic Seas. By E. K. KANE. Made during the Second Grinnell Expedition in search of Sir John Franklin, in 1853-'54-'55, at Van Rensselaer Harbor, and other points on the west coast of Greenland. Reduced and discussed by Charles A. Schott. November, 1859. 4to., pp. 120, 10 woodcuts. (S. C. XI.)
- 105. Catalogue of North American Mammals, chiefly in the Museum of the Smithsonian Institution. By Spencer F. Baird. July, 1857. 4to., pp. 22.
- 106. Catalogue of North American Birds, chiefly in the Museum of the Smithsonian Institution. By Spencer F. Baird. October, 1858. 4to., pp. 42.
- 107. Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1857. 35th Congress, 1st Session, Senate Mis. Doc. No. 272, House of Representatives, No. 135. 1858. 8vo., pp. 438, 100 woodcuts.

107. Report for 1857—Continued.

CONTENTS.

Report of Prof. J. HENRY, and Proceedings of the Board.

BAIRD, S. F. Report on publications, exchanges, museum, and experiences.

List of meteorological stations and observers.

STANLEY, J. M. Report of Committee of Regents on gallery of Indian portraits.

CONTAXAKI, MISS ELIZA B. Present of ornamental album from Greece.

CHASE, S. P. On telegraph.

GALE, L. D. On telegraph.

HALL, J. On telegraph.

Mason, C. On telegraph.

HENRY, J. On telegraph.

HENRY, J. On telegraph, deposition of, Sept. 1849.

HENRY, J. Communication relative to a publication by Prof. Morse.

FELTON, C. C. Report of special committee of Board of Regents on the communication of Prof. Henry relative to the electro-magnetic telegraph.

HENRY, J. History of the electro-magnetic telegraph.

LECONTE, J. Lecture on coal.

ALEXANDER, S. Lecture on vastness of the visible creation.

FENDLER, A. Meteorology and ethnology, Colonia Tovar, Venezuela. S. A.

LOGAN, T. M. Meteorology of Sacramento, California.

DEWEY, C. On best hours to find mean temperatures.

WISSNER, J. Meteorology of the District of Columbia.

MASTERMAN, S. Observations on natural phenomena, shooting staraurora, etc.

MÜLLER, J. Report on recent progress in physics. (Electricity, gar-vanism.)

- 108. Catalogue of North American Birds, chiefly in the Museum of the Smithsonian Institution. By Spencer F. Baird. 1859. 8vo. pp. 24. (M. C. II.)
- 108*. Same Title, (printed for labelling, with one side of each leaf blank.
- 109. Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1858. 35th Congress, 2d Session, Senate Mis. Doc. No. 49. House Rep., No. 57. 1859. 8vo., pp. 448, 48 woodcuts.

CONTENTS.

Report of Prof. J. HENRY, and Proceedings of the Board.

Correspondence:

Schleiden, R. Free freight between Germany and United States in Smithsonian Institution by the North German Lloyd.

SABINE, E. Or continuance of magnetic observations.

BAIRD, S. F. Report on publications, exchanges, museum, and explorations.

List of meteorological stations and observers.

CASWELL, A. Lecture on astronomy.

CUVIER, M. Memoir of Priestley.

BAIRD, S. F. Instructions for collecting nests and eggs of American birds.

BAIRD, S. F. Instructions for collecting insects.

169. Report for 1858—Continued.

LECONTE, J. L. Instructions for collecting coleoptera.

CLEMENS, B. Instructions for collecting hymenoptera.

UHLER, P. R. Instructions for collecting orthoptera.

UHLER, P. R. Instructions for collecting hemiptera.

UHLER, P. R. Instructions for collecting neuroptera.

LOEW, H.; OSTEN SACKEN, B. Instructions for collecting diptera.

CLEMENS, B. Instructions for collecting lepidoptera.

TAYLOR, A. S. Grasshoppers and locusts of America.

MOTSCHULSKY, V. On means of destroying the grasshopper.

MARTINS, C. Vegetable colonization of the British Isles of Shetland, Faroe, and Iceland.

DECANDOLLE, A. Causes which limit vegetable species towards the north in Europe and similar regions.

COOPER, J. G. Distribution of the forests and trees of North America, and catalogue of the native trees of the United States.

BLACKISTON; BLAND; WILLIS, J. R. Birds of Nova Scotia.

BLAND; WILLIS J. R. Birds of Bermuda.

DUPREZ, M. F. Atmospheric electricity.

MÜLLER, J. Recent progress in physics. (Galvanism.)

HENRY, J. Meteorological stations, cost of establishment of.

Hodgins, J. G. Meteorological stations of Upper Canada.

DUDLEY, T. Earthquake at New Madrid, Missouri.

NAILL, D. W. Dispersion of a cloud by an electrical discharge.

HARE, R. Method of forming small weights.

FRIEDLÄNDER, J. Plan of a Bibliography.

LYON, S. S. Antiquities from Kentucky.

GARDINER, R. H. Barometer, rain and snow gauges.

GUEST, W. E. Snow gauge.

GARDINER, R. H. Opening and closing of Kennebec river, Maine.

CANUDAS, A. Earthquakes in Guatemala.

Humphreys, A. A. Method of ascertaining the amount of water in rivers.

110. Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1859. 36th Congress, 1st Session, House of Representatives, Mis. Doc. No. 90. 1860. 8vo., pp. 450, 57 woodcuts.

CONTENTS.

Report of Prof. J. HENRY, and Proceedings of the Board.

BAIRD, S. F. Report of publications, exchanges, museum, and explorations.

List of meteorological stations and observers.

PEARCE, J. A. Notice of Richard Rush.

FELTON, C. C. Notice of W. W. Turner.

FELTON, C. C. Notice of Washington Irving.

BACHE, A. D. Notice of James P. Espy.

BACHE, A. D. Notice of G. Würdemann.

HENRY, J. Notice of Parker Cleaveland.

Correspondence:

DUKE OF NORTHUMBERLAND. Presentation of books.

2 HENRY, J. Account of Priestley's lens.

110. Report for 1859—Continued.

CUNARD, E. Free freight to Smithsonian Institution, between United States and England.

LOGAN, W. E. Request for duplicate shells.

Ross, B. R. Observations in Hudson's Bay Territory.

Johnson, S. W. Lectures on agricultural chemistry.

CARPENTER, P. P. Lectures on the shells of the Gulf of California.

MÆDLER, M. Movement of the stars around a central point.

DE LA RIVE, A. Report on the transactions of the Society of Physics and Natural History of Geneva, from July, 1858, to June, 1859.

RETZIUS, A. Present state of ethnology in relation to the form of the human skull.

FLOURENS, M. Memoir of Pyramus de Candolle.

AIRY, G. B. On the means which will be available for correcting the measures of the sun's distance in the next twenty-five years.

Powell, B. Report on the state of knowledge of radiant heat, made to the British Association in 1882, 1840, and 1854.

HILGARD, J. E. Description of the magnetic observatory at the Smithsonian Institution.

Poggendorff, J. C. On the use of the galvanometer as a measuring instrument.

MALLET, R. On observations of earthquake phenomena.

CASELLA, L. Description of meteorological instruments.

GREEN, J.; WÜRDEMANN, W. On filling barometer tubes.

Welsh, J. The construction of a standard barometer, and apparatus and processes employed in the verification of barometers at the Kew Observatory.

111. Smithsonian Contributions to Knowledge. Vol. XI. 1859. 4to., pp. 502, 20 woodcute, 23 plates.

CONTENTS.

Brewer, T. M. North American Oölogy. Part 1. Raptores and Fissionstres. No. 89.

GILLISS, J. M. Total eclipse of the sun, September 7, 1858, in Peru. No. 100.

BACHE, A. D. Magnetic and meteorological observations. Part I. No. 113.

KANE, E. K. Meteorological observations in the Arctic seas. No. 104 LECONTE, J. L. Coleoptera of Kansas and Eastern New Mexico. No. 126.

Sonntag, A. Observations on terrestrial magnetism in Mexico. No. 114. Loomis, E. On certain storms in Europe and America, December, 1806. No. 127.

112. Smithsonian Contributions to Knowledge. Vol. XII. 1860. 4to., pp. 538, 15 woodcuts, 3 plates.

CONTENTS.

KANE, E. K. Astronomical observations in the Arctic seas. No. 129. WHITTLESEY, C. Fluctuations of level in North American lakes. No. 119.

- 112. Contributions to Knowledge. Vol. XII—Continued.
 - CASWELL, A. Meteorological observations at Providence, Rhode Island, 28½ years. No. 103.
 - SMITH, N. D. Meteorological observations near Washington, Arkansas, 20 years. No. 181.
 - MITCHELL, S. W. Researches upon venom of the rattlesnake. No. 135.
- 113. Discussion of the Magnetic and Meteorological Observations made at the Girard College Observatory, Philadelphia, in 1840-'41-'42-'43-'44-'45. Part I. Investigation of the eleven year period in the amplitude of the solar-diurnal variation and of the disturbances of the magnetic declination. By A. D. BACHE. November, 1859. 4to., pp. 22, 5 woodcuts. (S. C. XI.)
- 114. Observations on Terrestrial Magnetism in Mexico. Conducted under the direction of Baron Von Müller, with notes and illustrations of an examination of the volcano Popocatepetl and its vicinity. By August Sonntag. February, 1860. 4to., pp. 92, 4 woodcuts, one plate. (S. C. XI.)
- 115. Proceedings of the Board of Regents of the Smithsonian Institution, in relation to the Electro-Magnetic Telegraph. 1861. 8vo., pp. 40, 7 woodcuts. (M. C. 11.)
- 116. List of Public Libraries, Institutions, and Societies in the United States and British Provinces of North America. By WILLIAM J. RHEES. 1859. 8vo., pp. 84.
- 117. Catalogue of Publications of Societies and of other Periodical Works in the Library of the Smithsonian Institution, July 1, 1858. Foreign Works. 1859. 8vo., pp. 264. (M. C. III.)
- 118. Catalogue of the Described Lepidoptera of North America. By John G. Morris. May, 1860. 8vo., pp. 76. (M. C. III.)
- 119. On Fluctuations of Level in the North American Lakes. By Charles Whittlesey. July, 1860. 4to., pp. 28, 2 plates of 4 figures. (S. C. XII.)
- 120. Results of Meteorological Observations made at Marietta, Ohio, between 1826 and 1859, inclusive. By S. P. HILDRETH. To which are added results of observations taken at Marietta, by Joseph Wood, between 1817 and 1823. Reduced and discussed by Charles A. Schott. September, 1867. 4to., pp. 52, 14 woodcuts. (S. C. xvi.)
- 121. Discussion of the Magnetic and Meteorological Observations made at the Girard College Observatory, Philadelphia, in 1840-'41'-42'-43-'44-'45. By A. D. Bache. Part II.—Investigation of the solar-diurnal variation in the magnetic declination and its annual inequality. June, 1862, 4to., pp. 28, 8 woodcuts. (S. C. XIII.)

122. Smithsonian Miscellaneous Collections. Vol. I. 1862. 8vo., pp. 733. 23 woodcuts.

CONTENTS.

Directions for meteorological observations. No. 148.

Coffin, J. H. Psychrometrical tables. No. 87.

GUYOT, A. Meteorological and physical tables. No. 153.

123. Smithsonian Miscellaneous Collections. Vol. II. 1862. 8vo., pp. 714, 33 woodcuts.

CONTENTS.

BOOTH, J. C.; MORFIT, C. Recent improvements in chemical arts. No. 27.

Proceedings of Board of Regents in relation to the electro-magnetic telegraph. No. 115.

STANLEY, J. M. Catalogue of portraits of North American Indians. No. 53.

BAIRD, S. F. Catalogue of North American birds. No. 108.

BAIRD, S. F.; GIRARD, C. Catalogue of North American reptiles. Serpents. No. 49.

Check-list shells North America. No. 128.

Directions for collecting specimens of natural history. No. 34.

HENRY, J. Circular to officers Hudson Bay Company. No. 137

Instructions for collecting nests and eggs. No. 189.

North American grasshoppers. No. 163.

North American shells. No. 176.

Morgan, L. H. Circular respecting relationship. No. 138.

124. Smithsonian Miscellaneous Collections. Vol. III. 1862. 8vo.. pp. 776, 49 woodcuts.

CONTENTS.

OSTEN SACKEN, R. Catalogue diptera North America. No. 102.

MORRIS, J. G. Catalogue described lepidoptera North America. No. 115.

LE CONTE, J. L. Classification coleoptera. I. No. 136.

Catalogue publications of societies in Smithsonian library. No. 117.

125. Smithsonian Miscellaneous Collections. Vol. IV. 1862. 8vo., pp. 762, 30 woodcuts.

CONTENTS.

HAGEN, H. Synopsis of North American neuroptera. No. 134. Morris, J. G. Synopsis of North American lepidoptera. No. 133.

- 126. The Coleoptera of Kansas and Eastern New Mexico. By John L LE Conte. December, 1859. 4to., pp. 64, 2 plates of 33 figures. 1 map. (S. C. xi.)
- 127. On Certain Storms in Europe and America, December, 1836. By ELIAS LOOMIS. February, 1860. 4to., pp. 28, 13 plates. (S. C. XI.

- 128. Check-lists of the Shells of North America. By Isaac Lea, P. P. Carpenter, Wm. Stimpson, W. G. Binney, and Temple Prime. June, 1860. 8vo., pp. 52. (M. C. 11.)
- 129. Astronomical Observations in the Arctic Seas. By E. K. KANE. Made during the Second Grinnell Expedition in search of Sir John Franklin, in 1853, 1854, and 1855, at Van Rensselaer Harbor, and other points in the vicinity of the northwest coast of Greenland. Reduced and discussed by Charles A. Schott. May, 1860. 4to., pp. 56, 3 woodcuts, 1 map. (S. C. xii.)
- 130. Tidal Observations in the Arctic Seas. By E. K. KANE. Made during the Second Grinnell Expedition in search of Sir John Franklin, in 1853, 1854, and 1855, at Van Rensselaer Harbor. Reduced and discussed by Charles A. Schott. October, 1860. 4to., pp. 90, 3 woodcuts, 4 plates. (S. C. XIII.)
- 131. Meteorological Observations made near Washington, Arkansas, from 1840 to 1859, inclusive. By NATHAN D. SMITH. October, 1860. 4to., pp. 96. (S. C. XII.)
- 132. Discussion of the Magnetic and Meteorological Observations made at the Girard College Observatory, Philadelphia, in 1840-'41-'42-'43-'44-'45. By A. D. Bache. Part III.—Investigation of the influence of the moon on the magnetic declination. June, 1862. 4to., pp. 16, 3 woodcuts. (S. C. XIII.)
- 133. Synopsis of the Described Lepidoptera of North America. By John G. Morris. Part I.—Diurnal and crepuscular lepidoptera. February, 1862. 8vo., pp. 386, 30 woodcuts. (M. C. IV.)
- 134. Synopsis of the Neuroptera of North America. With a list of the South American species. By Hermann Hagen. July, 1861. 8vo., pp. 368. (M. C. IV.)
- 135. Researches upon the Venom of the Rattlesnake. With an investigation of the anatomy and physiology of the organs concerned. By S. Weir Mitchell. December, 1860. 4to., pp. 156, 12 woodcuts. (S. C. XII.)
- 136. Classification of the Coleoptera of North America. By John L. Le Conte. Part I. March, 1862. 8vo., pp. 312, 49 woodcuts. (M. C. III.)
- 137. Circular to Officers of the Hudson's Bay Company. 1860. 8vo., pp. 6. (M. C. II; M. C. VIII.)
- 138. Circular in reference to the Degrees of Relationship among different Nations. By Lewis H. Morgan. January, 1860. 8vo., pp. 34. (M. C. 11.)

- 139. Instructions in reference to Collecting Nests and Eggs of North American Birds. January, 1860. 8vo., pp. 22, 20 woodcuts. (M. C. 11.)
- 140. List of the Coleoptera of North America. Prepared for the Smithsonian Institution. By John L. Le Conte. Part I. April, 1866. 8vo., pp. 82. (M. C. vi.)
- 141. Monographs of the Diptera of North America. By H. Loew. Part I. Edited, with additions, by R. Osten Sacken. April, 1862. 8vo., pp. 246, 15 woodcuts, 2 plates of 42 figures. (M. C. vi.)
- 142. Bibliography of North American Conchology, previous to the year 1860. By W. G. Binney. Part I.—American Authors. March, 1863. 8vo., pp. 658. (M. C. v.)
- 143. Land and Fresh-Water Shells of North America. By W. G. Binney. Part II.—Pulmonata Limnophila and Thalassophila. September, 1865. 8vo., pp. 172, 372 woodcuts. (M. C. vii.)
- 144. Land and Fresh-Water Shells of North America. By W. G. BINNEY. Part III.—Ampullariidæ, Valvatidæ, Viviparidæ, fresh-water Rissoidæ, Cyclophoridæ, Truncatellidæ, fresh-water Neritidæ, Helicinidæ. September, 1865. 8vo., pp. 128, 253 woodcuts. (M. C. VII.)
- 145. Monograph of American Corbiculadæ, (recent and fossil.) By Temple Prime. December, 1865. 8vo., pp. 92, 86 woodcuts. (M. C. VII.)
- 146. Meteorological Observations in the Arctic Seas. By Francis Lederold McClintock. Made on board the Arctic searching yacht "Fox," in Baffin Bay and Prince Regent's Inlet, in 1857, 1858, and 1859. Reduced and discussed by Charles A. Schott. May, 1862. 4to., pp. 164, 15 woodcuts, 1 map. (S. C. XIII.)
- 147. Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1860. 36th Congress, 2d Session. Senate Mis. Doc., No. 21, 1861. 8vo., pp. 448, 73 woodcuts.

HENRY, J. Secretary's report of operations.

BOARD OF REGENTS, Proceedings of

BAIRD, S. F. Report on publications, exchanges, museum, and explora-

List of meteorological stations and observers.

List of meteorological material contributed to Smithsonian Institution.

147. Report for 1860—Continued.

Correspondence:

CITIZENS OF PHILADELPHIA. Memorial relative to Lowe's aeronautic voyages across the Atlantic.

LEPSIUS, R. Presentation of books on Egypt.

MÜLLER, F. On Smithsonian exchanges.

HINCKS, WM. On Smithsonian exchanges.

HENRY, J. Reply to memorial on Lowe's aeronautic voyage.

HENRY, J. On currents of the atmosphere and aerial navigation.

ROGERS, F. Lectures on roads and bridges.

CARPENTER, P. P. Lectures on mollusca, or shell fish, and their allies.

Morlot, A. General views on archæology.

The microscope.

NICKLÉS, M. J. Scientific Congress of Carlsruhe, 1858.

CUVIER, M. Memoir of René Just Hauy.

SABINE, E. Magnetic storms.

GARDINER, R. H. Disappearance of ice.

FENDLER, A. Temperature of St. Louis, Missouri.

DEWEY, C. Best hours for temperature observations.

HENRY, J. Description of Smithsonian anemometer.

NEWTON, A. Suggestions for saving parts of the skeleton of birds.

Vollum, E. P. On the wingless grasshopper of California.

WURDEMAN, G. Specimens of flamingo, etc., from South Florida.

GESNER, W. Habits of pouched rat, or salamander of Georgia.

BARNARD, V. Birds of Chester county, Pennsylvania.

COOPER, J. G. Forests and trees of Florida and the Mexican Boundary.

- 148. Directions for Meteorological Observations and the Registry of Periodical Phenomena. 1860. 8vo., pp. 72, 23 woodcuts. (M. C. 1.)
- 149. Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1861. 37th Congress, 2d Session, House of Representatives Mis. Doc., No. 77. 1862. 8vo., pp. 464, 25 woodcuts.

CONTENTS.

HENRY, J. Secretary's report of operations.

BOARD OF REGENTS, Proceedings of

BAIRD, S. F. Report on publications, exchanges, museum, and explorations.

List of meteorological stations and observers.

List of meteorological material contributed to Smithsonian Institution.

KING, C. B. Catalogue of engravings presented to the Smithsonian Institution.

Correspondence:

FLADGATE, CLARKE, and FINCH. Smithson's residuary legacy.

KUNHARDT & Co. Free freight to Smithsonian Institution, between United States and Germany.

TROYON, F. Lacustrian habitations.

MORLOT, A. Archeology.

HILL, A. J. Map of antiquities of United States.

ROYAL ACADEMY OF SCIENCE, MADRID. Exchanges.

149. Report for 1861—Continued.

MEXICAN Soc. of Geog. and Statistics. Exchanges.

HAGEN, H. Neuroptera.

HAMILTON COLLEGE, CLINTON, N. Y. Exchange of specimens.

CODD, J. A. Acknowledgment for books.

SHEPARD, C. U. Arrangement of mineralogical collection.

University of Tobonto. Scientific co-operation.

ROYAL HORTICULTURAL SOCIETY, LONDON. Exchange of publications.

BETHUNE, C. J. Acknowledgment for books.

PEASE, W. H. Acknowledgment for shells, etc.

Circular of the Institute of Rupert's Land.

CRUMMELL, A: Facts respecting Liberia College.

GIBBS, G.; AND OTHERS. Recommendation of Shea's Indian guistics.

SHEA, J. G. Account of library of Indian linguistics.

Woolsey, T. D. Eulogy on Cornelius C. Felton.

Cox, S. S. Eulogy on Stephen A. Douglas.

ROGERS, F. Lecture on bridges.

ALEXANDER, S. Lecture on the relations of time and space.

HAYES, I. I. Lecture on Arctic explorations.

FLOURENS, M. Memoir of Geoffroy Saint Hilaire.

LAUGEL, A. The sun, its chemical analysis.

LEE, Dr. Progress of astronomical photography.

LESPIAULT, PROF. Small planets between Mars and Jupiter.

DUFOUR, C.; KÄMTZ. Scintillation of the stars.

DAUBRÉE, M. Metamorphism and the formation of crystalline rocks.

CRAIG, B. F. Nitrification.

HUNT, T. S. History of petroleum or rock-oil.

ALLEN, Z. Explosibility of coal oils.

Destructive effect of iron-rust.

TROYON, F. Lacustrian cities of Switzerland.

RUTIMEYER, A. Fauna of Middle Europe during the Stone Age.

TROYON, F. Report on ethnological collections of Museum at Lausal.nt.

TROYON, F. Archæological researches made at Concise.

PEALE, T. R. Ancient mound in St. Louis, Missourt.

GIBBS, G. Instructions for archæological investigations

HENRY, J. Circular, ancient mining in Lake Superior Copper Region

Morgan, L. H. Suggestions relative to ethnological map of Normal America.

Cours, E.; Prentiss, S. S. List of birds of the District of Columbia.

Prize Questions:

Holland Society of Science, Harlem.

Batavian Society of Experimental Philosophy of Rotterdam.

Society of Arts and Sciences of Utrecht.

Royal Academy of Netherlands.

150. Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1862. 37th Congress, 3d session, House of Representatives Mis. Doc., No. 25. 1863. 8vo., pp. 446, 94 woodcuts.

150. Report for 1862—Continued.

CONTENTS.

HENRY, J. Secretary's report of operations.

BOARD OF REGENTS, Proceedings of

BAIRD, S. F. Report on publications, exchanges, museum, and explorations.

List of meteorological stations and observers.

List of meteorological material contributed to Smithsonian Institution. Correspondence:

STANTON, E. M. Deposit of Beaufort library.

CANBY, E. Deposit of Bishop Johns' library.

SIBLEY, H. Meteorological telegrams.

ENGLEMANN, G. Heights of mountains in Colorado.

Guyor, A. On mountain measurements.

HUBBARD, J. S. Investigations of Biela's comet.

DARLINGTON, W. History of Chester county, Pa.

LYMAN, THEODORE. Ethnographical collections.

DE SCHLAGINTWEIT, H. Ethnographical collections.

DAA, L. K. Ethnological Museum of Norway.

DILLE, I. Antiquities in Missouri and Tennessee.

REID, A. Skulls and mummy from Patagonia.

GIBBS, G. Philological circular.

GIBBS, G. Ethnological map of United States.

GIBBS, G. Indian vocabularies.

WATKINS, E. A. Systems of relationship of Cree Indians.

FLACHENECKER, G. Indian languages.

PEASE, W. H. Natural history of Pacific islands.

WIRTZ, H. R. Herbarium captured in Tennessee.

METTENIUS, G. Acknowledgment for ferns.

DE SAUSSURE, H. Hymenoptera.

Romero, M. Explorations of John Xantus in Mexico.

BACHE, A. D. Eulogy on James A. Pearce.

BARNARD, F. A. P. Lectures on the undulatory theory of light.

WILSON, D. Lectures on physical ethnology.

MORLOT, A. Lecture on the study of high antiquity.

LUBBOCK, J. North American archæology.

· FLOURENS, M. Historical sketch of the Academy of Science of Paris.

FLOURENS, M. Memoir of Leopold von Buch.

FLOURENS, M. Memoir of Louis Jacques Thenard.

QUATREFAGES, M. Memoir of Isidore Geoffroy St. Hilaire.

PHIPSON, T. L. The catalytic force, or studies on the phenomena of contact.

HERSCHEL, J. On atoms.

LESLEY, J. P. On the classification of books.

RIED, A. Human remains from Patagonia.

PRIZE QUESTIONS:

London Institution of Civil Engineers.

Provincial Society of Arts and Sciences, Utrecht.

Royal Danish Society of Science.

Academy of Sciences of the Institute of Bologna.

151. Smithsonian Contributions to Knowledge. Vol. XIII. 1863. 4to., pp. 558; 80 woodcuts, 4 plates, 3 maps.

CONTENTS.

KANE, E. K. Tidal observations, Arctic seas. No. 180.

McClintock, F. L. Meteorological observations, Arctic seas. No. 145.

WHITTLESEY, C. Ancient mining on shores of Lake Superior. No. 155.

BACHE, A. D. Discussion, Girard College observations. Part II. No. 121.

BACHE, A. D. Discussion, Girard College observations. Part III. No. 182.

BACHE, A. D. Discussion, Girard College observations. Parts IV, V. VI. No. 162.

BACHE, A. D. Magnetic survey of Pennsylvania, &c. No. 166.

MITCHELL, S. W.; Morehouse, G. R. Researches upon anatomy and physiology of chelonia. No. 159.

- 152. Lectures on Mollusca or "Shell-fish," and their Allies. Prepared for the Smithsonian Institution. By Philip P. Carpenter. 1861. 8vo., pp. 140.
- 153. Tables, Meteorological and Physical, Prepared for the Smithsonian Institution. By Arnold Guyot. Third edition, revised and enlarged. 1859. 8vo., pp. 638. (M. C. I.)

CONTENTS.

- I. Thermometrical tables, A.
- II. Hygrometrical tables, B.
- III. Barometrical tables, C.
- IV. Hypsometrical tables, D.
- V. Meteorological corrections, E.
- VI. Miscellaneous tables, F.
- 154. List of Foreign Correspondents of the Smithsonian Institution. Corrected to January, 1862. May, 1862. 8vo., pp. 56. (M. C. v.)
- 155. Ancient Mining on the Shores of Lake Superior. By CHARLES WHITTLESEY. April, 1863. 4to., pp. 34, 30 woodcuts, 1 map. S. C. XIII.)
- 156. Catalogue of Minerals, with their Formulas, etc. By T. Egleston. June, 1863. 8vo., pp. 56. (M. C. vii.)
- 157. Results of Meteorological Observations made under the direction of the United States Patent Office and the Smithsonian Institution, from the year 1854 to 1859, inclusive, being a Report of the Commissioner of Patents made at the first session of the 36th Congress. Vol. I. 1861. 36th Congress, 1st session, Senate Ex. Doc. 4to., pp. 1270.
- 158. Smithsonian Miscellaneous Collections. Vol. V. 1864. 8vo., pp. 774.

158. Miscellaneous Collections. Vol. V—Continued. CONTENTS.

BINNEY, W. G. Bibliography of North American conchology. No. 142. Catalogue of publications of the Smithsonian Institution to June, 1862. No. 74.

List of foreign correspondents of the Smithsonian Institution to January, 1862. No. 154.

- 159. Researches upon the Anatomy and Physiology of Respiration in the Chelonia. By S. Weir Mitchell and George R. Morehouse. April, 1863. 4to., pp. 50, 10 woodcuts. (S. C. XIII.)
- 160. Instructions for Research Relative to the Ethnology and Philology of America. By George Gibbs. March, 1863. 8vo., pp. 56. (M. C. vii.)
- 161. A Dictionary of the Chinook Jargon or Trade Language of Oregon. By George Gibbs. March, 1863. 8vo., pp. 60. (M. C. VII.)
- 162. Discussion of the Magnetic and Meteorological Observations made at the Girard College Observatory, Philadelphia, in 1840-'41-'42-'43-'44-'45. Second section, comprising Parts IV, V, and VI. Horizontal force. Investigation of the eleven (or ten) year period and of the disturbances of the horizontal component of the magnetic force, with an investigation of the solar diurnal variation, and of the annual inequality of the horizontal force, and of the lunar effect on the same. By A. D. BACHE. November, 1862. 4to., pp. 78, 11 woodcuts. (S. C. XIII.)
- 163. Circular in Reference to the History of North American Grasshoppers. January, 1860. 8vo., pp. 4. (M. C. 11.)
- 164. Smithsonian Museum Miscellanea. 1862. 8vo., pp. 88. (M. C. VIII.)

CONTENTS.

Abbreviations of names of States and Territories of North America, for labelling insects, shells, &c.

A series of small figures, from 1-1648.

A series of medium figures, from 1-2747.

A series of large figures, from 1-2599.

Blank check-list of specimens.

- No. 5. consists of columns of figures from 1 to 1,000, and of two series, 25 and 50, to the 8vo. column. All these are stereotyped and printed with from one to eight columns on each page, with blank spaces of greater or less extent, as may be required.
- 185. Monograph of the Bats of North America. By H. Allen. June, 1864. 8vo., pp. 110, 73 woodcuts. (M. C. vii.)

- 166. Records and Results of a Magnetic Survey of Pennsylvania and parts of Adjacent States, in 1840 and 1841, with some additional records and Results of 1834, 1835, 1843 and 1862, and a map. By A. D. BACHE. October, 1863. 4to., pp. 88, 1 map. (S. C. XIII.)
- 167. New Species of North American Coleoptera. By John L. Le Conte. Part I. March, 1863. April, 1866. 8vo., pp. 180. (M. C. vi.)
- 168. Circular Relative to Collections of Birds from Middle and South America. December, 1863. 8vo., pp. 2. (M. C. VIII.)
- 169. Smithsonian Miscellaneous Collections. Vol. VI. 1867. 8vo., pp. 888. 15 woodcuts, 7 plates of 106 figures.

LOEW, H. Monograph of diptera. Part I. No. 141.

LOEW, H. Monograph of diptera. Part II. No. 171.

LE CONTE, J. L. List of coleoptera of North America. No. 140.

LE CONTE, J. L. New species of North American coleoptera. No. 167.

- 170. Comparative Vocabulary. May, 1863. 4to., pp. 20.
- 171. Monographs of the Diptera of North America. By H. LOEW.

 Part II. Edited by R. Osten Sacken. January, 1864. 8vo.,

 pp. 372, 5 plates of 44 figures. (M. C. vi.)
- 172. Palæontology of the Upper Missouri; a Report upon Collections made principally by the Expeditions under command of Lieutenant G. K. Warren, United States Topographical Engineers, in 1855 and 1856. Invertebrates. By F. B. Meek and F. V. Hayden. Part I. April, 1865. 4to., pp. 158, 48 woodcuts, 5 plates of 41 figures. S. C. xiv.)
- 173. The Gray Substance of the Medulla Oblongata and Trapezium. By John Dean. February, 1864. 4to., pp. 80, 5 woodcuts, 16 plates of 65 figures. (S. C. xvi.)
- 174. Bibliography of North American Conchology previous to the year 1860. By W. G. Binney. Part II. Foreign authors. June, 1864. 8vo., pp. 302. (M. C. ix.)
- 175. Discussion of the Magnetic and Meteorological Observations made at the Girard College Observatory, Philadelphia, in 1840, 1841, 1842, 1843, 1844 and 1845. Third section, comprising Parts VII, VIII. and IX. Vertical force. Investigations of the eleven (or ten) year period and of the disturbances of the vertical component of the magnetic force, and appendix on the magnetic effect of the aurora borealis; with an investigation of the solar diurnal variation, and of the

- annual inequality of the vertical force; and of the lunar effect on the vertical force, the inclination, and total force. By A. D. BACHE. April, 1864. 4to., pp. 72, 14 woodcuts. (S. C. XIV.)
- 176. Circular in Reference to Collecting North American Shells. January, 1860. 8vo., pp. 4. (M. C. 11.)
- 177. Check-list of the Invertebrate Fossils of North America. Cretaceous and Jurassic. By F. B. Meek. April, 1864. 8vo., pp. 42. (M. C. VII.)
- 178. Circular to Entomologists. 1860. 8vo., pp. 2. (M. C. VIII.)
- 179. Catalogue of Publications of Societies and of Periodical Works, belonging to the Smithsonian Institution, January 1, 1866. 1866. 8vo., pp. 596. (M. C. 1x.)
- 180. On the Construction of a Silvered Glass Telescope, fifteen and a half inches in aperture, and its use in Celestial Photography. By Henry Draper. July, 1864. 4to., pp. 60, 53 woodcuts. (S. C. xiv.)
- 181. Review of American Birds in the Museum of the Smithsonian Institution. By S. F. Baird. Part I.—North and Middle America. June, 1864—June, 1866. 8vo., pp. 484, 80 woodcuts. (M. C. xii.)
- 182. Results of Meteorological Observations made under the Direction of the United States Patent Office and the Smithsonian Institution, from the year 1854 to 1859, inclusive, being a Report of the Commissioner of Patents made at the first session of the 36th Congress. Vol. II. Part I. 36th Congress, 1st session. Senate Ex. Doc. 1864. 4to., pp. 546.
- 183. Check-list of the Invertebrate Fossils of North America. Miocene. By F. B. Meek. November, 1864. 8vo., pp. 34. (M. C. VII.)
- 184. Smithsonian Contributions to Knowledge. Vol. XIV. 1865. 4to., pp. 490, 158 woodcuts, 25 plates.

- BACHE, A. D. Discussion Girard College observations. Parts VII, VIII, IX. No. 175.
- BACHE, A. D. Discussion Girard College observations. Parts X, XI, XII. No. 186.
- DRAPER, H. Construction of silvered glass telescope and its use in celestial photography. No. 180.
- MEEK, F. B.; HAYDEN, F. V. Palsontology of the Upper Missouri. No. 172.
- LEIDY, J. Cretaceous reptiles of the United States. No. 192.

- 185. List of the Described Birds of Mexico, Central America, and the West Indies not in the Collection of the Smithsonian Institution.

 January 1, 1863. 8vo., pp. 8.
- 186. Discussion of the Magnetic and Meteorological Observations made at the Girard College Observatory, Philadelphia, 1840—'41—'42-'43-'4-4' '45. Fourth section, comprising Parts X, XI and XII. Dip and total force; analysis of the disturbances of the dip and total force; discussion of the solar diurnal variation and annual inequality of the dip and total force; and discussion of the absolute dip, with the final values for declination, dip and force between 1841 and 1845. By A. D. Bache. April, 1865. 4to., pp. 42, 8 woodcuts. (S. C. XIV.)
- 187. Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1863. 38th Congress, 1st Session. House of Representatives, Mis. Doc. No. 83. 1864. 8vo., pp. 420, 56 woodcuts.

HENRY, J. Secretary's report of operations.

BOARD OF REGENTS, Proceedings of

BAIRD, S. F. Report of publications, exchanges, museum, and explora-

List of meteorological stations and observers.

Meteorological material contributed to Smithsonian Institution. Correspondence:

GOULD, B. A. Discussion of Piazzi's astronomical observations.

PACKARD, F. A. Project of an outline history of education in the United States.

CHAMBER OF COMMERCE OF BORDEAUX. Exchange of publications.

AGRICULTURAL ASSOCIATION OF MILAN. Exchange of publications.

IRWIN, B. J. D. Tucson meteorite.

SANTIAGO AINSA. Tucson meteorite.

Poole, H. Cone-in-cone.

HUNGARIAN NATIONAL MUSEUM. Acknowledgment for birds.
UNIVERSITY OF CHRISTIANIA, NORWAY. Ethnological specimens presented.

IMP. GEOLOG. INSTITUTE, VIENNA. Tertiary fossils presented.

BRITISH MUSEUM. Electrotypes of engravings of shells granted.

FISHER, J. G. Acknowledgment of perrennibranchiates.

HAIDINGER, W. Honorary medal to Von Martius.

Hudson's Bay Co. Kennicott's explorations.

WHITNEY, W. D. Lectures on the principles of linguistic science.

DEBEAUMONT, ELIE. Memoir of C. F. Bautemps-Beaupré.

ALEXANDER, C. A. Origin and history of the Royal Society of London.

WETHERILL, C. M. Modern theory of chemical types.

DE LA RIVE, A. Phenomena which accompany the propagation of elatricity in highly rarefled elastic fluids.

MARCET, PROF. Report on Society of Physics and Natural History of Geneva, from July, 1862, to June, 1863.

187. Report for 1863—Continued.

PLATEAU, J. Experimental and theoretical researches on the figures of equilibrium of a liquid mass withdrawn from the action of gravity. Part I.

History of discovery relative to magnetism.

GAUTIER, PROF. Researches relative to the nebulæ.

MERINO, St. M. Figure of the earth.

ARAGO, F. Æronautic voyages, etc.

GLAISHER, JAS. Account of balloon ascensions.

BAEGERT, JAC. Aboriginal inhabitants of the California peninsula.

Jones, J. M. Kjækken-mædding in Nova Scotia.

MORLOT, A. Abstract of Dr. Keller's report on lacustrian settlements.

RAU, C. Agricultural implements of the North American stone period.

TROWBRIDGE, D. Ancient fort and burial ground in Tompkins county, New York.

KELLEY, O. H. Ancient town in Minnesota.

FOSTER, J. W. Ancient relics in Missouri.

DANILSEN, A. F. Mound in East Tennessee..

Purple dyeing, ancient and modern.

PEALE, T. R. Method of preserving lepidoptera.

FIGANIERRE, M. Account of remarkable accumulation of bats.

Tables of English and French weights and measures.

Table for conversion of centigrade degrees to Fahrenheit's scale.

188. Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1864. 38th Congress, 2d Session. House of Representatives, Mis. Doc. 1865. 8vo., pp. 450, 50 woodcuts.

CONTENTS.

HENRY, J. Secretary's report of operations.

BOARD OF REGENTS, Proceedings of

WEBB, T. W. Account of Prof. Henry Draper's telescope.

LEIDY, Jos. Review of "Cretaceous reptiles of United States."

BAIRD, S. F. Report on publications, exchanges, museum, and explorations.

List of foreign institutions from which donations to the library have been received in exchange. 1860-64.

List of meteorological stations and observers.

Meteorological material contributed to Smithsonian Institution.

WALLACH, R.; HENRY, J. Report of Committee of Regents in relation to the fire, January, 1865.

FOURIER, Jos. Memoir of Delambre.

DELAUNAY, M. Essay on the velocity of light.

WETHERILL, C. M. Ozone and antozone.

JAMIN, J. Vegetation and the atmosphere.

BECQUEREL, M. Preservation of copper and iron in salt water.

Prescrvation of wood.

Caoutchouc and gutta percha.

VON KAROLYI; CRAIG, B. F. Products of the combustion of gun-cotton and gun-powder.

PETTENKOFER, MAX. Apparatus for testing the results of perspiration and respiration.

LAMONT, J. The solar eclipse of July 18, 1860.

188 Report for 1864—Continued.

DEPEADOS, M. LE BARON. Eclipse of the sun, April 25, 1865.

DUBY, M. Report on the transactions of the Society of Physics and Natural History of Geneva, 1861.

DECANDOLLE, A. Report on the transactions of the Society of Physics and Natural History, 1862.

TROYON, FRED. On the crania helvetica.

PLATEAU, J. Experimental and theoretical researches on the figures of equilibrium of a liquid mass withdrawn from the action of gravity, etc. Parts II, III, IV.

RAU, C. Artificial shell deposits in New Jersey.

GIBBS, G. The intermixture of races.

BARGERT, J. Aboriginal inhabitants of the California peninsula. Part II.

MORLOT, A. The study of high antiquity in Europe.

PRIZE QUESTIONS:

Holland Society of Sciences at Harlem.

Imperial Society of Natural Sciences of Cherbourg.

Royal Prussian Academy of Sciences.

Imperial Academy of Sciences at Vienna.

EXPLORATIONS:

Scientific expedition to Mexico. A report addressed to the Emperor of France by the Minister of Public Instruction.

KIRBY, W. W. A journey to the Yukon, Russian America.

FEILNER, J. Explorations in Upper California in 1860. •

Hoy, P. R. Journal of an exploration of Western Missouri in 1854. Tables of English and French weights and measures.

Table of chemical equivalents of sixty-three elements.

- 189. Catalogue of the Orthoptera of North America described previous to 1867. By Samuel H. Scudder. October, 1868. 8vo., pp. 110. (M. C. VIII.)
- 190. Queries Relative to Tornadoes. By Joseph Henry. 8vo., pp. 4, 1 woodcut. (M. C. x.)
- 191. Smithsonian Miscellaneous Collections. Vol. VII. 1867. 8vo., pp. 878, 813 woodcuts.

CONTENTS.

ALLEN, H. Monograph of bats of North America. No. 165.

BINNEY, W. G. Land and fresh-water shells of North America. Part II. No. 143.

BINNEY, W. G. Land and fresh-water shells of North America. Part III. No. 144.

STIMPSON, W. Hydrobiinæ. No. 201.

PRIME, T. Monograph of American corbiculadæ. No. 145.

CONRAD, T. A. Check-list of fossils—eocene and oligocene. No. 200.

MEEK, F. B. Check-list of fossils-miocene. No. 188.

MEEK, F. B. Check-list of fossils—cretaceous and jurassic. No. 177.

EGLESTON, T. Catalogue of minerals. No. 156.

GIBBS, G. Dictionary of Chinook jargon. No. 161.

GIBBS, G. Instructions for ethnological and philological research. No. 160.

List of works published by the Smithsonian Institution to January. 1874. No. 208.

- 192. Cretaceous Reptiles of the United States. By Joseph Leidy. May, 1865. 4to., pp. 142, 35 woodcuts, 20 plates of 309 figures. (S. C. XIV.)
- 193. Duplicate Shells Collected by the United States Exploring Expedition under Captain C. Wilkes. 8vo., pp. 4.
- 194. Land and Fresh-water Shells of North America. Part I.—Pulmonata geophila. By W. G. Binney and T. Bland. February, 1869. 8vo., pp. 328, 723 woodcuts. (M. C. VIII.)
- 195. Discussion of the Magnetic and Meteorological Observations made at the Girard College Observatory, Philadelphia, in 1840-'41-'42-'43-'44-'45. Parts I to XII inclusive. (Nos. 113x1, 121x111, 132x111, 162x111, 175x117, 186x117). 4to., pp. 262, 49 woodcuts.
- 196. Physical Observations in the Arctic Seas. By Isaac I. Haves. Reduced and discussed by Charles A. Schott. June, 1867. 4to., pp. 286, 15 woodcuts, 3 maps, 3 plates. (S. C. xv.)
- 197. On the Fresh-water Glacial Drift of the Northwestern States. By Charles Whittlesey. May, 1866. 4to., pp. 38, 14 woodcuts, 1 map, 1 plate. (S. C. xv.)
- 198. Physical Observations in the Arctic Seas. By ELISHA KENT KANE. Made during the Second Grinnell Expedition in Search of Sir John Franklin, in 1853, 1854, and 1855, at Van Rensselaer Harbor and other points on the West Coast of Greenland. Reduced and discussed by Charles A. Schott. Part I.—Magnetism. II.—Meteorology. III.—Astronomy. IV.—Tides. (Nos. 97^x, 104^{xi}, 129^{xii}, 130^{xiii}.) 1859-'60. 4to., pp. 340, 17 woodcuts, 1 map, 6 plates.
- 199. An Investigation of the Orbit of Neptune, with General Tables of its Motion. By Simon Newcomb. January, 1866. 4to., pp. 116. (S. C. xv.)
- 200. Check-list of the Invertebrate Fossils of North America, Eccene and Oligocene. By T. A. CONRAD. May, 1866. 8vo., pp. 46. (M. C. VII.)
- 201. Researches upon the Hydrobiinæ and Allied Forms; chiefly made upon materials in the Museum of the Smithsonian Institution. By WILLIAM STIMPSON. August, 1865. 8vo., pp. 64, 32 woodcuts. (M. C. VII.)
- 202. Geological Researches in China, Mongolia, and Japan, during the years 1862 to 1865. By RAPHAEL PUMPELLY. August, 1866. 4to., pp. 173, 18 woodcuts, 3 plates of 10 figures, 6 plates of 15 maps. (S. C. xv.)

- 203. List of Works published by the Smithsonian Institution. January, 1866. 8vo., pp. 12. (M. C. vii.)
- 204. Results of Meteorological Observations made at Brunswick, Maine, between 1807 and 1859. By Parker Cleaveland. Reduced and discussed by Charles A. Schott. May, 1867. 4to., pp. 60, 8 woodcuts. (S. C. xvi.)
- 205. Circular Relating to Collections in Archæology and Ethnology. By JOSEPH HENRY. January, 1867. 8vo., pp. 2. (M. C. VIII.)
- 206. Smithsonian Contributions to Knowledge. Vol. XV. 1867. 40., pp. 620, 47 woodcuts, 13 plates, 4 maps.

NEWCOMB, S. Orbit of Neptune. No. 199.

WHITTLESEY, C. Fresh-water glacial drift of Northwestern States. No. 197.

Pumpelly, R. Geological researches in China, Mongolia, and Japan. No. 202.

HAYES, I. I. Physical observations in the Arctic seas. No. 196.

- 207. Suggestions Relative to Objects of Scientific Investigation in Russian America. By Joseph Henry. May, 1867. 8vo., pp. 10. (M. C. VIII.)
- 208. The Gliddon Mummy Case in the Museum of the Smithsonian Institution. By Charles Pickering. June, 1867. 4to., pp. 6, 1 plate. S. C. xvi.)
- 209. Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1865. 39th Congress, 1st Session, House of Representatives Mis. Doc. No. 102. 1866. 8vo., pp. 496, 139 woodcuts.

CONTENTS.

HENRY, J. Secretary's report of operations.

BOARD OF REGENTS. Proceedings of

BAIRD, S. F. Report of publications, exchanges, museum, and explorations.

List of addresses of foreign institutions since 1862.

List of meteorological stations and observers.

List of meteorological material contributed.

Correspondence:

HENRY, J. Letter to Secretary of Treasury on payment of interest in coin.

LEIDY, J.; TRYON, G. W. Report on shells presented to Academy of Natural Sciences.

MEXICAN Soc. of GEOG. AND STATISTICS. Exchange system.

209. Report for 1865—Continued.

EVANS, J. Climate of Colorado.

GIVEN, W. F. Remarkable electric phenomenon.

HALE, C. Explorations of the Nile.

MINING DEPT., MELBOURNE. Exchange system.

CARLETON, J. H. Meteorites in Mexico.

CONRAD, T. A. Chalk found in United States.

REGENTS OF UNIVERSITY OF STATE OF NEW YORK. Acknowledgment for specimens.

McMasters, S. Y. Language of Navajos said to resemble the Welsh.

GIBBS, G. Indian languages.

LISBOA, M. M. Books on Brazil presented.

BRAZILIAN NAT. HIST. MUSEUM. Specimens from Brazil presented.

BERN MUSEUM. Request for Bison.

IMPERIAL LIBRARY OF VIENNA. Books presented.

BRUCE, F. W. A., British Minister. Introducing and recommending Dr. Berendt.

DE IRISARRI, A. T., Guatemalan Minister. Introducing and recommending Dr. Berendt.

Molina, L., Costa Rican Minister. Introducing and recommending Dr. Berendt.

Rosing, J. Exchange system.

GOVERNMENT OF BREMEN. Exchange system.

Hodgins, J. G. Metebrological system of Canada.

BRITISH MUSEUM. Acknowledgment of specimens.

Toronto University. Objects of the Museum.

PETITOT, E. Account of the Indians of British America.

Act of Congress to transfer Smithsonian Library to Library of Congress.

BARNARD, J. G. Eulogy on Gen. Jos. G. Totten.

FLOURENS, M. Memoir of Ducrotay de Blainville.

CHOSSAT, Dr. Report on the transactions of the Society of Physics and Natural History of Geneva, from July, 1863, to June, 1864.

PLANTAMOUR, M. E. Report on the transactions of the Society of Physics and Natural History of Geneva, from July, 1864, to June, 1865.

LOOMIS, E. Aurora Borealis, or Polar Light; its phenomena and laws. The Senses. Sense of feeling; sense of smell.

MATTEUCCI, C. Lectures on electro-physiology.

DESOR, E. Palafittes or Lacustrian constructions of the Lake of Neuchatel.

PLATEAU, J. Experimental and theoretical researches on the figures of equilibrium of a liquid mass withdrawn from the action of gravity, etc. Part V.

LILLJEBORG, W. Outline of a systematic review of the classification of birds.

PRIZE QUESTIONS:

Royal Danish Society of Sciences.

Imperial Academy of Sciences of Vienna.

Pontifical Academy of the Nuovi Lincei.

Royal Scientific and Literary Institute of Lombardy.

Imperial Society of Science, Agriculture, and Arts of Lille.

Dunkirk Society for the encouragement of sciences, letters, and arts.

NEWTON, H. A. Metric system of weights and measures, with tables.

- 210. Arrangement of Families of Birds, adopted provisionally by the Smithsonian Institution. By S. F. BAIRD. June, 1866. 8vo., pp. 8. (M. C. VIII.)
- 211. Smithsonian Contributions to Knowledge. Vol. XVI. 1870. 4to., pp. 498, 76 woodcuts, 18 plates.

DEAN, J. Gray substance of the medulla oblongata and trapezium. No. 178.

CLEAVELAND, P. Meteorological observations, Brunswick, Maine, 52 years. No. 204.

Hildreth, S. P. Meteorological observations, Marietta, O. No. 120.

Pickering, C. Gliddon mummy case. No. 208.

COFFIN, J. H. Orbit and phenomena of a meteoric fire ball. No. 221.

Gould, B. A. Transatlantic longitude. No. 223.

SWAN, J. G. Indians of Cape Flattery. No. 220.

212. Smithsonian Miscellaneous Collections. Vol. VIII. 1869. 8vo., pp. 921, 730 woodcuts, 4 plates.

CONTENTS.

OSTEN SACKEN, R. Monograph of the diptera of North America. Part IV. No. 219.

Scudder, S. H. Catalogue of the orthopters of North America. No. 189.

BINNEY, W. G.; BLAND, T. Land and fresh-water shells of North America. Part I. No. 194.

BAIRD, S. F. Arrangement of families of birds. .No. 210.

HENRY, J. Circular to officers of the Hudson's Bay Company. No. 137.

HENRY, J.; GIBBS, G.; BAIRD, S. F. Suggestions relative to scientific investigations in Russian America. No. 207.

HENRY, J. Circular relative to archeology and ethnology. No. 205.

HENRY, J. Circular to entomologists. No. 178.

HENRY, J. Circular relative to collections of birds from Middle and South America. No. 168.

BAIRD, S. F. Smithsonian Museum miscellanea. No. 164.

213. Smithsonian Miscellaneous Collections. Vol. IX. 1869. 8vo., pp. 914.

BINNEY, W. G. Bibliography of North American conchology. Foreign authors. Part II. No. 174.

Catalogue of publications of societies and of periodicals in Smithsonian Library, 1866. No. 179.

214. Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1866. 39th Congress, 2d Session, House of Representatives, Mis. Doc. No. 83. 1867. 8vo., pp. 470, 70 woodcuts.

CONTENTS.

HENRY, J. Secretary's report of operations. BOARD OF REGENTS. Proceedings of

214. Report for 1866—Continued.

BAIRD, S. F. Report on publications, exchanges, museum, and explorations.

List of meteorological stations and observers.

Act of Congress to receive into the United States Treasury the residuary legacy of Smithson, etc.

Memorial of Board of Regents to Congress relative to the fund.

HENRY, J. Memoir of W. W. Seaton.

Report on system of accounts.

FLOURENS, M. Memoir of Magendie.

THE SENSES. Sense of taste; sense of hearing; sense of sight.

HUGGINS, W. Results of spectrum analysis applied to the heavenly bodies. External appearance of the sun's disk.

Moigno, Abbe. Accidental or subjective colors: Persistence of images, contrast, irradiation, daltonism, etc.

PLATEAU, J. Figures of equilibrium of a liquid mass withdrawn from the action of gravity. Part VI.

Gosse, Dr. Report on the transactions of the Society of Physics and Natural History of Geneva, from July, 1865, to June, 1866.

GIBBS, G. Notes on the Tinneh or Chepewyan Indians of British and Russian America. 1. The Eastern Tinneh, by Bernard R. Ross, Esq. 2. The Loucheux Indians, by William L. Hardisty. 3. The Kutchin Tribes, by Strachan Jones, Esq.

Von Hellwald, F. The American migration, with notes by Prof. Henry.

RAU, C. Indian pottery.

BRINTON, D. G. Artificial shell deposits of the United States.

DILLE, I. Sketch of ancient earth-works of Ohio.

Pile-work antiquities of Olmutz.

Estes, L. C. Antiquities on the banks of the Mississippi river and Lake Pepin.

GIBBS, G. Physical Atlas of North America.

DAVIS, E. H. On ethnological research.

SCHERZER; SCHWARZ. Table of anthropological measurements.

International Archæological Congress, organized by the Archæological Academy of Belgium, in concert with the French Society of Archæology. Antwerp, 1866.

HIGGINS, H. H. On vitality, with notes by Prof. Henry.

LEWIS, J. Instructions for collecting land and fresh-water shells.

WOOD, H. C. Instructions for collecting myriapods, phalangidse, etc.

WETHERILL, C. M. Plan of a research upon the atmosphere.

LEWIS; QUALE. Account of the cryolite of Greenland.

EXTRACTS FROM THE METEOROLOGICAL CORRESPONDENCE OF THE INSTI-

HENRY, J. Remarks by

DENNIS, W. C. Evaporation in Florida.

DENNIS, W. C. Fresh-water in the ocean.

DEWEY, C. The winds.

BALTZELL, J. Winds in Florida.

WARD, L. F. Barometer tube breaking suddenly.

Andrews, S. L. Meteors.

214. Report for 1866—Continued.

BALFOUR, J. The wind and fog.

Huntington, G. C. Climate of Kelley's Island, Ohio.

MALLINIKROOT, C. Changes of wind.

CAPEN, F. L. Meteorological discovery.

FENDLER, A. Meteorology of Colonia Towar.

HILLIER, S. L. Effect of lightning.

BANNISTER, H. M. Formation of clouds over Gulf stream.

BANNSSTER, H. M. Climate of Alaska.

VAILLANT, M. Horary variations of the barometer, with notes by Prof. Henry.

ENGLEHARDT, M. Formation of ice at the bottom of water.

SARTORIUS, C. Earthquake in Eastern Mexico, January, 1866.

DREUTZER, O. E. Statistics relative to Norwegian mountains, lakes, and the snow-line.

215. Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1867. 40th Congress, 2d Session, Senate Miss. Doc. No. 86. 1868. 8vo., pp. 506, 10 woodcuts.

CONTENTS.

HENRY, J. Secretary's report of operations.

BOARD of REGENTS. Proceedings of

BAIRD, S. F. Report of exchanges and museum.

BAIRD, S. F. List of expeditions and other sources from which the specimens in the Government Museum have been mainly derived.

List of meteorological stations and observers.

Meteorological material contributed in addition to the regular observations.

Cluss, A. Report of the architect.

Correspondence:

AMER. ACAD. OF ARTS AND SCIENCES. Exchange system.

KNIGHT, G. H. New system of weights and measures with 8 as the metrical number.

Bolles, E. C. Account of Portland Society of Natural History.

BATH AND WEST OF ENG. Soc. FOR ENCOUR. OF AGRIC., ARTS. ETC. Exchange system.

HAMBURG ZOOL. GARDENS. Exchange of specimens.

CHICAGO ACAD. OF SCIENCES. Acknowledgment for specimens.

TAYLOR, W. B. Report on improved system of numeration.

ST. PETERSBURG ACAD. OF SCIENCES. Exchange system.

ZISGENBALS, H. Schlagintweit ethnographic collection.

LLOYD, W. A. Sparrows sent to United States.

LLOYD, W. A. Exchange of specimens.

MUSEUM OF NAT. UNIV. OF GREECE. Exchange of specimens.

LABOULAYE, M. Acknowledgment for books on education.

ACAD. OF SCIENCES, STOCKHOLM. Acknowledgment of birds.

Gould, J. Acknowledgment of birds.

MAYBERRY, S. P. Gradual approach of sea upon land.

University of Costa Rica. Exchange of specimens.

Guild, R. A. Biographical notice of Chas. C. Jewett.

GRAY, A, Biographical notice of W. H. Harvey.

Agassiz, L. Report on use of new hall in Smithsonian Building.

215. Report for 1867—Continued.

Memorial of Regents to Congress asking appropriations for Museum.

DEBEAUMONT, E. Memoir of Legendre.

PELTIER, F. A. Memoir of J. C. A. Peltier.

Scientific researches of Peltier.

MAILLY, E. History of Royal Institution of Great Britain.

DE LA RIVE, Prof. Michael Faraday, his life and works.

FLOUBENS, M. The Jussieus and the natural method.

MARKY, M. Natural history of organized bodies.

MATTEUCCI, C. On the electrical currents of the earth.

Considerations on electricity.

Electricity—Account of lightning discharges, with notes by Prof. Henry.

G. W. Dodge, O. M. Poe, J. C. Cresson, H. Haas, H. J. Kron, B. F.

Mudge, New Haven Journal, S. D. Martin, G. Wright, C. G.

Boerner, W. S. Gilman, C. W. Dean, J. A. Osborne.

DARWIN, C. Queries about expression for anthropological inquiry.

PETTIGREW, J. B. Modes of flight in relation to aeronautics.

Man as the contemporary of the mammoth and reindeer in Middle Europe.

JAMIN, M. Photo-chemistry.

ABBE, C. Description of the observatories at Dorpat and Poulkova.

TYLOR, E. B. On traces of the early mental condition of man.

ETHNOLOGY:

GUNN, D. Indian remains, Red River Settlement, Hudson's Bay Territory.

READ, M. C. Ancient mound near Chattanooga, Tenn.

PIDGEON, W. Ancient burial mound in Indiana.

BERTHOUD, E. L. Ancient remains in Colorado.

AGNEW, S. A. Mounds in Mississippi.

WHITNEY, J. D. Cave in Calaveras county, California.

Ethnological Department of the French Exposition, 1867.

HAYDEN, F. V. Notes on Indian history.

MEIGS, J. A. Description of a human skull from Rock Bluff, Ill.

KABIS, M. Introduction to the study of the Coptic language.

SMART, C. Notes on the Tonto Apaches.

BERENDT, C. H. Explorations in Central America.

GUNN, D. Notes of an egging expedition to Shoal Lake, Lake Winnipeg.

ROTHBOCK, J. T. Sketch of the flora of Alaska.

METEOROLOGY:

Hurricane in the island of St. Thomas.

LATIMER, G. A. Earthquakes in St. Thomas.

Maritime disasters of the Antilles.

Dickinson, A. B. Eruption of a volcano in Nicaragua.

Young, W. J. Cloud-bursts.

WOODWORTH, A. Meteorite in Mexico.

Simson, R. Meteorite in Mexico.

ERNST, G. A. Meteorology of Caracas, South America.

TUCKETT, F. F. On barometer tables.

Rico.

Creat hurricane at Tortola, St. Thomas, and Porto

PIKE, N. Cyclone in the Indian Ocean.

215. Report for 1867—Continued.

PRIZE QUESTIONS:

Royal Danish Society of Sciences.

Pontifical Academy of the Nuovi Lincei.

DE LA RUE, W. Abbreviations used in England, 1867.

- 216. List of Photographic Portraits of North American Indians in the Gallery of the Smithsonian Institution. 1867. 8vo., pp. 42. (M. C. xiv.)
- 217. Letter by M. Hoek in reference to the Meteoric Shower of November 13th, 1867. 8vo., pp. 4.
- 218. Systems of Consanguinity and Affinity of the Human Family. By Lewis H. Morgan. 1869. 4to., pp. 616, 6 woodcuts, 14 plates. (S. C. xvii.)
- 219. Monographs of the Diptera of North America. By R. Osten Sacken. Part IV. 1869. 8vo., pp. 358, 7 woodcuts, 4 plates of 92 figures. (M. C. VIII.)
- 220. The Indians of Cape Flattery, at the entrance to the Strait of Fuca. Washington Territory. By James G. Swan. March, 1870. 4to., pp. 118, 44 woodcuts. (S. C. xvi.)
- 221. The Orbit and Phenomena of a Meteoric Fire Ball, seen July 20, 1860. By James H. Coffin. May, 1869. pp. 56, 3 woodcuts, 1 map. (S. C. xvi.)
- 222. Tables and Results of the Precipitation, in Rain and Snow, in the United States, and at some stations in adjacent parts of North America, and in Central and South America. By Charles A. Schott. March, 1872. 4to., pp. 178, 8 woodcuts, 5 plates, and 3 double charts. (S. C. XVIII.)
- 223. The Transatlantic Longitude, as determined by the Coast Survey Expedition of 1866. By Benjamin Apthorp Gould. October, 1869. 4to., pp. 110, 2 woodcuts. (S. C. xvi.)
- 224. Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1868. 40th Congress, 3d Session. House of Representatives, Ex. Doc. No. 83. 1869. 8vo., pp. 474, 40 woodcuts.

CONTENTS.

HENRY, J. Secretary's report of operations.

BOARD OF REGENTS, Proceedings of

List of meteorological stations and observers.

List of meteorological material contributed to Smithsonian Institution.

DELAFIELD, R. Report of Committee of Regents on Washington canal. Flourens, M. Memoir of Cuvier.

FLOURENS, M. History of the works of Cuvier.

DE BEAUMONT, E. Memoir of Oersted.

Notice of Christian Frederic Schænbein, the discoverer of ozone.

234. Report for 1868—Continued.

HENRY, J. Appendix to notice of Schoenbein.

HAGEN, G. Memoir of Encke.

RAWSON, R. Memoir of Eaton Hodgkinson.

CAZIN, A. Recent progress in relation to the theory of heat.

MULLER, J. Principles of the mechanical theory of heat.

MAGRINI, L. Continuous vibratory movement of all matter, ponderable and imponderable.

TYNDALL, J. Radiation.

DAUBRÉE, M. Synthetic experiments relative to meteorites.

Brush, G. J. Catalogue of meteorites in the mineralogical collection of Yale College.

DE SAUSSURE, H. Observations on the electric resonance of mountains.

STEWART, B. Experiments on aneroid barometers made at the Kew Observatory.

ELLERY, R. L. J. Address of the president of the Royal Society of Victoria.

WARTMANN, E. Report on the transactions of the Society of Physics and Natural History of Geneva, from July, 1867, to June, 1868.

Broca, P. History of the transactions of the Anthropological Society of Paris, from 1865 to 1867.

RAU, C. Drilling in stone without metal.

RAU, C. Agricultural flint implements in southern Illinois.

Notice of the Blackmore Museum, Salisbury, England.

PRIZE QUESTIONS:

Holland Society of Sciences of Harlem.

Imperial Academy of Sciences, Belles Lettres, and Arts, of Bordeaux.

Pollock, J. Assay of gold and silver coins at the Mint of the United States.

Table of foreign gold and silver coins.

List of publications of the Smithsonian Institution up to July, 1869, with systematic and alphabetical index.

- 225. List of Foreign Correspondents of the Smithsonian Institution; corrected to January, 1870. April, 1870. 8vo., pp. 56.
- 226. A List of the Smithsonian Publications, from 1846 to 1869. November, 1869. 8vo., pp. 34.
- 227. Arrangement of the Families of Mollusks. By Theodore Gill. February, 1871. 8vo., pp. 65. (M. C. x.)
- 228. Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1869. 41st Congress, 3d Session. House of Representatives, Ex. Doc. No. 153. 1871. 8vo., pp. 430, 38 woodcuts, 1 map.

CONTENTS.

HENRY, J. Secretary's report of operations.

BOARD OF REGENTS, Proceedings of

List of meteorological stations and observers.

List of meteorological material contributed to Smithsonian Institution.

228. Report for 1869—Continued.

BERTHRAND, M. Kepler: his life and works.

ARAGO, M. Eulogy on Thomas Young.

DE BEAUMONT, E. Memoir of Auguste Bravais.

RAU, C. Memoir of C. F. P. Von Martius.

MATTEUCCI, C. Life and scientific labors of Stefano Marianini.

HUNT, T. S. Chemistry of the earth.

MATTEUCCI, C. Electrical currents of the earth.

MAREY, M. Phenomena of flight in the animal kingdom.

BABINET, M. The northern seas.

LOMBARD, H. C. Report on the transactions of the Society of Physics and of Natural History of Geneva, July, 1868, to June, 1869.

Simpson, J. H. Coronado's march in search of the "Seven Cities of Cibola."

LUBBOCK, J. Social and religious condition of the lower races of man.

HUXLEY, T. H. Principles and methods of palsontology.

SCHOTT, A. Remarks on the "Cara Gigantesca" of Yzamal, in Yucatan.

BECQUEREL, M. Forests and their climatic influence.

Brenndecke, F. Meteorites.

ABICH, S. Remarkable forms of hail-stones in Georgia.

SARTORIUS, C. Eruption of the volcano of Colima.

229. Smithsonian Contributions to Knowledge. Vol. XVII. 1871. 4to., pp. 616, 6 woodcuts, 14 plates.

CONTENTS.

MORGAN. Systems of consanguinity and affinity of the human family. No. 218.

- 230. Arrangement of the Families of Mammals. By Theodore Gill. November, 1872. 8vo., pp. 104. (M. C. xi.)
- 231. Memoranda of Inquiry relative to the Food Fishes of the United States. By S. F. BAIRD. 1871. 8vo., pp. 8. (M. C. x.)
- 232. The Secular Variations of the Elements of the Orbits of the Eight Principal Planets, Mercury, Venus, the Earth, Mars, Jupiter, Saturn. Uranus, and Neptune, with tables of the same; together with the obliquity of the ecliptic and the precession of the equinoxes in both longitude and right ascension. By John N. Stockwell. 1872. 4to., pp. 220. (S. C. XVIII.)
- 233. Converging Series expressing the Ratio between the Diameter and the Circumference of a Circle. By Wm. Ferrel. April, 1871. 4to., pp. 6. (S. C. xvIII.)
- 234. Questions Relative to the Food Fishes of the United States. By S. F. BAIRD. 1871. 8vo., pp. 14. (M. C. x.)
- 235. Instructions for Observations of Thunder Storms. By JOSEPH HENRY. 1871. 8vo., p. 1. (M. C. x.)

- 236. Circular Relative to Heights. By JOSEPH HENRY. 1871. 8vo., pp. 2. (M. C. x.)
- 237. Directions for constructing Lightning Rods. By JOSEPH HENRY. 1871. 8vo., pp. 4. (M. C. x.)
- 238. List of the Institutions, Libraries, Colleges, and other Establishments in the United States in correspondence with the Smithsonian Institution. By Wm. J. Rhees. July, 1872. 8vo., pp. 256. (M. C. x.)
- 239. Observations on Terrestrial Magnetism, and on the Deviations of the Compasses of the United States Iron Clad Monadnock during her cruise from Philadelphia to San Francisco, in 1865 and 1866. By WM. HARKNESS. 1872. 4to., pp. 226, 2 woodcuts. (S. C. XVIII.)
- 240. Problems of Rotary Motion presented by the Gyroscope, The Precession of the Equinoxes, and The Pendulum. By J. G. BARNARD. 1872. 4to., pp. 56, 6 woodcuts. (S. C. xix.)
- 241. A Contribution to the History of the Fresh Water Algo of North America. By Horatio C. Wood, Jr. 1872. 4to., pp. 272, 21 plates of 340 figures. (S. C. xix.)
- 242. Lucernariæ and their Allies. A Memoir on the Anatomy and Physiology of Haliclystus Auricula and other Lucernarians, with a Discussion of their Relations to other Acalephæ, to Beroids and Polypi. By H. J. Clark. 1878. 4to., pp. 138, 11 plates of 149 figures. (S. C. XXII.)
- 243. List of Foreign Correspondents of the Smithsonian Institution; corrected to January, 1872. 1872. 8vo., pp. 66. (M. C. x.)
- 244. Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1870. 42d Congress, 1st Session. House of Representatives, Ex. Doc. No. 20. 1871. 8vo., pp. 494, 28 woodcuts.

HENRY, J. Secretary's report of operations.

BOARD OF REGENTS, Proceedings of

List of meteorological stations and observers.

List of meteorological material contributed to Smithsonian Institution.

HENRY, J. Eulogy on Alexander Dallas Bache.

BACHE, A. D. Lecture on Switzerland.

HENRY, J. On a physical observatory.

ARAGO, M. The History of my Youth, an autobiography.

ARAGO, M. Eulogy on Herschel.

Life and labors of Henry Gustavus Magnus.

Anderson, M. B. Life of Prof. Chester Dewcy.

TAYLOR, W. B. Thoughts on the nature and origin of force.

244. Report for 1870—Continued.

VON LIEBIG, J. Induction and deduction.

HAUGHTON, S. Relation of food to work, and its bearing on medical practice.

REYNOLDS, J. E. Hydrogen as gas and as a metal.

WISEMAN, N. Identification of the artisan and artist.

BABINET, M. The diamond and other precious stones.

ETHNOLOGY:

GIBBS, G. On the language of the aboriginal Indians of America.

WILLIAMS, H. C. On antiquities in some of the Southern States.

GARDNER, W. H. Ethnology of the Indians of the valley of the Red River of the North.

FINCK, H. Account of antiquities in the State of Vera Cruz, Mexic

DUNNING, E. O. Account of antiquities in Tennessee.

STEPHENSON, M. F. Account of ancient mounds in Georgia.

DAYTON, E. A. Explorations in Tennessee.

HARWOOD, A. A. Account of the sarcophagus in the National Museum, now in charge of the Smithsonian Institution.

GRANT, E. M. Account of the discovery of a stone image in Tensessee, now in possession of the Smithsonian Institution.

BLYDEN, E. D. On mixed races in Liberia.

Fowler, J. On shell-heaps.

PEALE, T. R. On the uses of the brain and marrow of animas among the Indians of North America.

Lyon, S. S. Report of an exploration of ancient mounds in Univ. county, Kentucky.

BARRANDT, A. Sketch of ancient earthworks on the Upper Missour.

STELLE, J. P. Account of aboriginal ruins at Savannah, Tennessee.

STELLE, J. P. Account of aboriginal ruins in Hardin Co., Tennes-er.

TERRESTRIAL PHYSICS:

CAMPBELL, J. V. The earthquake in Peru, August 13, 1868.

PALMIERI, Prof. The electro-magnetic seismograph.

Johnson, W. W. On the distribution of forest trees in Montans. Idaho, and Washington.

SARGENT, W. D. Influence of the aurora on the telegraph.

METEOROLOGY:

Poër, A. New classification of clouds.

TACCHINI, P. On the evaporation observed at Palermo, in 1865 and 1866.

ZANTEDESCHI, F. On the electricity of induction in the zerial strate of the atmosphere, which, in the shape of a ring, surround a cloud that is resolving into rain, snow, or hail.

PALMIERI, Prof. On the presence of electricity during the fall of rain.

ELLIOTT, R. S. Climate of Kansas.

PORTER, Commodore. Account of a hail storm on the Bosphorus.

BACHE, G. M. Account of a hail storm in Texas.

245. Check List of Publications of the Smithsonian Institution to July. 1872. 1872. 8vo., pp. 21. (M. C. x.)

246. Smithsonian Contributions to Knowledge. Vol. XVIII. 1872. 4to., pp. 646, 10 woodcuts, 5 plates, 3 charts.

CONTENTS.

SCHOTT, C. A. Tables of rain and snow. No. 222.

STOCKWELL, J. N. Secular variations of the orbits of planets. No. 282.

HARKNESS, W. Observations on terrestrial magnetism. No. 239.

FERREL, W. Converging series, expressing the ratio between the diameter and the circumference of a circle. No. 238.

- 247. Arrangement of the Families of Fishes, or Classes Pisces, Marsipobranchii, and Leptocardii. By Theodore Gill. November, 1872. 8vo., pp. 96. (M. C. XI.)
- 248. On the Geology of Lower Louisiana and the Salt Deposit on Petite Anse Island. By Eugene W. Hilgard. June, 1872. 4to., pp. 38, 4 woodcuts, 2 plates. (S. C. XXIII.)
- 249. Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1871. 42d Congress, 1st Session. Senate Mis. Doc. No. 149. 1873. 8vo., pp. 473, 3 woodcuts.

CONTENTS.

HENRY, J. Secretary's report of operations.

BOARD OF REGENTS, Proceedings of

List of meteorological stations and observers.

Meteorological material contributed to Smithsonian Institution.

Meteorological articles received by the Institution, and deposited in the Library of Congress. Auroras, earthquakes, etc.

Dodge, N. S. Memoir of Sir John Frederick William Herschel.

ARAGO, M. Eulogy on Joseph Fourier.

Odling, W. Professor Thomas Graham's scientific work.

HELMHOLTZ, H. On the relation of the physical sciences to science in general.

KORNHUBER, G. A. Alternate generation and parthenogenesis in the animal kingdom.

REICHARDT, W. Present state of our knowledge of cryptogamous plants.

STOCKWELL, J. N. Recent researches on the secular variations of the planetary orbits.

DE FOREST, E. L. Methods of interpolation applicable to the graduation of irregular series, such as tables of mortality. Part I.

DE SAUSSURE, H. Report on the transactions of the Society of Physics and Natural History of Geneva, from July, 1870, to June, 1871.

Expedition toward the North Pole:

HENRY, J.; HILGARD, J. E.; NEWCOMB, S.; BAIRD, S. F.; MEEK, F. B.; AGASSIZ, L. Scientific instructions to Captain Hall.

ETHNOLOGY:

COMFORT, A. J. Indian mounds near Fort Wadsworth, Dakota.

BERTHOUD, E. L. Antiquities on the Cache la Poudre River, Weld county, Colorado Territory.

Lyon, W. B. Antiquities in New Mexico.

249. Report for 1871—Continued.

SPAINHOUR, J. M. Antiquities in Lenoir county, North Carolina.

McConnell, E. M. Account of the old Indian village, Kushkushkee, near Newcastle, Pennsylvania.

GROSSMANN, F. E. Pima Indians of Arizona.

CROOK, G. Indian mode of making arrow-heads and obtaining fire.

PETER, R. Ancient mound near Lexington, Kentucky.

Brown, D. Shell-heap in Georgia.

SCHOTT, A. Remarks on ancient relic of Maya sculpture.

MUCH, M. Ancient history of North America.

ROTHER, F. L. O. On the language of the Dakota or Sioux Indians.

METEOROLOGY, with notes by Professor Henry:

LATIMER, G. Meteorology of Porto Rico.

Collins, Colonel. Meteorology of the Green river country.

LAMARK. Distinction between tornadoes and tempests.

MEEK, J. B. Account of a tornado which occurred in Spruce Creek Valley, Centre county, Pennsylvania.

HENRY, J. Effect of the moon on the weather.

KNIGHT, R. T.; HENRY, J. Connection of gales of wind and appearance of the aurora.

HARRISON, W. Account of a storm in Butler county, Kansas. June 23, 1871.

250. Smithsonian Miscellaneous Collections. Vol. X. 1873. 8vo., pp. 913, 5 woodcuts.

CONTENTS.

CABPENTER, P. P. Mollusks of western North America. No. 252.

GILL, T. Arrangement of the families of mollusks. No. 227.

HENRY, J. Instructions for observations of thunder storms. No. 235.

HENRY, J. Circular relative to heights. No. 236.

HENRY, J. Directions for constructing lightning rods. No. 287.

HENRY, J. Queries relative to tornadoes. No. 190.

BAIRD, S. F. Questions relative to the food fishes of the United States. No. 284.

BAIRD, S. F. Memoranda of inquiry relative to food fishes. No. 234.

RHEES, W. J. List of institutions, etc., in the United States in correspondence with the Smithsonian Institution. No. 238.

List of foreign correspondents of the Smithsonian Institution, for 1872.

Check list of publications of the Smithsonian Institution, 1872. No. 245.

251. Memoir of C. F. P. Von Martius. By C. Rau. 1871. 8vo., pp. 12.

252. The Mollusks of Western North America. By P. P. CARPENTER. December, 1872. 8vo., pp. 446, 4 woodcuts. (M. C. x.)

CONTENTS.

Supplementary report on the present state of our knowledge with regard to the mollusca of the west coast of North America.

Review of Professor C. B. Adams' "Catalogue of the Shells of Panama" from the type specimens.

252. Mollusks of Western America—Continued.

Diagnoses of new forms of mollusks collected at Cape St. Lucas, Lower California.

Contributions towards a monograph of the pandoridæ.

Diagnoses of new forms of mollusca from the Vancouver district.

Diagnoses of new species and a new genus of mollusks, from the Reigen Mazatlan collection; with an account of additional specimens presented to the British Museum.

Descriptions of new species and varieties of chitonide and acmeide, from the Panama collection of the late Prof. C. B. Adams.

Diagnoses of new species of mollusks from the west tropical region of North America.

Diagnoses of new forms of mollusca from the west coast of North America, first collected by Col. E. Jewett.

Diagnoses of new forms of mollusca, collected by Col. E. Jewett, on the west tropical shores of North America.

Diagnoses des mollusques nouveaux provenant de Californie et faisant partie du Musée de l'Institution Smithsonienne.

On the pleistocene fossils collected by Col. E. Jewett, at Santa Barbara, California; with descriptions of new species.

- 253. Land and Fresh Water Shells of North America. Part IV. Strepomatidæ (American Melanians). By Geo. W. Tryon, Jr. December, 1873. 8vo., pp. 490, 871 woodcuts. (M. C. xvi.)
- 254. Synopsis of American Wasps. Solitary Wasps. By Henry De Saussure. December, 1875. 8vo., pp. 430, 4 plates of 31 figures. (M.C. xiv.)
- 255. The Constants of Nature. Part I. Specific Gravities: Boiling and Melting Points; and Chemical Formulæ. By Frank Wigglesworth Clarke. December, 1873. 8vo., pp. 272. (M. C. XII.)
- 256. Monographs of the Diptera of North America. Part III. By H. Loew. December, 1873. 8vo., pp. 381, 4 plates of 116 figures. (M. C. xi.)
- 257. Systematic Index to List of Foreign Correspondents of the Smithsonian Institution. 1872. 8vo., pp. 30. (M. C. x.)
- 258. Bibliographical Index to North American Botany; or, Citations of Authorities for all the recorded Indigenous and Naturalized Species of the Flora of North America; with a Chronological Arrangement of the Synonymy. Part I. Polypetalæ. By Sereno Watson. March, 1878. 8vo., pp. 484. (M. C. xv.)
- 259. Explorations of the Aboriginal Remains of Tennessee. By Joseph Jones. October, 1876. 4to., pp. 181, 110 woodcuts. (S. C. XXII.)
- 260. Regulations of the Smithsonian Institution. January, 1872. 8vo., pp. 41, 1 woodcut.

- 261. Directions for Collecting and Preserving Insects. By A. S. PACKARD, Jr. 1873. 8vo., pp. 60, 55 woodcuts. (M. C. XL.)
- 262. An Investigation of the Orbit of Uranus, with General Tables of its Motion. By Simon Newcomb. August, 1873. 4to., pp. 296. (S. C. XIX.)
- 263. Circular of Instructions to Observatories relative to Telegraphic Announcements of Astronomical Discoveries. By Joseph Henry. May, 1873. 8vo., pp. 4. (M. C. XII.)
- 264. New Species of North American Coleoptera. Part II. By John L. Le Conte. 1873. 8vo., pp. 74. (pp. 169-240.) (M. C. XI.)
- 265. Classification of the Coleoptera of North America. Part II. By John L. Le Conte. June, 1873. 8vo., pp. 72. (pp. 279-348. (M. C. XI.)
- 266. On the Structure of Cancerous Tumors and the mode in which Adjacent Parts are Invaded. Toner Lecture No. I. Delivered March 28, 1873. By J. J. WOODWARD. November, 1873. 8vo., pp. 44. 5 woodcuts. (M. C. xv.)
- 267. The Haidah Indians of Queen Charlotte's Islands, British Columbia, with a brief description of their Carvings, Tattoo Designs, etc. By James G. Swan. July, 1874. 4to., pp. 22, 7 plates of 24 figures (S. C. xxi.)
- 268. The Winds of the Globe; or, the Laws of Atmospheric Circulation over the Surface of the Earth. By James Henry Coffin. The tables completed and maps drawn by S. J. Coffin, with a discussion and analysis of the tables and charts by Alexander Weikof. December, 1875. 4to., pp. 781, 4 woodcuts, 26 plates of 221 figures. (S. C. xx.)
- 269. The Sculptures of Santa Lucia Cosumalwhuapa, in Guatemala, with an account of travels in Central America and on the Western Coast of South America. By S. Habel. 1878. 4to., pp. 94, 8 plates of 25 figures. (S. C. xxII.)
- 270, Catalogue of the described Diptera of North America. By C. R. OSTEN SACKEN. 1878. 8vo., pp. 324. (M. C. XVI.)
- 271. Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1872. 42d Congress, 3d Session. House of Representatives, Mis. Doc. No. 107. 1873. 8vo., pp. 456, 109 woodcuts.

HENRY, J. Secretary's report of operations. BOARD OF REGENTS, Proceedings of

1. Report for 1872—Continued.

List of meteorological stations and observers.

HENRY, J. Notes relative to George Catlin.

Report of Committee of Regents on Corcoran Art Gallery.

Agassiz, L. Narrative of the Hassler expedition.

BACHE, A. D. Bequest to National Academy of Sciences.

CORCORAN, W. W. Deed of gift of Art Gallery.

TONER, J. M. Deed of foundation of Toner lectures.

TYNDALL, J. Trust fund for the promotion of science in the U.S.

HAMILTON, J. Bequest of \$1,000 to Smithsonian Institution.

HENRY, J. Circular sent with specimens presented.

ARAGO, M. Eulogy on Ampère.

FISCHER, F. Scientific labors of Edward Lartet.

PEABODY, A. P. Scientific education of mechanics and artisans.

BAUER, A. Organic bases.

KLETZINSKY, Prof. Nitrogen bodies of modern chemistry.

EGLESTON, T. Scheme for the qualitative determination of substances by the blowpipe.

Blowpipe apparatus of Hawkins and Wale.

Suess, E. Boundary line between geology and history.

BREZINA, A. Explanation of the principles of crystallography and crystallophysics.

WŒIKOFF, A. Meteorology in Russia.

DONATI, G. B. Phenomena in telegraphic lines during the aurora borealis.

ETHNOLOGY:

Broca, P. The Troglodytes, or cave dwellers, of the valley of the Vézere.

RAU, C. Ancient aboriginal trade in North America.

RAU, C. North American stone implements.

BRUFF, J. G. Indian engravings on the face of rocks along Green River valley in the Sierra Nevadas.

LEE, J. C. Y. Ancient ruin in Arizona.

BARRANDT, A. Haystack mound, Lincoln county, Dakota.

BREED, E. E. Earthworks in Wisconsin.

DEAN, C. K. Mound in Wisconsin.

WARNER, J. Big elephant mound in Grant county, Wisconsin.

CUTTS, J. B. Ancient relics in northwestern Iowa.

PERRINE, T. M. Mounds near Anna, Union county, Illinois.

PETER, R. Ancient mounds in Kentucky.

STEPEENSON, M. F. Mounds in Bartow county, Georgia.

McKinley, W. Mounds in McIntosh and Early counties, Georgia.

HOTCHKISS, T. P. Indian remains in Caddo parish, Louisiana.

LOCKETT, S. H. Mounds in Louisiana.

PEALE, T. R. Prehistoric remains in vicinity of City of Washington, D. C.

DEVEREUX, J. H. Catalogue of cabinet of Indian relics presented to Smithsonian Institution.

DEVEREUX, J. H. Ancient pottery from Phillips County, Arkansas.

KIPP, J. On the accuracy of Catlin's account of the Mandan ceremonies.

272. Smithsonian Contributions to Knowledge. Vol. XIX. 1874. 4to., pp. 640, 6 woodcuts, 21 plates.

CONTENTS.

BARNARD, J. G. Problems of rotary motion. No. 240.

Wood, H. C. Fresh-water alge of North America. No. 241.

NEWCOMB, S. Orbit of Uranus. No. 262.

273. Smithsonian Miscellaneous Collections. Vol. XI. 1874. 8vo., pp. 790, 55 woodcuts, 4 plates.

CONTENTS.

GILL, T. Arrangement of the families of mammals. No. 230.

GILL, T. Arrangement of the families of fishes. No. 247.

Loew, H. Monograph of the diptera of North America. Part III. No. 256.

PACKARD, A. S. Directions for collecting and preserving insects. No. 261.

LE CONTE, J. L. New species of North America coleoptera. Part I No. 264.

LE CONTE, J. L. Classification of North America coleoptera. Part II. No. 265.

274. Smithsonian Miscellaneous Collections. Vol. XII. 1874. 8vo., pp. 767, 86 woodcuts.

CONTENTS.

BAIRD, S. F. Review of American birds. Part I. No. 181.

CLARKE, F. W. The constants of nature. Part I.—Specific gravities. No. 255.

HENRY, J. Circular on Telegraphic announcements of astronomical discoveries. No. 268.

275. Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1873. 43d Congress, 1st Session. Senate Mis. Doc. No. 130. 1874. 8vo., pp. 452, 33 woodcuts.

CONTENTS.

HENRY, J. Secretary's report of operations.

BOARD OF REGENTS, Proceedings of

BAIRD, S. F. Report on Museum and exchanges.

ENDLICH, F. M. List of minerals in the National Museum.

Classified record of monthly meteorological reports preserved in the Smithsonian Institution.

Classified list of meteorological publications, and meteorological articles in periodicals deposited in Library of Congress in 1873.

GARFIELD, J. A. Biographical notice of S. P. Chase and L. Agassiz.

HAMLIN, H. Biographical notice of S. P. Chase.

PARKER, P. Biographical notice of L. Agassiz.

Hamilton, J. Bequest deposited in United States Treasury.

Dodge, N. S. Memoir of Charles Babbage.

275. Report for 1873—Continued.

HENRY, J. On the "Moon-Hoax."

BABBAGE, C. Extracts from writings of

STEBBINS, R. P. Memoir of Louis Agassiz.

GRAY, A. Memoir of John Torrey.

STEVENS, J. A. Memoir of George Gibbs.

DALTON, J. C. Origin and propagation of disease.

HELMHOLZ H.; MAXWELL, J. C. Later views of the connection of electricity and magnetism.

Gould, B. A. Account of the astronomical observatory at Cordoba, Argentine Republic.

MAILLY, E. Estimate of the population of the world.

Morin, A. Warming and ventilating buildings. Part I.

DEFOREST, E. Additions to a memoir on methods of interpolation.

Part II.

ETHNOLOGY:

SCHUMACHER, P. Remarks on the Kjökken-Möddings on the North-west Coast of America.

BERENDT, C. H. On a grammar and dictionary of the Carib or Karif language, with some account of the people by whom it is spoken.

GILLMAN, H. The mound-builders and platyenemism in Michigan.

MASON, O. T. The Leipsic "Museum of Ethnology."

PERRINE, T. M. Antiquities of Union county, Illinois.

PATTON, A. Antiquities of Knox county, Indiana, and Lawrence County, Illinois.

Miscellaneous Correspondence:

Dall, W. H. Explorations on the Western coast of North America.

PIERSON, W. M. Discovery of a large meteorite in Mexico.

BRUNOT, F. R. On the habits of the beaver.

JEVONS, W. S. On a national library.

PRIZE QUESTIONS OF SCIENTIFIC SOCIETIES:

Society for the Encouragement of Science, Literature, and Art, Dan-kirk, France.

Society of Science, Art, and Literature of Hainaut, Mons. Belgium.

Royal Institute for the Encouragement of the Natural; Economical, and Technological Sciences, Naples, Italy.

Royal Academy of Science, Literature, and the Fine Arts, Brussels, Belgium.

Society of Sciences of Haarlem, Holland.

- 276. The Constants of Nature. Part II. A Table of Specific Heats for Solids and Liquids. By Frank W. Clarke. April, 1876. 8vo., pp. 58. (M. C. XIV.)
- 277. Tables, Distribution, and Variations of the Atmospheric Temperature in the United States and some adjacent parts of America. By Charles A. Schott. April, 1876. 4to., pp. 360, 10 woodcuts, 2 plates, 3 maps. (S. C. xxi.)
- 278. Check-list of Publications of the Smithsonian Institution. July, 1874. 8vo., pp. 24.

- 279. On Strain and Over-action of the Heart. Toner Lecture No. III. Delivered May 14th, 1874. By J. M. Da Costa. August. 1874. 8vo., pp. 32, 2 woodcuts. (M. C. xv.)
- 280. Statement and Exposition of certain Harmonies of the Solar System. By Stephen Alexander. March, 1875. 4to., pp. 104, 20 woodcuts. (S. C. xxi.)
- 281. On the General Integrals of Planetary Motion. By Simon Newcomb. December, 1874. 4to., pp. 40. (S. C. xxi.)
- 282. A Study of the Nature and Mechanism of Fever. Toner Lecture No. IV. Delivered January 20, 1872. By Horatio C. Wood. February, 1875. 8vo., pp. 50. (M. C. xv.)
- 283. Catalogue of the Fishes of the East Coast of North America. By Theodore Gill. 1875. 8vo., pp. 56. (M. C. xiv.)
- 284. Smithsonian Contributions to Knowledge. Vol. XX. 1876. 4to., pp. 794, 4 woodcuts, 26 plates.

COFFIN, J. H. The winds of the globe. No. 268.

285. Smithsonian Contributions to Knowledge. Vol. XXI. 1876. 4to., pp. 543, 30 woodcuts, 9 plates, 3 maps.

CONTENTS.

ALEXANDER, S. Harmonies of the solar system. No. 280.

NEWCOMB, S. Integrals of planetary motion. No. 281.

Swan, J. G. Haidah Indians of Queen Charlotte's Islands, British Columbia. No. 267.

Schott, C. A. Tables, atmospheric temperature in the United States. No. 277.

286. Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1874. 43d Congress, 2d Session. House Doc. No. 56. 1875. 8vo., pp. 416, 46 woodcuts.

CONTENTS.

HENRY, J. Secretary's report of operations.

BOARD of REGENTS. Proceedings of

List of articles deposited by the Smithsonian Institution in the Corcoran Art Gallery.

List of Indian vocabularies received from the Wheeler Expedition.

MITCHELL, S. W. Inquiries relative to the disease known as "Chorea." Classified record of meteorological material preserved in the Smithsonian Institution. Auroras, instruments, meteors, rain, wind, general meteorology, local meteorology.

GRAY, A.; COPPÉE, H. Report of Committee of Regents on the Smithsonian Museum.

286. Report for 1874—Continued.

ARAGO, M. Eulogy on La Place.

MAILLY, M. Eulogy on Quetelet.

DUMAS, M. Eulogy on A. A. de la Rive.

HILGARD, J. E. Tides and tidal action in harbors.

LEMSTRÖM, S.; DE LA RIVE, A. A. Electricity of the atmosphere and the Aurora Borealis.

DE CANDOLLE, A.; GRAY, J. E. On a dominant language for science.

SCHOTT, C. A.; EVERETT, J. D. Underground temperature.

DUPRE, W.; HENRY, J. Earthquakes in North Carolina, 1874.

DE LA RIVE, A. A. Report on the transactions of the Society of Physics and Natural History of Geneva, from July, 1872, to June, 1878.

MORIN, A. Warming and ventilating buildings. Part II.

ETHNOLOGY:

Schumacher, P. Ancient graves and shell-heaps of California.

KING, W. M. Account of the burial of an Indian squaw, San Bernardino county, California.

McWhorter, T. Ancient mounds of Mercer county, Illinois.

PRATT, W. H. Antiquities of Whitesides county, Illinois.

FARQUHARSON, R. J. A study of skulls and long bones, from mounds near Albany, Illinois.

TIFFANY, A. S. The shell-bed skull.

HILL, G. W. Antiquities of northern Ohio.

ROBERTSON, R. S. The age of stone, and the troglodytes of Breck-inridge county, Kentucky.

DAVIS, A. C. Antiquities of Isle Royale, Lake Superior.

SMITH, J. W. C. Antiquities of Yazoo county, Mississippi.

WRIGHT, D. F. Antiquities of Tennessee.

LAW, A. E. Antiquities of Blount county, Tennessee.

Cushing, F. H. Antiquities of Orleans county, New York.

ROBERTSON, R. S. Antiquities of La Porte county, Indiana.

ROBERTSON, R. S. Antiquities of Allen and De Kalb counties, Ind.

HAILE, J.; McHENRY, J. W. Antiquities of Jackson county, Tennessee.

ANDERSON, W. Antiquities of Perry county, Ohio.

BRYAN, O. N. Antiquities of Charles county, Maryland.

Kron, F. J. Antiquities of Stanley and Montgomery counties, North Carolina.

MITCHELL, A. Antiquities of Florida.

BARTRAM, J. Antiquities of Florida.

287. The Archæological Collection of the United States National Museum, in charge of the Smithsonian Institution, Washington, D. C. By Charles Rau. 1876. 4to., pp. 118, 340 woodcuts. (S. C. xxII.)

288. The Constants of Nature. First supplement to Part I. Specific Gravities, Boiling Points, and Melting Points. By Frank W. Clarke. April, 1876. 8vo., pp. 62. (M. C. XIV.)

289. The Constants of Nature. Part III. Tables of Expansion by Heat for Solids and Liquids. By Frank W. Clarke. April, 1876. 8vo., pp. 58. (M. C. xiv.)

- 290. Circular for Distribution at the Centennial Exhibition, 1876, containing List of Smithsonian Publications and Rules of Distribution; List of Foreign Agents; Number of Foreign and Domestic Institutions in Correspondence; List of Regents, Officers, and Assistants of the Institution. 1876. 12mo., pp. 12.
- 291. Dual Character of the Brain. Toner Lecture No. II. Delivered April 22, 1874. By C. E. Brown-Séquard. January, 1877. 8vo., pp. 26. (M. C. xv.)
- 292. Check-list of North American Batrachia and Reptilia; with a Sytematic List of the Higher Groups, and an Essay on Geographical Distribution based on the specimens contained in the United States National Museum. By EDWARD D. COPE. 1875. 8vo., pp. 108. (M. C. XIII.) Bulletin of the National Museum No. 1.
- 293. Contributions to the Natural History of Kerguelen Island, made in Connection with the American Transit-of-Venus Expedition, 1874—75.

 By J. H. Kidder. I.—Ornithology, by Elliott Coues. 1875. 8vo., pp. 61. (M. C. XIII.) Bulletin of the National Museum, No. 2.
- 294. Contributions to the Natural History of Kerguelen Island, made in connection with the United States Transit-of-Venus Expedition, 1874-75. By J. H. Kidder. II.—Oology, Botany, &c. 1876. 8vo., pp. 124. (M. C. XIII.) Bulletin of the National Museum, No. 3.

KIDDER, J. H.; Coues, E. Oology.

BOTANY:

GRAY, ASA. A.—Phænogamia, filices, et lycopodiacem.

JAMES, T. P. B.-Musci.

TUCKERMAN, E. C.—Lichones.

FARLOW, W. G. Algæ.

. CROZET FLOBA.

ENDLICH, F. M. Geology.

KIDDER, J. H. Mammals.

GILL, T. N. Fish.

DALL, W. H. Mollusks.

INSECTS:

OSTEN-SACKEN, C. C. Diptera.

HAGEN, H. A. Pseudo-neuroptera.

SMITH, S. I. Crustaceans.

VERRILL, A. E. Annelids, echinoderms, and anthozos.

KERSHNER, E. Collection.

KIDDER, J. H.; COUES, E. A study of chionis minor.

295. Birds of Southwestern Mexico, collected by Francis E. Sumichrast for the United States National Museum. By George N. Lawrence.

1875. 8vo., pp. 56. (M. C. XIII.) Bulletin of the National Museum, No. 4.

- 296. Catalogue of the Fishes of the Bermudas. Based chiefly upon the collections of the United States National Museum. By G. Brown Goode. 1876. 8vo., pp. 84. (M. C. XIII.) Bulletin of the National Museum, No. 5.
- 297. Classification of the Collection to illustrate the Animal Resources of the United States. A List of Substances derived from the Animal Kingdom, with Synopsis of the Useful and Injurious Animals and a Classification of Methods of Capture and Utilization. By G. Brown Goode. 1876. 8vo., pp. 140. (M. C. XIII.) Bulletin of the National Museum, No. 6.
- 298. Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1875. 44th Congress, 1st Session. Senate Mis. Doc. No. 115. 1876. 8vo., pp. 422, 354 woodcuts.

CONTENTS.

HENBY, J. Secretary's report of operations.

BOARD OF REGENTS, Proceedings of

BAIRD, S. F. Report on Museum and explorations.

BAIRD, S. F. Report on the proposed plan of exhibit by the Smithsonian Institution at the International Centennial Exhibition at Philadelphia.

PARKER, P. Eulogy on Henry Wilson.

ARAGO, M. Eulogy on Alexander Volta.

DE CANDOLLE, A. Probable future of the human race.

DE CANDOLLE, A. Report on the transactions of the Society of Physics and Natural History of Geneva, from July, 1873, to June, 1874.

PRESTWICH, J. The past and future of geology.

WEX, H. G. Diminution of the water of rivers and streams.

TAYLOR, W. B. Refraction of sound.

HENRY, J. On the organization of local scientific societies.

ETHNOLOGY:

MORTILLET, G. DE; CHANTRE, E. International code of symbols for charts of pre-historic archæology.

GILLMAN, H. Characteristics pertaining to ancient man in Michigan. Abbott, C. C. The stone age in New Jersey.

299. Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1876. 44th Congress, 2d Session. Senate Mis. Doc. No. 46. 1877. 8vo., pp. 488, 73 woodcuts.

CONTENTS.

HENRY, J. Secretary's report of operations.

BOARD OF REGENTS, Proceedings of

BAIRD, S. F. Report on Museum and explorations.

BAIRD, S. F. Report on Centennial Exhibition of 1876.

Centennial awards to Smithsonian Institution.

299. Report for 1876—Continued.

GRAY, A.; SARGENT, A. A.; CLYMER, H. Report of Committee on the Museum.

Bancroft, G. Memorial to Congress, in behalf of the Regents, for new museum.

List of collections presented by foreign commissioners to the United States.

ARAGO, M. Eulogy on Gay Lussac.

FIALHO, A. Biographical sketch of Dom Pedro II.

TAYLOR, W. B. Kinetic theories of gravitation.

PILAR, G. The revolutions of the crust of the earth.

KIRKWOOD, D. The asteroids between Mars and Jupiter.

ETHNOLOGY:

MASON, O. T. The Latimer collection of antiquities from Porto Rico in the National Museum at Washington, D. C.

ROMER, F. F. Pre-bistoric antiquities of Hungary.

BLONDEL, S. Jade. A historical, archeological, and literary study of the mineral called Yu by the Chinese.

WILLIAMSON, G. Antiquities in Guatemala.

BERENDT, C. H. Collection of historical documents in Guatemala.

STRONG, MOSES. Observations on the pre-historic mounds of Grant county, Wisconsin.

SNYDER, J. F. Deposits of flint implements, Illinois.

SMITH, C. D. Ancient mica mines in North Carolina.

PEET, S. D. Double-walled earthwork in Ashtabula county, Ohio.

ELLSWORTH, E. W. Ancient implement of wood, from Connecticut.

Powers, S. Centennial mission to the Indians of Western Nevada and California.

DOYLE, W. E. Indian forts and dwellings, Indian Territory.

BRACKETT, A. G. The Sioux or Dakota Indians.

- 300. On the Surgical Complications and Sequels of the Continued Fevers. With a Bibliography of works on Diseases of the Joints, Bones, Larynx, the Eye, Gangrene, Haematoma, Phlegmasia, etc. Tones Lecture No. V. Delivered February 17, 1876. By William W. Keen. March, 1876. 8vo., pp. 72, 5 woodcuts. (M. C. xv.)
- 301. List of Publications of the Smithsonian Institution, July, 1877. 1877. 8vo., pp. 72. (M. C. xiv.)
- 302. Subcutaneous Surgery, its Principles, and its recent Extension in Practice. Toner Lecture No. VI. Delivered September 13, 1876. By William Adams. April, 1877. 8vo., pp. 20. (M. C. xv.)
- 303. Contributions to the Natural History of the Hawaiian and Fanning Islands and Lower California, made in connection with the United States North Pacific Surveying Expedition, 1873-'75. By Thos. H. Streets. 1877. 8vo., pp. 172. (M. C. XIII.) Bulletin of the National Museum, No. 7.

303. Natural History of Hawaiian Islands—Continued. CONTENTS.

ORNITHOLOGY.

HERPETOLOGY.

ICHTHYOLOGY:

- I. Fishes of Upper and Lower California.
- II. Fishes of the Hawaiian Islands.
- III. Fishes of the Fanning Islands.
- IV. Fishes from the Samoan Islands.

CRUSTACEA.

BOTANY.

- 304. Index to the Names which have been appplied to the Subdivisions of the Class Brachiopoda, excluding the Rudistes, previous to the year 1877. By W. H. Dall. 1877. 8vo., pp. 88. (M. C. XIII.) Bulletin of the National Museum, No. 8.
- 305. Contributions to North American Ichthyology. Based primarily on the collections of the United States National Museum. I.—Review of Rafinesque's Memoirs on North American Fishes. By David S. Jordan. 1877. 8vo., pp. 56. (M. C. XIII.) Bulletin of the National Museum, No. 9.
- 306. Contributions to North American Ichthyology. Based primarily on the Collections of the United States National Museum. II. A.— Notes on Cottidæ, Etheostomatidæ; Percidæ, Centrarchidæ, Aphododeridæ, Dorysomatidæ, and Cyprinidæ, with revisions of the genera and descriptions of new or little known species. B.—Synopsis of the Siluridæ of the Fresh Waters of North America. By David S. Jordan. 1877. 8vo., pp. 124, 45 plates of 100 figures. (M. C. XIII.) Bulletin of the National Museum, No. 10.
- 307. Report on the Centennial Exhibition, Philadelphia. By S. F. BAIRD. 1877. 8vo., pp. 22.
- 308. Contributions to North American Ichthyology, based primarily on the collections of the United States National Museum. III. A.—On the Distribution of the Fishes of the Alleghany Region of South Carolina, Georgia, and Tennessee, with descriptions of new or little known species. By David S. Jordan and Alembert W. Brayton. B.—A Synopsis of the Family Catostomidæ. By David S. Jordan. 1878. 8vo., pp. 237. (M. C. XXIII.) Bulletin of the National Museum, No. 12.
- 309. List of Foreign Correspondents of the Smithsonian Institution. Corrected to January, 1878. March, 1878. 8vo., pp. 120. (M. C. xv.)
- 310. On the Internal Structure of the Earth, considered as affecting the Phenomena of Precession and Nutation; supplementary to article under the above head, Smithsonian Contributions to Knowledge, Vol. XIX, No. 240, being the third of the Problems of Rotary Mo-

- tion. By J. G. BARNARD. August, 1877. 4to., pp. 19, 4 woodcuts. (S. C. XXIII.)
- 311. Index Catalogue of Books and Memoirs relating to Nebulse and Clusters, etc. By EDWARD S. HOLDEN. November, 1877. 8vo., pp. 126. (M. C. xiv.)
- 312. Smithsonian Miscellaneous Collections. Vol. XIII. 1878. 8vo., pp. 982, 45 plates. Bulletins of the National Museum, Nos. 1-10.

COPE, E. D. North American batrachia and reptilia. No. 292.

KIDDER, J. H.; Cours, E. Birds of Kerguelen Island. No. 293.

Kidder, J. H., and others. Oology, botany, &c., of Kerguelen Island. No. 294.

LAWRENCE, G. N. Birds of Mexico. No. 295.

GOODE, G. B. Fishes of Bermuda. No. 296.

GOODE, G. B. Classification of animal resources. No. 297.

STREETS, T. H. Natural history of Hawaiian Islands, Fanning Islands, and Lower California. No. 808.

DALL, W. H. Index of brachiopoda. No. 304.

JORDAN, D. S. North American Ichthyology. Nos. 805 and 806.

- 313. The Flora of St. Croix and the Virgin Islands. By Baron H. F. A. EGGERS. 1879. 8vo., pp. 139. (M. C. XXIII.) Bulletin of the National Museum, No. 13.
- 314. Smithsonian Miscellaneous Collections. Vol. XIV. 1878. 8vo., pp. 911, 4 plates.

CONTENTS.

DE SAUSSURE, H. Synopsis of American wasps. No. 254.

GILL, T. Catalogue of fishes. No. 288.

CLARKE, F. W. Specific gravity tables. Supp. I. No. 288.

CLARKE, F. W. Specific heat tables. No. 276.

CLARKE, F. W. Tables of expansion by heat. No. 289.

Catalogue of photograph portraits of North American Indians. No. 216. Check list of Smithsonian publications to July, 1877. No. 301.

8vo.,

HOLDEN, E. S. Catalogue of books relative to nebulæ. No. 311.

315. Smithsonian Miscellaneous Collections. Vol. XV. 1878.

pp. 880, 53 woodcuts.

CONTENTS.

WATSON, S. Botanical index. No. 258..

WOODWARD, J. J. Toner Lecture I. Cancerous tumors. No. 266.

Brown-Séquard, C. E. Toner Lecture II. The brain. No. 291.

DA COSTA, J. M. Toner Lecture III. The heart. No. 279.

Wood, H. C. Toner Lecture IV. Study of fever. No. 282.

KEEN, W. W. Toner Lecture V. Continued fevers. No. 800.

ADAMS, W. Toner Lecture VI. Subcutaneous surgery. No. 302.

List of foreign correspondents of the Smithsonian Institution to January, 1878. No. 809.

- 315. Miscellaneous Collections. Vol. XV—Continued.
 - Circular in reference to American archmology. No. 816.
 - Circular. Inquiries relative to crawfish and crustaces. No. 819.
 - Circular relating to collections of living reptiles. No. 820.
- 316. Circular in reference to American Archæology. 1878. 8vo., pp. 15, 38 woodcuts. (M. C. xv.)
- 317. A Classification and Synopsis of the Trochilidæ. By Daniel Giraud Elliot. 1879. 4to., pp. 289, 127 woodcuts. (S. C. xxiii.)
- 318. On the Remains of Later Pre-historic Man, obtained from Caves in the Catherina Archipelago, Alaska Territory, and especially from the Caves of the Aleutian Islands. By W. H. Dall. 1878. 4to., pp. 44, 10 plates of 39 figures. (S. C. xxII.)
- 319. Circular of Inquiries relative to the Natural History of the American Crawfish and other Fresh Water Crustacea. 1878. 8vo., pp. 8, 2 woodcuts. (M. C. xv.)
- 320. Circular relating to Collections of Living Reptiles. 1878. 8vo., pp. 2. (M. C. xv.)
- 321. The Nature of Reparatory Inflammation in Arteries after Ligatures, Acupressure, and Torsion. Toner Lecture No. VII. Delivered June 27, 1878. By Edw. O. Shakespeare. March, 1879. 8vo., pp. 74, 7 plates of 11 figures. (M. C. xvi.)
- 322. Smithsonian Miscellaneous Collections. Vol. XVI. 1880. 8vo., pp. 950, 871 woodcuts, 7 plates.

TRYON, G. W. Land and fresh water shells. Strepomatides. No. 253.

OSTEN SACKEN, C. R. Catalogue of diptera. No. 270.

SHAKESPEARE, E. O. Nature of reparatory inflammation. No. 821.

Circular relative to scientific and literary exchanges. No. 824.

Business arrangements of the Smithsonian Institution. No. 825.

ELLIOT, D. G. List of species of humming birds. No. 884.

List of the principal scientific and literary institutions in the United States. No. 885.

List of publications of the Smithsonian Institution. No. 844.

323. Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1877. 45th Congress, 2d Session. Senate Mis. Doc. No. 35. 1878. 8vo., pp. 500, 49 woodcuts.

CONTENTS.

HENRY, J. Secretary's report of operations.

BOARD OF REGENTS, Proceedings of

BAIRD, S. F. Report on Museum and explorations.

BAIRD, S. F. List of explorations furnishing collections to the National Museum, 1888 to 1877.

323. Report for 1877—Continued.

HOLMGREN, F. Color blindness in its relation to accidents by rail and sea.

HENRY, J. Color blindness.

PLANTAMOUR, E. Report on the transactions of the Geneva Society of Physics and Natural History, from June, 1874, to June, 1875.

MÜLLER, J. Report on the transactions of the Geneva Society of Physics and Natural History, from June, 1875, to June, 1876.

FAVRE, A. Report on the transactions of the Geneva Society of Physics and Natural History, from June, 1876, to June, 1877.

ETHNOLOGY:

Cannon, G. L. Antiquities of Jefferson and Clear Creek counties, Colorado.

STRONG, Moses. Antiquities in Wisconsin.

HART, J. N. DE. The mounds and osteology of the mound builders of Wisconsin.

Breed, E. E. Pits at Embarrass, Wisconsin.

MOULTON, M. W. Mounds in Delaware county, Iowa.

Knapp, Mrs. G. Earthworks on the Arkansas river, sixteen miles below Little Rock.

LYKINS, W. H. R. Antiquities of Kansas City, Missouri.

SHAW, J. The mound builders in the Rock river valley, Illinois.

COCHRANE, J. Antiquities of Mason county, Illinois.

HILL, G. W. Ancient earthworks of Ashland county, Onio.

CASE, H. B. Flint implements in Holmes county, Ohio.

MILLER, F. Mound in Trumbull county, Ohio.

FRIEL, J. Antiquities of Hancock county, Kentucky.

Robertson, R. S. Antiquities of Nashville, Tennessee.

CLARK, W. M. Antiquities of Tennessee.

Jones, Jr., C. C. Aboriginal structures in Georgia.

BAILEY, W. B. F. Antiquities of Spalding county, Georgia.

GAINES, A. S.; CUNNINGHAM, K. M. Shell-heaps on Mobile river.

RAU, C. The stock in trade of an aboriginal lapidary.

RAU, C. Observations on a gold ornament from a mound in Florida.

HALDEMAN, S. S. On a polychrome bead from Florida.

HARRISON, A. M. Colored bead dug from a mound on the eastern coast of Florida.

MAYBERRY, S. P. Shell-heaps at the mouth of Saint John's river, Florida.

TAYLOR, W. M. Ancient mound in western Pennsylvania.

SHEPARD, E. M. Deposit of arrow heads near Fishkill, New York.

GIBBS, G. J. Stone celts in the West Indies and Africa.

GALT, F. L. The Indians of Peru.

Bowers, S. History and antiquities of Santa Rosa Island, California McParlin, T. A. Notes on the history and climate of New Mexico.

WEISMANN, A. On the change of the Mexican axolotl to an amblystoms.

ABBE, C. Short memoirs on meteorological subjects, translated by, viz:

HANN, J. On the diminution of aqueous vapor with increasing altitude in the atmosphere.

HANN, J. On the influence of rain upon the barometer and upon the formation of precipitation in general.

333. Report for 1877—Continued.

HANN, J. Atmospheric pressure and rain-fall.

HANN, J. The laws of the variation of temperature in ascending currents of air.

Sohncke, L. The law of the variation of temperature in ascending moist currents of air.

REYE, T. Rain-fall and barometric minima.

HANN, J. On the relation between the difference of pressure and the velocity of the wind according to the theories of Ferrel and Colding.

FERREL, W. Reply to the criticisms of J. Hann.

Colding, A. Some remarks concerning the nature of currents of air.

Colding, A. On the whirlwind at St. Thomas on the 21st of August, 1871.

PESLIN, M. On the relation between barometric variations and the general atmospheric currents.

- 324. Circular relative to Scientific and Literary Exchanges. 1879. 8vo., pp. 2. (M. C. xvi.)
- 325. Business Arrangements of the Smithsonian Institution. January, 1879. 8vo., pp. 7. (M. C. xvi.) Free.
- 326. Catalogue of the Collection to illustrate the Animal Resources and the Fisheries of the United States, exhibited at Philadelphia, in 1876, by the Smithsonian Institution and the United States Fish Commission, and forming a part of the United States National Museum. By G. Browne Goode. 1879. 8vo., pp. 367. (M. C. XXIII.) Bulletin of the National Museum, No. 14.
- 327. The Scientific Writings of James Smithson. Edited by W. J. Rhees. 1879. 8vo., pp. 166, 32 woodcuts, 1 plate. (Portrait.) (M. C. xxi.)

CONTENTS.

SMITHSON, J. Scientific writings.

Johnson, W. R. Memoir on the scientific character and researches of James Smithson.

IRBY, J. R. McD. Works and character of James Smithson.

328. The Smithsonian Institution. Documents relative to its Origin and History. By WILLIAM J. RHEES. 1879. 8vo., pp. 1027. (M. C. XVII.)

CONTENTS.

SMITHSON, J. Will of

Correspondence between attorneys in England, Department of State, Richard Rush, and others relative to the bequest of Smithson.

RUSH, R. The case stated.

Opinion of English counsel.

Decree of Chancery Court awarding bequest to the United States.

Account of case of the United States.

Bill of costs of the case of the United States.

RUSH, R. Accounts of

SMITHSON, J. Schedule of personal effects.

338. Smithsonian Institution. Documents—Continued.

Smithson, J. Residuary legacy of

Congressional proceedings in relation to the bequest. 1885 to 1877.

Twenty-fourth to Forty-fourth Congress.

HENRY, J. Digest of act of Congress to establish the Smithsonian Institution.

ADAMS, J. Q. Extracts from Memoirs of, relative to Smithson bequest.

TREASURY OF U. S. Accounts of, with Smithson fund.

Proposed applications of Smithson bequest.

Report of Organization Committee of the Board of Regents of the Smithsonian Institution.

HENRY, J. Programme of organization of Smithsonian Institution. Letters relative to the programme of organization.

329. The Smithsonian Institution. Journals of the Board of Regents, Reports of Committees, Statistics, etc. By WILLIAM J. RHEEL 1879. 8vo., pp. 851. (M. C. XVIII.)

CONTENTS.

Journal of Proceedings of the Board of Regents from 1846 to 1876.

Report of the Special Committee of the Board of Regents on the distribution of the income.

Report of the Special Committee of the Board of Regents—Prof. Felton—on the present of the Greek album from Miss E. Contaxaki.

Report of the Special Committee of the Board of Regents—Prof. Feltonon the purchase of the Stanley Indian gallery.

HENRY, J. Communication relative to a publication by Prof. S. F. B. Morse.

Report of the Special Committee of the Board of Regents on the invention of the electro-magnetic telegraph.

Report of the Special Committee of the Board of Regents relative to the Smithsonian fire of January 24, 1865.

Act of Congress to transfer custody of library to Library of Congress.

Act of Congress to receive into Treasury of the United States residuary legacy of Smithson, and to authorize increase of fund to a million dollars.

Report of the Special Committee of the Board of Regents on best use of new hall of Institution.

Report of Executive Committee of the Board of Regents on the Washington city canal.

Report of the Executive Committee of the Board of Regents relative to the Corcoran Art Gallery.

Report of the Special Committee of the Board of Regents on the museum. Eulogies and Biographical Sketches:

Agassiz, L.; Bache, A. D.; Chase, S. P.; Cleaveland, P.; Douglas, S. A.; Espy, J. P.; Felton, C. C.; Harvey, W. H.; Irving. W.; Jewett, C. C.; Pearce, J. A.; Priestley, J.; Rush, R.; Seaton, W. W.; Totten, J. G.; Turner, W. W.; Wilson, H.; Wurdeman, G.

AGASSIZ, L. Narrative of expedition through Straits of Magellan to California.

BACHE, A. D. Will of, establishing Bache Scientific Fund.

CORCORAN, W. W. Deed of gift and trust of Corcoran Art Gallery.

329. Smithsonian Institution. Journals, &c.—Continued.

TONER, J. M. Deed of, establishing Toner Lectures for advance of medical science.

TYNDALL, J. Deed of, establishing trust for promotion of science in the United States.

HAMILTON, J. Bequest of, to Smithsonian Institution.

Circular sent with specimens of natural history, etc., presented to institutions.

Journal of the Executive Committee from Sept. 12, 1846, to Dec. 21, 1849. Report of the Executive Committee, from 1847 to 1875.

Journal of the Building Committee, from Feb. 17, 1847, to Dec. 1, 1847.

Report of the Building Committee, 1847-1867.

General financial and statistical statements, receipts and expenditures, classified, 1846 to 1877.

Statistics of literary and scientific exchanges, additions to the library, etc., 1846 to 1877.

Appropriations from National Treasury by Congress for Smithsonian Institution and National Museum, 1846-1876.

List of Regents of the Institution from 1846 to 1879, according to mode of appointment, residence, etc.

Act of Congress to establish the Smithsonian Institution, August 10, 1846.

The same, according to the Revised Statutes.

By-Laws of the "Establishment."

The "Establishment" of the Smithsonian Institution, organization and journal of proceedings, 1846 to 1877.

Examination of Professor Henry by the English Government Scientific Commission, June 28, 1870.

330. Smithson and his Bequest. By WILLIAM J. RHEES. 1880. 8vo., pp. 76, 9 plates. (M. C. xxi.)

CONTENTS.

Life of Smithson.

Legislation of Congress in relation to the bequest.

List of first Board of Regents.

Obituary notice of Smithson, March, 1880.

DUTENS, L. Account of the First Duke of Northumberland.

Notice of the First Duke in The Gentleman's Magazine, 1786.

Coffin-plate inscription of Hugh Smithson.

Account of Earl Percy.

Notice of Smithson's paper on "Tabasheer."

DAVY, SIR H. Notice of Smithson's paper on "Calamines."

Illustrations of presentations of books to Smithson.

BERZELIUS. Notice of Smithson's researches.

Extracts from Smithson's scientific writings.

Catalogue of the library of Smithson.

Notice of the city of Washington, in Harriott's travels.

Notice of the city of Washington, in Weld's travels.

331. The Palenque Tablet, in the United States National Museum, Washington, D. C. By Charles Rau. 1879. 4to., pp. 90, 2 plates. (S. C. XXII.)

332. Proceedings of the U. S. National Museum for 1878. Vol. I. 1879. 8vo., pp. 524, 8 cuts, 8 plates. (M. C. xix.)

CONTENTS.

BEAN, T. H. Description of a new sparoid fish, Sargus holbrookii, from Savannah bank.

On the occurrence of Stichæus punctatus (Fabr.) Kröyer, at St. Michael's, Alaska.

On the identity of Euchalarodus Putnami Gill, with Pleuronectes glaive-(Storer) Gill, with notes on the habits of the species.

Description of a species of Lycodes (L. Turneri) from Alaska. See also under Goode and Bean.

BELDING, L. A partial list of the birds of Central California.

Cook, C. The manufacture of porpoise-oil.

DALL, W. H. Description of new forms of mollusks from Alaska contained in the collections of the National Museum.

Postpliocene fossils in the Coast Range of California.

Fossil mollusks from later tertiaries of California.

Note on shells from Costa Rica kitchenmidden, collected by Drs. Flint and Bransford.

Distribution of Californian tertiary fossils.

Descriptions of new species of shells from California in the collections of the National Museum.

Report on the limpets and chitons of the Alaskan and Arctic regions. descriptions of genera and species believed to be new.

EDWARDS, V. N. On the occurrence of the Oceanic Bonito, Orcyans pelamys (Linné) Poey, in Vineyard Sound, Massachusetts.

GILL, T. Synopsis of the pediculate fishes of the Eastern Coast of Extratropical North America.

Note on the Antennariidæ.

On the proper specific name of the common pelagic antennariid Pterophryne.

Note on the Ceratiidæ.

Note on the Maltheidæ.

GOODE, G. B. The Clupea tyrannus of Latrobe.

The occurrence of Belone latimanus in Buzzard's bay, Massachusetts. The voices of Crustaceans.

A revision of the American species of the genus Brevoortia, with a description of a new species from the Gulf of Mexico.

The occurrence of Hippocampus antiquorum, or an allied form, on St. George's Banks.

The occurrence of the Canada porcupine in West Virginia.

On two fishes from the Bermudas mistakenly described as new by Dr. Günther.

GOODE, G. B.; BEAN, T. H. The Craig flounder of Europe, Glyptory-halus cynoglossus, on the coast of North America.

The Oceanic Bonito on the coast of the United States.

Description of Caulolatilus microps, a new species of fish from the Gulf coast of Florida.

- 332. Proceedings of National Museum. Vol. I—Continued.
 - GOODE, G. B.; BEAN, T. H. On a new serranoid fish, Epinephelus Drum-mond-Hayi, from the Bermudas and Florida.
 - Descriptions of two new species of fishes, Lutjanus Blackfordii and Lutjanus stearnsii, from the Coast of Florida.
 - A note upon the Black Grouper (Epinephelus nigritus (Holbrook) Gill) of the Southern coast.
 - Descriptions of two gadoid fishes, Phycis Chesteri and Haloporphyrus viola, from the deep-sea fauna of the Northwestern Atlantic.
 - Description of Argentina syrtensium, a new deep-sea fish from Sable Island Bank.
 - The identity of Rhinonemus caudacuta, (Storer) Gill with Gadus cimbrius Linn.
 - Note on Platessa ferruginea D. H. Storer, and Platessa rostrata H. R. Storer.
 - On the identity of Brosmius brosme americanus Gill, with Brosmius brosme (Müller) White.
 - JACKSON, J. B. S. Arsenic acid for protecting anatomical preparations from insects.
 - JEFFERSON, J. P. On the mortality of fishes in the Gulf of Mexico in 1878.
 - JEFFERSON, J. P.; PORTER, J. Y.; Moore, T. On the destruction of fish in the vicinity of the Tortugas during the months of September and October, 1878.
 - JORDAN, D. S. Notes on a collection of fishes from Clackamas river, Oregon.
 - JORDAN, D. S.; GILBERT, C. H. Notes on the fishes of Beaufort harbor, North Carolina.
 - LAWRENCE, G. N. Catalogue of the birds of Dominica, from collections made for the Smithsonian Institution, by Frederick A. Ober, together with his notes and observations.
 - Catalogue of the birds of St. Vincent, from collections made by Fred.

 A. Ober, under the directions of the Smithsonian Institution.

 with his notes thereon.
 - Catalogue of the birds of Antigua and Barbuda, from collections made for the Smithsonian Institution, by Fred. A. Ober, with his observations.
 - Catalogue of the birds of Grenads, from a collection made by Fred.

 A. Ober for the Smithsonian Institution, including others seen by him, but not obtained.
 - Catalogue of the birds collected in Martinique, by Fred. A. Ober for the Smithsonian Institution.
 - Catalogue of a collection of birds obtained in Guadeloupe for the Smithsonian Institution, by Fred. A. Ober.
 - A general catalogue of the birds noted from the islands of the Lesser Antilles visited by Fred. A. Ober; with a table showing their distribution, and those found in the United States.
 - LUPTON, N.T. On the breeding habits of the sea-catfish (Ariopsis milberti?)

 MERRILL, J. C. Notes on the ornithology of Southern Texas, being a list of birds observed in the vicinity of Fort Brown, Texas, from February, 1876, to June, 1878.

- 382. Proceedings of National Museum. Vol. I-Continued.
 - POEY, F. Notes on the American species of the genus Cybium.
 - PRATT, R. H. Catalogue of casts taken by Clark Mills of the heads of sixty-four Indian prisoners of various western tribes, and held at Fort Marion, St. Augustine, Florida, in charge of R. H. Pratt.
 - RIDGWAY, R. On a new humming bird (Atthis Ellioti) from Guatemala.

 A review of the American species of the genus Scops, Savigny.
 - Descriptions of several new species and geographical races of birds contained in the collection of the United States National Museum.
 - Description of two new species of birds from Costa Rica, and notes on other rare species from that country.
 - Descriptions of new species and races of American birds, including a synopsis of the Genus Tyrannus, Cuvier.
 - STEARNS, S. A note on the Gulf menhaden, Brevoortia patronus, Goode. STEINDACHNER, F. Note on Perca flavescens.
 - WILMOT, S. Notes on the western gizzard shad, Dorosoma cepedianum heterurum (Raf.) Jordan.
- 8vo., pp. 503, 2 wood cuts, 7 plates. (M. C. xix.)

- BEAN, T. H. A list of European fishes in the collection of the United States National Museum.
 - On the species of Astroscopus of the Eastern United States.
 - On the occurrence of *Hippoglossus vulgaris*, Flem., at Unalashka and St. Michael's, Alaska.
 - Description of an apparently new species of Gasterosteus (G. atkinsii) from the Schoodic Lakes, Maine.
 - Description of a new fish from Alaska (Anarrhichas lepturus), with notes upon other species of the genus Anarrhichas.
 - Notes on collection of fishes from eastern Georgia.
 - Description of a new species of Amiurus (A. ponderosus) from the Mississippi river.
 - Descriptions of two species of fishes, collected by Prof. A. Dugès in Central Mexico.
 - Descriptions of some genera and species of Alaskan fishes. See also under Goode and Bean.
- Brewer, T. M. Notes on the nests and eggs of the eight North American species of empidonaces.
- Cooper, J. G. On the migrations and nesting habits of west coast birds.
- Cours, E. Fourth instalment of ornithological bibliography, being a list of faunal publications relating to British birds.
- GARKE, H. On the birds of Heligolaud.
- GOODE, G. B. A study of the trunk fishes (Ostraciontides), with notes upon the American species of the family.
 - A preliminary catalogue of the fishes of the St. John's river and the east coast of Florida, with descriptions of a new genus and three new species.
 - Description of a new species of amber fish (Seriola stearnsii) obtained near Pensacola, Fla., by Mr. Silas Stearns.

- 333. Proceedings of National Museum. Vol. II-Continued.
 - GOODE, G. B.; BEAN, T. H. Description of Alepocephalus bairdii, a new species of fish from the deep-sea fauna of the western Atlantic.
 - Description of a species of Lycodes (L. paxillus), obtained by the United States Fish Commission.
 - Description of a new species of Liparis (L. ranula), obtained by the United States Fish Commission off Halifax, Nova Scotia.
 - Catalogue of a collection of fishes sent from Pensacola, Fla., and vicinity, by Mr. Silas Stearns, with descriptions of six new species.
 - Description of a new genus and species of fish, Lopholatilus chamæleonticeps, from the south of New England.
 - On the occurrence of Lycodes vahlii, Reinhardt, on La Have and Grand Banks.
 - Catalogue of a collection of fishes obtained in the Gulf of Mexico by Dr. J. W. Velie, with description of seven new species.
 - HARGER, O. Notes on New England Isopoda.
 - JORDAN, D. S. Notes on certain typical specimens of American fishes in the British Museum and in the Museum d'Histoire Naturelle, at Paris.
 - Description of new species of North American fishes.
 - Notes on a collection of fishes obtained in the streams of Guanajuato and in Chapala lake, Mexico, by Prof. A. Dugès.
 - KIDDER, J. H. Report of experiments upon the animal heat of fishes, made at Provincetown, Mass., during the summer of 1879, in connection with operations of the United States Fish Commission.
 - Lockington, W. N. Review of the *Pleuronectidæ* of San Francisco.

 Descriptions of new genera and species of fishes from the coast of California.
 - MERRILL, J. C. On the habits of the Rocky Mountain goat.
 - PRATT, R. H. List of names, ages, tribe, &c., of Indian boys and girls at Hampton Normal and Agricultural Institute, Virginia, plaster casts of whose heads were taken by Clark Mills, Esq., March, 1879.
 - SMITH, S. I. Occurrence of *Chelura terebrans*, a crustacean destructive to the timber of submarine structures, on the coast of the United States.
 - Notice of a new species of the Willemæsia group of crustacea (recent Eryontidæ.)
 - VERRILL, A. E. Notice of recent additions to the marine invertebrata of the northeastern coast of America, with descriptions of new genera and species and critical remarks on others.
 - VERRILL, A. E.; RATHBUN, R. List of marine invertebrata from the New England coast, distributed by the United States Commission of Fish and Fisheries.
 - WHITE, C. A. Descriptions of new species of carboniferous invertebrate fossils.
 - Descriptions of new cretaceous invertebrate fossils from Kansas and Texas.
 - Note on Endothyra ornata.
 - Note on Criocardium and Ethmocardium.

- 334. List of described species of Humming Birds. By Daniel Giraud Elliot. 1879. 8vo., pp. 22. (M. C. xvi.)
- 335. List of the Principal Scientific and Literary Institutions in the United States, May, 1879. 1879. 8vo., pp. 6. (M. C. xvi.)
- 336. Smithsonian Miscellaneous Collections. Vol. XVII. 1880. 8vo., pp. 1034.

Documents relative to the origin and history of the Smithsonian Institution. No. 828.

337. Smithsonian Miscellaneous Collections. Vol. XVIII. 1880. 8vo., pp. 851.

CONTENTS.

Journals of the Board of Regents, Reports of Committees, Statistics, etc. No. 329.

- 338. Notes on the Life and Character of Joseph Henry. Read before the Philosophical Society of Washington, October 26, 1878. By James C. Welling. 1880. 8vo., pp. 30, 1 plate. (Portrait.) (M. C. XXL-
- 339. A Memoir of Joseph Henry: a Sketch of his Scientific Work. Read before the Philosophical Society of Washington, October 26, 1878. By William B. Taylor. 1880. 8vo., pp. 225, 1 plate. (Portrait.) (M. C. XXI.)
- 340. Smithsonian Contributions to Knowledge. Vol. XXII. 1880. 4to., pp. 537, 474 woodcuts, 20 plates.

CONTENTS.

Jones, J. Explorations of aboriginal remains of Tennessee. No. 259. HABEL, S. Sculptures of Santa Lucia Cosumaluhuapa, in Guatemaa. No. 269.

RAU, C. Archæological collection of U.S. National Museum. No. 287 RAU, C. Palenque tablet in the U.S. National Museum. No. 381. Dall, W.H. Remains of later pre-historic man in Alaska. No. 318.

341. Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1878. 45th Congress, 3d Session. Senate, Mis. Doc. No. 59. 1879. 8vo., pp. 575, 12 woodcuts.

CONTENTS.

BAIRD, S. F. Secretary's report of operations.

Report of Government explorations and surveys.

Acts and resolutions of Congress relative to the Smithsonian Institution and the National Museum. Forty-fifth Congress, 2d and 3d Ses., 1878, 1879.

BOARD OF REGENTS, Proceedings of

GRAY, A. Biographical memoir of Joseph Henry.

ARAGO, F. Biography of Condorcet.

FAVRE, E. Biographical notice of Louis Agassiz.

341. Report for 1878—Continued.

TAYLOR, W. B. Henry and the telegraph.

LAUTENBACH, B. F. Irritation of a polarized nerve.

Wood, H. C. Researches upon fever.

LECONTE, J. Constants of nature.

List of apparatus available for scientific researches.

OBER, F. A. Ornithology of the Carribee Islands.

KUMLIEN, L. Report of explorations in Greenland.

HENRY, J. Researches in sound.

342. Contributions to the Natural History of Arctic America, made in connection with the Howgate Polar Expedition, 1877-'78. By LUDWIG KUMLIEN. 1879. 8vo., pp. 179. (M. C. XXIII.) Bulletin of the National Museum, No. 15.

CONTENTS.

KUMLIEN, L. Ethnology, mammals, and birds.

BEAN, T. H. Fishes.

VERRILL, A. E. Annelides, molluscoids, and radiates.

DALL, W. H. Mollusks.

INSECTS:

EDWARDS, W. H. Diurnal lepidoptera.

SCUDDER, S. H. and others. Hymenoptera, nocturnal lepidoptera, diptera, coleoptera, neuroptera, and arachnida.

GRAY, A. Plants.

TUCKERMAN, E. Lichens.

FARLOW, W. G. Algæ.

- 343. Annual Reports of the Secretary of the Smithsonian Institution, JOSEPH HENRY, 1865 to 1877. 1880. 8vo., pp. 548.
- 344. Check-list of Publications of the Smithsonian Institution, July, 1879. 1879. 8vo., pp. 16. (M. C. xvi.)
- 345. Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1879. 1880. 46th Congress, 2d Session. Senate Mis. Doc. No. 54. 8vo., pp. 631, 216 woodcuts.

CONTENTS.

BAIRD, S. F. Secretary's report of operations.

BOARD OF REGENTS, Proceedings of

Acts and resolutions of Congress relative to the Smithsonian Institution and the National Museum. 45th Congress, 8d Session; 46th Congress, 2d Session. 1878-1880.

Report of the National Museum Building Commission.

Report of the Architects.

RHEES, W. J. James Smithson and his bequest.

KNIGHT, E. H. A study of the savage weapons at the Centennial Exhibition, Philadelphia, 1876.

ANTHROPOLOGY:

Worsaar, J. J. A. The preservation of antiquities and national monuments in Denmark.

HAVARD, V. The French half-breeds of the Northwest.

845. Report for 1879—Continued.

NORRIS, P. W. Prehistoric remains in Montana, between Fort Ellis and the Yellowstone river.

BRACKETT, A. G. The Shoshone or Snake Indians: their religion. superstitions, and manners.

BURR, R. T. Ruins in White river canon, Pima county, Arizona.

ARMSTRONG, T. Mounds in Winnebago county, Wisconsin.

Anderson, W. G. Mounds near Quincy, Ill., and in Wisconsin.

EVANS, S. B. Notes on some of the principal mounds in Des Moires valley.

DAHLBERG, R. N. and C. Composition of ancient pottery found near the mouth of Chequest creek, at Pittsburg, on the Des Moines river.

BROADHEAD, G. C. Prehistoric evidences in Missouri.

Thompson, T. Mounds in Muscatine county, Iowa, and Rock Island county, Illinois.

TŒLLNER, A. Antiquities of Rock Island county, Illinois.

OEHLER, A. Stone cists near Highland, Madison county, Illinois.

MITCHELL, B. Mounds in Pike county, Illinois.

ADAMS, W. H. Mounds in the Spoon River valley, Illinois.

Quick, E. R. Mounds in Franklin county, Indiana.

JACKMAN, F. Mounds and earthworks of Rush county, Indiana.

Jones, Jr., C. C. Primitive manufacture of spear and arrow points along the line of the Savannah river.

GESNER, W. Mica beds in Alabama.

Hough, J. Mounds in Washington county, Mississippi.

BRODNAX, B. H. Mounds in Moorehouse Parish, Louisiana.

BEAUCHAMP, W. M. Wampum belts of the Six Nations.

Andrews, F. D. Indian relics from Schoharie, New York.

WALKER, S. T. Preliminary explorations among the Indian mounds in Southern Florida.

WALKER, S. T. Report on the shell heaps of Tampa bay, Florida.

NUTTER, F. H. Mounds on Gideon's farm, near Excelsior, Hennepin county, Minnesota.

MASON, O. T. Summary of correspondence of the Smithsonian Institution previous to January 1, 1880, in answer to circular No. 316.

MASON, O. T. Anthropological investigations during the year 1879.

BOEHMER, G. H. Index to papers on anthropology published by the Smithsonian Institution, 1847 to 1878.

Pisko, F. J. On the present fundamental principles of physics.

Von Baumhauer, E. H. A universal meteorograph, designed for detached observatories.

Holden, E. S. Reports of American observatories.

346. Smithsonian Contributions to Knowledge. Vol. XXIII. 1881. 4to., pp. 766, 160 woodcuts, 18 plates of 165 figures.

CONTENTS.

CLARK, H. J. Lucernarise and their allies. No. 242.

HILGARD, E. W. Geology of Lower Louisiana and Petite Anse Island. No. 248.

BARNARD, J. G. Internal structure of the earth. No. 310.

ELLIOT, D. G. Classification and synopsis of the trochilidæ. No. 317.

WOOD, H. C. Fever; a study in physiology. No. 357.

- 347. Nomenclature of Clouds. Two lithographic plates. 1851. 8vo.
- 348. Report on the Fishes of the New Jersey Coast, as observed in the Summer of 1854. By Spencer F. Baird. 1855. 8vo., pp. 40.
- 349. Suggestions for the Sanitary Drainage of Washington City. Toner Lecture No. VIII. Delivered May 26, 1880. By George E. Waring, Jr. June, 1880. 8vo., pp. 26.
- 350. A Map of the Stars near the North Pole; for observations on the Aurora. Copied from the Map used in the Toronto Observations. 1856. 15 inches by 12 inches.
- 351. On the Distribution of the Forests and Trees of North America; with notes on its Physical Geography. By J. G. Cooper. 1859. 8vo., pp. 36, 1 woodcut.
- 352. Brief Abstract of a series of six Lectures on the Principles of Linguistic Science. Delivered at the Smithsonian Institution in March, 1864. By WILLIAM D. WHITNEY. 1864. 8vo., pp. 22.
- 353. Tables and Results of the Precipitation, in Rain and Snow, in the United States, and at some stations in adjacent parts of North America, and in Central and South America. Collected by the Smithsonian Institution, and discussed under the direction of Joseph Henry and Spencer F. Baird, Secretaries. By Charles A. Schott. May, 1881. 4to., pp. 269, 8 woodcuts, 5 plates, 5 charts.
- 354. Essay on the Velocity of Light. By M. Delaunay. Translated for the Smithsonian Institution by Alfred M. Mayer. 1864. 8vo., pp. 31.
- 355. Ozone and Antozone. By CHARLES M. WETHERILL. 1864. 8vo., pp. 12.
- 356. A Memorial of Joseph Henry. (Published by order of Congress.) 1880. 8vo., pp. 532, 1 plate.

INTRODUCTION. Proceedings in Congress relative to public commemoration.

PART I.—OBSEQUIES OF JOSEPH HENRY.

WAITE, M. R. Mortuary announcement.

Proceedings of the Board of Regents.

The funeral, May 16, 1878.

Hodge, C. Prayer at the funeral.

MITCHELL, S. S. Funeral sermon.

PART II.—MEMORIAL EXERCISES AT THE CAPITOL.

Announcement by Executive Committee of the Regents.

McCosh, J. Introductory prayer.

HAMLIN, H. Address.

356.—Memorial of Joseph Henry—Continued.

WITHERS, R. E. Address.

GRAY, A. Address.

CLYMER, H. Reading of telegrams.

ROGERS, W. B. Address.

GARFIELD, J. A. Address.

Cox, S. S. Address.

SHERMAN, W. T. Address.

SUNDERLAND, B. Concluding prayer.

PART III.—MEMORIAL PROCEEDINGS OF SOCIETIES.

Proceedings of the Philosophical Society of Washington.

Proceedings of the Albany Institute.

MEADS, O. Memorial minute.

Proceedings of the United States Light-House Board.

Dod, S. B. Memorial discourse.

CAMERON, H. C. Reminiscences.

WELLING, J. C. Methorial address.

TAYLOR, W. B. Memorial address.

LOVERING, J. Obituary memoir.

NEWCOMB, S. Biographical memoir.

MAYER, A. M. Memorial address.

APPENDIX. Proceedings in Congress regarding the erection of a monument to Joseph Henry.

- 357. Fever: A Study in Morbid and Normal Physiology. By Horatio C. Wood. January, 1878. 8vo., pp. 263, 16 woodcuts, 5 plates of 16 figures. (S. C. xxIII.)
- 358. The Constants of Nature. Part IV. Atomic Weight Determinations: A Digest of the Investigations published since 1814. By George F. Becker. August, 1880. 8vo., pp. 152.
- 359. A Planisphere of the Visible Heavens, extending to 40° of south declination, for observations on Meteoric displays. Prepared by a Committee of the Connecticut Academy of Arts and Sciences. 1864. 30 inches by 24½ inches.
- 360. Palafittes, or Lacustrian Constructions of the Lake of Neuchstel By E. Desor. Translated for the Smithsonian Institution. 1865. 8vo., pp. 53, 118 woodcuts.
- 361. An account of the Aboriginal Inhabitants of the California Peninsula, as given by Jacob Baegert, a German Jesuit Missionary, who lived there seventeen years during the second half of the last century. Translated and arranged for the Smithsonian Institution, by Charles Rau. 1865. 8vo., pp. 41.
- 362. Artificial Shell-deposits in New Jersey. By Charles Rau. 1865. 8vo., pp. 6, 1 woodcut.
- 363. Instructions for collecting Land and Fresh Water Shells. By James Lewis. 1866. 8vo., pp. 8.

- 364. Outline of a Systematic Review of the Class of Birds. By W. LILLJE-BORG. Translated for the Smithsonian Institution. 1866. 8vo., pp. 16.
- 365. Notes on the Tinneh or Chepewyan Indians of British and Russian America. 1.—The Eastern Tinneh; by Bernard R. Ross. 2.—The Loucheux Indians; by William L. Hardisty. 3.—The Kutchin Tribes; by Strachan Jones. Compiled by George Gibbs. 1866. 8vo., pp. 25, 11 woodcuts.
- 366. Directions for Collecting, Preserving, and Transporting Specimens of Diatomacea, and other Microscopic Organisms. By ARTHUR M. EDWARDS. 1867. 8vo., pp. 7.
- 367. Sketch of the Flora of Alaska. By J. T. ROTHROCK. 1867. 8vo., pp. 33.
- 368. Indian Pottery. By Charles Rau. 1867. 8vo., pp. 11, 9 woodcuts.
- 369. Dorpat and Poulkova. By CLEVELAND ABBE. 1867. 8vo., pp. 23.
- 370. A Deposit of Agricultural Flint Implements found in Southern Illinois. By Charles Rau. 1868. 8vo., pp. 9, 5 woodcuts.
- 371. The Metric System of Weights and Measures, with Tables. Prepared for the Smithsonian Institution. By H. A. Newton. 1868. 8vo., pp. 23.
- 372. Drilling in Stone without the use of Metal. By CHARLES RAU. 1869. 8vo., pp, 11, 12 woodcuts.
- 373. Meteorological Stations, and Observers of the Smithsonian Institution, in North America and adjacent Islands, from the year 1849, to the end of the year 1868. 1869. 8vo., pp. 42.
- 374. Three Rain-Charts of the United States, showing the distribution by Isohyetal lines of the mean precipitation in rain and melted snow. 1, for the Summer months; 2, for the Winter months; 3, for the Year. 1870. 20 inches by 14 inches.
- 375. Thoughts on the Nature and Origin of Force. By WILLIAM B. TAYLOR. 1870. 8vo., pp. 19.
- 376. On the Chemistry of the Earth. By T. STERRY HUNT. 1871. 8vo., pp. 26.
- 377. The Diamond and other Precious Stones. By M. Babinet. Translated for the Smithsonian Institution, by John Stearns. 1872. 8vo., pp. 33.
- 378. The Language of the Dakota or Sioux Indians. By F. L. O. RŒHRIG. 1872. 8vo., pp. 19.

- 379. Eulogy on Prof. Alexander Dallas Bache, late Superintendent of the U.S. Coast Survey, President of the National Academy of Sciences, etc. By Joseph Henry. 1872. 8vo., pp. 28.
- 380. The Scientific Education of Mechanics and Artizans. By Andrew P. Peabody. 1873. 8vo., pp. 13.
- 381. Temperature-Chart of the United States, showing the distribution, by Isothermal lines, of the mean temperature for the Year. 1873. 16! inches by 10! inches.
- 382. North American Stone Implements. By Charles Rau. 1873. 8vo., pp. 16, 7 woodcuts.
- 383. Archæological Researches in Nicaragua. By J. F. Bransford. 1881. 4to., pp. 100, 202 woodcuts, 2 plates of 40 figures.
- 384. Circular in reference to Shipping Fresh Fish and other Animals. By S. F. Baird. 1881. 8vo., pp. 4.
- 385. Ancient Aboriginal Trade in North America. By CHARLES RAU. 1873. 8vo., pp. 49.
- 386. Explanation of the Principles of Crystallography and Crystallophysics. By Aristides Brezina. Translated for the Smithsonian Institution, by T. Egleston. 1874. 8vo., pp. 36.
- 387. Three Temperature-Charts of the United States, showing the distribution, by Isothermal curves, of the mean temperature of the lower atmosphere. 1, for the Summer months; 2, for the Winter months: 3, for the Year. 1874. 28½ inches by 19¾ inches.
- 388. Temperature Chart of the United States, showing the distribution, by Isothermal lines, of the mean temperature for the year. 1874. 20 inches by 13½ inches.
- 389. An Account of Investigations relative to Illuminating Materials. (From the Light-House Board Report for 1875.) By JOSEPH HENRY. 1881. 8vo., pp. 25.
- 390. On Tides and Tidal Action in Harbors. By Julius E. Hilgard, 1875. 8vo., pp. 22, 15 woodcuts.
- 391. An act to establish the Smithsonian Institution, etc. Revised Statutes: Title LXXIII. 1875. 8vo., pp. 10.
- 392. The Prehistoric Antiquities of Hungary. An Address delivered by F. F. Romer, at the Opening of the International Anthropological Congress, held at Budapest, September, 1876. Translated for the Smithsonian Institution, by Charles Rau. 1877. 8vo., pp. 9.

- 393. The Mound Builders, and Platycnemism in Michigan. By Henry Gillman. 1874. 8vo., pp. 28, 12 woodcuts. Also, Certain Characteristics pertaining to Ancient Man in Michigan. By Henry Gillman. 1877. 8vo., pp. 13, 13 woodcuts.
- 394. The Stone Age in New Jersey. By C. C. Abbott. 1877. 8vo. pp. 136, 58 plates of woodcuts, containing 223 figures.
- 395. Kinetic Theories of Gravitation. By WILLIAM B. TAYLOR. 1877. 8vo., pp. 80.
- 396. Notes on the History and Climate of New Mexico. By Thomas A. McParlin. 1877. 8vo., pp. 30.
- 397. The Latimer Collection of Antiquities from Porto Rico, in the National Museum at Washington, D. C. By Otis T. Mason. 1877. 8vo., pp. 23. 14 plates of woodcuts, containing 60 figures.
- 398. Short Memoirs on Meteorological Subjects: By Julius Hann; L. Sohncke; Theodore Reye; William Ferrel; A. Colding; and M. Peslin. Compiled and Translated for the Smithsonian Institution by CLEVELAND ABBE. 1878. 8vo., pp. 104.
- 399. Color Blindness in its Relation to Accidents by Rail and Sea. By FRITHIOF HOLMGREN. Translated for the Smithsonian Institution, by M. L. Duncan. 1878. 8vo., pp. 72, 5 woodcuts.
- 400. Aboriginal Structures in Georgia. By Charles C. Jones, Jr. 1878. 8vo., pp. 13, 5 woodcuts.
- 401. On the change of the Mexican Axolotl to an Amblystoma. By AUGUST WEISMANN. Translated for the Smithsonian Institution, by Henry M. Douglass. 1878. 8vo., pp. 29.
- 402. The Stock-in-Trade of an Aboriginal Lapidary. By CHARLES RAU. 1878. 8vo., pp. 9, 16 woodcuts.
- 403. Observations on a Gold Ornament found in a Mound in Florida. By Charles Rau. 1878. 8vo., pp. 6, 1 woodcut.
- 404. On a Polychrome Bead from Florida. By S. S. HALDEMAN. 1878. 8vo., pp. 6, 2 woodcuts.
- 405. An Historical Sketch of Henry's Contribution to the Electro-Magnetic Telegraph; with an account of the origin and development of Prof. Morse's invention. By WILLIAM B. TAYLOR. 1879. 8vo., pp 103.
- 406. Henry on Sound. A Summary of Researches in Sound conducted in the service of the United States Light-House Board, by Joseph Henry, during the years 1865 to 1877. 1879. 8vo., pp. 106, 12 woodcuts.

- 407. Biographical Memoir of Joseph Henry. Prepared in behalf of the Board of Regents. By Asa Gray. 1879. 8vo., pp. 35.
- 408. Report of the Secretary of the Smithsonian Institution, for the year 1878. By S. F. Baird. 1879. 8vo., pp. 60.
- 409. Report of the National Museum Building Commission, and of the Architects, for 1879. 1880. 8vo., pp. 18, 2 folding plates.
- 410. Reports of Astronomical Observatories for 1879. By E. S. HOLDEN. 1880. 8vo., pp. 60.
- 411. The effect of Irritation of a Polarized Nerve. By B. F. LAUTEN-BACH. 1880. 8vo., pp. 59.
- 412. On the Zoological Position of Texas. By Edward D. Cops. 1880. 8vo., pp. 55. Bulletin of the National Museum No. 17.
- 413. Exhibit of the Fisheries and Fish Culture of the United States of America, at the Internationale Fischerei Austsellung, at Berlin, April 20, 1880, and forming a part of the collection of the National Museum, made by the United States Fish Commission. Prepared under the direction of G. Brown Goode. 1880. 8vo., pp. 279. Bulletin of the National Museum, No. 18.
- 414. Base Chart of the United States. Prepared by Charles A. Schott.
 1880. (281 x 191 inches.)
- 415. A Study of the Savage Weapons at the Centennial Exhibition, Philadelphia, 1876. By Edward H. Knight. 1880. 8vo., pp. 90, 147 woodcuts.
- 416. Smithsonian Miscellaneous Collections. Vol. XIX. 1880. 8vo.. pp. 1034.

Proceedings of the U.S. National Museum. Vol. I. 1878. No. 332. Proceedings of the U.S. National Museum. Vol. II. 1879. No. 333.

- 417. Henry as a Discoverer. A Memorial Address, delivered before the American Association for the Advancement of Science, August 26, 1880. By Alfred M. Mayer. 1880. 8vo., pp. 36.
- 418. Report of the Secretary of the Smithsonian Institution, for the year 1879. By Spencer F. Baird. 1880. 8vo., pp. 76.
- 419. Report of the Secretary of the Smithsonian Institution, for the year 1880. By Spencer F. Baird. 1881. 8vo., pp. 88.
- 420. Anthropological Investigations in 1879. By Otts T. Mason. 1881. 8vo., pp. 30.
- 421. Index to Anthropological Articles in Publications of the Smithsonian Institution. By George H. Bœhmer. 1881. 8vo., pp. 10.

- 422. Nomenclature of North American Birds chiefly contained in the United States National Museum. By Robert Ridgway. 1881. 8vo., pp. 94. Bulletin of the National Museum, No. 21.
- 423. Smithsonian Miscellaneous Collections. Vol. XX. 1881. 8vo. pp. 846.

Bulletin of the Philosophical Society of Washington:

Vol. I, March, 1871, to June, 1874, pp. 218.

Vol. II, October 10, 1874, to November 2, 1878, pp. 452.

· Vol. III, November 9, 1878, to June 19, 1880, pp. 169.

424. Smithsonian Miscellaneous Collections. Vol. XXI. 1881. 8vo., pp. 773, 42 woodcuts, 13 plates.

CONTENTS.

RHEES, W. J. James Smithson and his bequest. No. 380.

SMITHSON, J. Scientific writings. No. 827.

Johnson, W. R. Scientific character and researches of Smithson. No. 327.

1RBY, J. R. McD. The works and character of Smithson. No. 327. A memorial of Joseph Henry. No. 856.

425. Proceedings of the United States National Museum for 1880. Vol. III. 1881. 8vo., pp. 595, 5 cuts, 2 plates.

CONTENTS.

- BEAN, T. H. Description of a new hake (*Phycis earlii*), from South Carolina, and a note on the occurrence of *Phycis regius* in North Carolina.
 - Check-list of duplicates of North American fishes distributed by the Smithsonian Institution in behalf of the United States National Museum, 1877-'80.
- CATTIE, S. T. On the genitalia of male eels and their sexual characters. ENDLICH, F. M. List of species and varieties of minerals in the National Museum of the United States in 1879.
- GARMAN, S. Synopsis and descriptions of the American Rhinobatidæ.
- GILL, T. On the identity of the genus Leurynnis Lockington, with Lycodopsis Collett.
- GOODE, G. B. Descriptions of seven new species of fishes from deep soundings on the Southern New England coast, with diagnoses of two undescribed genera of flounders and a genus related to Merlucius.
 - Fishes from the deep waters on the South coast of New England obtained by the United States Fish Commission in the summer of 1880.
 - The frigate mackerel (Auxis Rochei) on the New England coast.
 - Notacanthus phasganorus, a new species of Notacanthidæ from the Grand Banks of Newfoundland.
- HAY, O. P. On a collection of fishes from Eastern Mississippi.

425. Proceedings of National Museum. Vol. III-Continued.

HEILPRIN, A. On some new species of eocene mollusca from the Southern United States.

JORDAN, D. S. Notes on a collection of fishes from East Florida, obtained by J. A. Henshall.

Notes on a collection of fishes from St. John's river, Florida, obtained by A. H. Curtiss.

Note on a forgotten paper of Dr. Ayres, and its bearing on the numenclature of the cyprinoid fishes of the San Francisco markets.

Note on "Sema" and "Dacentrus."

Description of a new species of Caranx (Caranx Beani), from Beaufor'.

North Carolina.

JORDAN, D. S.; GILBERT, C. H. Notes on a collection of fishes from San Diego, California.

Description of new flounder (Xystreurys liolepis), from Santa Catalina Island, California.

Description of a new ray (Platyrhina triseriata), from the coast of California.

Description of a new species of "rock cod" (Sebastichthys serricepent from the coast of California.

On the occurrence of Cephaloscyllium laticeps (Duméril) Gill on the coast of California.

On the oil shark of Southern California (Galeorhinus galeus.)

Description of a new flounder (*Pleuronichthys verticalis*), from the coast of California, with notes on other species.

Notes on sharks from the coast of California.

On the generic relations of Platyrhina exasperata.

Description of a new species of Schastichthys (Schastichthys miniatus . from Monterey bay, California.

Description of a new species of "rock fish" (Sebastichthys carnatus. from the coast of California.

Description of a new species of ray (Raia stellulata), from Monterey. California.

Description of a new species of Xiphister and Apodichthys, from Monterey, California.

Description of two new species of Sebastichthys (Sebastichthys entomelas and Sebastichthys rhodochloris), from Monterey bay, Cal.

Description of a new agonoid fish. (Brachiopsis xyosternus), from Monterey bay, California.

Description of a new flounder (Hippoglossoides exilis), from the coset of California.

Description of a new species of ray (Raia rhina) from the coast of California.

Description of two new species of fishes (Ascelicthys rhodorus and Scytalina cerdale), from Neah Bay, Washington Territory.

Description of two new species of scopeloid fishes (Sudis ringers and Myctophum crenulare), from Santa Barbara channel, California.

Description of two new species of flounders (Parophrys ischyurus and Hippoglossoides elassodon), from Puget Sound.

Description of seven new species of sebastoid fishes, from the coast of California.

425. Proceedings of National Museum. Vol. III-Continued.

JORDAN, D. S.; GILBERT, C. H. Description of a new embiotocoid (Abeona aurora), from Monterey, California, with notes on a related species.

Description of a new flounder (*Platysomatichthys stomias*), from the coast of California.

Description of a new embiotocoid fish (Cymatogaster rosaceus), from the coast of California.

Description of a new species of deep-water fish (Icichthys Lockingtoni), from the coast of California.

Description of a new embiotocoid fish (Ditrema atripes), from the coast of California.

Description of a new scorpsenoid fish (Sebastichthys maliger), from the coast of California.

Description of a new scorpænoid fish (Sebastichthys proriger), from Monterey Bay, California.

Description of a new agonoid (Agonus vulsus), from the coast of California.

Description of a new species of Hemirhamphus (Hemirhamphus Rosæ), from the coast of California.

Description of a new species of notidanoid shark (Hexanchus corinus), from the Pacific coast of the United States.

Description of a new species of Nemichthys (Nemichthys avocetta), from Puget Sound.

Description of a new species of Paralepis (Paralepis coruscans), from the Straits of Juan de Fuca.

List of the fishes of the Pacific coast of the United States, with a table showing the distribution of the species.

On the generic relations of Belone exilis (Girard).

Notes on a collection of fishes from Utah Lake.

Description of a new species of rockfish (Sebastichthys chrysomelas), from the coast of California.

LAWRENCE, G. N. Description of a new species of bird of the family Turdidæ, from the island of Dominica, W. I.

Description of a new species of parrot of the genus Chrysotis, from the island of Dominica.

Description of a new species of Icterus, from the West Indies.

Lockington, W. N. Remarks on the species of the genus Chirus found in San Francisco market, including one hitherto undescribed.

Description of a new fish from Alaska (Uranidea microstoma).

Description of a new species of Agonides (Brachyopsis verrucosus), from the coast of California.

Description of a new genus and some new species of California fishes (Icosteus ænigmaticus and Osmerus attenuatus).

Description of a new chiroid fish (Myriolepis zonifer), from Monterey Bay, California.

Description of a new sparoid fish (Sparus brachysomus), from Lower California.

Note on a new flat fish (Lepidopsetta isolepis), found in the markets of San Francisco.

Description of a new species of Prionotus (Prionotus stephanophrys), from the coast of California.

RATHBUN, R. The littoral marine fauna of Provincetown, Cape Cod, Massachusetts.

- 495. Proceedings of National Museum. Vol. III-Continued.
 - RIDGWAY, R. Revisions of nomenclature of certain North American birds.
 - A catalogue of the birds of North America.
 - Catalogue of Trochilidæ in the collection of the United States National Museum.
 - RYDER, J. A. On Camaraphysema, a new type of sponge.
 - List of the North American species of myriapods belonging to the family of the Lysiopetalidæ, with a description of a blind form from Luray Cave, Virginia.
 - SMITH, R. On the occurrence of a species of *Cremnobates* at San Diego. California.
 - SMITH, S. I. Preliminary notice of the crustacea dredged in 64 to 325 fathoms off the south coast of New England by the United States Fish Commission, in 1880.
 - SWAN, J. G. The surf smelt of the northwest coast, and the method of taking them by the Quillehute Indians, West Coast of Washington Territory.
 - The eulachon or candle fish of the northwest coast.
 - VERRILL, A. E. Notice of recent additions to the marine invertebrata of the northeastern coast of America, with descriptions of new genera and species, and critical remarks on others.
 - Part II.—Mollusca, with notes on Annelida, Echinodermata, etc... collected by the United States Fish Commission.
 - Part III.—Catalogue of mollusca recently added to the fauna of Southern New England.
 - WHITE, U. A. Note on the occurrence of *Productus giganteus* in Cal. Note on Acrothele.
 - Description of a new cretaceous Pinna from New Mexico.
 - Note on the occurrence of Stricklandia salteri and S. davidsoni in Georgia.
 - Description of a very large fossil gasteropod, from the State of Puebla.

 Mexico.
 - Descriptions of new invertebrate fossils from the mesozoic and cenozoic rocks of Arkansas, Wyoming, Colorado, and Utah.
- 426. A Synopsis of the Scientific Writings of Sir William Herschel. By Edward S. Holden and Charles S. Hastings. 1881. 8vo., pp. 118.
- 427. Record of Recent Progress in Science, 1879 and 1880. Astronomy. By Edward S. Holden. 1881. 8vo., pp. 39.
- 428. Record of Recent Progress in Science, 1879 and 1880. Geology and Mineralogy. By George W. Hawes. 1881. 8vo., pp. 30.
- 429. Record of Recent Progress in Science, 1879 and 1880. Physics and Chemistry. By George F. Barker. 1881. 8vo., pp. 65.
- 430. Record of Recent Progress in Science, 1879 and 1880. Botany. By WM. G. FARLOW. 1881. 8vo., pp. 19.

- 431. Record of Recent Progress in Science, 1879 and 1880. Zoology. By Theodore Gill. 1881. 8vo., pp. 62.
- 432. Record of Recent Progress in Science, 1880. Anthropology. By Otis T. Mason. 1881. 8vo., pp. 51.
- 433. Report of a visit to the Luray Cavern, in Page County, Virginia, under the auspices of the Smithsonian Institution, July 13 and 14, 1880. 1881. 8vo., pp. 12, 6 woodcuts.
- 434. Report of the National Museum Building Commission and of the Architects for 1880. 1881. 8vo., pp. 12.
- 435. Discussion of the Barometric Observations of E. S. Snell. By F. H. Loud. 1881. 8vo., pp. 23, 7 woodcuts.
- 436. List of Periodicals received by the Smithsonian Institution. 1881. 8vo., pp. 9.
- 437. Check-list of Publications of the Smithsonian Institution to December, 1881. 1881. 8vo., pp. 22.
- 438. Reports of Astronomical Observatories for 1880. By E. S. Holden and George H. Bæhmer. 1881. 8vo., pp. 128.
- 439. Warming and Ventilating Occupied Buildings. By ARTHUR MORIN. Translated for the Smithsonian Institution by Clarence B. Young. 1882. 8vo., pp. 92, 32 woodcuts.
- 440. Articles on Anthropological Subjects contributed to the Annual Reports of the Smithsonian Institution from 1863 to 1877. By Charles Rau. 1882. 8vo., pp. 180, 53 woodcuts.

RAU, C., Preface by. (February, 1882.)

BAEGERT, J. Account of the aboriginal inhabitants of the Californian peninsula. Translated and arranged for the Smithsonian by Chas. Rau. From Smithsonian Report for 1868-64. No. 861.

Agricultural implements of the North American stone period. From Report for 1863.

Artificial shell deposits in New Jersey. From Report for 1864. No. 862. Indian pottery. From Report for 1866. No. 868.

Drilling in stone without metal. From Report for 1868. No. 872.

Deposit of agricultural flint implements in Southern Illinois. From Report for 1868. No. 870.

Memoir of C. F. P. Von Martius. From Report for 1869. No. 251.

Ancient aboriginal trade in North America. From Report for 1872. No. 385.

North American stone implements. From Report for 1872. No. 382.

ROMER, F. F. Prehistoric antiquities of Hungary. Translated by Chas. Rau. From Report for 1876. No. 392.

Stock in trade of an aboriginal lapidary. From Report for 1877. No. 402. Observations on a gold ornament from a mound in Florida. From Report for 1877. No. 403.

- 441. The Constants of Nature. Part V. A Recalculation of the Atomic Weights. By Frank Wigglesworth Clarke. 1882. 8vo., pp. 293.
- 442. Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1880. Forty-sixth Congress, 3d Session, Senate Mis. Doc. No. 31. 1881. 8vo., pp. 782, 13 woodcuts.

BAIRD, S. F. Secretary's report of operations.

List of periodicals received by the Institution.

Rules for the examination of specimens submitted to the Institution.

TAYLOR, F. W. Report of the chemist.

Additions to the collections of the National Museum.

Additions to Museum by the Berlin International Fisheries Exhibition.

Stevenson, J. Report of explorations in New Mexico and Arizons. Receipts and distribution of specimens.

GOODE, G. B. The first decade of the U.S. Fish Commission.

Report of the Executive Committee on the Henry statue.

Report of the Executive Committee for 1880.

BOARD OF REGENTS, Proceedings of

Report of the National Museum Building Commission for 1880.

Report of the Superintending Architects of the Museum Building.

RECORD OF RECENT SCIENTIFIC PROGRESS:

BAIRD, S. F. Introduction.

HOLDEN, E. S. Astronomy.

HAWES, G. W. Geology.

BARKER, G. F. Physics.

BARKER, G. F. Chemistry.

HAWES, G. W. Mineralogy.

FARLOW, W. G. Botany.

GILL, T. Zoology.

MASON, O. T. Anthropology.

MASON, O. T. Bibliography of anthropology.

Mason, O. T. Abstracts of anthropological correspondence.

MASON, O. T. Report on Luray Cavern.

Loud, F. H. Discussion of Prof. Snell's barometric observations.

HENRY, J. Investigation of illuminating materials.

Holden, E. S.; Hastings, C. S. Synopsis of the scientific writings of William Herschel.

HOLDEN, E. S.; BŒHMER, G. H. Reports of astronomical observatories.

- 443. Results of Meteorological Observations made at Providence, Rhode Island, extending over a period of forty-five years, from December, 1831, to December, 1876. By ALEXIS CASWELL. 1882. 4to, pp. 40.
- 444. Guide to the Flora of Washington and Vicinity. By LESTER F. WARD. 1881. 8vo., pp. 265. One map. Bulletin of the National Museum, No. 22.

- 445. Plan of Organization and Regulations of the United States National Museum. By G. Brown Goode. 1882. 8vo., pp. 58. Two woodcuts. (Circular of U. S. National Museum, No. 1.)
- 446. Circular addressed to Friends of the Museum. By Spencer F. Baird. 1882. 8vo., pp. 2. (Circular of U. S. National Museum, No. 2.)
- 447. Circular in reference to Petroleum Collections. By Spencer F. Baird. 1882. 8vo., pp. 4. (Circular of U.S. National Museum, No. 3.)
- 448. Circular concerning the Department of Insects. By Spencer F. Baird. 1882. 8vo., pp. 2. (Circular of U. S. National Museum, No. 4.)
- 449. Establishment and Officers of the Smithsonian Institution and National Museum, January 1, 1882. 1882. 8vo., pp. 2. (Circular of U. S. National Museum, No. 5.)
- 450. Classification and Arrangement of the Materia Medica Collection. By James M. Flint. 1882. 8vo., pp. 2. (Circular of U.S. National Museum, No. 6.)
- 451. A Classification of the Forms in which Drugs and Medicines appear, and are administered. By James M. Flint. 1882. 8vo., pp. 8. (Circular of U. S. National Museum, No. 7.)
- 452. Memoranda for Collectors of Drugs for the Materia Medica Section of the National Museum. By James M. Flint. 1882. 8vo., pp. 2. (Circular of U. S. National Museum, No. 8.)
- 453. Circular in reference to the Building-Stone Collection. By Spencer F. Baird. 1882. 8vo., pp. 6. (Circular of U. S. National Museum, No. 9.)
- 454. Two letters on the work of the National Museum. By BARNET PHILLIPS. 1882. 8vo., pp. 10. (Circular of U.S. National Museum, No. 10.)
- 455. A Provisional Classification of the Food Collections. By G. Brown Goode. 1882. 8vo., pp. 18. (Circular of U. S. National Museum, No. 11.)
- 456. Classification of the Collections of the U. S. National Museum to illustrate the Art of Taxidermy. By W. T. Hornaday. 1882. 8vo., pp. 2. (Circular of U. S. National Museum, No. 12.)
- 457. Outline of a Scheme of Classification for the Collections in the United States National Museum. By G. Brown Goode. 1882. 8vo., pp. 4. (Circular of U. S. National Museum, No. 13.)

- 458. Circular requesting material for the Library of the United States National Museum. By Spencer F. Baird. 1882. 8vo., pp. 4. (Circular of U. S. National Museum, No. 14.)
- 459. The Organization and Objects of the United States National Museum. By G. Brown Goode. 1882. 8vo., pp. 4. (Circular of U. S. National Museum, No. 15.)
- 460. Directions for Collecting and Preserving Plants. By LESTER F. WARD. 1882. 8vo., pp. 32.
- 461. Check-List. Flora of Washington and Vicinity. By LESTER F. WARD. 1882. 8vo., pp. 62.
- 462. Catalogue of Old World Birds in the U.S. National Museum. By ROBERT RIDGWAY. 1882. 8vo., pp. 20.
- 463. Bibliography of the Fishes of the Pacific Coast of the United States. to the end of 1879. By Theodore Gill. 1882. 8vo., pp. 77. (M. C. XXIII.) Bulletin of National Museum, No. 11.
- 464. Directions for Collecting and Preserving Fish. By Tarleton H. Bean. 1881. 8vo., pp. 6.
- 465. List of Marine Invertebrates, mainly from the New England Coast, distributed by the U.S. National Museum. Series II. By RICHARD RATHBUN. 1881. 8vo., pp. 6.
- 466. Directory of Officers, Collaborators, Employés, etc., of the Smithsonian Institution, National Museum, Geological Survey, Bureau of Ethnology, and Fish Commission. 1882. 8vo., pp. 8.
- 467. Proceedings of the United States National Museum for 1881. Vol. IV. 1882. 8vo., pp. 600, 13 cuts, 2 plates. (M. C. xxII.)

- BAIRD, S. F. Notes on certain aboriginal shell mounds on the coast of New Brunswick and of New England.
- BEAN, T. H. Descriptions of new species of fishes (Uranidea marginata.

 Potamocottus Bendirei) and of Myctophum crenulare J. and G.

Notes on some fishes from Hudson's Bay.

Descriptions of new fishes from Alaska and Siberia.

Directions for collecting and preserving fish.

A preliminary catalogue of the fishes of Alaskan and adjacent waters.

A partial bibliography of the fishes of the Pacific coast of the United States and of Alaska, for the year 1880.

Notes on a collection of fishes made by Capt. Henry E. Nichols. U.S. N., in British Columbia and Southern Alaska, with descriptions of new species and a new genus (*Delolepis*).

See also under Goode and Bean.

BENDIRE, C. Notes on Salmonidae of the upper Columbia.

467. Proceedings of National Museum. Vol. IV—Continued.

BOYD, C. H. Remains of the walrus (?) in Maine.

CARLIN, W. E. Observations of Siredon lichenoides.

DALL, W. H. On the genera of chitons.

On certain limpets and chitons from the deep waters off the eastern coast of the United States.

ENDLICH, F. M. An analysis of water destructive to fish in the Gulf of Mexico.

FARLOW, W. G. Report on the contents of two bottles of water from the Gulf of Mexico, forwarded by the Smithsonian Institution.

GILBERT, C. H. See under Jordan and Gilbert.

GILL, T. Note on the Latiloid genera.

GLAZIER, W. C. W. On the destruction of fish by polluted waters in the Gulf of Mexico.

GOODE, G. B. The Taxonomic relations and geographical distribution of the members of the swordfish family, (Xiphiidæ.)

GOODE, G. B.; BEAN, T. H. Description of a new species of fish, Apogon pandionis, from the deep water off the mouth of Chesapeake Bay.

Benthodesmus, a new genus of deep-sea fishes, allied to Lepidopus.

HAWES, G. W. On the mineralogical composition of the normal meso-zoic diabase upon the Atlantic border.

On the determination of feldspar in thin sections of rocks.

INGERSOLL, E. On the fish mortality in the Gulf of Mexico.

JAPANESE LEGATION, Washington. Catalogue of a collection of Japanese cotton fibre presented to the U.S. National Museum by the Government of Japan, together with the amount of the annual crop of Japan and the price of cotton.

JOHNSON, S. H. Notes on the mortality among fishes of the Gulf of Mexico.

JORDAN, D. S.; JOUY, P. L. Check-list of duplicates of fishes from the Pacific coast of North America, distributed by the Smithsonian Institution in behalf of the United States National Museum, 1881.

JORDAN, D. S.; GILBERT, C. H. Notes on the fishes of the Pacific coast of the United States.

Description of Sebastichthys mystinus.

Description of a new species of Ptychochilus (Ptychochilus Harfordi) from Sacramento river.

Note on Raia inornata.

Notes on a collection of fishes, made by Lieut. Henry E. Nichols, U. S. N., on the west coast of Mexico, with descriptions of new species.

List of fishes collected by Lieut. Henry E. Nichols, U. S. N., in the Gulf of California and on the west coast of Lower California, with descriptions of four new species.

Descriptions of thirty-three new species of fishes from Mazatlan, Mexico.

Description of a new species of *Pomadasys* from Mazatlan, with a key to the species known to inhabit the Pacific coasts of tropical America.

467. Proceedings of National Museum. Vol. IV—Continued.

Description of a new species of Xenichthys (Xenichthys ocyurus) from the west coast of Central America.

Descriptions of five new species of fishes from Mazatlan, Mexico.

Jour, P. L. Description of a new species of Squalius (Squalius alicia) from Utah Lake.

LAWRENCE, G. N. Description of a new sub-species of Loxigilla from the island of St. Christopher, West Indies.

Lockington, W. N. Description of a new genus and species of Cottida.

LUGGER, O. The occurrence of the Canada porcupine in Maryland.

McKAY, C. L. A review of the genera and species of the family Contrarchidoe, with a description of one new species.

MOORE, M. A. Fish mortality in the Gulf of Mexico.

Pirz, A. Methods of making and preserving plaster casts.

PLATEAU, F. The rapid preparation of large myological specimens.

PORTER, J. Y. On the destruction of fish by poisonous water in the Gulf of Mexico.

RATHBUN, R. List of marine invertebrates, mainly from the New England coast, distributed by the United States National Museum. (Series II.)

List of Marine invertebrates from the New England coast, distributed by the United States National Museum. (Series III.)

RAU, C. List of anthropological publications by

RIDGWAY, R. On a duck new to the North American fauna.

On Amazilia Yucatanensis, (Cabot.) and A. cerviniventris, Gould.

A review of the genus Centurus, Swainson.

List of species of Middle and South American birds not contained in the United States National Museum.

List of special desiderata among North American birds.

Catalogue of Old World birds in the United States National Museum.

Notes on some Costa Rican birds.

Description of a new fly-catcher, and a supposed new petrel, from the Sandwich Islands.

Description of a new owl from Porto Rico.

Descriptions of two new thrushes from the United States.

On two recent additions to the North American bird fauna, by L. Belding.

RYDER, J. A. On Semper's method of making dry preparations.

SHUFELDT, R. W. Remarks upon the osteology of Opheosaurus ventralis.

SMITH, R. Description of a new Gobioid fish, (Othonops cos.) from San Diego, California.

Description of a new species of Gobiesox, (Gobiesox rhessodon,) from San Diego, California.

SMITH, S. B. On the Chinook names of the salmon in the Columbia River.

STEJNEGER, L. Description of two new races of Myadestes obscurus, Lafr.
TRANSLATION. Metallic castings of delicate natural objects.

TRUE, F. W. On the North American land tortoises of the genus Xero-bates.

On the rare rodent Cricetodipus parvus (Baird) Coues.

- 467. Proceedings of National Museum. Vol. IV-Continued.
 - WARD, L. F. Catalogue of a collection of Japanese woods presented to the U.S. National Museum by the University of Tokio, Japan.
 - WHITE, C. A. On certain cretaceous fossils from Arkansas and Colorado. Anonymous. The comparative action of dry heat and sulphurous acid upon putrefactive bacteria.

APPENDICES—Circulars of the U.S. National Museum:

- No. 1. Plan of organization and regulations of the Museum.
- No. 2. Circular addressed to the friends of the Museum.
- No. 3. Circular in reference to petroleum collections.
- No. 4. Circular concerning the department of insects.
- No. 5. Establishment and officers of the Smithsonian Institution and Museum.
- No. 6. Classification and arrangement of the materia medica collection.
- No. 7. A classification of the forms in which drugs and medicines appear and are administered.
- No. 8. Memoranda of collectors of drugs for the materia medica section of the National Museum.
- No. 9. Circular in reference to the building-stone collection.
- No. 10. Two letters on the work of the National Museum.
- No. 11. A provisional classification of the food collections.
- No. 12. Classification of the collections to illustrate the art of taxidermy.
- No. 13. Outline of a scheme of Museum classification
- No. 14. Circular requesting material for the library of the Museum.
- No. 15. The organization and objects of the National Museum.
- No. 16. Plans for the installation of collections.
- No. 17. Contributions and their acknowledgment.
- No. 18. List of publications of the United States National Museum.
- 468. Smithsonian Miscellaneous Collections. Vol. XXII. 1882. 8vo., pp. 1200, cuts 18, plates 4.

CONTENTS.

Proceedings of the U.S. National Museum. Vol. III. 1880. No. 425. Proceedings of the U.S. National Museum. Vol. IV. 1881. No. 467.

- 469. List of Foreign Correspondents of the Smithsonian Institution. Corrected to January, 1882. 1882. 8vo., pp. 174.
- 470. Nomenclator Zoologicus. An Alphabetical List of all Generic Names that have been employed by Naturalists for Recent and Fossil Animals from the earliest times to the close of the year 1879. In two parts. Part I.—List of Generic Names employed in Zoology and Paleontology to the close of the year 1879, chiefly supplemental to those catalogued by Agassiz and Marschall, or indexed in the Zoological Record. By Samuel H. Scudder. 1882. 8vo., pp. 398. Bulletin of the National Museum, No. 19.
- 471. List of Marine Invertebrates from the New England Coast, distributed by the U.S. National Museum. Series III.—Educational Series. By Richard Rathbun. 1881. 8vo., pp. 4.
- 472. Plans for the Installation of Collections in the U.S. National Museum. By G. Brown Goode. 1882. 8vo., pp. 2. (Circular of U.S. National Museum, No. 16.)

- 473. Contributions to the United States National Museum, and their Acknowledgement. 1882. 8vo., pp. 2. (Circular of U.S. National Museum, No. 17.)
- 474. List of Publications of the United States National Museum. 1882. 8vo., pp. 12. (Circular of U. S. National Museum, No. 18.)
- 475. Smithsonian Miscellaneous Collections. Vol. XXIII. 1882. 8vo., pp. 1003. Bulletins of the National Museum, Nos. 11 to 15.

- GILL, T. Bibliography of fishes of Pacific coast. Bulletin No. 11. No. 468.
- JORDAN, D. S.; BRATTON, A. W. Fishes of Alleghany region of South Carolina, Georgia, and Tennessee, and synopsis of Catostomidæ. Bulletin No. 12. No. 808.
- EGGERS, H. F. A. Flora of St. Croix and the Virgin Islands. Bulletin No. 18. No. 318.
- GOODE, G. B. Catalogue of collection illustrating animal resources and fisheries of the United States. Bulletin No. 14. No. 326.
- Kumlien, L. Natural history of Arctic America. Bulletin No. 15. No. 342.
- 476. First Annual Report of the Bureau of Ethnology to the Secretary of the Smithsonian Institution, 1879-80. By J. W. Powell, Director. 1881. Royal 8vo., pp. 638. 343 cuts, 54 plates, 1 folded map.

CONTENTS.

POWELL, J. W. Report of the Director of the Bureau.

Powell, J. W. On the evolution of language.

POWELL, J. W. Mythology of the North American Indians.

POWELL, J. W. Wyandot government.

Powell, J. W. Limitations to the use of some Anthropological data.

YARROW, H. C. Study of the mortuary customs of the North American Indians.

HOLDEN, E. S. Studies in Central American picture-writing.

ROYCE, C. C. Cessions of land by Indian tribes to the United States.

MALLERY, GARRICK. Sign language among North American Indians, compared with that among other peoples and deaf-mutes.

PILLING, J. C. Catalogue of linguistic manuscripts in the library of the Bureau of Ethnology.


Dorsey, J. O. Illustration of the method of recording Indian languages. How the rabbit caught the sun in a trap. (An Omaha myth.)

GATSCHET, A. S. Details of an Indian conjurer's practice. (In the Klamath Lake dialect.)

GATSCHET, A. S. The relapse. (In the Klamath Lake dialect.)

Riggs, S. R. . A dog's revenge. (A Dakota Fable.)

- 477. Report on International Exchange, with list of Official Publications of the U.S. Government, between 1868 and 1881. By George H. Bæmmer. 1882. 8vo., pp. 113.
- 478. Catalogue of Publications of the Smithsonian Institution up to July 1, 1882, with Index to all Articles in the "Smithsonian Contributions to Knowledge," "Miscellaneous Collections," "Annual Reports," "Bulletins and Proceedings of the U. S. National Museum," etc. By William J. Rhees. 1882. 8vo., 1882, pp.



CLASSIFIED LIST OF SEPARATE PUBLICATIONS

OF THE

SMITHSONIAN INSTITUTION.

SUBJECTS.

- I. ANATOMY, PHYSIOLOGY, MEDICINE, and SURGERY.
- II. ANTHROPOLOGY. (See also Philology.)
- III. ARCHÆOLOGY. (See Anthropology.)
- IV. ARCHITECTURE.
 - V. ASTRONOMY.
- VI. BIBLIOGRAPHY.
- VII. BIOGRAPHY.
- VIII. BIOLOGY. (See also Anatomy, Botany, Microscopy, Palsontology, and Zoology.)
 - IX. BOTANY.
 - X. CHEMISTRY and TECHNOLOGY.
 - XI. ELECTRICITY and MAGNETISM.
 - XII. ETHNOLOGY. (See Anthropology, also Philology.)
- XIII. GEOLOGY. (See also Palsontology.)
- XIV. MAGNETISM. (See Electricity, etc.)
 - XV. MATHEMATICS.
- XVI. MEDICINE and SURGERY. (See Anatomy, etc.)
- XVII. METEOROLOGY.
- XVIII. MICROSCOPY.
 - XIX. MINERALOGY.
 - XX. MISCELLANEOUS.
 - XXI. NATURAL HISTORY. (See Biology.)
- XXII. PALÆONTOLOGY.
- XXIII. PHILOLOGY.
- XXIV. PHYSICAL GEOGRAPHY.
 - XXV. Physics. (See also Terrestrial Physics.)
- XXVI. Physiology. (See Anatomy, etc.)
- XXVII. TECHNOLOGY. (See Chemistry, etc.)
- XXVIII. TERRESTRIAL PHYSICS.
 - XXIX. Zoology.—Birds.

Fishes.

Insects.

Mammals.

Mollusks.

Radiates.

Reptiles, and Batrachians.

Shells. (See Mollusks.)

I.—ANATOMY, PHYSIOLOGY, MEDICINE, AND SURGERY.

	No. Catalo
ADAMS. Subcutaneous surgery (Toner Lecture, No. 6)	
Brown-Séquard. Dual character of the brain (Toner Lecture, No. 2)	
DACOSTA. The heart (Toner Lecture, No. 8)	
DEAN. Medulla oblongata (s. c.)	
FLINT. Classification of drugs and medicines	
FLINT. Classification of materia medica collection	
FLINT. Memoranda for collection of drugs	
Holmgren. Color-blindness	
Jones. Chemical and physiological investigations (s. c.)	
KEEN. Continued fevers (Toner Lecture, No. 5)	
LAUTENBACH. Effect of irritation of a polarized nerve	
LEIDY. Flora and fauna within living animals (s. c.)	
MITCHELL and Morehouse. Chelonia (s. c.)	
MITCHELL. Venom of rattlesnake (s. c.)	
SHAKESPEARE. Reparatory inflammation (Toner Lecture, No. 7)	
WARING. Sanitary drainage of Washington (Toner Lecture, No. 8)	
WEISMANN. Change of Mexican axolotl	
Wood. Morbid and normal physiology of fever (s. c.)	
Wood. Study of fever (Toner Lecture, No. 4)	
WOODWARD. Cancerous tumors (Toner Lecture, No. 1)	
WYMAN. Rana pipiens (s. c.)	
	
II.—ANTHROPOLOGY. (See also Philology.)	
ABBOTT. Stone age in New Jersey	
Archæology, circular on2	
BEGERT. Aboriginal inhabitants of California	
BEHMER. Index to archeological articles in Smithsonian publications	
Bransford. Archæological researches in Nicaragua (s. c.)	
DALL. Prehistoric man, Aleutian Islands (s. c.)	
DESOR. Palafittes of the Lake of Neuchatel	
GIBBS. Ethnological and philological instructions	
GIBBS. Tinneh, or Chepewyan Indians	
GILLMAN. Characteristics of ancient man in Michigan	

HABEL. Sculptures in Guatemala (s. c.)

HALDEMAN. Polychrome bead from Florida.....

269

404

ANTHROPOLOGY—Continued.	Cata
HARDISTY. Loucheux Indians	
Haven. American archæology (s. c.)	
Indians, photographic portraits of, list of	
Jones. Aboriginal structures in Georgia	
ONES. Antiquities of Tennessee (s. c.)	
ones. Kutchin Indians	
Knight. Savage weapons	
LAPHAM. Antiquities of Wisconsin (s. c.)	
Mason. Anthropological investigations in 1879	
Mason, " 1880	
Mason. Latimer antiquities from Porto Rico	
MASON. Recent progress in anthropology	
MAYER. Mexican archæology (s. c.)	
Morgan. Circular relative to relationship	
Morgan. Systems of consanguinity (s. c.)	
Pickering. Gliddon mummy case (s. c.)	
Powell. First annual report of Bureau of Ethnology	
RAU. Ancient aboriginal trade in North America	
RAU. Archæological collection in Smithsonian Institution (s. c.)	••
RAU. Articles on anthropological subjects, (from Reports S. I.)	
Rau. Artificial shell deposits in New Jersey	·
RAU. Deposit of agricultural flint implements	
RAU. Drilling in stone without metal	
RAU. Gold ornament in mound in Florida	
RAU. Indian Pottery	
RAU. North American stone implements	
Rau. Palenque tablet (s. c.)	
RAU. Stock-in-trade of aboriginal lapidary	
ROMER. Prehistoric antiquities of Hungary (translated by RAU)	
Ross. Eastern Tinneh Indians	
Squire and Davis. Ancient monument of Mississippi valley (s. c.)	
SQUIER and DAVIS. Aboriginal monuments of New York (s. c.)	
STANLEY. Catalogue of Indian Portraits	
Swan. Indians of Cape Flattery (s. c.)	
Swan. Haidah Indians (s. c.)	
WHITTLESEY. Ancient mining at Lake Superior (s. c.)	
WHITTLESEY Ancient works in Ohio (s. C.)	

IV.—ARCHITECTURE.

Cata	dogue.
Building-stones, circular relative to	453
OWEN. Hints on public architecture	P
Report of National Museum building commission and architects for 1879	409
Report of National Museum building commission and architects for 1880	434

V.—ASTRONOMY.

ABBE. Obser	rvatories of Dorpat and Poulkova	
Alexander.	Harmonies of the solar system (s. c.)	
Astronomical	telegram circular	
Downes. Oc	ccultations 1848 (s. c.)	
Downes.	" 1849 (s. c.)	
Downes.	" 1850 (s. c.)	
Downes.	" 1851 (s. c.)	
Downes.	" 1852 (s. c.)	
Downes.	" 1858 (s. c)	
Draper. Sil	lvered-glass telescope (s. c.)	
Eclipse, annu	lar, 1854	
GILLISS. Sol	lar eclipse, Peru (s. c.)	
Gould. Hist	tory of Neptune	
Gould. Trai	nsatlantic longitude (s. c.)	
HILL. Map	solar eclipse	
HOLDEN and	BCHMER. Reports of observatories for 1880	
Holden and	HASTINGS. Synopsis of Sir William Herschel's writings	B
Holden. In	dex catalogue of Nebulæ	
Holden. Re	ecent progress in astronomy	·
Holden. Re	ports of observatories for 1879	
KANE. Astro	onomical observations arctic seas (s. c.)	129
NEWCOMB. I	Integrals of planetary motion (s. c.)	
Newcomb. (Orbit of Neptune (s. c.)	
Newcomb. (Orbit of Uranus (s. c.)	
RUNKLE. Pl	anetary tables (s. c.)	
RUNKLE. Pla	anetary tables, supplement	
Star-map, circ	eumpolar	
Star-map, of t	he visible heavens	
STOCKWELL.	Orbits of planets (s. c.)	
WALKER. RO	esearches Neptune (s. c.)	

	ASTRONOMY—Continued.	No. in Catalogue.
WALKER.	Ephemeris Neptune, 1848 (s. c.)	
WALKER.		
WALKER.	. 1850 (s. c.)	6
WALKER.		
WALKER.		
		
	VI.—BIBLIOGRAPHY.	
BINNEY.	Bibliography of conchology, part I	142
BINNEY.	46 66 II	174
Exchanges,	s, list of Smithsonian, part I	 73
Exchanges,	s, " II	
Exchanges,	•	
GILL. Bil	ibliography of fishes of Pacific coast	 463
GIRARD.	Bibliography of natural history, 1851	48
Holden an	nd Hastings. Synopsis of Sir William Herschel's writ	tings 426
Holden.	Index catalogue of books and memoirs on Nebulæ	311
Institutions	ns, principal literary and scientific in United States	335
JEWETT.	Catalogue system	47
JEWETT.	Public libraries	25
Publication	ns, Smithsonian Institution, list of (successive issues,)	74, 203, 226,
	245, 278, 290, 30	1, 344, 437, 478
Periodicals,	s, received by the Institution, list of	Q, 436
Publication	ns of learned societies in library Smithsonian Inst	73, 85, 179
RHEES. C	Datalogue and index of Smithsonian publications	479
RHEES. P	Public libraries	116, 238
WATSON.	Bibliography of North American botany	258
	VII.—BIOGRAPHY.	•
_		
	emoir of Joseph Henry	
	Eulogy on Alex. Dallas Bache	
• •	emorial of (By numerous contributors.)	
	nd Hastings. Scientific writings of Sir William Hersel	
	Henry as a discoverer	
	moir of Von Martius	
	Life and bequest of James Smithson	
	Memoir of and scientific work of Joseph Henry	
Welling.	Life and character of Joseph Henry	

HARVEY.

HARVEY.

"

VIII.—BIOLOGY.	(See also Anatomy, Microscopy, Palmontology, I	t eatoe
	and Zoology.)	

and Zoology.)
No. in Catalog: -
BAIRD. Circular for shipping fresh-fish and other animals
Circular for collecting birds
Circular for collecting craw-fish
Circular for collecting grasshoppers 163
Circular for collecting shells176, 363
Circular for observations in Russian America
Circular to officers of the Hudson Bay Co137
Circular to entomologists178
COPE. Zoology of Texas 412
Directions for collecting and preserving specimens 34
Directions for collecting and preserving specimens of nests and eggs 139
EDWARDS. Directions for collecting diatomacea 366
GILL. Recent progress in zoology 431
GIRARD. Bibliography of natural history 48
Goode. Animal resources and fisheries exhibit, 1876 326
Goode. Classification of animal resources 297
KIDDER and others. Natural history of Kerguelen Island 294
Kumlien. Natural history of Arctic America 342
LEIDY. Flora and fauna within animals (s. c.) 44
Museum miscellanea, numbers, labels, etc164
PACKARD. Directions for collecting insects
Proceedings of the U.S. National Museum I, 332; II, 333; III, 425; IV, 467
Periodical phenomena register65, 148
Scudder. List of genera of animals
STIMPSON. Marine Invertebrata, Grand Manan, (s. c.) 50
STREETS. Natural history of Hawaiian, Fanning Islands, etc 303
WEISMANN. Change of Mexican axolotl 401
· · · · · · · · · · · · · · · · · · ·
IX.—BOTANY.
Change Thomas and Anna of Manch Assessed
Cooper. Forests and trees of North America
EGGERS. Flora of St. Croix
FARLOW. Recent progress in botany 430
GRAY. Plantæ Wrightianæ, I (s. c.)
GRAY. " II (8. C.)
HARVEY. North American marine algae, I (s. c.)

II (8. C.)....

III (s. c.)____

95

OF THE SMITHSONIAN INSTITUTION.

BOTANY—Continued	No. in Catalogue.
HARVEY. North American marine algee, complete (s	
LEIDY. Flora within animals (s. c.)	-
Periodical phenomena, register of	
ROTHBOCK. Flora of Alaska	
Torrey. Batis maritima (s. c.)	
Torrey. Darlingtonia Californica (s. c.)	
HARVEY. Plantæ Frémontianæ (s. c.)	
WARD. Check list of flora of Washington	
WARD. Directions for collecting and preserving pla	
WARD. Guide to the flora of Washington	•
WATSON. Index to North American botany	
Wood. Fresh water algæ (s. c.)	
WOOD. I Took water algo (6. C.)	
X.—CHEMISTRY AND TECH	NOLOGY.
BARKER. Recent progress in chemistry	429
BECKER. Atomic weight determinations	358
BOOTH and MORFIT. Chemical arts	
CLARKE. Expansion tables	
CLARKE. Recalculation of atomic weights	
CLARKE. Specific gravity tables	
CLARKE. " " " 1st supplement	
CLARKE. Specific heat tables	
GIBBS and GENTH. Ammonia-cobalt bases (s. c.)	
HARE. Explosiveness of nitre (s. c.)	
HENRY. Investigation of illuminating materials	
HUNT. Chemistry of the earth	
Jones. Chemical and physiological investigations (s	
MOBIN. Warming and ventilating occupied building	-
NEWTON. Metric system of weights and measures	
SMITHSON. Scientific writings	
TAYLOR. Henry and the telegraph	
WETHERILL. Ozone and antozone	355
XI.—ELECTRICITY AND MAC	ANTON
BACHE. Magnetic discussion, part I (s. c.)	
	121
	132
7	

ELECTRICITY AND MAGNETISM—Continued. No. in Catalogue.
BACHE. Magnetic discussion, parts IV to VI (s. c.) 162
BACHE. " parts VII to IX (s. c.) 175
BACHE. " parts X to XII (s. c.) 186
BACHE. " parts I to XII, complete (s. c.) 195
BACHE. Magnetic survey of Pennsylvania (s. c.) 166
Gould. Transatlantic longitude by electro-magnetic signals (s. c.) 223
HARKNESS. Magnetic observations on the iron-clad Monadnock (s. c.) 239
HAYES. Arctic observations (s. c.) 196
HENRY. Communication to Regents on the electro-magnetic telegraph 115
KANE. Arctic observations (s. c.)97, 198
Locke. Observations on terrestrial magnetism (s. c.) 35
SECCHI. Researches on electrical rheometry (s. c.) 36
Sonntag. Terrestrial magnetism, Mexico (s. c.)
TAYLOR. Henry and the telegraph 405
XII.—ETHNOLOGY. (See Anthropology and Philology.) ———————————————————————————————————
Cooper. Physical geography of North America
HAWES. Recent progress in geology
HILGARD. Geology of lower Louisiana (s. c.)248
HITCHCOCK. Surface geology (s. c.)
MASON. Luray Cavern in Virginia
Pumpelly. Geology of China, Mongolia, and Japan (s. c.) 202
WHITTLESEY. Fluctuations of lakes (s. c.) 119
WHITTLESEY. Glacial drift (s. c.) 197
XIV.—MAGNETISM. (See Electricity.)
XV.—MATHEMATICS.
ALVORD. Tangencies of circles (s. c.)
FERREL. Converging series (s. c.) 233
NEWCOMB. General integrals of planetary motion (s. c.) 281

XVI.—MEDICINE. (See Anatomy, etc.)

XVII.—METEOROLOGY.

Ca	No. in talogue
ABBE. Meteorological memoirs	
BACHE. Girard College observations (s. c.) 113, 121, 132, 162, 175, 186	•
Caswell. Observations, Providence, Rhode Island, 1881-1860 (s. c.)	103
CASWELL. Observations, Providence, Rhode Island, 1831-1876 (s. c.)	443
CHAPPELSMITH. Tornado (s. c.)	. 59
Clouds, nomenclature of	347
Coffin. Meteoric fire-ball (s. c.)	221
Coffin. Psychrcmetrical tables	87
COFFIN. Winds, northern hemisphere (s. c.)	52
COFFIN. Winds of the globe (s. c.)	268
Colding. The nature of air currents	398
COLDING. The whirlwind of St. Thomas in 1871	398
COOPER, J. G. Forests and trees of North America	351
Directions for meteorological observations1), 148
FERREL. Reply to strictures of J. Hann	398
Force. Record of auroras (s. c.)	84
Guyor. Meteorological directions19), 148
Guyor. Meteorological and physical tables3:	
HANN. Diminution of vapor with altitude	
HANN. Influence of rain on the barometer	
HANN. Atmospheric pressure and rain-fall	
HANN. Laws of temperature in ascending currents	
HANN. Relation between pressure and wind	
HAYES. Arctic observations (s. c.)	
Henry. Circular, altitudes	
HENRY. " directions for constructing lightning rods	
HENRY. "thunder-storms	
Henry. " tornadoes	
HILDRETH AND WOOD. Observations, Marietta, Ohio (s. c.)	120
Hork. Letter on meteoric shower	
Loud. Discussion of Snell's barometric observations	435
Loomis. Storms in 1886 (s. c.)	127
Map of the stars near the north pole for aurora	350
McClintock. Arctic observations (s. c.)	146 206
	396
MEECH. Intensity sun's heat (s. c.).	83
CLEAVELAND. Meteorological observations, Brunswick, Maine (s. c.)	
KANE. Meteorological arctic observations (s. c.)104	
Meteorological observations, 1855	93.

METEOROLOGY—Continued.	no. Catal	. IN Ogue.
Meteorological results, 1854-59, vol. 1		157
Meteorological results, 1854-59, vol. 2		182
Meteorological stations and observers 1849-1868		373
Meteors, November		217
OLMSTEAD. Aurora (s. c.)		81
Periodical phenomena, registry of	65,	148
PESLIN. Relation between barometric pressure and air currents		398
Planisphere of the visible heavens, for observation of meteors		359
Rain charts for summer, winter, and year		374
REYE. Rainfall and barometric minima		398
SCHOTT. Base chart of the United States		414
SCHOTT. Tables of rain and snow in the United States (s. c.)	222,	353
Schott. Tables of temperature (s. c.)		
SMITH. Observations, Washington, Arkansas (s. c.)		131
Sounces. Variation of temperature in ascending currents of moist air_		398
Temperature chart of U.S. for year	•	
Temperature charts of the U.S. for summer, winter, and year	-	387
BAILEY. Microscopic observations, Southern States (s. c.)		23
Bailey. New microscopic organisms (s. c.)		63
BAILEY. Soundings, coast survey (8. C.)		20
DEAN. Medulla oblongata (S. C.)		173
EDWARDS. Directions for collecting microscopic organisms		366
LEIDY. Flora and fauna within living animals (s. c.)		44
WOODWARD. Cancerous tumors		26 6
		
XIX.—MINERALOGY.		
BABINET. Diamond and precious stones		377
Brezina. Crystallography and crystallophysics		38 6
EGLESTON. Catalogue of minerals		1.56
HAWES. Recent progress in mineralogy		428
		
XX.—MISCELLANEOUS.		
Relative to the Smithsonian Institution.		
Act of Congress to establish the Smithsonian Institution	 0,	3 91
BORHMER. Report on international exchanges	-	477

		MISC	ELLANEOUS	—Cont	inued.	No. 1n C atal ogu
Busines	e arrange	ments of the S	Smithsonian Ins	stitutio	n	•
Commit	itee on O	rganization, re	port of			В,
Correspo	ondence 8	Equier and Day	vis			1
Circular	relative	to exchanges.				32
Circular	r relative	to Smithsonia	n Institution			29
DALLAS	. Addre	ss, laying cor	ner-stone of Sm	ithsonia	an Institution	
Digest o	of act of	Congress relati	ive to Smithsoni	ian Ins	titution	
Director	y of offic	ers, collaborat	ors and employe	68	### +	- 4 6
Establis	hment an	d officers of S	mithsonian Inst	titution	and Museum	44
HENRY.	Exposi	tion, Smithson	n's bequest			
_	-	•	-			
_						
	•				64, 154,	
		_				
					1	
		ganization				
_		_				26
BAIRD.		-	•		va	
BAIRD.						
AIRD.	_				1, 1876	
ircular	of the U	Inited States N	Vational Museum	•	1	
"	44		44	66	2	
44	46	44	44	11	8	
* 6	4.6	4.6	74	"	4	44
• 6	66	66	4.	"	5	44
4.6	£6 ·	4.6	44	"	6	
"	44	66	4.6	"	7	45
"	if	44	44	"	8	
44	"	44	46	46	9	45
16	"	44	44	46	10	45
• 6	44	46	4.6	"	11	45
44	44	4.6	44	66	12	45
14	44	44	66	"	18	45
"	"	4.6	4.6	"	14	45
44	46	• "	66		15	
16	16	48	44		16	

MISCELLANEOUS—Continued.), in logue,
Circular of the United States National Museum, No. 17	473
18	474
Circular requesting material for Museum library	458
Classification of animal resources	297
Contributions to Museum, and acknowledgement	473
GOODE. Animal resources and fisheries	326
Goode. Classification of Museum collections	457
GOODE. Classification of the food collections	455
Goode. Installation of collections in Museum	472
Goode. Organization and objects of the Museum	459
Goode. Organization and regulations of the Museum	445
HORNADAY. Classification of the collections in taxidermy	456
Hudson's Bay circular	137
PHILLIPS. Letters on the work of the National Museum	454
Publications of National Museum, list of	474
XXII.—PALÆONTOLOGY. Conbad. Check list of eocene fossils	200
GIBBES. Mosasaurus (s. c.)	14
LEIDY. Ancient fauna, Nebraska (s. c.)	36
LEIDY. Cretaceous reptiles (s. c.)	192
	192
LEIDY. Cretaceous reptiles (s. c.)	192 72
LEIDY. Cretaceous reptiles (s. c.) LEIDY. Extinct sloths (s. c.)	192 72 41
LEIDY. Cretaceous reptiles (s. c.) LEIDY. Extinct sloths (s. c.) LEIDY. Fossil ox (s. c.)	192 72 41 177
LEIDY. Cretaceous reptiles (s. c.) LEIDY. Extinct sloths (s. c.) LEIDY. Fossil ox (s. c.) MEEK. Check list of cretaceous and jurassic fossils	192 72 41 177 183
LEIDY. Cretaceous reptiles (s. c.) LEIDY. Extinct sloths (s. c.) LEIDY. Fossil ox (s. c.) MEEK. Check list of cretaceous and jurassic fossils MEEK. Check list of miocene fossils MEEK and HAYDEN. Palseontology of Nebraska (s. c.) XXIII.—PHILOLOGY.	192 72 41 177 183 172
LEIDY. Cretaceous reptiles (s. c.) LEIDY. Extinct sloths (s. c.) LEIDY. Fossil ox (s. c.) MEEK. Check list of cretaceous and jurassic fossils MEEK. Check list of miocene fossils MEEK and HAYDEN. Palsontology of Nebraska (s. c.) XXIII.—PHILOLOGY. Bowen. Yoruba grammar and dictionary (s. c.)	192 72 41 177 183 172
LEIDY. Cretaceous reptiles (s. c.) LEIDY. Extinct sloths (s. c.) LEIDY. Fossil ox (s. c.) MEEK. Check list of cretaceous and jurassic fossils MEEK. Check list of miocene fossils MEEK and HAYDEN. Palseontology of Nebraska (s. c.) XXIII.—PHILOLOGY. BOWEN. Yoruba grammar and dictionary (s. c.) GIBBS. Chinook jargon	192 72 41 177 183 172 98 161
LEIDY. Cretaceous reptiles (s. c.) LEIDY. Extinct sloths (s. c.) LEIDY. Fossil ox (s. c.) MEEK. Check list of cretaceous and jurassic fossils MEEK. Check list of miocene fossils MEEK and HAYDEN. Palsontology of Nebraska (s. c.) XXIII.—PHILOLOGY. BOWEN. Yoruba grammar and dictionary (s. c.) GIBBS. Chinook jargon GIBBS. Comparative Vocabulary	192 72 41 177 183 172 98 161 170
LEIDY. Cretaceous reptiles (s. c.) LEIDY. Extinct sloths (s. c.) LEIDY. Fossil ox (s. c.) MEEK. Check list of cretaceous and jurassic fossils MEEK. Check list of miocene fossils MEEK and HAYDEN. Palseontology of Nebraska (s. c.) XXIII.—PHILOLOGY. BOWEN. Yoruba grammar and dictionary (s. c.) GIBBS. Chinook jargon	192 72 41 177 183 172 98 161

OF THE SMITHSONIAN INSTITUTION.

PHILOLOGY—Continued.	. in logue
Morgan. Systems of consanguinity (s. c.)	_
Riggs. Dakota grammar and dictionary (s. c.)	4
REHRIG. Language of the Dakota or Sioux Indians	37
WHITNEY. Lectures on Linguistics	35
XXIV.—PHYSICAL GEOGRAPHY.	
Cooper. Physical geography of North America	35
ELLET. Physical geography of the Mississippi valley (s. c.)	13
HAYES. Observations in the Arctic seas (s. c.)	19
HENRY. Circular relative to heights	23
Mason. Luray Cavern in Virginia	
XXV.—PHYSICS. (See also Terrestrial Physics.)	
BARKER. Recent progress in physics	42
BARNARD. Internal structure of the earth (s. c.)	31
BARNARD. Problems of rotary motion (s. c.)	24
Brezina. Crystallography and crystallophysics	38
Coffin. Winds of the globe (s. c.)52,	26
DAVIS. Law of deposit of flood-tide (s. c.)	3
DELAUNAY. Essay on velocity of light	35
DRAPER. Telescope in photography (s. c.)	18
Gould. Transatlantic longitude (s. c.)	
Guyor. Physical and meteorological tables31,	
HAYES. Physical observations, Arctic seas (8. C.)	
HENRY. Electro-magnetic telegraph	
HENRY. Investigation of illuminating materials	
Henry. Lightning rods	
Henry. Researches on sound.	
HILGARD. Tides and tidal action	39
Holmgren. Color-blindness	39
	37
KANE. Physical observations, Arctic seas (S. C.)	
MERCH. Heat and light of the sun (s. c.)	8
Newton. Metric weights and measures	37
TAYLOR. Henry and the telegraph	40
TAYLOR. Kinetic theories of gravitation	39
TATION Noture and origin of force	37

PHYSICS—Continued.

No. in Catalogue

SCHOTT. Base chart of the United States	41
SECCHI. Electrical rheometry (s. c.)	3
WHITTLESEY. Fluctuations, level of lakes (s. c.)	
XXVI.—PHYSIOLOGY. (See Anatomy, etc.)	
XXVII.—TECHNOLOGY. (See CHEMISTRY, etc.)	
XXVIII.—TERRESTRIAL PHYSICS.	
BACHE. Discussion of magnetic observations (s. c.)I, 113; II, 121; III,	132
IV-VI, 162; VII-IX, 175; X-XII, 186; I-XII	-
Bache. Magnetic survey of Pennsylvania (s. c.)	16
Barnard. Problems of rotary motion, and precession of equinoxes (s. c.)	
COFFIN. Winds of the globe (s. c.)	26
COFFIN. Winds of the northern hemisphere (s. c.)	5
DAVIS. Dynamical action, etc., of the flood tide (s. c.)	3
Force. Record of auroral phenomena (s. c.)	
HAYES. Physical observations in Arctic seas (s. c.)	
HENRY. Circular relative to altitudes	
HILGARD. Tides and tidal action in harbors	390
HUNT. Chemistry of the earth	37
	91
KANE. Physical observations in the Arctic seas.(8. C.)	196
KANE. Tidal observations in the Arctic seas (s. c.)	
LOCKE. Observations on terrestrial magnetism in United States (s. c.)	35
MÜLLER. Observations on terrestrial magnetism in Mexico (s. c.)	114
	81
Sonntag. Observations on terrestrial magnetism in Mexico (s. c.)	114
WHITTLESEY. Fluctuations of level in the North American lakes (s. c.)	
•	
XXIX.—ZOOLOGY.	
. Birds.	
BAIRD. Arrangement of birds	210
BAIRD. Catalogue of birds, 4to	106
BAIRD. " 8vo	108

ZOOLOGY—Continued. No. No. 1	o. in Llogue.
BAIRD. Instructions for collecting bird's nests and eggs	_
BAIRD. Review of birds, part I	181
Brewer. North American oology, part I (s. c.)	89
Circular, birds of South America	168
Desiderata of birds of Mexico, Central America, etc.	185
ELLIOT. List of Trochilids, (humming birds)	334
ELLIOT. Monograph of Trochilids (s. c.)	317
KIDDER and Cours. Birds of Kerguelen Island	293
LAWRENCE. Birds of Mexico	295
LILLJEBORG. Outline of classification of birds	364
RIDGWAY. Catalogue of old world birds in Museum	462
RIDGWAY. Nomenclature of North American birds	422
Fishes.	
BAIRD. Circular for shipping fresh fish	384
BAIRD. Circular relative to food fishes	
BAIRD. Fishes of New Jersey coast	348
BAIRD. Inquiry relative to food-fishes	231
BEAN. Directions for collecting and preserving fish	464
GILL. Arrangement of fishes	247
GILL. Bibliography of fishes of Pacific coast	463
GILL. Catalogue of fishes	283
GILL. Fishes of Western North America.	307
GIRARD. Monograph, cottoids (s. c.)	30
Goode. Berlin exhibit of fisheries and fish culture	413
Goode. Fishes of Bermuda	296
JORDAN and BRAYTON. Fishes of South Carolina, Georgia, and Tennessee	308
JORDAN. Notes on cottide	306
JORDAN. Notes on Rafinesque's memoirs, North American fishes	305
Insects.	
AGASSIZ. Classification of insects (8. C.)	16
Circular to entomologists	178
Circular concerning department of insects	448
DE SAUSSURE. Monograph of wasps, hymenoptera	254
Hagen. Synopsis of neuropteta	134
LE CONTE. Classification of coleoptera, part I	136
LE CONTE. " part II	265
Le Conte. Kancas and New Mexico coleoptera (s. c.)	126

	ZOOLOGY—Continued.	alo		
LE CONTE.	List of coleoptera			
LE CONTE.	New species of coleoptera, part I			
LE CONTE.	ee part II	. :		
LOEW. Moi	ograph of diptera, part I			
JOEW.	" part II			
JOEW.	" part III	•		
Aelsneime	R. Catalogue of coleoptera			
Morris. C	talogue of lepidoptera	•		
dorris. Sy	nopsis of lepidoptera.	•		
. •	EN. Monograph of diptera, part IV			
	EN. Catalogue of diptera10			
	Directions for collecting insects			
	Catalogue of orthoptera			
	,			
	Mammals.			
	126////406.			
	onograph, North American bats			
BAIRD. Catalogue, mammals				
	of families of mammals			
lit. List	of families of mammals	•		
FILL, List	Mollusks. ibliography North American conchology, part I			
BINNET. B	Mollusks. ibliography North American conchology, part I	•		
BINNEY. B BINNEY. L	Mollusks. ibliography North American conchology, part I " " " II and and fresh-water shells, part I. Pulmonata geophila	•		
BINNEY. BINNEY. LINNEY. LINNEY.	Mollusks. ibliography North American conchology, part I			
BINNEY. BINNEY. LINNEY. LINNEY. LINNEY.	Mollusks. ibliography North American conchology, part I			
SINNEY. BINNEY. LINNEY. LINNEY. SINNEY. SINNEY.	Mollusks. ibliography North American conchology, part I			
SINNEY. BINNEY. LINNEY. LINNEY. SINNEY. SINNEY. CARPENTER.	Mollusks. ibliography North American conchology, part I " " " II and and fresh-water shells, part I. Pulmonata geophila " " " II. Pulmonata Limnophila " " " III. Ampullariidæ, etc Mollusks of Western North America Chitonidæ			
SINNEY. BINNEY. LINNEY. LINNEY. LINNEY. CARPENTER. CARPENTER.	Mollusks. ibliography North American conchology, part I			
SINNEY. BINNEY. LINNEY. LINNEY. LINNEY. CARPENTER. CARPENTER.	Mollusks. ibliography North American conchology, part I " " " II and and fresh-water shells, part I. Pulmonata geophila " " " II. Pulmonata Limnophila " " " III. Ampullariidæ, etc Mollusks of Western North America Chitonidæ			
SINNEY. BINNEY. LINNEY. LINNEY. LINNEY. CARPENTER. CARPENTER. CARPENTER. CARPENTER. Check list of	Mollusks. ibliography North American conchology, part I			
BINNEY. BINNEY. BINNEY. BINNEY. BINNEY. CARPENTER. CARPENTER. CARPENTER. Check list of	Mollusks. ibliography North American conchology, part I			
SINNEY. SINNEY. SINNEY. SINNEY. SINNEY. CARPENTER. CARPENTER. Check list of Circular for Call. Index	Mollusks. ibliography North American conchology, part I			
SINNEY. SINNEY. SINNEY. SINNEY. SINNEY. CARPENTER. CARPENTER. CARPENTER. Check list of Circular for Call. Index	Mollusks. ibliography North American conchology, part I			
SINNEY. BINNEY. L. SINNEY. L. SINNEY. L. SINNEY. CARPENTER. CARPENTER. CARPENTER. Check list of Circular for Call. Index Exploring explo	Mollusks. ibliography North American conchology, part I			
SINNEY. SINNEY. SINNEY. SINNEY. SINNEY. CARPENTER. CARPENTER. CARPENTER. Check list of Circular for Carploring ex Supporting ex S	Mollusks. ibliography North American conchology, part I			
SINNEY. BINNEY. L. SINNEY. L. SINNEY. L. SINNEY. CARPENTER. CARPENTER. Check list of Circular for Call. Indexploring exploring	Mollusks. ibliography North American conchology, part I			

OF THE SMITHSONIAN INSTITUTION.

	ZOOLOGY-	-Continue		lo, in alogue.	
	Rad	iates.			
BAIRD. Circular fo	r collecting cray-fish .			319	
CLARK. Lucernaria	æ (s. c.)		**	242	
			, series II	465	
RATHBUN. "	"	46	" III	471	
STIMPSON. Marine	invertebrata of Grand	d Manan (s	s. c.)	50	
	Rep	tiles.		•	
BAIRD and GIRARD.	Catalogue of serper	nts		49	
BAIRD. Circular or	BAIRD. Circular on collecting reptiles				
Cope. North American batrachia and reptilia					
MITCHELL and MOE	EHOUSE. Chelonia (s. C.)		159	
•	Shells. (Se	e Mollusks.	.)		

SMITHSUNIAN CONTRIBUTIONS TO KNOWLEDGE. (Quarto.)

Vol.	DATE.	Pages.	Wood- cuts.	PLATES.	Maps.	No. in Series.	CONTAINS NOS. OF CATALOGUE.
I	1848	860	207	48		2	1.
. II	1851	572	.89	24		26	8, 12, 20, 18, 14, 16, 17, 23, 15,
III	1852	562		85		88	4, 5, 6, 7, 11. 85, 86, 30, 82, 22, 83, 37, 24, 29.
IV	1852	426				89	4 0.
v	1858	588	4	45		55	44, 41, 45, 43, 42.
VI	1854	484	9	53		56	46, 60, 61, 50, 52, 58, 54.
VII	1855	260	74	72	2	76	59, 63, 70, 72, 78.
VIII	1856	564	52	9		78	71, 81, 80, 82, 84, 85.
IX	1857	480	45	22		92	88, 90, 86, 88, 79, 9 4 .
X	1858	462	1	16	1	99	95, 97, 98.
XI	1859	502	20	28		111	89, 100, 118, 104, 126, 114, 127.
XII	1860	58 8	15	8		112	129, 119, 103, 131, 185.
XIII	1868	558	80	4	8	151	180, 146, 155, 121, 132, 162, 166.
XIV	1865	490	158	25		184	159. 175, 186, 180, 172, 192.
X∇	1867	620	47	18	4	206	199, 197, 202, 196.
XVI	1870	498 .	76	18		211	173, 204, 120, 208, 221, 223, 220.
XVII	1871	616	6	14		229	218.
IIIVX	1872	6 4 6	10	5	8,	24 6	222, 282, 289, 233.
XIX	1874	640	6	21		272	240, 241, 262.
XX	1876	79 4	4	26		284	268.
XXI	1876	5 4 8	80	9	8	285	280, 281, 267, 277.
XXII	1880	587	474	20		840	259, 269, 287, 331, 318.
XXIII	1881	766	160	18		346	242, 248, 310, 317, 357.

SMITHSONIAN MISCELLANEOUS COLLECTIONS. (Octavo.)

Vol	DATE.	Pages.	Coxs.	PLATES.	No. in Series.	CONTAINS NOS. OF CATALOGUE.
I	1862	788	28		122	148, 87, 158.
II	1862	714	88		128	27, 115, 58, 108, 49, 128, 84, 187, 189,
ш	1862	776	49		124	163, 176, 188. 102, 118, 186, 117.
IV	1862	762	80		125	18 4 , 188.
V	1864	774			158	142, 74, 15 4 .
VI	1867	888	15	7	169	141, 171, 140, 167.
VII	1867	878	818		191	165, 148, 144, 201, 145, 200, 188, 177, 156, 161, 160, 208.
VIII	1869	921	780	4	212	219, 189, 194, 210, 187, 207, 205, 178, 168, 164.
IX	1869	914			218	174, 179.
X	1878	918	5		250	252, 227, 235, 236, 287, 190, 284, 288, 248, 245.
XI	1874	790	55	4	278	280, 247, 256, 261, 264, 265.
XII	1874	767	86		274	181, 255, 268.
XIII	1878	982	45		812	292, 298, 294, 295, 296, 297, 808, 804, 305, 806.
XIV	1878	911		4	814	254, 283, 288, 276, 289, 216, 801, 811.
XV	1878	880	58		815	258, 266, 291, 279, 282, 800, 802, 809, 816, 819, 820.
XVI	1880	950	871	7	822	258, 270, 321, 824, 825, 884, 885, 844.
XVII	1880	1084			886	828.
XVIII	1880	851			887	329.
XIX	1880	1084			416	882, 888.
XX	1881	8 4 6			428	
XXI	1881	778	42	18	424	880, 827, 856.
XXII	1882	1200	18	4	4 68	425, 467.
IXIII	1882	1008			475	463, 808, 818, 826, 842.

ANNUAL REPORTS OF THE SMITHSONIAN INSTITUTION. (Octavo.)

	Pages.	Woodcuts.	No. in Smithsonian Catalogue.	When Published
1st, for 1846	88		G	1847
2d, for 1847	208		H	1848
8d, for 1848	64		I	1849
4th, for 1849	272		21	1850
5th, for 1850	826		28	1851
6th, for 1851	10 4		51	1852
7th, for 1852	96		57	1853
8th, for 1858	810		67	1854
9th, for 1854	464	4	75	1855
10th, for 1855	440	79	77	1856
11th, for 1856	468	69	91	1857
12th, for 1857	438	100	107	1858
18th, for 1858	448	48	109	1859
14th, for 1859	4 50	57	110	1860
15th, for 1860	448	78	147	1861
l6th, for 1861	464	25	149	1862
17th, for 1862	446	94	150	1868
18th, for 1868	420	56	187	1864
19th, for 1864	450	50	188	1865
20th, for 1865	496	189	209	1866
21st, for 1866	470	70	214	1867
22d, for 1867	506	10	215	1868
28d, for 1868	474	40	224	1869
24th, for 1869	480	88	. 228	1871
25th, for 1870	494	28	244	1871
26th, for 1871	478	8	249	1873
27th, for 1872	456	119	271	1878
28th, for 1878	452	88	275	1874
29th, for 1874	416	46	286	1875
80th, for 1875	422	854	298	1876
Blst, for 1876	488	78	299	1877
32d, for 1877	500	49	828	1878
88d, for 1878	575	12	841	1879
34th, for 1879	681	216	845	1880
35th, for 1880	782	18	442	1881

PUBLICATIONS OF THE U. S. NATIONAL MUSEUM. (Octavo.)

No. of Museum Series.		TITL	€.		In Vol. of Miscellaneous Collections.	No. in Smithsonian Catalogue.
1	Bulletin of th	ne U. S. Nati	onal Muse	ım, No. 1	XIII	292
2	"	"	16	No. 2	XIII	298
8	44	"	46	No. 8	XIII	294
4	46	16	4.6	No. 4	XIII	295
5	44	"	44	No. 5	XIII	296
6	• 66	16	"	No. 6	XIII	297
7	66	"	1.6	No. 7	XIII	808
8	66	"	46	No. 8	XIII	804
9	66	4.6	"	No. 9	XIII	805
10		"	66	No. 10	XIII	806
11	66	• "	8.6	N o. 11	XXIII	463
12	86	44	66	No. 12	XXIII	808
13	44	. 46	44	No. 18	XXIII	818
14	44	46	66	No. 14	XXIII	826
15	44	"	66	No. 15	XXIII	842
16	44	"	4.6	No. 16	<i>j</i> ,	
17	Proceedings	Nat. Mus.,	Vol. I		XIX	882
18	Bulletins Na	t. Mus., Vol	. I, Nos. 1	to 10	XIII	812
19	Proceedings	Nat. Mus.,	V ol. II		XIX	883
20	Bulletin of th	ne U.S. Nati	ional Muser	um, No. 17		412
21	"	"	4.6	No. 18		418
22	"	"	44	No. 19		470
28	"	"	46	No. 20		
24	"	"	44	No. 21		422
25	Proceedings	Nat. Mus.,	Vol. III		XXII	425
26		ne U.S. Nati				444
27	Proceedings	Nat. Mus.,	Vol. IV		XXII	467
28	Bulletins Na	•			XXIII	4 75

PUBLICATIONS OF THE U.S. NATIONAL MUSEUM—Continued. SEPARATE LIST OF BULLETINS.

No.	When Published.	Pages.	Cuts.	Plates.	Maps.	In Vol. of Miscellaneous Collections.	No. in Museum Series.	No. in Smithsonism Catalogue.
1	1875	108				XIII.	1	292
2	1875	61				XIII	2	293
8	1876	124				XIII	8	294
4	1875	56				XIII	4	295
5	1876	84				XIII	5	296
6	1876	140				XIII	6	297
7	1877	172				XIII	7	908
8	1877	88				XIII	8	304
9	1877	56				XIII	9	305
10	1877	124		45		XIII	10	306
11	1882	77				ıfıxx	11	463
12	1878	287				XXIII	12	308
18	1879	189				XXIII	13	813
14	, 1879	867				XXIII	14	326
15	1879	179				IIIXX	15	842
16							16	
17	1880	55					20	412
18	·1880	279					21	418
19	1882	898					22	470
20							28	
21	1881	94					24	423
22	1881	265			1		.26	444

THE BULLETINS COLLECTED IN VOLUMES.

Vol.	Contents.	Published.	Pages.	Outs.	Plates.	In Vol. of Miscellaneous Collections.	No. in Museum Series.	No. in Smithsonian Catalogue.
I	Nos. 1-10	1878	982		45	XIII	18	812
II	Nos. 11-15	1882	1008	****		XXIII	28	475

PUBLICATIONS OF THE U.S. NATIONAL MUSEUM—Continued.

SEPARATE LIST OF PROCEEDINGS.

Vol.	For Year.	Pages.	Cuts.	Plates.	When Published.	In Vol. of Miscellaneous Collections.	No. in Museum Series.	No. in Smithsonian Catalogue.
I	1878	524	8	8	1879	XIX	17	882
11	1879	503	2	7	1880	XIX	19	388
111	1880	594	5	2	1881	XXII	25	425
17	1881	600	18	2	1882	XXII	27	4 67

PUBLICATIONS OF THE BUREAU OF ETHNOLOGY OF THE SMITHSONIAN INSTITUTION. (Imperial Octavo.)

•	Title.	Published.	Pages.	Cuts.	Plates.	Maps.	No. in Smithsonian Catalogue.
First Annual	Report, for 1879-80	1881	688	348	54	1	476

PUBLICATIONS . HAVING SEPARATE NUMBERS IN THE SMITH-SONIAN SERIES, BUT INCLUDED IN THE ANNUAL REPORTS OF THE INSTITUTION.

		Vel. Repa
A	Journal of Regents	18
B	Report of Organization Committee	18
D	Address at laying corner-stone. DALLAS	18
E	Exposition of Bequest. HENRY	18
F	First Report of Secretary. HENRY	18
J	Programme of Organization. HENRY	18
K	Correspondence, Squier and Davis	18
L	First Report of Organization Committee	15
M	Reports of Institution to 1849	19
N	Officers and Regents	19
25	Public Libraries. JEWETT	18
152	Lectures on Mollusca. CARPENTER.	18
251	Memoir of Von Martius. RAU	18
307	Report on Centennial. BAIRD	18
343	Annual Reports. HENRY	1865-
3 4 8	Fishes of New Jersey. BAIRD	18
351	Forests of North America. Cooper	
352	Lectures on Linguistics. WHITNEY	18
354	Essay on Velocity of Light. DELAUNAY	18
355	Ozone and antozone. WETHERILL	15
36 0	Palafittes. Desor.	18
361	Aborigines of California. BAEGERT.	1863
362	Artificial Shell Deposits in New Jersey. RAU	18
364	Classification of Birds. LILLJEBORG	18
365	Tinneh Indians. Ross and others	18
367	Flora of Alaska. Rothrock	186
368	Indian Pottery. RAU	
369	Dorpat and Poulkova. ABBE	180
37 0	Flint Implements in Illinois. RAU	186
371	Metric Tables. Newton	186
372	Drilling in Stone without Metal. RAU	_ 186
373	Meteorological Stations and Observers	
375	Origin and Nature of Force. TAYLOR	
376	Chemistry of the Earth. HUNT	
377	Diamond and Precious Stones. BABINET	
378	Dakota Language. RŒHRIG	
379	Eulogy on A. D. Bache. HENRY.	1870

	PUBLICATIONS HAVING SEPARATE NUMBERS—Continued.	ol. of
380		eport
		1872
385		1872
386	Crystallography. Brezina	1872
389	Investigation of Illuminants. HENRY	1880
390	Tides. HILGARD	1874
392		1876
393	Mound Builders and Ancient Man in Michigan. GILLMAN1873 and	
394		1878
395	Kinetic Theories of Gravitation. TAYLOR	1876
396	History and Climate of New Mexico. McParlin	187
397	Latimer Antiquities. Mason	1876
398	Meteorological Memoirs. ABBE	187
399	Color-blindness. Holmgren	187
400	Aboriginal Structures in Georgia. Jones	187
401		187
402	Stock-in-trade of Aboriginal Lapidary. RAU	187
403	Gold Ornament from Florida. RAU	187
404	Polychrome Bead from Florida. *HALDEMAN	187
405	Henry and the Telegraph. TAYLOR	1878
406	Researches in Sound. HENRY	1878
407	Memoir of Joseph Henry. GRAY	187
408	Report of Secretary for 1878. BAIRD.	1878
409	Report of Museum Building Commission and Architects	1879
410	Reports of Observatories. Holden	1879
411	Irritation of Polarized Nerve. LAUTENBACH	1878
415	Savage Weapons. KNIGHT	1879
418	Report of Secretary for 1879. BAIRD.	1879
419	Report of Secretary for 1880. BAIRD	1880
420	Anthropological Investigations. MASON	1879
421	Index to Anthropological Articles. BŒHMER	1879
426	Synopsis of Herschel's Writings. Holden and Hastings	1880
427	Recent Progress in Astronomy. Holden	1880
428	"Geology and Mineralogy. HAWES	1880
4 29	" Physics and Chemistry. BARKER	1880
430	. Botany. FARLOW	1880
431	" Zoology. GILL	1880
432	" Anthropology. MASON	1880
433	Report of Visit to Luray Cave. MASON	1880
134	Report of Museum Building Commission and Architects	1880

PUBLICATIONS NOT IN REGULAR SERIES OF "CONTRIBUTIONS," "COLLECTIONS," OR "REPORTS.

- C Digest of Act of Congress.
- P Hints on Public Architecture. OWEN.
- O Check-list of Periodicals.
- 18 Report on the Discovery of the Planet Neptune. Gould.
- 47 On Construction of Catalogues of Libraries. JEWETT.
- 48 Bibliography of American Natural History for 1851. GIRARD.
- 62 Catalogue of Described Coleoptera of the United States. MELSHEIMER.
- 64 List of Foreign Institutions.
- 65 Registry of Periodical Phenomena.
- 66 The Annular Eclipse of May 26, 1854.
- 68 Vocabulary of the Jargon. MITCHELL and TURNER.
- 69 List of Domestic Institutions.
- 93 Meteorological Observations for 1855.
- 101 Map of the Solar Eclipse.
- 105 Catalogue of North American Mammals. BAIRD.
- 106 Catalogue of North American Birds. BAIRD.
- 116 List of Public Libraries. RHEES.
- 157 Results of Meteorological Observations, 1854-1859. Vol. I.
- 170 Comparative Vocabulary.
- 182 Results of Meteorological Observations, 1854-1859. Vol. II.
- 185 List of Birds of Mexico, &c. BAIRD.
- 193 List of Duplicate Shells collected by Wilkes.
- 217 Letter of M. Hoek on Meteoric Shower.
- 225 List of Foreign Correspondents.
- 226 List of Smithsonian Publications.
- 260 Regulations of the Smithsonian Institution.
- 278 Check-list of Publications of Smithsonian Institution.
- 290 Circular for distribution at Centennial.
- 347 Nomenclature of Clouds.
- 350 Map of Stars near the North Pole.
- 359 Planisphere of the Visible Heavens.
- 366 Directions for collecting Diatomacea, &c. EDWARDS.
- 414 Base Chart.
- 437 Check-list of Smithsonian Publications.
- 466 Directory of Officers and Employés.

LIST OF PAPERS PUBLISHED IN THE "SMITHSONIAN CONTRIBU-TIONS TO KNOWLEDGE."

Sama.	Author.	Title of Work.	Commissions of Reference.
1	SQUIER, E. G., AND DAVIS, E. H.	Ancient Monuments of Mississippi Valley.	E. Robinson, D. D., J. R. Bart- lett, Prof. W. W. Turner, S. G. Morton, M. D., G. P. Marsh.
3	WALKER, S. C	Researches, Planet Neptune.	Diolong Par Di, G. Ar Bankan.
4	Walker, S. C	Ephemeris of Neptune for 1848.	•
5	Walker, 8. C	Ephemeris of Neptuno for 1849.	
6	Walker, S. C	Ephemeris of Neptune for 1850.	
7	WALKER, 8. C	Ephemeris of Neptune for 1851.	
8	Downes, John	Occultations in 1848.	
9	Downes, John	Occultations in 1849.	
10	Downes, John	Occultations in 1830.	
11	Downes, John	Occultations in 1851.	•
12	Lieber, Francis	Vocal Sounds of L. Bridgman	Col. W. W. S. Bliss, Miss D. L. Dix.
13	Ellet, Charles	Physical Geography of U.S	Maj. A. Mordecai, Capt. F. A. Smith, Lieut. M. L. Smith.
14	GIBBES, R. W	Memoir on Mosasaurus	Prof. Louis Agassiz, Prof. Henry D. Rogers.
15	SQUIER, E. G	Aboriginal Monuments of New York	Brantz Mayer, Wm. W. Turner.
16	Agassiz, Louis,	Classification of Insects	Prof. C. D. Meigs, Thaddeus W. Harris, M. D.
17	HARE, ROBERT	Explosiveness of Nitre	John Torrey, M. D., Col. J. J. Abert.
20	BAILEY, J. W	Microscopic Examination of Soundings	
22	GRAY, AFA	Plantæ Wrightianæ. Part I	
Ľ;	Bailey, J. W	Microscopic Observations in South Caro- lina, Georgia, and Florida.	Prof. Wm. B. Rogers, Prof. Lewis R. Gibbes.
24 ;	WALKER, S. C	Ephemeris of Neptune, 1852. Appendix I.	
29	Downes, John	Occultations in 1852.	•
:.0	GIRARD, CHARLES	Fresh-Water Fishes of North America	Dr. Jared P. Kirtland, Dr. J. E. Holbrook.
32	HARVEY, WM. H	Marine Algæ of North America. Part I	Prof. J. W. Bailey, Dr. Asa Gray.
33	DAVIS, CHAS. H	Law of Deposit of Flood Tide	Prof. L. Agassiz, Prof. A. Guyot.
35	LOCKE, JOHN	Observations on Terrestrial Magnetism	Prof. Elias Loomis, Prof. Julius E. Hilgard.
26	SECCHI, A	Researches on Electrical Rhoometry	J. H. Lane, Prof. Jas. Curley.
37		Ancient Works in Ohio	Brantz Mayer, E. G. Squier.
40	Riggs, S. R	Dakota Grammar and Dictionary	Minnesota Historical Society, Prof. W. W. Turner, Prof. C. C. Felton.
41	LEIDY, JOSEPH	Extinct American Ox	Paul B. Goddard, M. D., Joseph Pancoast, M. D.
42	GRAY, ASA	Plantæ Wrightianæ. Part II	Prof. John Torrey, John Carey.
43	HARVEY, WM. H	Marine Alge of North America. Part II	Prof. J. W. Bailey, Dr. Asa Gray.
44	LEIDY, JOSEPH	Flora and Fauna within Living Animals	Prof. J. W. Bailey, Charles F. Beck, M. D.
4 5	WYMAN, JEFFRIES	Anatomy of Rana Pipiens	J. B. S. Jackson, M. D., Joseph
4 6	TORREY, JOHN	Plantæ Fremontianæ	Dr. W. Darlington, Dr. Asa Gray-

PAPERS PUBLISHED IN SMITHSONIAN CONTRIBUTIONS TO KNOWLEDGE-Continued.

SERIAL NUMBER.	Author.	TITLE TO WORK.	COMMISSIONS OF REFERENCE
5 0	STIMPSON, WM	Marine Invertebrata of Grand Manan	A. A. Gould, M. D., Prof. J. F.
52	Coffin, Jas. H	Winds of the Northern Hemisphere	Dana. Prof. W. B. Rogers, Prof E.
54	Downes, John	Occultations in 1853.	Loomis.
5 8	LEIDY, JOSEPH	Ancient Fauna of Nebraska	Prof. J. Hall, J. L. Leconte, M.
5 9	Chappelsmith, J	Tornado in Indiana	Prof. J. H. Coffin, Prof. A. (3-
60	Torrey, John	Batis Maritima	well. Prof. A. Gray, John Carey.
61	Torrey, John	Darlingtonia Californica	Prof. A. Gray, John Carey.
63	Bailey, J. W	New Species of Microscopic Organisms	Prof. C. R. Gilman, M. D., W.
70	Lapham, I. A	Antiquities of Wisconsin	I. Burnett, M. D. American Antiquarian Socret
71	Haven, S. F	Archæology of the United States	Peter Force, Brantz Mayer.
72	LEIDY, JOSEPH	Extinct Sloth Tribe of North America	Isaac Hays, M. D., Prof. W. L.
73	••••••	Publications of Societies in Smithsonian Library.	Horner.
79	Runkle, John D	Tables for Planetary Motion	Prof. Benj. Peirce, Com. Char E. Davis.
80	ALVORD, BENJ	Tangencies of Circles and Spheres	
81	OLMSTED, D	Secular Period of Aurora Borealis	
82	Jones, Joseph	Investigation on Amer. Vertebrata	S. Jackson, M. D., J. Leidy, M.D. J. Wyman, M. D.
.8 3	Мексн, L. W	Relative Intensity of Heat and Light of the Sun.	Prof. B. Peirce, Dr. B. A. Godil. Jr.
· 84	Force, Peter	Auroral Phenomena in North Latitudes.	
85	***************************************	Publications of Societies in Smithsonian Library. Part II.	
86	Mayer, Brantz	Mexican History and Archæology	S. F. Haven, E. H. Davis, M. P.
8 8	GIBBS, W., AND GENTH, F. A.	Ammonia Cobalt Bases	Prof. John F. Frazer, Prof. Jel Torrey.
89	BREWER, TH. M	North American Oology. Part I	John Gould, John Cassin.
90	Нітенсоск, Е	Illustrations of Surface Geology.	T
94	Runkle, John D	Asteroid Supplement to New Tables for $b = \frac{(i)}{s}$.	Prof. Benj. Peirce, Com. Cl.as. H. Davis.
95	HARVEY, WM. H	Marine Algæ of North America. Part III.	
96	HARVEY, WM. H	Marine Algæ of North America. 3 parts complete.	Prof. J. W. Bailey, Dr. Asa Gray
97	KANE, E. K	Magnetic Observations in the Arctic Seas	Prof. A. D. Bache.
98	Bowen, T. J	Yoruba Grammar and Dictionary	Prof. J. W. Gibbs, Prof. W. D. Whitney, R. Anderson.
100	Gilliss, J. M	Eclipse of the Sun, Sept. 7, 1858	Prof. J. S. Hubbard, J. H. Lane.
103	CASWELL, A	Meteorological Observations, Providence, R. I.	
104	KANE, E. K	Meteorological Observations in Arctic Seas.	Prof. A. D. Bache.
113	Васни, А. Д	Magnetic and Meteorological Observa- tions at Girard College. Part I.	Prof. Benjamin Peirce, Prof. W. Chauvenet.

PAPERS PUBLISHED IN SMITHSONIAN CONTRIBUTIONS TO KNOWLEDGE-Continued.

Stutat.	AUTHOR.	Title of Work.	Commissions of Reference.	
114	SONNTAG, A	Terrestrial Magnetism in Mexico	Prof. Stephen Alexander, Pro	
119	Whittlesey, Ch	Fluctuations of Level in N. A. Lakes	Arnold Guyot. Capt. A. A. Humphreys, Capt. A.	
120	HILDRETH, S. P., and Wood, J.	Meteorological Observations, Marietta, Ohio.	W. Whipple.	
121	Васпе, А. D	Magnetic and Meteorological Observa- tions at Girard College. Part II.		
126	Le Conte, John L.	Coleoptera of Kansas and New Mexico	C. Zimmerman, F.E. Melsheimer.	
127	Lookis, E	Storms in Europe and America, Dec. 1836.	Dr. S. P. Hildreth, Prof. A. Caswell.	
129	KANE, E. K	Astronomical Observations in the Arctic Seas.	Prof. A. D. Bache.	
130	KANE, E. K	Tidal Observations in the Arctic Seas	Prof. A. D. Bache.	
131	Smith, N. D	Meteorological Observations in Arkansas from 1840 to 1859.		
132	Bacne, A. D	Magnetic and Meteorological Observa- tions at Girard College. Part III.		
135	MITCHELL, S. W	Venom of the Rattlesnake	Franklin Bache, M. D., Robley	
146	M'CLINTOCK, SIR F. L.	Meteorological Observations in the Arctic Seas.	Dunglison, M. D.	
155	WHITTLESEY, CH	Ancient Mining on Lake Superior	D. Wilson, LL. D., E. H. Davis, M. D.	
159	MITCHELL, S. W., & MOREHOUSE, G. R.	Anatomy and Physiology of Respira- tion in Chelonia.	Prof. J. Wyman, Prof. J. Leidy.	
162	Васне, А. D	Magnetic and Meteorological Observa- tions at Girard Coll. Pt. IV, V, & VI.		
163	BACHE, A. D	Magnetic Survey of Pennsylvania.		
172	MEEK, F. B., AND HAYDEN, F. V.	Palmontology of the Upper Missouri. Part I.	Isaac Lea, Prof. Jas. D. Dana.	
173	DEAN, JOHN	Gray Substance of the Medulla Oblongata.	Dr. W. A. Hammond, Prof. Jeff- ries Wyman.	
175	Васив, А. D	Mag. and Met. Observ. at Girard College. Parts VII, VIII, and IX.		
180	DRAPER, H	Construction of a Silvered Glass Telescope.	Prof. Wolcott Gibbs, Com. J. M. Gilliss.	
186	Васпи, А. D	Mag. and Met. Observ. at Girard College. Parts X, XI, and XII.		
192	LEIDY, JOSEPH	Cretaceous Reptiles of the U.S	Prof. L. Agassiz, Prof. E. D. Cope.	
195	Васив, А. D	Girard College Observations. Complete. Parts I to XII.	Prof. Benj. Peirce, Prof. W. Chauvenet.	
1146	HATES, I. I	Physical Observations in the Arctic Seas.		
197	WHITTLESEY, CH	Glacial Drift of Northwestern States	Prof. L. Agassiz, Prof. J. P. Lesley.	
198	KANE, E. K	Physical Observations in the Arctic Seas. Complete.	Prof. A. I). Bache.	
199	Newcomb, S	Orbit of Neptune	Adm'l C. H. Davis, Prof. Stephen Alexander.	
202	PUMPELLY, R	Geological Researches in China, Mongolia, &c.	National Academy of Sciences.	
3H '	CLEAVELAND, P	Meteorological Observations, Brunswick, Me., 1807–1859.		

PAPERS PUBLISHED IN SMITHSONIAN CONTRIBUTIONS TO KNOWLEDGE—Continued.

			<u> </u>
SEELAL NUMBER.	Author.	Title of Work.	Commissions of Reference.
208	Pickering, Chas	Gliddon Mummy Case in Smithsonian Institution.	
218	Morgan, L. H	Systems of Consanguinity and Affinity	Prof. J. A. Hadley, J. H. Fran- bull, Prof. W. D. Whitney.
220	Swan, Jas. G	Indians of Cape Flattery	George Gibbs, Jeffries Wymar
221	Coppin, James H	Orbit, &c., of Meteoric Fire Ball, July 20, 1860.	Prof. H. A. Newton, C. A. Schert
222	SCHOTT, CHAS. A	Tables of Rain and Snow in United States.	
223	GOULD, B. A	Transatlantic Longitude.	
232	STOCKWELL, J. N	Secular Variations of Orbits of Planets	Prof. J. H. C. Coffin, Prof. S. Newcomb.
233	FERREL, WM	Converging Series, Ratio of Diameter, and Circumference of Circles.	Newcomb.
239	Harrness, Wm	Magnetic Observations on the Monad- nock.	Pres. Acad. Sciences. Prof. J. H. C. Coffin, Prof. F. Rogers
240	BARNARD, J. G	Problems of Rotary Motion.	
241	W00D, H. C	Fresh-Water Algæ of North America	Dr. J. Torrey, Dr. F. A. P. Farnard.
242	CLARK, H. J	Lucernariæ.	natu.
248	Hilgard, E. W	Geology of Lower Louisiana.	
259	Jones, Jos	Antiquities of Tennessee	Dr. G. A. Otis, Prof. O. T. Ma-
262	Newcomb, S	Orbit of Uranus	Prof. J. H. C. Coffin, Prof. Ass. Hall.
267	Swan, J. G	Haidah Indians	J. C. Welling, LL. D., Dr. 6 3. Otis.
268	Coffin, J. H	Winds of the Globe.	Otio.
269	Habel, Simeon	Sculptures of Santa Lucia Cosumal- whuapa in Guatemala.	Prof. W. D. Whitney, J. H. Trumbull.
277	Зснотт, С. А	Temperature Tables.	•
280	Alexander, S	Harmonies of Solar System.	
281	Newcomb, S	Planetary Motion	Prof. H. A. Newton, G. W. Hill.
287	Rau, Charles	Archæological Collection, Nat. Museum.	
310	Barnard, J. G	Internal Structure of the Earth.	
317	ELLIOT, D. G	Classification and Synopsis of Trochilidæ.	
318	Dall, Wm. H	Remains of Man from Caves in Aleutian Islands.	
331	Rau, Charles	The Palenque Tablet	S. F. Haven, H. H. Bancroft.
353	Зснотт, С. А	Tables of Rain Fall, (2d edition.)	
357	W00в, H. C	Researches on Fever	S. Weir Mitchell, M. D., J. Woodward, M. D.
383	Bransford, J. F	Archæological Researches in Nicaragua	Garrick Mallery, H. C. Yarr 4. M. D
443	Caswell, A	Meteorological Observations, Providence, R. I., 1831 to 1876.	

ALPHABETICAL INDEX OF ARTICLES

IN THE

SMITHSONIAN CONTRIBUTIONS TO KNOWLEDGE, Vols. 1-23. 1848-1881.

SMITHSONIAN MISCELLANEOUS COLLECTIONS, Vols. 1-23. 1862-1882.

> SMITHSONIAN ANNUAL REPORTS, Vols. 1-35. 1846-1880.


BULLETINS OF THE U. S. NATIONAL MUSEUM,
Nos. 1-22, (except 16 and 20,) 1875-1881,

PROCEEDINGS OF THE U. S. NATIONAL MUSEUM, Vols. 1-4. 1878-1881.

AND IN THE

FIRST ANNUAL REPORT OF THE BUREAU OF ETHNOLOGY OF THE SMITHSONIAN INSTITUTION,

Vol. 1. 1879–1880.



ALPHABETICAL INDEX.

- NOTE.—The heavy-faced figures indicate the specific number in the Catalogue of a separate publication; the ordinary, or light-faced figures indicating the number of a publication of which the paper is only a constituent part.
- "R." represents the annual Smithsonian Report, immediately followed by the year of the report, and the number given to it in the Catalogue.
- "P." represents the Proceedings of the National Museum, followed by the year of publication and the number given to the volume in the Catalogue.
- "E." represents Report of the Bureau of Ethnology.

A.

ABBE, U. Description of Observatories at Dorpat and Poulkova. (R, 1867) 215, 369
ABBE, C. Meteorological memoirs, translated by. (R. 1877)
Аввотт, С. С. Stone age in New Jersey. (R. 1875) 298, 394
Abbreviations of names of States, etc., for labelling collections 164
Abbreviations used in England in 1867—W. DE LA RUE. (R. 1867) 215
Abeona aurora—Jordan and Gilbert. (P. 1880) 425
ABICH, S. Remarkable forms of hail-stones. (R. 1869) 228
Aboriginal indians of America, language of-G. GIBBS. (R. 1870) 244
Aboriginal inhabitants of Cal J. BAEGERT. (R. 1868; R. 1864) _ 187, 188, 440, 361
Aboriginal lapidary, Stock-in-trade of-C. RAU. (R. 1877) 823, 440, 402
Aboriginal monuments of State of New York—E. G. Squier 15
Aboriginal remains of Tennessee—J. Jones 259
Aboriginal ruins at Savannah, Tenn.—J. P. STELLE. (R. 1870) 244
Aboriginal ruins in Hardin Co., Tenn.—J. P. STELLE. (R. 1870) 244
Aboriginal shell-mounds, New Brunswick & New EngS. F. BAIRD. (P. 1881) 467
Aboriginal structures in Georgia—C. C. Jones, Jr. (R. 1877) 828, 400
Abstracts of anthropological correspondence—O. T. MASON. (R. 1879; R.
1880) 845, 442
Academy, Pontifical, Rome—See Prize questions.
Academy of Sciences—See National.
Academy of Sciences, Bordeaux—See Prize questions.
Academy of Sciences, Chicago, acknowledging specimens. (R. 1867) 215
Academy of Sciences, St. Petersburg, exchange system. (R. 1867) 215
Academy of Sciences, Vienna—See Prize questions.

Acalephæ, anatomy and physiology of-H. J. CLARK	
Accidental or subjective colors—ABBE MOIGNO. (R. 1866)	
Account of—	
aboriginal ruins at Savannah, Tenn.—J. P. STELLE. (R. 1	870)
aboriginal ruins in Hardin Co., Tenn.—J. P. STELLE. (R	. 1870)
aborigines of California-J. BAEGERT. (R. 1868; R. 1864).	187, 188, 440,
ancient mounds, Georgia-M. F. STEPHENSON. (R. 1870)	
antiquities, Tennessee-E. O. DUNNING. (R. 1870)	
antiquities, Vera Cruz, Mexico—H. Finck. (R. 1870)	
astronomical observatory at Cordoba—B. A. GOULD. (R.	1878)
balloon ascensions—J. GLAISHER. (R. 1863)	••••••
barometer, new-J. Green. (R. 1855)	
. burial of Indian squaw, California-W. M. King. (R. 18	74)
Carib or Karif language and people. (R. 1878)	
discovery of stone image in Tennessee-E. M. GRANT. (R	
Draper's telescope—T. W. WEBB. (R. 1864)	
eclipse of sun in Peru, 1858-J. M. GILLISS	
hail storm in Texas—G. M. BACHE. (R. 1870)	
hail storm on Bosphorus-Com. PORTER. (R. 1870)	
Indians of British America—E. PETITOT. (R. 1865)	
lightning discharges. (R. 1867)	
Lindheimer's botanical explorations—A. GRAY. (R. 1849))
marble used in extension of U.S. Capitol—J. HENRY. (R	. 1856)
"occultator" for indicating moon's position—T. HILL	
old Indian village, Kushkushkee, Pa.—E. M. McConnell	(R. 1871)_
Portland Society of Natural History—E. C. Bolles. (R.	1867)
Priestley's lens—J. HENRY. (R. 1859)	
remarkable accumulation of bats-M. FIGANIERRE. (R. 1	863)
sarcophagus in National Museum—A. A. HARWOOD. (R.	1870)
storm in Butler Co., Kans., June 28, 1871—W. HARRISON.	(R. 1871)
tornado in Spruce Creek Valley, Pa.—J. B. MEEK. (R. 1	871)
tornado near New Harmony, Indiana—J. CHAPPELSMITH	
travels in Central and South America—S: HABEL	
Acid, sulphurous, action of, upon putrefactive bacteria. (P. 1881	.)
Acknowledgment of birds—Academy of Sciences, Stockholm. (R. 1867)
Acknowledgment of birds—J. Gould. (R. 1867)	
Acknowledgment of contributions to National Museum. (P. 188	1) 467
Acknowledgment of ferns-G. METTENIUS. (R. 1862)	
Acknowledgment of perennibranchiates—J. G. FISHER. (R. 186	i8)
Acknowledgment of shells, &cW. H. PEASE. (R. 1861)	
Acknowledgment of specimens—British Museum. (R. 1865)	h 41720 00000 -4
Acknowledgment of specimens—Chicago Academy of Sciences.	(R. 1867)

Acknowledgment of specimens—Regents of Univ. of State of N. Y. (R. 1865).	209
Acmæidæ, from Panama, new species of—P. P. CARPENTER	252
Acoustics applied to public buildings—J. HENRY. (R. 1856)	91
Acoustics, different kinds of refraction—W. B. TAYLOR. (R. 1875)	298
Acoustics, lectures on architecture in relation to-D. B. REID. (R. 1856)	91
Acrothele, note on—C. A. WHITE. (P. 1880)	425
Acts of Congress—See Congress.	
ADAMS, C. B. Catalogue of shells of Panama, review of-P. P. CARPENTER.	252
ADAMS, C. B. Shells collected by, description of	252 .
Adams, W. Subcutaneous surgery. Toner lecture No. VI	302
ADAMS, W. H. Mounds in Spoon River Valley, Illinois. (R. 1879)	345
Additions to Museum by Berlin Fisheries Exhibition. (R. 1880)	442
Address at laying of corner-stone of Smithsonian building-G. M. Dallas. D,	329
Address on the Smithsonian Institution—J. HENRY. (R. 1858) E	67
Address, Pres. Royal Society of Victoria-R. L. J. ELLERY. (R. 1868)	224
Addresses at memorial services of J. Henry—See Henry.	
Addresses, list of, foreign institutions, since 1862. (R. 1865)	209
Administration of drugs and medicines—J. M. FLINT. (P. 1881) 467,	
	147
	147
	147
	147
	187
	187
	215
Affinity, systems of—L. H. Morgan	
Africa, stone celts in—G. J. GIBBS. (R. 1877)	
Africa, Yoruba, account of, and grammar and dictionary	
AGASSIZ, L., biographical notice of—J. A. GARFIELD. (R. 1873) 275,	
AGASSIZ, L., biographical notice of—P. PARKER. (R. 1878) 275,	
Agassiz, L., biography of—E. FAVRE. (R. 1878)	
Agassiz, L., memoir of—R. P. STEBBINS. (R. 1873)	
Agassiz, L., supplemental list of generic names employed by—S. H. Scudder.	
Agassiz, L., papers by—	270
Classification of insects from embryological data	16
Narrative of the Hassler expedition. (R. 1872) 271,	
On the formation of a museum. (R. 1849)	
Report on use of new hall in Smithsonian building. (R. 1867) 215,	
	249
	286
•	215
Agonidæ, new species of, from California—W. N. Lockington. (P. 1880)	
·	

Agonus vulsus—Jordan and Gilbert. (P. 1880)
Agricultural Association of Milan, on exchanges. (R. 1863)
Agricultural chemistry, lectures on—S. W. Johnson. (R. 1859) 110
Agricultural flint implements in Illinois—C. RAU. (R. 1868) 224, 440, 370
Agricultural flint implements, Southern Illinois—C. RAU. (R. 1868) 224, 370
Agricultural implements of N. American stone period—C. RAU. (R. 1863) 187. 441
AINSA, SANTIAGO. Tucson meteorite. (R. 1863)
Air, temperature in ascending currents of—J. HANN. (R. 1877) 323.
Air, temperature in ascending moist currents of—L. Sohncke. (R. 1877) 325.
Air-currents, nature of—A. Colding. (R. 1877)
AIRY, G. B. Correcting measures of the sun's distance. (R. 1859)
Alabama, mica beds in—W. GESNER. (R. 1879)
Alabama, shell-heaps—A. S. Gaines and K. M. Cunningham. (R. 1877)
Alaska, bibliography of fishes of, for 1880—T. H. BEAN. (P. 1881)
Alaska, climate of—H. M. BANNISTER. (R. 1866)
Alaska, description of Lycodes Turneri from—T. H. BEAN. (P. 1878)
Alaska, descriptions of new fishes from—T. H. BEAN. (P. 1881)
Alaska, flora of—J. T. ROTHROCK. (R. 1867)
Alaska, limpets and chitons of, report on—W. H. Dall. (P. 1878)
Alaska, new fish from—T. H. BEAN. (P. 1879)
Alaska, new fish from—W. N. Lockington., (P. 1880)
Alaska, new forms of mollusks from—W. H. DALL. (P. 1878)
Alaska, notes on Nichols' collection of fishes of—T. H. BEAN. (P. 1881) "
Alaska, occurrence of Hippoglossus vulgaris at—T. H. BEAN. (P. 1879)
Alaska, occurrence of Stichæus punctatus at—T. H. BEAN. (P. 1878)
Alaska, preliminary catalogue of fishes of—T. H. Bean. (P. 1881)
Alaska, remains of prehistoric man from caves in—W. H. Dall
Alaska, some genera and species of fishes from—T. H. Bean. (P. 1879)
Albany, Illinois, mound skull and bones—R. J. FARQUHARSON. (R. 1874)
Albany Institute, memorial proceedings relative to Joseph Henry
Album from Greece, presented by Miss E. B. Contaxaki. (R. 1857)
Aleutian Islands, someins of prohistoric man from W. H. Davids
Aleutian Islands, remains of prehistoric man from—W. H. Dall
ALEXANDER, B. S. Report of architect. (R. 1854)
ALEXANDER, C. A. Origin and history of Royal Society of London. (B. 1863)
ALEXANDER, C. A., (translations by)—
Accidental or subjective colors—ABBE MOIGNO. (R. 1866)
American migration—F. von Hellwald. (R. 1866)
Biography of Peltier by his Son. (R. 1867)
Carlsruhe Scientific Congress—M. J. NICKLES. (R. 1860)
Catalytic force—T. L. Phipson. (R. 1862)

Electri	cal currents of the earth—C. MATTEUCCI. (R. 1867)
Electri	city. (R. 1867)
Electro	-physiology—C. MATTEUCCI. (R. 1865)
Extern	al appearance of sun's disk. (R. 1866)
	s of equilibrium of a liquid mass—J. Plateau. (R. 1865; R. 36) 209
Forma	tion of ice at bottom of water-M. Englehardt. (R. 1866)
Histor	of Academy of Sciences of Paris-M. Flourens. (R. 1862)
Histor	of Royal Institution of Great Britain—E. MAILLY. (R. 1867)
Horar	variations of barometer—M. VAILLANT. (R. 1866)
Intern	ational Archæological Congress. (R. 1866)
	s, the, and the natural method—M. Flourens. (R. 1867)
	Faraday—De LA RIVE. (R. 1867)
	Kepler-M. Berthrand. (R. 1869)
Man tl	e contemporary of the mammoth. (R. 1867)
_	r of A. Bravais-E. DE BEAUMONT. (R. 1869)
Memo	r of D. de Blainville-M. FLOURENS. (R. 1865)
	r of von Buch-M. Flourens: (R. 1862)
	r of A. de Candolle-M. FLOURENS. (R. 1859)
	r of Delambre—J. Fourier. (R. 1864)
	r of R. J. Häuy-M. Cuvier. (R. 1860)
	r of Legendre—E. DE BEAUMONT. (R. 1867)
	r of L. J. Thenard—M. FLOURENS. (R. 1862)
	r of Magendie—M. Flourens. (R. 1866)
	r of Priestley-M. Cuvier. (R. 1858)
	r of Geoffroy St. Hilaire-M. FLOURENS. (R. 1861)
	cope. (R. 1860)
	ent of the stars—M. MAEDLER. (R. 1859)
	l History of organized beings-M. MARKY. (R. 1867)
	es or lacustrian constructions—E. DESOR. (R. 1865)
	chemistry—M. Jamin. (R. 1867)
Pile-w	ork antiquities of Olmutz. (R. 1866)
	vation of wood. (R. 1864)
	fic Congress of Carlsrube—M. J. Nickles. (R. 1860)
	ation of the stars—C. D. Kämtz. (R. 1861)
_	f feeling: sense of smell. (R. 1865)
_	f hearing. (R. 1866)
	f sight. (R. 1866)
	f taste. (R. 1866)
~	planets between Mars and Jupiter—LESPIAULT. (R. 1861)

ALEXANDER, C. A., (translations by)—Continued.
State of ethnology in relation to form of human skull—A. Pelzius. (R. 1859)
Transactions of Geneva Society of Physics, &c. (R. 1859; R. 1864; R. 1865; R. 1866) 110, 188, 209. 21
ALEXANDER, S. Harmonies of the solar system 28
ALEXANDER, S. Lecture on relations of time and space. (R. 1861) 14
ALEXANDER, S. Lecture on vastness of visible creation. (R. 1857) 10
Algæ—J. W. BAILEY
Algæ, Arctic, list of-W. H. HARVEY
Algæ, Arctic America—W. G. FARLOW
Algæ, bibliography of—H. C. Wood
Algæ, Kerguelen Island-W. G. FARLOW
Algæ, lecture on marine—W. H. HARVEY. (R. 1855)
Algæ, marine. Part 1, Melanospermeæ-W. H. HARVEY 3
Algæ, marine. Part 2, Rhodospermeæ—W. H. HARVEY
Algæ, marine. Part 8, Chlorospermeæ-W. H. HARVEY 9
Algæ, marine. Parts 1, 2, 8, complete—W. H. HARVEY 9
Algæ, North American fresh water, history of-H. C. Wood 24
Algæ, Rhode Island, list of—S. T. OLNEY
ALLEN, H. Monograph of bats of North America
ALLEN, Z. Explosibility of coal oils. (R. 1861)
Alphabetical index of genera of birds—S. F. BAIRD
Alphabetical index to Smithsonian publications 470
Alternate generation and parthenogenesis-G. A. KORNHUBER. (R. 1871) 24
Altitude, diminution of vapor with increasing-J. HANN. (R. 1877) 323. 34
Altitudes, circular relative to—J. HENRY 23
ALVORD, B. Tangencies of circles and of spheres
Amazilia cerviniventris-R. RIDGWAY. (P. 1881) 46
Amazilia yucatanensis—R. RIDGWAY. (P. 1881)
Amblystoma, change of axolotl to—A. Weismann. (R. 1877) 323, 403
Amblystoma, luridum—P. R. Hoy. (R. 1854)
America—See Arctic, British, Central, Middle, North, Russian, South.
America, ancient history of—M. Much. (R. 1871)
America, collection of charts and maps of—J. G. Kohl. (R. 1856)
America, grasshoppers and locusts of—A. S. TAYLOR. (R. 1858)
America, language of aboriginal Indians of-G. GIBBS. (R. 1870) 24
America, storms in, December, 1836—E. Loomis
American Academy of Arts and Sciences-
On Smithsonian exchange system. (R. 1867)213
Report on organization of Smithsonian. (R. 1858)
Thanks for exchanges. (R. 1855)

of smithsonian publications.	128
	77.6
American Antiquarian Society. Antiquities of Wisconsin	
American Association for Advancement of Science—	
Combined meteorological observations proposed. (R. 1851)	51
Memorial address on Joseph Henry. A. M. MAYER	
American authors, North American conchology—W. G. BINNEY	142
American Board of Commissioners for Foreign Missions. Grammar and dictionary of Dakota language	
American corbiculadæ, monograph of-T. PRIME	145
American correspondents, list of69,	238
Amerićan Ethnological Society, report on Squier and Davis' work. (R. 1847) I	I, K
American migration—F. Von Hellwald; Jos. Henry. (R. 1866)	214
American natural history, bibliography for 1851. C. GIRARD	48
Amiurus ponderosus, from Mississippi river—T. H. BEAN. (P. 1879)	888
Ammonia-cobalt bases—W. GIBBS; F. A. GENTH.	88
Ampère, eulogy on-F. Arago. (R. 1872)	27
Amphorosteus, memoir on—R. W. GIBBES	14
Ampullariid W. G. BINNEY	14
Analyses of Chinese and Japanese coals—J. A. MACDONALD	203
Analysis, chemical, of the sun—A. LAUGEL. (R. 1861)	149
Analysis of blood—J. Jones	82
Analysis of water destructive to fish in Gulf of Mexico—F. M. Endlich. (P. 1881)	46'
Analysis, spectrum—W. Huggins. (R. 1866)	214
Anarrhichas lepturus, from Alaska, and other species—T. H. BEAN. (P. 1878)	388
Anatomical preparations, arsenic acid for protecting, from insects—J. B. S. JACKSON. (P. 1878)	38
Anatomy and physiology of chelonia—S. W. MITCHELL; G. R. MOREHOUSE.	159
Anatomy and physiology of lucernarians—H. J. CLARK	242
Anatomy and physiology of the rattlesnake—S. W. MITCHELL	13
Anatomy of the nervous system of Rana pipiens—J. WYMAN	45
Ancient aboriginal trade in North America—C. RAU. (R. 1872) 271, 440,	385
Ancient and modern purple dyeing. (R. 1863)	18
Ancient burial mound in Indiana—W. Pidgeon. (R. 1867)	21
Ancient earthworks of Ashland county, Ohio-G. W. HILL. (R. 1877)	82
Ancient earthworks of Ohio, sketch of—I. DILLE. (R. 1866)	214
Ancient earthworks on Upper Missouri—A. BARBANDT. (R. 1870)	24
Ancient fauna of Nebraska—J. Leidy	58
Ancient fort and burial ground, Tomp. Co., N. Y.—D. TROWBRIDGE. (R. 1863)	18
Ancient graves and shell-heaps of California—P. Schumacher. (R. 1874)	28
Ancient history of North America—M. Much. (R. 1871)	24

Ancient implement of wood found in Conn.—E. W. Ellsworth. (R. 1876)__ 299

Ancient man in Michigan, characteristics of-H. GILLMAN. (R. 1875) 298, 3	393
Ancient mica mines North Carolina—C. D. SMITH. (R. 1876)	299
Ancient mining on Lake Superior-C. WHITTLESEY	155
Ancient mining on Lake Superior, circular on—J. HENRY. (R. 1861)	149
Ancient monuments in the Miss. val., correspondence relative to. (R. 1847) H	i, K
Ancient monuments in the Mississippi valley—E. G. SQUIER; E. H. DAVIS	1
Ancient mound in western Pennsylvania-W. M. TAYLOR. (R. 1877)	823
Ancient mound near Chattanooga, Tennessee—M. C. READ. (R. 1867)	215
Ancient mound, near Lexington, Ky.—R. PETER. (R. 1871)	249
Ancient mound, St. Louis, MoT. R. PEALE. (R. 1861)	149
Ancient mounds in Georgia—M. F. STEPHENSON. (R. 1870)	244
Ancient mounds in Kentucky—R. Peter. (R. 1872)	271
Ancient mounds of Mercer county, Illinois—T. McWhorter. (R. 1874)	286
Ancient mounds of Union county, Kentucky—S. S. Lyon. (R. 1870)	244
Ancient pottery from Phillips county, Arkansas—J. H. DEVEREUX. (R. 1872)	271
Ancient pottery on Des Moines river—R. N. and C. Dahlberg. (R. 1879)	34 5
Ancient relic of Maya sculpture—A. Schott. (R. 1871)	249
Ancient relics in Missouri—J. W. FOSTER. (R. 1868)	187
Ancient relics in northwestern Iowa—J. B. Cutts. (R. 1872)	271
Ancient remains in Canada West, near Prescott—W. E. Guest. (R. 1856)	91
Ancient remains in Colorado—E. L. BERTHOUD. (R. 1867)	215
Ancient ruin in Arizona—J. C. Y. LEE. (R. 1872)	271
Ancient town in Minnesota—O. H. KELLEY. (R. 1868)	187
Ancient works in Ohio, description of—C. WHITTLESEY	37
Anderson, M. B. Life of Professor Chester Dewey. (R. 1870)	244
ANDERSON, W. Antiquities of Perry county, Ohio. (R. 1874)	286
Anderson, W. G. Mounds in Illinois and Wisconsin. (R. 1879)	34 5
Andrews, F. D. Indian relics from Schoharie, New York. (R. 1879)	845
Andrews, S. L. Meteors. (R. 1866)	214
Anemometer, description of Smithsonian—J. HENRY. (R. 1860)	147
Aneroid barometers, experiments—B. STEWART. (R. 1868)	274
Animal kingdom, list of substances derived from—G. B. Goode	297
Animal kingdom, parthenogenesis in—G. A. KORNHUBER. (R. 1871)	540
Animal resources of United States, classification of collection—G. B. Goode	297
Animal resources of United States, collection at Centennial—G. B. Goods	326
Animals, circular for shipping—S. F. BAIRD	384
Animals, flora and fauna within living—J. LEIDY	44
Animals, list of all generic names of—S. H. SCUDDER	
Animals, synopsis of useful and injurious—G. B. Goode	
Anna, Illinois, mounds near—T. M. PERRIN. (R. 1872)	
Annals of Philosophy, list of Smithson's papers in. (R. 1858) 67	
Annelide of Arctic America A R VERRITT	342

Annelids of northeastern coast of America—A. E. VERRILL. (P. 1879;	
Annular eclipse of May 26, 1854	
Anthozoa of Kerguelen Island—A. E. VERBILL	
Anthozoa of northeast coast of America—A. E. VERRILL. (P. 1879)	
Anthropological articles, Smithsonian, index to—G. H. Bornmer. (R. 1879)	
Anthropological Congress, address before—F. F. Romen. (R. 1876) 2	
Anthropological correspondence, abstracts of—O. T. Mason. (R. 1880)	
Anthropological correspondence, summary of—O. T. MASON. (R. 1879)	
Anthropological data, limitations to use of—J. W. Powell. (E. 1880)	
Anthropological inquiry, queries for—C. DARWIN. (R. 1867)	
Anthropological investigations in 1879—O. T. MASON. (R. 1879) 8	
Anthropological measurements, table of—SCHERZER; SCHWARZ. (R. 1866)	
Anthropological publications, list of C. RAU's. (P. 1881)	
Anthropological Society of Paris, transactions of—P. Broca. (R. 1868)	
Anthropological Society of Vienna, address on ancient history of Nor	
America—M. Much. (R. 1871)	
Anthropology, articles on. (R. 1879)	
Anthropology, bibliography of—O. T. MASON. (R. 1880)4	_
Anthropology, index to Smithsonian papers on—G. H. Boehmer. (R. 1879)	
Anthropology, progress of, in 1880—O. T. Mason. (R. 1880) 4	
Anthropology, recent progress in—O. T. MASON. (R. 1880)	•
Antigua, catalogue of Ober's collection of birds of—G. N. LAWRENCE. (P. 187	•
Antilles, maritime disasters of the. (R. 1867)	
Antilles—See Lesser Antilles.	
Antiquarian Society, American. Antiquities of Wisconsin	
Antiquities, California, Santa Rosa Island—S. Bowers. (R. 1877)	
Antiquities, Col., Jefferson and Clear Creek Cos.—G. L. CANNON. (R. 1877)	•
Antiquities, Colorado, Weld county—E. L. BERTHOUD. (R. 1871)	
Antiquities, Denmark, preservation of—J. J. A. Worsaae. (R. 1879)	
Antiquities, Florida—J. BARTRAM. (R. 1874)	
Antiquities, Florida—A. MITCHELL. (R. 1874)	
Antiquities, Georgia, Spalding county—W. B. F. BAILEY. (R. 1877)	
Antiquities, Guatemala—G. WILLIAMSON. (R. 1876)	
Antiquities, Hungary, prehistoric—F. F. ROMER. (R. 1876) 299, 4	4 0, 3
Antiquities, Illinois, Lawrence county—A. PATTON. (R. 1878)	
Antiquities, Illinois, Mason county—J. Cochrane. (R. 1877)	
Antiquities, Illinois, Rock Island county—A. Toellner. (R. 1879)	
Antiquities, Illinois, Union county-T. M. PERRINE. (R. 1878)	
Antiquities, Illinois, Whitesides county—W. H. PRATT. (R. 1874)	

Antiquities, Indiana, Allen and DeKalb counties—R. S. Robertson. (R. 1874) 286
Antiquities, Indiana, Knox county—A. PATTON. (R. 1878) 275
Antiquities, Indiana, Laporte county—R. S. Robertson. (R. 1874) 280
Antiquities, Kentucky—S. S. Lyon. (R. 1858)
Antiquities, Kentucky, Hancock county—J. FRIEL. (R. 1877) 323
Antiquities, Maryland, Charles county-O. N. BRYAN. (R. 1874) 286
Antiquities, Mexico, Vera Cruz—H. FINCK. (R. 1870) 244
Antiquities, Michigan, Isle Royale—A. C. Davis. (R. 1874) 286
Antiquities, Mississippi river and Lake Pepin-L. C. Estes. (R. 1866) 216
Antiquities, Mississippi, Yazoo county—J. W. C. Smith. (R. 1874) 29
Antiquities, Missouri and Tennessee—I. DILLE. (R. 1862)
Antiquities, Missouri, Kansas City—W. H. R. LYKINS. (R. 1877)
Antiquities, New Mexico—W. B. Lyon. (R. 1871) 249
Antiquities, New York, Orleans county—F. H. Cushing. (R. 1874) 28
Antiquities, Nicaragua—E. G. SQUIER. (R. 1850)
Antiquities, North Carolina, Lenoir county—J. M. Spainhour. (R. 1871) 24:
Antiquities, N. C., Stanley and Montgomery counties—F. J. Kron. (R. 1874)
Antiquities, Ohio—G. W. HILL. (R. 1874)
Antiquities, Ohio, Perry county—W. Anderson. (R. 1874)
Antiquities, Olmutz pile-work. (R. 1866)
Antiquities, Porto Rico, Latimer collection—O. T. MASON. (R. 1876) 299, 397
Antiquities, Southern States—H. C. WILLIAMS. (R. 1870) 249
Antiquities, Tennessee-W. M. CLARK. (R. 1877)
Antiquities, Tennessee—E. O. Dunning. (B. 1870)
Antiquities, Tennessee—J. Jones 259
Antiquities, Tennessee—R. S. Robertson. (R. 1877)
Antiquities, Tennessee—D. F. WRIGHT. (R. 1874)
Antiquities, Tennessee, Blount county-Miss A. E. Law. (R. 1874)
Antiquities, Tennessee, Jackson county—J. Haile; J. W. McHenry. (B. 1874)
Antiquities, United States, proposed map of—A. J. HILL. (R. 1861)
Antiquities, Wisconsin—I. A. LAPHAM
Antiquities, Wisconsin—M. STRONG. (R. 1877)
Antiquity in Europe, the study of high—A. Morlot. (R. 1862; R. 1864) 150. 140
Antiquity in United States, vestiges of—S. F. HAVEN
Antozone and ozone—C. M. WETHERILL. (R. 1864)
Apaches, Tonto, notes on the—C. SMART. (R. 1867)
Aphododeridæ, notes on—D. S. JORDAN
Apodichthys-Jordan and Gilbert. (P. 1880)
Apogon pandionis, description of-G. B. GOODE; T. H. BEAN. (P. 1881) 45
Apparatus available for research, list of. (R. 1878)
Apparatus, blow-pipe—Hawkins and Walk. (R. 1872)
Apparatus for testing perspiration and respiration—Petrenkoves. (R. 1864: 1864)

Apparatus, letter relative to gift of—R. HARE. (R. 1848)
Appropriations for Museum, memorial of regents asking for. (R. 1867) 215, 329
Aqueous vapor, diminution of, with altitude—J. HANN. (R. 1877) 823, 898
Arachnida of Arctic America—S. H. Scudder
Arago, F. Aeronautic voyages. (R. 1868) 187
Autobiography. (R. 1870) 244
Biography of Condorcet. (R. 1878) 341
Eulogy on Ampère. (R. 1872) 271
Eulogy on Joseph Fourier. (R. 1871) 249
Eulogy on Gay-Lussac. (R. 1876) 299
Eulogy on Herschel. (R. 1870) 244
Eulogy on La Place. (R. 1874) 286
Eulogy on Alex. Volta. (R. 1875) 298
Eulogy on Thomas Young. (R. 1869) 228
Archæological Academy of Belgium, Archæological Congress organized by.
(R. 1866) 214
Archæological collection of the U.S. National Museum—C. RAU 287
Archæological Congress, International, Antwerp. (R. 1866) 214
Archæological investigations, instructions for-G. GIBBS. (R. 1861) 149, 160
Archæological researches at Concise—F. TROYON. (R. 1861) 149
Archæological researches in Nicaragua-J. F. Bransford 383
Archæological study of jade—S. BLONDEL. (R. 1876) 299
Archæology, circular relative to American
Archæology, French Society of. (R. 1866)214
Archæology, general views on-A. Morlot. (R. 1860; R. 1861) 147, 149
Archæology, Mexican—B. MAYER 86
Archæology, North American—J. Lubbock. (R. 1862) 150
Archæology of the United States—S. F. HAVEN
Archæology, prehistoric, international code of symbols for charts of. (R. 1875) 298
Archæology and ethnology, circular for collecting specimens of 205
Architect's report—B. S. ALEXANDER. (R. 1854) 75, 829
Architect's report—A. Cluss. (R. 1867) 215, 329
Architects' report for 1879—Cluss & Schulze. (R. 1879)
Architects' report for 1880—Cluss & Schulze. (R. 1880) 442, 434
Architecture, hints on public—R. D. Owen
Architecture in relation to ventilation, warming, lighting, fire-proofing, acoustics,
and the general preservation of health—D. B. REID. (R. 1856) 91
Arctic algæ, list of-W. H. HARVEY 95
Arctic America, natural history of—L. Kumlien
Arctic atmospheric pressure—I. I. HAYES
Arctic auroras—P. Force84
Arctic expedition, Howgate-L. Kumlikn

Arctic explorations, lecture on—I. I. HAYES. (R. 1861)	149
Arctic latitudes and longitudes—E. K. KANE	129
Arctic observations-See Force, Hall, Hayes, Kane, Kumlien, McClintock, Scho	tt.
Arctic regions, limpets and chitons of, report on-W. H. Dall. (P. 1878)	882
Arctic regions, record of voyage of yacht "Fox"	146
Arctic seas, astronomical observations—E. K. KANE	
Arctic seas, magnetical observations—E. K. KANE	
Arctic seas, meteorological observations—E. K. KANE	
Arctic seas, meteorological observations—F. L. McClintock	
Arctic seas, physical observations in—I. I. HAYES	
Arctic seas, physical observations in—E. K. KANE	
Arctic seas, tidal observations—E. K. KANE	
Arctic temperature—I. I. HAYES	
Arctic winds—I. I. HAYES	
Argentina syrtensium, new deep-sea fish-Goode and Bean. (P. 1878)	
Argentine Republic, astronomical observatory—B. A. Gould. (R. 1873)	
Ariopsis Milberti, breeding habits of—N. T. Lupton. (P. 1878)	
Arizona, ancient ruin in—J. C. Y. LEE. (R. 1872)	
	442
Arizona, Pima county, ruins in—R. T. Burr. (R. 1879)	34 5
Arizona, Pima Indians of—F. E. Grossman. (R. 1871)	
Arizona, Tonto Apaches of—C. SMART. (R. 1867)	
Arkansas, cretaceous fossils from—C. A. WHITE. (P. 1881)	
Arkansas, earthworks in—Mrs. G. KNAPP. (R. 1877)	
Arkansas, invertebrate fossils from—C. A. WHITE. (P. 1880)	
Arkansas, meteorological observations near Washington-N. D. SMITH	
Arkansas, Phillips county, ancient pottery—J. H. DEVEREUX. (R. 1872)	
Armstrong, T. Mounds in Winnebago county, Wisconsin. (R. 1879)	
Arrangement of families of birds—S. F. BAIRD	
Arrangement of families of fishes—T. GILL	
Arrangement of families of mammals—T. GILL	
Arrangement of families of mollusks—T. GILL	
Arrangement of mineralogical collection—C. U. Shepard. (R. 1861)	
Arrangement of the materia medica collection-J. M. FLINT. (P. 1881) 467,	
	325
Arrow-heads, deposit of, near Fishkill, New York-E. M. SHEPARD. (R. 1877)	323
Arrow-heads, Indian mode of making and obtaining fire—G. CROOK. (R. 1871)	249
Arrows, lectures on wounds of poisoned—D. Brainard. (R. 1854)	
Arsenic acid for protecting anatomical preparations—J. B. S. JACKSON. (P. 1878)	
Art, on tables of constants of nature and—C. BABBAGE. (R. 1856)	
Art Gallery, Corcoran, deed of gift of. (R. 1872) 271,	
Art Gallery, Corcoran, deposits in, by Smithsonian Institution. (R. 1874)	

Art Gallery, Corcoran, report of Committee of Regents on. (R. 1872) 271, 329
Arteries, reparatory inflammation in. Toner lecture No. VII-E. O. SHAKE-
SPEARE 321
Artificial shell deposits in New Jersey—C. RAU. (R. 1864) 188, 440, 362
Artificial shell deposits of the United States-D. G. BRINTON. (B. 1866) 214
Artisan and artist, identification of the-N. WISEMAN. (R. 1870) 244
Artisans, scientific education of mechanics and—A. P. PEABODY. (R. 1872) 271, 380
Arts, chemical, recent improvements in-J. C. Booth; C. Morfit
Ascelicthys rhodorus—Jordan and Gilbert. (P. 1880) 425
Ascensions—See Balloons.
Assay of coins at mint of the United States-J. Pollock. (R. 1868) 224
Asteroid supplement to new tables—J. D. Runkle 94
Asteroids between Mars and Jupiter—D. Kirkwood. (R. 1876) 299
Astronomical discoveries, telegraphic announcements of-J. Henry 263
Astronomical observations—W. HARKNESS
Astronomical observations, Arctic—I. I. HAYES
Astronomical observations, discussion of Piazzi's—B. A. Gould. (R. 1868) 187
Astronomical observations in the Arctic Seas—E. K. KANE129
Astronomical observatories, reports of—E. S. Holden; G. H. Boehmer.
(R. 1880)
Astronomical observatories, reports of, 1879—E. S. Holden. (R. 1879) 345, 410
Astronomical observatories, reports of, 1880—E. S. HOLDEN; G. H. BOEHMER.
(R. 1880)
Astronomical observatory at Cordoba, Argentine Republic, account of—B. A.
Gould. (R. 1878)
Astronomical photography, progress of—Dr. Lee. (R. 1861) 149
Astronomical tables—J. D. RUNKLE
Astronomical telegram circular—J. HENRY 263
Astronomy, lecture on—A. CASWELL. (R. 1858) 109
Astronomy, progress of, in 1879 and 1880—E. S. Holden. (R. 1880) 442, 427
Astronomy, recent progress in—E. S. Holden. (R. 1880) 442, 427
Astroscopus, species of, in Eastern United States-T. H. BEAN. (P. 1879) 388
Atlantic coast, microscopic examination of soundings-J. W. BAILEY 20
Atlantic, Lowe's aeronautic voyage across. (R. 1860) 147
Atlas, physical, of North America, proposed—G. Gibbs. (R. 1866) 214
Atmosphere, currents of the, and aerial navigation—J. HENRY. (R. 1860) 147
Atmosphere, diminution of aqueous vapor with altitude—J. HANN. (R. 1877) 323, 898
Atmosphere, plan of research upon—C. M. WETHERILL. (R. 1866) 214
Atmosphere, vegetation and the—J. Jamin. (R. 1864) 188
Atmospheric circulation over the earth, laws of—J. H. Coffin 268
Atmospheric currents, relation between barometric variations and—M. PESLIN.
(R. 1877)

Atmospheric electricity—M. F. Duprez. (R. 1858)	109
Atmospheric pressure and rain fall—J. HANN. (R. 1877)	, 398
Atmospheric pressure, Arctic—I. I. HAYES	196
Atmospheric pressure, Arctic—E. K. KANE	104
Atmospheric pressure, Arctic—F. L. McCLINTOCK	146
Atomic volume, bibliography of—F. W. CLARKE	2.5)
Atomic weight determinations—G. F. BECKER	358
Atomic weights, recalculation of—F. W. CLARKE	441
Atoms—J. Herschel. (R. 1862)	150
Atthis Ellioti, a humming bird from Guatemala—R. RIDGWAY. (P. 1878)	302
Aurora and electricity—S. Lemström; A. A. De La Rive. (R. 1874)	286
Aurora, connection of gales of wind and—R. T. KNIGHT. (R. 1871)	249
Aurora, directions for observing—J. HENRY. (R. 1855)	77
Aurora, effect of, on magnetic declination—A. D. BACHE	175
Aurora, influence of, on the telegraph—W. D. SARGENT. (R. 1870)	244
Aurora, instructions for observations of—Toronto Observatory	148
Aurora, magnetic effect of—A. D. BACHE	175
Aurora, map of stars for observations on	35 0
Aurora, observations on—S. Masterman. (R. 1857)	107
Aurora, or Polar light, its phenomena and laws—E. Loomis. (R. 1865)	209
Aurora, phenomena in telegraphic lines during—G. B. Donatt. (R. 1872)	271
Aurora, recent secular period of—D. Olmsted	81
Auroras, articles on, received by the Institution. (R. 1871; R. 1874) 249,	236
Auroras, records of, in high northern latitudes—P. Force	84
Autobiography, the history of my youth—F. ARAGO. (R. 1870)	244
Auxis Rochei, frigate mackerel, on New England coast—G. B. Goode. (P. 1880)	425
Awards to Smithsonian Institution at the Centennial Exhibition. (R. 1876; R. 1878)	341
Axolotl changed to amblystoma—A. Weismann. (R. 1877) 323,	401
AYRES, Dr. Notes on fishes. (P. 1880)	425

B.

BABBAGE, C., extracts from writings of. (R. 1873)	275
BABBAGE, C., memoir of, by N. S. Dodge. (R. 1878)	275
BABBAGE, C. On constants of nature and art. (R. 1856)	91
BABINET, J. The diamond and other precious stones. (R. 1870) 244,	377
BABINET, J. The northern seas. (R. 1869)	228
BACHE, A. D.—	
Bequest to National Academy of Sciences. (R. 1872) 271	, 329
Discussion of magnetic and meteorological observations. Part I	113
Discussion of magnetic and meteorological observations. Part 11	121
Discussion of magnetic and meteorological observations. Part III	132
Discussion of magnetic and meteorological observations. Pts. IV, V, VI	162
Discussion of magnetic and meteorological observations. Pts. VII, VIII, IX.	175
Discussion of magnetic and meteorological observations. Pts. x, x1, x11	186
Discussion of magnetic and meteorological observations. Parts I-XII	195
Eulogy on James A. Pearce. (R. 1862)	150
Lecture on Switzerland. (R. 1870)	244
Magnetic survey of Pennsylvania	166
Notice of James P. Espy. (R. 1859)	110
Notice of G. Würdemann. (R. 1859)	110
BACHE, A. D., eulogy on, by J. HENRY. (R. 1870) 244,	379
BACHE, A. D., list of scientific papers of—B. A. Gould. (R. 1870) 244, 329	, 376
BACHE, G. M. Hail storm in Texas. (R. 1870)	244
Bacteria, putrefactive, action of dry heat and sulphurous acid upon. (P. 1881)	467
BAEGERT, J. Aboriginal inhabitants of California peninsula. (R. 1868; R.	
1864) 187, 188, 440,	361
Baffin's Bay, meteorological observations in—F. L. McCLINTOCK	146
BAILEY, J. W.—	
Algæ	28
Diatomacea 2	8, 68
Infusoria	28
Microscopical examination of soundings, Atlantic coast	20
Microscopical observations in South Carolina, Georgia, Florida	23
New species and localities of microscopical organisms	63
BAILEY, W. B. F. Antiquities of Spalding county, Georgia. (R. 1877)	323
Baird, S. F.—	
•Alphabetical index of genera of birds	108
Arrangement of families of birds	
Catalogue of North American birdsoctavo, 108; quarto,	106
Catalogue of North American mammals	105

BAIRD, S. F.—Continued.
Circular concerning department of insects. (P. 1881) 467, 446
Circular in reference to building-stone collection. (P. 1881) 467, 453
Circular in reference to petroleum collections. (P. 1881) 467, 447
Circular in reference to shipping fresh fish and other animals 384
Circular relative to scientific and literary exchanges 329
Circular requesting material for the library of the National Museum. (P. 1881) 467, 458
Circular to friends of the Museum. (P. 1881) 467, 446
Directions for collecting, preserving, and transporting specimens of natu-
ral history. (R. 1856) 91, 34
Fishes on the coasts of New Jersey and Long Island. (R. 1854) 75, 348
Instructions for collecting insects. (R. 1858)
Instructions for collecting nests and eggs of North American birds. (R. 1858)
List of explorations furnishing collections to National Museum, 1838 to 1877. (R. 1877) 32
List of sources from which specimens have been derived, 1888 to 1867. (R. 1867)
Memoranda of inquiry relative to food fishes 231
Museum miscellanea
Notes on aboriginal shell-mounds on coast of New Brunswick and of
New England. (P. 1881) 467
Notes on Henry's researches on sound. (R. 1878) 341, 406
On reports of recent scientific progress. (R. 1880)
Questions relative to food fishes 234
Register of periodical phenomena65, 148
Report of Secretary for 1878. (R. 1878) 341, 408
Report of Secretary for 1879. (R. 1879) 845, 418
Report of Secretary for 1880. (R. 1880) 442, 419
Report on Centennial Exhibition. (R. 1876) 299, 307
Report on and statistics of British Museum. (R. 1850)
Report on plan of exhibit at the International Exhibition at Philadelphia. (R. 1875)
Reports on publications, exchanges, explorations, and Museum, 1851-1877. (R. 1851-1877) 51, 57, 67, 75, 77, 91, 107, 109, 110, 147, 149, 150, 187, 188 209, 214, 215, 224, 228, 244, 249, 271, 275, 286, 298, 299, \$2,
Review of American birds. Part I. North and Middle America 181
Scientific instructions to Captain Hall. (R. 1871)249
Scientific investigations in Russian America. Natural history 207
BAIRD, S. F.; GIRARD, C. Catalogue of North American reptiles. Part I 49
Balfour, J. The wind and fog. (R. 1866)
Balloon ascensions, account of—J. GLAISHER. (R. 1868)

•
Balloon ascensions—See Aeronautics.
BALTZELL, J. Winds in Florida. (R. 1866)
BANCBOFT, G. Memorial to Congress for new Museum building. (R. 1876) 2
BANCROFT, G. On publication of Spanish works on New Mexico. (R. 1855)_
BANNISTER, H. M. Climate of Alaska. (R. 1866)
Bannister, H. M. Formation of clouds over Gulf stream. (R. 1866)
Baptist Missionary Society. Yoruba grammar and dictionary
Barbuda, catalogue of Ober's collection of birds of—G. N. LAWRENCE. (P. 1878)
BARKER, G. F. Recent progress in chemistry. (R. 1880) 442, 4
BARKER, G. F. Recent progress in physics. (R. 1880) 442, 4
BARNARD, F. A. P. Lectures on the undulatory theory of light. (R. 1862) 1
BARNARD, J. G. Eulogy on Gen. Joseph G. Totten. (R. 1865)
BARNARD, J. G. Internal structure of the earth 3
BARNARD, J. G. Problems of rotary motion2
BARNARD, V. Birds of Chester county, Pennsylvania. (R. 1860)
Barometer, account of Green's. (R. 1855) 77,
Barometer, Green's standard
Barometer, horary variations of the—F. VAILLANT; J. HENRY. (R. 1866) 2
Barometer, influence of rain upon the—J. HANN. (R. 1877) 823, 8
Barometer, rain and snow gauges—R. H. GARDINER. (R. 1858)
Barometer tables, on construction of—F. F. Tuckett. (R. 1867)
Barometer tube breaking suddenly—L. F. WARD. (R. 1866)
Barometer tubes, on filling—J. Green; W. Würdemann. (R. 1859) 1
Barometers at the Kew Observatory, verification of—J. WELSH. (R. 1859) 1
Barometers, experiments on aneroid—B. STEWART. (R. 1868)
Barometric minima and rainfall—J. T. REYE. (R. 1877)
Barometric observations, discussion of Snell's—F. H. Loud. (R. 1880) 442, 45
Barometric observations, discussion of Shell's—F. H. Houd. (10. 1000) == 412, 20
Barometric variations and atmospheric currents, relation between—M. PESLIN.
(R. 1877) 328, 8
Barometrical tables—A. GUYOT
,
BARRANDT, A. Haystack mound, Dakota. (R. 1872)
BARTLETT, J. R. On publication of Squier and Davis' Ancient Monuments.
(R. 1847)
BARTRAM, J. Antiquities of Florida. (R. 1874)
Base chart of the United States—C. A. SCHOTT
Bases, ammonia-cobalt, researches on—W. GIBBS; F. A. GENTH
Bases, organic—A. BAUER. (R. 1872)
Batavian Society of Experimental Philosophy, Rotterdam. Prize questions.
(R. 1861)

Bath and West of England Society for encouragement of Agriculture, Arts, etc. Exchanges. (R. 1867) 215
Batis maritima—J. Torrey
Batrachia, North American—E. D. Cope
Bats, account of remarkable accumulation of—M. FIGANIERRE. (R. 1863) 187
Bats of North America, monograph of—H. ALLEN 165
BAUER, A. Organic bases. (R. 1872) 271
Bay of Fundy, marine invertebrata of—W. STIMPSON
Bead, colored, from a mound at Black Hammock, Florida—A. M. HARRISON. (R. 1877)
Bead, polychrome, from Florida-S. S. HALDEMAN. (R. 1877) 323, 404
BEAN, T. H.—
Check-list of fishes distributed by Smithsonian Institution. (P. 1880) 425
Description of a new hake (Phycis Earlii) and note on Phycis regius. (P. 1880)
Description of new fish from Alaska, Anarrhichas lepturus, and notes on genus. (P. 1879)
Description of new sparoid fish, Sargus Holbrookii, from Savannah bank. (P. 1878)
Description of new species of Amiurus from the Mississippi. (P. 1879).
Description of new species of Gasterosteus from Schoodic Lakes, Maine. (P. 1879)
Description of species of Lycodes (L. Turneri) from Alaska. (P. 1878).
Descriptions of genera and species of Alaskan fishes. (P. 1879) 32
Descriptions of new fishes from Alaska and Siberia. (P. 1881) 467
Descriptions of new species of fishes (Uranidea marginata and Potamo-cottus Bendirei,) and of Myctophum crenulare. (P. 1881)
Descriptions of species of fishes collected by Dugès in Mexico. (P. 1879) 33
Directions for collecting and preserving fish. (P. 1881) 467, 464
Fishes of Arctic America. (Bull. 15, U.S. National Museum) 343
Identity of Euchalarodus Putnami with Pleuronectes glaber and notes on habits. (P. 1878)
List of European fishes in National Museum. (P. 1879) 32
Notes on collection of fishes from eastern Georgia. (P. 1879)
Notes on fishes collected by H. E. Nichols in British Columbia and Alaska, with descriptions of new species and new genus (Delolepis.) (P. 1881)
Notes on fishes from Hudson's Bay. (P. 1881)
Occurrence of Hippoglossus vulgaris in Alaska. (P. 1879)
Occurrence of Stichœus punctatus at St. Michael's, Alaska. (P. 1878) 3
Partial bibliography of fishes of Pacific coast of United States and Alaska for 1880. (P. 1881)
Preliminary catalogue of fishes of Alaskan waters. (P. 1881) #:
Species of Astroscopus of Eastern United States. (P. 1879)

BEAN, T. H.; GOODE, G. B.—	
Catalogue of Stearns's fishes of Florida and descriptions of new species. (P. 1879)	388
Catalogue of Velie's fishes in Gulf of Mexico and description of new species. (P. 1879)	388
Craig flounder of Europe (Glyptocephalus cynoglossus) on coast of North	
• America. (P. 1878)	382
Description of Alepocephalus Bairdii, new deep-sea fish, western Atlantic. (P. 1879)	888
Description of Argentina syrtensium, new deep-sea fish, Sable Island bank. (P. 1878)	882
Description of Caulolatilus microps from Gulf coast of Florida. (P. 1878)	832
Description of Lycodes paxillus. (P. 1879)	388
Description of new genus and species of fish, Lopholatilus chamæleonticeps, from New England. (P. 1879)	
Description of new species of fish, Apogon pandionis, from mouth of Chesapeake Bay. (P. 1881)	•
Description of new species of fish (Lutjanus Blackfordii and Lutjanus Stearnsii) from coast of Florida. (P. 1878)	8 8 2
Descriptions of gadoid fishes, Phycis chesteri and Haloporphyrus viola, from northwestern Atlantic. (P. 1878)	882
Identity of Brosmius brosme Americanus, Gill, with Brosmius brosme (Müller) White. (P. 1878)	882
Identity of Rhinonemus caudacuta with Gadus cimbrius. (P. 1878)	832
New genus of fishes, Benthodesmus. (P. 1881)	4 67
New serranoid fish, Epinephelus Drummond-Hayi, from Bermudas and Florida. (P. 1878)	882
New species of Liparis (L. ranula) from Halifax. (P. 1879)	888
Note on Platessa ferruginea and Platessa rostrata. (P. 1878)	882
Note upon black grouper (Epinephelus nigritus) of the Southern coast. (P. 1878)	382
Occurrence of Lycodes Vahlii on La Have and Grand Banks. (P. 1879.)	3 38
Oceanic bonito on coast of United States. (P. 1878)	882
BEAUCHAMP, W. M. Wampum belts of the Six Nations. (R. 1879)	3 4 5
Beaufort, (N. C.,) library, deposit of, by E. M. Stanton. (R. 1862)	150
Beaufort, N. C., notes on fishes of—D. S. JORDAN. (P. 1880)	425
Beaufort harbor, notes on fishes of—D. S. JORDAN; C. H. GILBERT. (P. 1878)	832
BEAUMONT—See De Beaumont.	
Beautemps-Beaupré, C. F., memoir of, by Elie de Beaumont. (R. 1863)	187
Beaver, on the habits of the—Felix R. Brunot. (R. 1878)	275
Beaver Island, Mich., natural history of-J. J. STRANG. (R. 1854)	75
BECKER, G. F. Atomic weight determinations	358

BECQUEREL, A. E. Forests and their climatic influence. (R. 1869) 228
BECQUEREL, A. E. Preservation of copper and iron in salt water. (R. 1864.) 188
Beirut, Syria, account of sarcophagus from—A. A. HARWOOD. (R. 1870) 244
BELDING, L. Additions to North American bird fauna by—R. RIDGWAY. (P. 1881) 467
Belding, L. Partial list of birds of central California. (P. 1878)
Belgium, Archæological Academy of. Archæological congress organized by. (R. 1866)
Belgium—See Prize questions.
Belgium, Society of Science, etc.—See Prize questions.
Belone exilis, generic relations of-D. S. JORDAN; C. H. GILBERT. (P. 1880) 425
Belone latimanus, occurrence of, in Buzzard's bay-G. B. Goods. (P. 1878) 32
BENDIRE, C. Notes on Salmonida of Columbia river. (P. 1881) 4.
Benthodesmus, a new genus of deep-sea fishes-G. B. Goode; T. H. Bran.
(P. 1881) 467
Bequest of A. D. Bache to National Academy of Sciences. (R. 1872) 271, 329
Bequest of S. Habel. (R. 1879)
Bequest of J. Hamilton to Smithsonian Institution. (R. 1872; R. 1873) 271, 273
Bequest of Smithson, act of Congress, July 1, 1886, accepting. (R. 1853) 67. M
Bequest of Smithson, exposition of—JHENRY
Bequest of Smithson, history of 328. 330
Bequest, Smithson and his-W. J. RHEES. (R. 1879) 345, 330
BEBENDT, C. H.—
Collection of historical documents in Guatemala. (R. 1876) 299
Explorations in Central America. (R. 1867) 215
On the Carib language and people. (R. 1878) 275
recommendation of, by British Minister. (R. 1865) 209
recommendation of, by Guatemalan Minister. (R. 1865) 209
recommendation of, by L. Molina, Costa Rican Minister. (R. 1865) 209
BERLANDIER, L. Catalogue of collection of historical and geographical manu-
scripts, maps, &c., of Mexico. (R. 1854) 75
Berlin Fisheries Exhibition, additions to museum from. (R. 1880) 442
Berlin Fisheries Exhibition, catalogue of United States' exhibit at—G. B. Goode 413
Bermuda, birds of—T. BLAND; J. R. WILLIS. (R. 1858)
Bermudas, catalogue of fishes of—G. B. Goode 296
Bermudas, fishes from, mistakenly described as new by Günther-G. B. Goods.
(P. 1878) 332
Bermudas, new serranoid fish from—G. B. Goode; T. H. BEAN. (P. 1878) 332
Bern Museum, request for bison. (R. 1865) 200
Beroids, relation of lucernarians to—H. J. CLARK242

BERRIEN, J. McPherson. Construction of act establishing Smithsonian. (R. 1858)67
BERTHOUD, E. L. Ancient remains in Colorado. (R. 1867) 215
BERTHOUD, E. L. Antiquities of Colorado. (R. 1871) 249
BERTHRAND, M. Life and works of Kepler. (R. 1869) 228
Best hours for temperature observations—C. DEWEY. (R. 1860) 147
BETHUNE, C. J. Acknowledgment of books. (R. 1861) 149
Bibliographia Americana Historico-Naturalis for 1851—C. GIRARD 48
Bibliographia Americana, prospectus of a-H. STEVENS. (R. 1848)
Bibliographical index of North American Botany—S. WATSON 258
Bibliographical sketches of vestiges of antiquity in United States—S. F. HAVEN 71
Bibliography of alge—H. C. Wood 241
Bibliography of American natural history for 1851—C. GIRARD 48
Bibliography of anthropology-O. T. MASON. (R. 1880) 442, 432
Bibliography of atomic volume and specific gravity, expansion, and boiling and
melting—F. W. CLARKE 255
Bibliography of Chinook jargon—G. GIBBS
Bibliography of Dakota language—S. R. Riggs 40
Bibliography of diptera—C. R. OSTEN SACKEN 270
Bibliography of diseases of joints, bones, larynx, eye, gangrene, etc.—W. W. KEEN
Bibliography of expansion—F. W. CLARKE 289
Bibliography of fishes—T. GILL247
Bibliography of fishes of Pacific coast of United States and Alaska for 1880—
T. H. BEAN. (P. 1881) 467
Bibliography of fishes of Pacific coast of United States-T. GILL 463
Bibliography of mammals—T. GILL 280
Bibliography of mollusks—T. GILL 227
Bibliography of nebulæ, clusters, milky way, nebular hypothesis, etc.—E. S. HOLDEN
Bibliography of North American conchology previous to 1860. Part 1. American authors—W. G. BINNEY
Bibliography of North American conchology previous to 1860. Part II. Foreign authors—W. G. BINNEY
Bibliography of ornithology: faunal publications relative to British birds—E. Cours. (P. 1879)
Bibliography of orthoptera—S. H. SCUDDER 189
Bibliography of rattlesnakes, serpent venom, etc.—S. W. MITCHELL 185
Bibliography of specific heat—F. W. CLARKE 276
Bibliography of Yoruba language—T. J. Bowen 98
Bibliography, plan of—J. FRIEDLÄNDER. (R. 1858)
Biela's comet, investigations of—J. S. HUBBARD. (R. 1862) 150
Big elephant mound in Grant county. Wiscopsin—J. WARNER. (R. 1872) 271

BINNEY, W. G.—	
Bibliography of North American conchology previous to 1860. Part I. American authors	142
Bibliography of North American conchology previous to 1860. Part II. Foreign authors	174
Check-list of shells of North America	128
Land and fresh-water shells of North America. Part I	194
Land and fresh-water shells of North America. Part II	143
Land and fresh-water shells of North America. Part III	144
Biographical memoir of Joseph Henry-A. GRAY. (R. 1878) 341, 856,	407
Biographical notice of L. Agassiz-E. FAVRE. (R. 1878)	341
Biographical notice of L. Agassiz-J. A. GARFIELD. (R. 1873)	277
Biographical notice of L. Agassiz-P. PARKER. (R. 1873)	275
Biographical notice of S. P. Chase—J. A. GARFIELD. (R. 1878)	475
Biographical notice of S. P. Chase—H. HAMLIN. (R. 1873)	27.0
Biographical notice of W. H. Harvey-A. GRAY. (R. 1867)	215
Biographical notice of Charles C. Jewett-R. A. Guild. (R. 1857)	215
Biographical sketch of Dom Pedro II-A. FIALHO. (R. 1876)	299
Biography of Condorcet—F. ARAGO. (R. 1878)	341
Biography—See Memoirs.	
Bird, a new humming, from Guatemala—R. RIDGWAY. (P. 1878)	332
Birds, acknowledgment of—Academy of Sciences, Stockholm. (R. 1867)	215
Birds, acknowledgment of—J. Gould. (R. 1867)	215
Birds, acknowledgment of—Hungarian National Museum. (R. 1863)	157
Birds, alphabetical index of genera of—S. F. BAIRD	108
Birds, American, instructions for collecting nests and eggs of—S. F. BAIRD. (R. 1858) 109.	139
Birds and mammals of Missouri river, list of—E. HARRIS. (R. 1850)	
Birds, arrangement of families of—S. F. BAIRD	210
Birds, bibliography of British—E. Cours. (P. 1879)	333
Birds, catalogue of North American—S. F. BAIRD. Octavo	106
Birds, catalogue of North American—S. F. BAIRD. Quarto	106
Birds, catalogue of Old World, in Nat. Museum -R. RIDGWAY. (P. 1881) 467.	462
Birds, circular relative to collections of	168
Birds, dates of first appearance of—F. B. Hough	182
Birds, description of new species of, from Costa Rica—R. RIDGWAY. (P. 1878)	332
Birds, descriptions of new-R. RIDGWAY. (P. 1881)	467
Birds, descriptions of several new species and races of, in National Museum— R. RIDGWAY. (P. 1878)	9 3-1
Birds, desiderata among North American—R. RIDGWAY. (P. 1881)	
Birds' eggs, suggestions for forming collections of—A. NEWTON	150
Birds, humming, list of—D. G. Elliot	

OF SMITHSONIAN PUBLICATIONS.

Birds, list of described, of Mexico, Central America, and the West Indies, not in Smithsonian collections
Birds' nests and eggs, instructions for collecting
Birds, new species and races of American, and synopsis of genus Tyrannus—R. BIDGWAY. (P. 1878)
Birds, new species of Turdidæ, from Dominica—G. N. LAWRENCE. (P. 1880)
Birds; nomenclature and catalogue of-R. RIDGWAY
Birds. North American oölogy—T. M. Brewer.
Birds, notes on some Costa Rican—R. RIDGWAY, (P. 1881)
Birds of Antigua, catalogue of Ober's collection of-G. N. LAWRENCE. (P. 1878)
Birds of Arctic America—L. Kumlien
Birds of Barbuda, catalogue of Ober's collection-G. N. LAWRENCE. (P. 1878)
Birds of Bermuda-T. BLAND; J. R. WILLIS. (R. 1858)
Birds of Caribbee Islands—F. A. OBER. (R. 1878)
Birds of Central California-L. BELDING. (P. 1878)
Birds of Chester county, Pennsylvania-V. BARNARD. (R. 1860)
Birds of District of Columbia, list of—E. Coues; S. S. Prentiss. (R. 1861)
Birds of Dominica, catalogue of Ober's collection of—G. N. LAWRENCE. (P. 1878)
Birds of Grenada, catalogue of Ober's collection-G. N. LAWRENCE. (P. 1878)
Birds of Guadeloupe, catalogue of Ober's collection—G. N. LAWRENCE. (P. 1878)
Birds of Heligoland-H. GÄRKE. (P. 1879)
Birds of Kerguelen Island-J. H. KIDDER; E. COURS
Birds of Lesser Antilles, catalogue of Ober's collection of—G. N. LAWRENCE, (P. 1878)
Birds of Martinique, catalogue of Ober's collection of—G. N. LAWRENCE. (P. 1878)
Birds of Middle and South America not in National Museum—R. RIDGWAY. (P. 1881)
Birds of Missouri river—E. HARRIS. (R. 1850)
Birds of North America, nomenclature and catalogue of—R. RIDGWAY. (P. 1880)
Birds of Nova Scotia—Blackiston; T. Bland; J. R. Willis. (R. 1858)
Birds of Saint Vincent, catalogue of Ober's collections of—G. N. LAWRENCE. (P. 1878)
Birds of South America, circular relative to collections of
Birds of Southern Texas—J. C. MERBILL. (P. 1878)
Birds of Southwestern Mexico, collected by F. E. Sumichrast—G. N. LAW-
Birds of West coast, migrations and nesting habits of—J. G. Cooper. (P. 1879)
Birds of West Indies, catalogue of Ober's collections from—G. N. LAWRENCE. (P. 1878)
Birds, review of American. Part I—S. F. BAIRD
Birds, revisions of nomenclature of—R. RIDGWAY. (P. 1880)

Birds, suggestions for saving skeletons of—A. NEWTON. (R. 1860)
Birds, systematic review of classification of—W. LILLJEBORG. (R. 1865.) 209, 3
Birds, two new North American, added by L. Belding—R. RIDGWAY. (P. 1881)
BISHOP, W. D. Commissioner of Patents, Meteorological observations I
Bison, request of Bern Museum for. (R. 1865)
Bites of serpents, nature and cure of-D. Brainard. (R. 1854)
Black bass of the Ohio, habits of the-J. Eoff. (R. 1854)
Black Hammock, colored bead from—A. M. HARRISON. (R. 1877)
BLACKISTON, -; BLAND, T.; WILLIS, J. R. Birds of Nova Scotia. (R. 1858)
Blackmore Museum, Salisbury, England, notice of the. (R. 1868)
Black Mountain, North Carolina, topography of-T. L. CLINGMAN. (R. 1855)
BLAINVILLE, D. DE, memoir of-M. FLOURENS. (R. 1865)
BLAND, T.; BINNEY, W. G. Land and fresh-water shells of North America. Part I.—Pulmonata geophila
BLAND; WILLIS, J. R. Birds of Bermuda. (R. 1858)
Bland, T.; Blackiston, —; Willis, J. R. Birds of Nova Scotia. (R. 1858)
Blank check-lists, labels, figures, &c
Blind deaf mute, vocal sounds of Laura Bridgman, the—F. LIEBER
Blodget, L. List of meteorological observers. (R. 1853)
Blondel, S. Historical, archæological, and literary study of jade. (R. 1876)
Blood, analysis of—J. Jones
Blossoming of plants, dates of—F. B. Hough
Blowpipe apparatus of Hawkins and Wale. (R. 1872)
Blowpipe, qualitative determination by the—T. EGLESTON. (R. 1872)
BLYDEN, E. D. Mixed races in Liberia. (R. 1870)
Board of Commissioners for Foreign Missions—See American Board, etc.
Board of Regents, Journals of. (See each annual report) 32
Board of Regents—See Regents.
BOEHMER, G. H. Index to anthropological articles in Smithsonian publications. (R. 1879) 345,
BOEHMER, G. H. Report on International Exchange
BOEHMER, G. H. Reports of astronomical observatories, 1880. (R. 1880) 442,
BOEHMER, G. H.; HOLDEN, E. S. Reports of observatories, 1880. (R. 1880) 442,
BOERNER, C. G. Lightning discharges. (R. 1867)
Boiling point, bibliography of—F. W. CLARKE
Boiling points, tables of—F. W. CLARKE 255
Bolles, E. C. Account of Portland Society of Natural History. (R. 1867)
Bologna, Academy of Sciences of the Institute of. Prize questions. (R. 1862)
Bones, bibliography of diseases of the—W. W. KEEN
Bonito, oceanic, description of—V. N. EDWARDS. (P. 1878)
Bonito, oceanic, on coast of United States-G. B. Goode; T. H. Bran. (P. 1878)

OF SMITHSONIAN PUBLICATIONS.	147
Books, acknowledgment of—C. J. BETHUNE. (R. 1861)	149
Books, acknowledgment of—J. A. Codd. (R. 1861)	149
Books, acknowledgment of-M. LABOULAYE. (R. 1867)	215
Books, copyright, from 1846 to 1849—C. C. JEWETT. (R. 1850)	28
Books on Brazil, presented—M. M. Lisboa. (R. 1865)	209
Books on Egypt, presentation of-R. Lepsius. (R. 1860)	147
Books, on the classification of—J. P. LESLEY. (R. 1862)	150
Books, presentation of—Duke of Northumberland. (R. 1859)	110
Books, presented—Imperial Library of Vienna. (R. 1865)	209
BOOTH, J. C.; MORFIT, C. Recent improvements in the chemical arts	27
Bordeaux, Chamber of Commerce of. Exchange of publications. (R. 1863)	187
Bordeaux, Imperial Academy of Sciences. Prize questions. (R. 1868)	224
Bosphorus, account of a hailstorm on the—Com. Porter. (R. 1870)	244
Boston, Mass., American Academy—See American Academy.	
Botanical explorations in New Mexico and California—A. GRAY. (R. 1849)	21
Botany, index to North American—S. WATSON	258
Botany of Hawaiian and Fanning Islands and California—T. H. STREETS	808
Botany of Kerguelen Island-A. GRAY; T. P. JAMES; E. TUCKERMAN; W.	
G. FARLOW	294
Botany, progress in, in 1879 and 1880—W. G. FARLOW. (R. 1880) 442, Botany—See Plants.	430
Boundary line between geology and history—E. Surss. (R. 1872)	271
Bown, T. J. Yoruba grammar and dictionary	98
Bowers, S. Explorations in Santa Rosa Island, California. (R. 1877)	828
Boyd, C. H. Remains of walrus (?) in Maine. (P. 1881)	467
Brachiopoda, index of—W. H. DALL	304
Brachyopsis verrucosus, from California-W. N. Lockington. (P. 1880)	425
Brachyopsis xyosternus—D. S. Jordan; C. H. GILBERT. (P. 1880)	425
Brackett, A. G. Shoshone or Snake Indians, religion, superstitions, manners, etc. (R. 1879)	345
Brackett, A. G. Sioux or Dakota Indians. (R. 1876)	299
Brain and marrow of animals, uses of, among Indians—T. R. PEALE. (R. 1870)	244
Brain, dual character of the—C. E. Brown-Séquard. Toner lecture No. II	
Brainard, D. Lecture on bites of serpents and wounds of poisoned arrows. (R. 1854)	75
Bransford, J. F. Archeological researches in Nicaragua	383
Bransford, J. F., note on shells from Costa Rica kitchen-midden, collected by—W. H. Dall. (P. 1878)	832
Bravais, Auguste, memoir of, by E. De Beaumont. (R. 1869)	228
Brayton, A. W. Distribution of the fishes of South Carolina, Georgia, and Tennessee	308
Brayton, A. W. North American ichthyology	808

Brazil, books on, presented—M. M. Lisboa. (R. 1865)
Brazil, specimens from, presented. Brazilian Nat. Hist. Museum. (R. 1865) 2%
Brazilian Nat. Hist. Museum, specimens from Brazil presented. (R. 1865) 209
Breed, E. E. Earthworks in Wisconsin. (R. 1872)
Breed, E. E. Pits at Embarrass, Wisconsin. (R. 1877) 325
Breeding habits of sea-catfish—N. T. Lupton. (P. 1878) 83
Bremen, government of, exchange system. (R. 1865) 29
Brenndecke, F. Meteorites. (R. 1869)
Brevoortia patronus, note on—S. STEARNS. (P. 1878)
Brevoortia, revision of American species of—G. B. Goode. (P. 1878) 325
Brewer, T. M. Instructions for collecting and preserving nests and eggs 135
Brewer, T. M. Nests and eggs of North American species of empidonaces. (P. 1879)
Brewer, T. M. North American oölogy. Part 1. Raptores and Fissirostres. 85
Brezina, A. Explanation of principles of crystallography and crystallophysics. (R. 1872)
Bridges, lectures on roads and—F. Rogers. (R. 1860; R. 1861) 147, 147
BRIDGMAN, L., vocal sounds of—F. LIEBER
BRINTON, D. G. Artificial shell deposits in the United States. (R. 1866) 218
British America, account of the Indians of—E. Petitot. (R. 1865) 35
British America, Tinneh or Chepewyan Indians of-G. GIBBS. (R. 1866.) 214. 365
British Association, report to, on radiant heat—B. Powell. (R. 1859) li
British birds, list of faunal publications relative to—E. Cours. (P. 1879) \$\frac{35}{2}\$
British Columbia, Haidah Indians of-J. G. SWAN 267
British Columbia, notes on fishes of—T. H. BEAN. (P. 1881) 47
British Minister, F. W. A. Bruce, recommending Dr. Berendt. (R. 1865) 26
British Museum, acknowledgment of specimens. (R. 1865) 36
British Museum, electrotypes of shells granted by. (R. 1863)
British Museum, notes on typical American fishes in—D. S. JORDAN. (P. 1879) 37
British Museum, statistics of—S. F. BAIRD. (R. 1850)
British Provinces of North America, list of libraries, institutions, and societies in—W. J. Rhees
Broadhead, G. C. Prehistoric evidences in Missouri. (R. 1879)
Broca, P. History of Anthropological Society of Paris, 1865 to 1867. (R. 1868)
Broca, P. Troglodytes, or cave dwellers, of the valley of the Vézère. (R. 1872)
BRODNAX, B. H. Mounds in Moorehouse parish, Louisiana. (R. 1879) 345
Brosmius brosme Americanus, Gill, identity of, with Brosmius brosme (Müller) White—G. B. GOODE; T. H. BEAN. (P. 1878)
Brown, D. Shell heap in Georgia. (R. 1871)
Brown-Séquard, C. E. Dual character of the brain. Toner lecture No. II. 291
BRUFF, J. G. Indian engravings along Green river valley. (R. 1872) 277

CE DELIES ON LONG OF THE CONTROL OF	
SRUNOT, F. R. Habits of the beaver. (R. 1878)	275
Brunswick, Maine, meteorological observations—P. CLEAVELAND	204
BRUSH, G. J. Catalogue of meteorites in Yale College. (R. 1868)	224
Brussels, Belgium, Royal Academy of Sciences, Literature, and the Fine Arts. See Prize questions.	
BRYAN, O. N. Antiquities of Charles county, Maryland. (R. 1874)	286
Buch, L. von, memoir of, by M. Flourens. (R. 1862)	150
Budapest, address before Anthropological Congress at—F. F. Romen. (R. 1876)	392
Building Commission, National Museum, report of, for 1879. (R. 1879) 845,	409
Building Commission, National Museum, report of, for 1880. (R. 1880) 442,	434
Building Committee of Regents of S. I., journal and reports of, 1847-1867	829
Building Committee of Smithsonian, report of, on architecture	P
Building materials—R. D. Owen	P
Building materials, mode of testing—J. HENRY. (R. 1856)	91
Building materials, report on. (R. 1847)	329
Building, new, for museum, memorial to Congress for. (R. 1876)	299
Building, report on use of new hall in-L. Agassiz. (R. 1867)	215
Building-stone collection, circular in reference to-S. F. BAIRD. (P. 1881) 467,	453
Building stones, report on—D. D. OWEN. (R. 1847)	9, H
Buildings, public, acoustics applied to—J. HENRY. (R. 1856)	91
Buildings, warming and ventilating—A. Morin. Part 1. (R. 1878) 275,	439
Buildings, warming and ventilating—A. Morin. Part II. (R. 1874) 286,	
Bulletin National Museum—	
No. 1. Batrachia and reptilia—E. D. Cope	292
. No. 2. Birds of Kerguelen Island-J. H. KIDDER; E. COUES	293
No. 8. Oölogy, plants, mammals, etc., Kerguelen Island—J. H. KIDDER	
and others	
No. 4. Birds of Mexico—G. N. LAWBENCE	
No. 5. Catalogue of fishes of Bermudas—G. B. Goode	296
No. 6. Classification of animal resources of United States—G. B. GOODE.	297
No. 7. Natural history, Hawaiian Islands, etc.—T. H. STREETS	303
No. 8. Index of brachiopoda—W. H. Dall	304
No. 9. Review of Rafinesque's fishes—D. S. JORDAN	305
No. 10. Notes on cottidæ, etc.; synopsis of siluridæ—D. S. JORDAN	306
No. 11. Bibliography of Pacific fishes—T. GILL	463
No. 12. Fishes of South Carolina, Georgia, Tennessee; synopsis of catostomidæ—D. S. Jordan; A. N. Brayton	308
No. 13. Flora of St. Croix and Virgin Islands—H. F. A. EGGERS	313
No. 14. Catalogue Centennial exhibit, animal resources and fisheries— G. B. Goode	326
No. 15. Natural history and ethnology, Arctic America—L. Kumlien.	342
No. 16. (Not published.)	

Bulletin National Museum—Continued.

No. 17. Zoölogical position of Texas—E. D. Cope	412
No. 18. Fishery exhibit of United States at Berlin-G. B. Goode	413
No. 19. List of genera in zoölogy and paleontology—S. H. SCUDDER	47 0
No. 20. (Not published.)	
No. 21. Nomenclature of North American birds—R. RIDGWAY	422
No. 22. Guide to flora of Washington, D. CL. F. WARD	444
Bulletin of Philosophical Society of Washington, Vols. I, II, III	
Bulletin, Smithsonian, No. 1. Vocabulary of the jargon	
Bulletins of the U.S. National Museum, catalogue and index of	
Bureau of Ethnology, directory of officers and employés of	
Burial caves—J. Jones	
Burial ground, ancient, in Tompkins Co., N. YD. TROWBRIDGE. (R. 1863)	
Burial, modes of—J. Jones	
Burial of an Indian squaw, California—W. M. KING. (R. 1874)	
BURB, R. T. Ruins in Pima county, Arizona. (R. 1879)	
Business arrangements of the Smithsonian Institution	
BUTLER, A. P. Report of Senate Judiciary Committee on management of Smithsonian. (R. 1855)	
Buzzard's bay, Mass., occurrence of Belone latimanus in—G. B. Goode. (P. 1878)	
C.	•
Cache la Poudre river, Colorado, antiquities on—E. L. BERTHOUD. (R. 1871) California—	249
account of botanical explorations in—A. GRAY. (R. 1849)	21
ancient graves and shell heaps of-P. Schumacher. (R. 1874)	
burial of Indian squaw in—W. M. KING. (R. 1874)	
cave in Calaveras county—J. D. WHITNEY. (R. 1867)	
climate of—A. Gibbons. (R. 1854)	
climate of	
description of Gobiesox rhessodon from San Diego—R. SMITH. (P. 1881)	
description of new gobioid fish from San Diego—R. SMITH. (P. 1881)	467
descriptions of new species of shells from—W. H. DALL. (P. 1878)	332
	33 2
fishes of—See D. S. JORDAN; C. H. GILBERT.	
	30\$

California—Continued.	
fossil mollusks from later tertiaries of—W. H. Dall. (P. 1878) 8	82
history and antiquities of Santa Rosa Island—S. Bowers. (R. 1877) 8	2
Indians of—S. Powers. (R. 1876) 2	98
list of fishes of, collected by Lt. H. E. Nichols-D. S. Jordan; C. H. Gilbert. (P. 1881)	67
meteorological observations at Sacramento—F. W. HATCH. (R. 1854)	78
meteorological observations at Sacramento—T. M. Logan. (R. 1854)	75
meteorology of Sacramento-T. M. Logan. (R. 1857) 1	07
natural history of—T. H. STREETS 30	03
new genera and species of fishes from coast of—W. N. Lockington. (P. 1879)	88
new pitcher plant from—J. Torrey	61
notes on a collection of fishes from San Diego—D. S. JORDAN; C. H. GILBERT. (P. 1880)	2
occurrence of Cremnobates at San Diego-R. SMITH. (P. 1880) 4	2
occurrence of Productus giganteus in—C. A. WHITE. (P. 1880) 4	28
plants collected by J. C. Frémont in -J. TORREY	46
pleistocene fossils from-P. P. CARPENTER 2	52
postpliocene fossils in coast range of—W. H. Dall. (P. 1878) 8	82
wingless grasshopper of—E. P. Vollum. (R. 1860)	47
California, central, partial list of birds of—L. Belding. (P. 1878) 8	82
California, Gulf of, lectures on shells of—P. P. CARPENTER. (R. 1859) 1	10
California peninsula, aboriginal inhabitants of—J. BAEGERT. (R. 1863; R. 1864) 187, 188, 440, 30	61
California, upper, explorations in, 1860—J. FEILNER. (R. 1864) 1	88
Camaraphysema, a new type of sponge—J. A. RYDER. (P. 1880) 4	25
Camel, lecture on the-G. P. Marsh. (R. 1854)	78
CAMEBON, H. C. Reminiscences of Joseph Henry 8	56
CAMPBELL, J. V. Earthquake in Peru, August 18, 1868. (R. 1870) 2	41
Canada, meteorological system of—J. G. Hodgins. (R. 1865) 2	909
Canada East, description of observatory at St. Martin, Isle Jesus—C. SMALL- WOOD. (R. 1856)	91
Canada porcupine, occurrence of, in Maryland—O. Lugger. (P. 1881) 4	67
Canada porcupine, occurrence of, in West Virginia—G. B. Goode. (P. 1878) 8	82
Canada West, ancient Indian remains near Prescott—W. E. Guest. (R. 1856)	91
Canada West, meteorological stations of—J. G. Hodgins. (R. 1858) 1	08
, , , , , , , , , , , , , , , , , , ,	24
CANBY, S. Deposit of Bishop John's library. (R. 1862) 1	
Cancerous tumors, structure of—J. J. WOODWARD 20	5€
Candle fish of northwest coast—J. G. Swan. (P. 1880)	2
CANDOLLE, P. DE, memoir of, by M. Flourens. (R. 1859)1	1(
CARRON G. I. Antiquities of Coloredo (R. 1877)	28

Canudas, A. Earthquakes in Guatemala. (R. 1858) 100
Caoutchoue and gutta percha. (R. 1864)
Cape Cod, littoral marine fauna of-R. RATHBUN. (P. 1880) 42
Cape Flattery, Indians of—J. G. SWAN 220
CAPEN, F. L. Meteorological discovery. (R. 1866)21
Capitol, U. S., memorial exercises at
Capitol, U. S., account of marble used in extension of-J. HENRY. (R. 1856) 9
Cara gigantesca of Yzamal, in Yucatan—A. Schott. (R. 1869) 22
Caracas, South America, meteorology of-G. A. Ernst. (R. 1867) 21
Caranx Beani, from Beaufort, N. C., description of-D. S. JORDAN. (P. 1880) 42
Carib language and people—C. H. BERENDT. (R. 1873) 274
Caribbee Islands, ornithology of-F. A. OBER (B. 1878) 34
CARLETON, J. H. Excursion to Abó, Quarra, and Gran Quivira, New Mexico. (R. 1854)
CARLETON, J. H. Meteorites in Mexico. (R. 1865) 205
CARLIN, W. E. Observations of Siredon lichenoides. (P. 1881) 46
Carlsruhe, Scientific Congress of, 1858-F. J. NICKLES. (R. 1860) 147
CARPENTER, P. P.—
Check-list of the shells of North America
Lectures on mollusca, or shell-fish, and their allies. (R. 1860) 147, 152
Lectures on shells of Gulf of California. (R. 1859) 110
Mollusks of western North America 252
CASE, H. B. Flint implements in Holmes county, Ohio. (R. 1877) 32-
CASELLA, L. Description of meteorological instruments. (R. 1859) 110
CASS, L., Secretary of State. Circular to diplomatic agents relative to L. H.
Morgan's research 13
Castings, metallic, of delicate natural objects. (P. 1881) 467
Casts, catalogue of, of Indian boys and girls at Hampton Normal Institute, Virginia—R. H. Pratt. (P. 1879)
Casts, catalogue of, of Indian prisoners at St. Augustine, Florida—R. H.
PRATT. (P. 1878)
Casts of statues, etc., proposed by W. J. Stone. (R. 1855)
Casts, plaster, methods of making and preserving—A. Pirz. (P. 1881) 457
Caswell, A.—
Lecture on astronomy. (R. 1858)
Meteorological observations at Providence, R. I., 1831-1860 103
Meteorological observations at Providence, R. I., 1831-1876 443
Catalogue, general stereotype, of public libraries, report of commission on. (R. 1850)
Catalogue of—
birds of Antigua and Barbuda, collected by F. A. Ober-G. N. LAW-BENCE. (P. 1878)
birds of Dominica, collected by F. A. Ober-G. N. LAWRENCE. (P. 1878) \$52

Catalogue of—Continued. birds of Grenada, collected by F. A. Ober—G. N. LAWRENCE. (P. 1878) 33	82
birds of Guadeloupe, collected by F. A. Ober—G. N. LAWRENCE. (P. 1878)	89
birds of Lesser Antilles, collected by F. A. Ober—G. N. LAWRENCE. (P. 1878)	
birds of Martinique, collected by F. A. Ober-G. N. LAWRENCE. (P. 1878)	
birds of North America—R. RIDGWAY. (P. 1880) 425, 42	12
birds of St. Vincent, collected by F. A. Ober-G. N. LAWRENCE. (P. 1878)	
books and memoirs relating to nebulæ and clusters—E. S. HOLDEN 31	.1
casts taken by Clark Mills of Indian boys and girls at Hampton, Virginia—R. H. Pratt. (P. 1879)	38
casts taken by Clark Mills of Indian prisoners at St. Augustine, Florida— R. H. Pratt. (P. 1878)———————————————————————————————————	
collection at International Exhibition, 1876. (R. 1876) 299, 32	
collection of Japanese cotton fibre presented by Government of Japan. (P. 1881)46	
collection of Japanese woods presented by University of Tokio-L. F. WARD. (P. 1881)	
collection to illustrate animal resources and fisheries of United States exhibited at Philadelphia, 1876—G. B. Goode 32	
described coleopters of the United States—F. E. Melsheimer	
described diptera of North America—C. R. OSTEN SACKEN 102, 27	
described lepidoptera of North America—J. G. Morris	
engravings presented to Smithsonian—C. B. King. (R. 1861) 14	
fishes from Pensacola, Florida, collected by S. Stearns—G. B. Goode; T. H. Bean. (P. 1879)	22
fishes obtained in the Gulf of Mexico by J. W. Velie—G. B. Goode; T. H. Bean. (P. 1879)	
fishes of Alaskan and adjacent waters—T. H. BEAN. (P. 1881) 46	
fishes of the Bermudas—G. B. Goode29	
fishes of the east coast of North America—T. GILL 28	3
fishes of St. John's river and east coast of Fla.—G. B. Goode. (P. 1879) 33	18
geological specimens—D. D. OWEN. (R. 1854)7	5
historical and geographical manuscripts, maps, etc.—L. Berlandier. (R. 1854) 7	75
Indian relics presented to Smithsonian—J. H. DEVEREUX. (R. 1872) 27	1
library of James Smithson	0
linguistic manuscripts in library of Bureau of Ethnology—J. C. Pilling. (E. 1879-80)47	'6
meteorites in Yale College—G. J. Brush. (R. 1868) 22	4
minerals, with their formulas—T. EGLESTON	F

Centennial Exhibition, catalogue of collection of animal resources and fisheries at—G. B. Goode 326
Centennial Exhibition, circular for distribution at 290
Centennial Exhibition, plan of proposed Smithsonian exhibit at—S. F. BAIRD. (R. 1875)
Centennial Exhibition, report on—S. F. BAIRD. (R. 1876) 299, 307
Centennial Exhibition, savage weapons at—E. H. KNIGHT. (R. 1879) 845, 415
Centennial mission to Indians of Western Nevada and California—S. Powers. (R. 1876)
Centigrade degrees, tables for conversion of, to Fahrenheit. (R. 1868) 187
Central America—
circular relative to collections of birds of 168
description of new fishes from—D. S. JORDAN; C. H. GILBERT. (P. 1881)
explorations in—C. H. BERENDT. (R. 1867) 215
list of birds of, not in Smithsonian collection 185
rain-fall in—C. A. Schott 222, 353
review of birds of—S. F. BAIRD 181
studies in picture writing of—E. S. HOLDEN. (E. 1879-80) 476
travels in—S. Habel269
Centrarchidse, notes on—D. S. JORDAN 806
Centrarchidse, review of family—C. L. McKAY. (P. 1881) 467
Centurus, review of genus—R. RIDGWAY. (P. 1881) 467
Cephaloscyllium laticeps-D. S. JORDAN; C. H. GILBERT. (P. 1880) 425
Ceratiidæ, note on—T. Gill. (P. 1878)
Ceremonies at laying of corner-stone of Smithsonian Institution. (R. 1847). H, 829
Ceremonies, Mandan, accuracy of Catlin's account of-J. KIPP. (R. 1872) 271
Cessions of land by Indian tribes to United States—C. C. Royce. (E. 1879-80) 476
CHACE, G. I. Lecture on oxygen and its combinations. (R. 1855) 77
Chalk found in United States—T. A. CONRAD. (R. 1865) 209
Chamber of Commerce of Bordeaux. Exchange of publications. (R. 1863) 187
Change of Mexican axolotl to amblystoma—A. Weismann. (R. 1877) 323, 401
CHANNING, W. F. Lecture on the American fire-alarm telegraph. (R. 1854) 75
CHANTEE, E. Code of symbols for prehistoric archæology. (R. 1875) 298
Chapala Lake, notes on Dugès' fishes from—D. S. JORDAN. (P. 1879) 838
CHAPPELSMITH, J. Tornado near New Harmony, Indiana, April 80, 1852 59
Characteristics of ancient man in Michigan—H. GILLMAN. (R. 1875) 298, 393
Chart, base, of the United States—C. A. SCHOTT 414
Chart, temperature, of the United States, year—C. A. Schott 381, 388
Charts and maps of America, on a collection of—J. G. Kohl. (R. 1856) 91
Charts of prehistoric archeology, international code of symbols for—G. DE MORTILLET; E. CHANTRE. (R. 1875)
Charts, rain, of the United States, summer, winter, year—C. A. Schott 374

Charts, temperature, of United States, summer, winter, year-C. A. Schott 387
Charts, wind—J. H. COFFIN; S. J. COFFIN
CHASE, S. P., biographical notice of, by J. A. GARFIELD. (R. 1878)
CHASE, S. P., biographical notice of, by H. HAMLIN. (R. 1873) 27
CHASE, S. P., statement of, relative to the telegraph. (R. 1857) 107, 11
Chattanooga, Tenn., ancient mound near-M. C. READ. (R. 1867) 21
Check-list of—
duplicates of fishes from Pacific coast, distributed by Smithsonian Insti-
tution in 1881—D. S. JORDAN; P. L. JOUY. (P. 1881) 46
fishes, distributed by Smithsonian Institution for National Museum-
T. H. BEAN. (P. 1880)
fossils of North America, miocene-F. B. MEEK 18
invertebrate fossils of North America, cretaceous and jurassic-F. B.
Meek
invertebrate fossils of North America, eocene and oligocene-T. A. Con-
RAD 200
North American batrachia and reptilia—E. D. Cope 292
periodicals received by Smithsonian Institution, 1858 (
plants of Washington and vicinity—L. F. WARD 460
publications of Smithsonian Institution to July, 1872 24
publications of Smithsonian Institution to July, 1874 278
publications of Smithsonian Institution to July, 1877 301
publications of Smithsonian Institution to July, 1879 34
publications of Smithsonian Institution to December, 1881 435
shells—Lea, Carpenter, and others 128
Smithsonian publications
Check-lists, blank164
Chelonia, anatomy and physiology of-S. W. MITCHELL; G. R. MOREHOUSE. 159
Chelonia, extinct, of Nebraska—J. LEIDY
Chelura terebrans, occurrence of, on coast of U.S.—S. I. SMITH. (P. 1879) 323
Chemical analysis of the sun—A. LAUGEL. (R. 1861)
Chemical and physiological investigations of vertebrata—J. Jones82
Chemical arts, recent improvements in—J. C. BOOTH; C. MORFIT 27
Chemical equivalents of sixty-three elements, table of. (R. 1864) 1864
Chemical formulæ, tables of—F. W. CLABKE255
Chemical types, modern theory of—C. M. WETHERILL. (R. 1863) 185
Chemist, report of—F. W. TAYLOB. (R. 1880) 422
Chemistry, lectures on agricultural—S. W. Johnson. (R. 1859) 110
Chemistry of the earth—T. S. Hunt. (R. 1869) 228, 376
Chemistry, photographic—M. Jamin. (R. 1867) 215
Chemistry, recent progress in, 1879 and 1880—G. F. BARKER. (R. 1880)_ 442, 429
Chepewvan Indians, notes on the-G. GIBBS and others. (R. 1866) 214, 365

Cherbourg, Imperial Society of Natural Sciences of. Prize questions. (R. 1864) Chesapeake Bay, new fish from mouth of—G. B. Goode; T. H. Bean. (P. 1881)	188 467
Chester county, Pennsylvania, birds of—V. BARNARD. (R. 1860)	147
Chester county, Pennsylvania, history of—W. DARLINGTON. (R. 1862)	150
Chicago Academy of Sciences. Acknowledging specimens. (R. 1867)	218
China, geological researches in—R. Pumpelly	202
Chinese coal-bearing rocks, fossil plants from—J. S. NEWBERRY	202
Chinese coals, analyses of-J. A. MACDONALD	202
Chinese mineral called yu, study of—S. BLONDEL. (R. 1876)	298
Chinook jargon—H. HALE	161
Chinook jargon-M. LIONNET; B. R. MITCHELL	68
Chinook jargon, bibliography of—G. GIBBS	161
Chinook jargon, dictionary of—G. GIBBS	161
Chinook names of salmon in Columbia river—S. B. SMITH. (P. 1881)	467
Chionis minor, study of-J. H. KIDDER; E. COUES-	294
Chiroid fish (Myriolepis zonifer) from California, new—W. N. Lockington. (P. 1880)	428
Chirus, remarks on genus-W. N. Lockington. (P. 1880)	428
Chitonidse, new species of, from Panama-P. P. CARPENTER	252
Chitons of Alaskan and Arctic regions, report on-W. H. Dall. (P. 1878)	882
Chitons from deep waters off eastern coast of U. SW. H. DALL. (P. 1881.)	467
Chitons, on genera of—W. H. Dall. (P. 1881)	467
Chlorospermeze, marine algze of North America—W. H. HARVEY	95
Chorea, inquiries relative to disease known as—S. W. MITCHELL. (R. 1874)	286
CHOSSAT, Dr. Transactions of Geneva Society of Physics and Natural History, 1863 to 1864. (R. 1865)	209
Christiania, Norway, University of, ethnological specimens presented. (R. 1868)	187
Chrysotis, new species of, from Dominica-G. N. LAWRENCE. (P. 1880)	425
Cibola, Coronado's march in search of cities of-J. H. SIMPSON. (R. 1869)	228
Circle, ratio between diameter and circumference of-W. Ferrel	233
Circles and spheres, tangencies of—B. ALVORD	80
Circular—	
ancient mining in Lake Superior copper region. (R. 1861)	149
for distribution at the Centennial Exhibition	290
giving directions for constructing lightning-rods—J. HENRY	237
in reference to collecting North American shells	176
in reference to history of North American grasshoppers—P. R. UHLER-	163
of inquiries relative to crawfish and other fresh water crustacea	319
of Institute of Rupert's Land. (R. 1861)	149
of instructions for observations of thunder storms—J. HENRY	235
of instructions to observatories, relative to telegraph announcements of	

Circular—Continued.	
philological—G. Gibbs. (R. 1862)	
relating to collections in archæology and ethnology	
relating to collections of living reptiles	
relative to American archæology—O. T. Mason. (R. 1879)	34 5, 31 6
relative to collections of birds from Middle and South America	168
relative to food fishes	234
relative to heights	236
relative to observations in Russian America	207
relative to scientific and literary exchanges	324
relative to shipping fresh fish and other animals	384
respecting copyright. (R. 1854)	57
respecting new report on libraries. (R. 1854)	75
respecting system of relationship—L. H. Morgan	138
sent with specimens presented. (R. 1872)	271
to entomologists	178
to officers of Hudson's Bay Company	137
Circulars of United States National Museum:	
No. 1. Plan of organization and regulations of the Museum—G.	
B. Goode. (P. 1881)	-
No. 2. To friends of the Museum—S. F. BAIRD. (P. 1881)	
No. 8. Petroleum collections—S. F. BAIRD. (P. 1881)	
No. 4. Department of insects—S. F. Baird. (P. 1881)	
No. 5. Establishment and officers of S. I. and Nat. Mus. (P. 1881)	
No. 6. Materia-medica collection—J. M. FLINT. (P. 1881)	467, 450
No. 7. Forms of drugs and medicines—J. M. FLINT. (P. 1881)	467, 451
No. 8. Collections of drugs for materia-medica section—J. M.	100 150
FLINT. (P. 1881)	
No. 9. Building-stone collection—S. F. BAIRD. (P. 1881)	
No. 10. Letters on the National Museum—B. PHILLIPS. (P. 1881)	467, 454
No. 11. Provisional classification of the food collections—G. B.	ART ARE
No. 12. Classification of the collections to illustrate taxidermy—	2011 200
W. T. Hornaday. (P. 1881)	467, 456
No. 18. Scheme of classification for the collections in the Museum—	,
G. B. GOODE. (P. 1881)	467, 457
No. 14. Asking for contributions to library of Museum-S. F.	
BAIRD. (P. 1881)	467, 458
No. 15. Organization and objects of the Museum—G. B. Goode.	100 ACC
(P. 1881)	
No. 16. Plans for installation of collections—G. B. Goods. (P. 1881)	507, 572
No. 17. Contributions to the Museum and their acknowledgment. (P. 1881)	467 472
No. 18. List of publications of the Museum. (P. 1881)	
TIA. TO. THE AT LANGUAGE AT PIG TERBORM. (T. 1001)	301, 3/3

OF SMITHSUNIAN PUBLICATIONS.
Cists, stone, Madison county, Illinois—A. OEHLER. (R. 1879) 846
Citizens of Philadelphia, memorial of, relative to aeronautic voyage of Lowe.
(R. 1860)
Civil Engineers, London Institution of. Prize questions. (R. 1862) 150
Clackamas river, Oregon, notes on fishes from—D. S. JORDAN. (P. 1878) 882
CLARK, H. J. Lucernarise and their allies, anatomy, physiology, and relations of 242
CLARK, H. J., memoir of, by A. S. PACKARD 242
CLARK, W. M. Antiquities of Tennessee. (R. 1877) 828
CLARKE, FLADGATE, and FINCH. Residuary legacy of Smithson. (R. 1861) 149, 828
CLARKE, F. W
Bibliography of atomic volume, specific gravity, &c 255
Recalculation of atomic weights41
Specific gravities, boiling and melting points, chemical formulæ, tables 255
Specific gravities, &c. Supplementary tables 288
Specific heat tables for solids and liquids 276
Tables of expansion by heat, for solids and liquids
Classification and arrangement of materia medica collection—J. M. FLINT.
(P. 1881)
Classification and synopsis of trochilide—D. G. Elliot
Classification of—
birds, review of the-W. LILLJEBORG. (R. 1865) 209, 364
books, on the—J. P. LESLEY. (R. 1862) 150
clouds-A. Pory. (R. 1870)244
coleoptera. Part I—J. L. LE CONTE 136
coleoptera. Part 11-J. L. LE CONTE 265
collection to illustrate animal resources of United States-G. B. Goode 297
collections of U.S. National Museum-G.B. Goode. (P. 1881) 467, 457
collections of United States National Museum to illustrate taxidermy—
W. T. Hornaday. (P. 1881) 467, 456
food collections—G. B. Goode. (P. 1881) 467, 455
forms in which drugs and medicines appear and are administered—J. M.
FLINT. (P. 1881) 467, 451
insects from embryological data—L. Agassız 16
methods of capture and utilization of animals—G. B. Goode 297
Classified list of meteorological publications and articles in periodicals received
by Smithsonian Institution. (R. 1873) 275
Classified record of meteorological material preserved in Smjthsonian. (R.
1874) 286
Classified record of monthly meteorological reports. (R. 1878) 275
CLEAVELAND, P. Meteorological observations at Brunswick, Maine 204
CLEAVELAND, P., notice of, by J. HENRY. (R. 1859) 110
CLEMENS, B. Instructions for collecting hymenopters. (R. 1858) 109

CLEMENS, B. Instructions for collecting lepidoptera. (R. 1858)	. 10
CLEMENS, B. On lepidoptera	. 13
Climate and history of New Mexico-T. A. McParlin. (B. 1877) 82	3, 39 (
Climate of Alaska—H. M. BANNISTER. (R. 1866)	. 21
Climate of California—T. M. LOGAN. (R. 1855)	
Climate of Colorado—J. Evans. (R. 1865)	. 200
Climate of Kansas—R. S. Elliott. (R. 1870)	. 24
Climate of Kelley's Island, Ohio-G. C. HUNTINGTON. (R. 1866)	. 2 1:
Climate of San Francisco—H. GIBBONS. (R. 1854)	. "
Climate—See Meteorology.	
Climatic influence of forests—M. BECQUEREL. (R. 1869)	. 23
CLINGMAN, T. L. Topography of Black Mountain, N. C. (R. 1855)	. 7
Clinton, N. Y.—See Hamilton College.	
Closing of Kennebec river, Maine-R. H. GARDINER. (R. 1858)	10
Cloud-bursts—W. J. Young. (R. 1867)	21
Cloud, dispersion of, by an electrical discharge—D. W. NAILL. (R. 1858)	
Clouds, electricity of induction in—F. ZANTEDESCHI. (R. 1870)	. 24
Clouds, formation of, over Gulf Stream-H. M. BANNISTER. (R. 1866)	. 21
Clouds, new classification of—A. Poey. (R. 1870)	. 24
Clouds, nomenclature of	
Clupes tyrannus of Latrobe-G. B. Goode. (P. 1878)	, 000 000
CLUSS, A. Report of architect. (R. 1867)	21.
Cluss & Schulze, report of, for 1879. (R. 1879) 343	, 40 9
Cluss & Schulze, report of, for 1880. (R. 1880) 442	434
CLYMER, H. Memorial of J. Henry	35
CLYMER, H., and others. Report on Museum. (R. 1876)	20
Coal, Chinese and Japanese, analyses of—J. A. MACDONALD	30
Coal, Chinese, fossil plants in—J. S. NEWBERRY	200
Coal, lecture on—J. LE CONTE. (B. 1857)	103
Coal oils, explosibility of—Z. Allen. (R. 1861)	143
Coast Survey expedition of 1866, to determine transatlantic longitude—B. A. Gould	9 04
Coast Survey soundings on Atlantic coast, examination of-J. W. BAILEY	A
Cobalt (ammonia-) bases-W. GIBBS; F. A. GENTH	88
COCHBANE, J. Antiquities of Mason county, Illinois. (R. 1877)	32:
CODD, J. A. Acknowledgment for books. (R. 1861)	149
Code of symbols for charts of archeology—G. DE MORTILLET; E. CHANTRE.	
(R. 1875)	50
Coffin, J. H.—	
Orbit and phenomena of meteoric fire ball	221
Psychrometrical tables	87
Storms of 1859	162

COFFIN, J. H.—Continued. Winds of the globe	269
Winds of the Northern Hemisphere	
COFFIN, S. J. Tables and maps of winds of the globe	
Cogswell, J. G. Report on Jewett's general stereotype catalogue of pul	
libraries	
Coins, assay of gold and silver, at United States mint—J. Pollock. (R. 18	
Coins, table of foreign, gold and silver. (R. 1868)	224
Colding, A. Meteorology. (R. 1877)	828, 898
Colding, A. Nature of currents of air. (R. 1877)	
Colding, A. Theory of relation between pressure and the velocity of wind. (R. 1877)	
Colding, A. Whirlwind at St. Thomas in August, 1871. (R. 1877)	828, 398
Coleoptera—	•
catalogue of the described, of the United States-F. E. MELSHEIMER	62
classification of. Part I-J. L. LE CONTE	136
classification of. Part II—J. L. LE CONTE	265
instructions for collecting—J. L. LE CONTE. (R. 1858)	109
new species of North American. Part I-J. L. LE CONTE	
new species of North American. Part II-J. L. LE CONTE	264
of Arctic America—S. H. SCUDDEB	842
of Kansas and eastern New Mexico—J. L. LE CONTE	126
of North America, list of-J. L. LE CONTE	140
Colima, eruption of the volcano of—C. SARTORIUS. (R. 1869)	 22 8
Collaborators of Smithsonian Institution, directory of	 4 66
Collecting—	
and preserving diptera, directions for-C. R. OSTEN SACKEN	102
and preserving fish, directions for-T. H. BEAN. (P. 1881)	467, 464
and preserving plants, directions for-L. F. WARD	 46 0
insects, directions for—A. S. PACKARD	261
insects, instructions for-S. F. BAIRD. (R. 1858)	109
land and fresh-water shells, instructions for—J. Lewis. (R. 1866)	21 4, 363
microscopic organisms, directions for	68
myriapods, phalangidse, etc.—H. C. Wood. (R. 1866)	214
nests and eggs, instructions for-T. M. Brewer	· 13 9
nests and eggs of birds—S. F. BAIRD. (R. 1858)	109
North American shells, circular in reference to	176
specimens, directions for—S. F. BAIRD. (R. 1856)	
specimens of diatomacea, directions for-A. M. EDWARDS	366
specimens of natural history—S. F. BAIRD	34
Collection—	
and preservation of marine invertebrates—W. STIMPSON	84
archæological, of United States National Museum—C. RAU	287

Collection—	
of historical documents in Guatemala—C. H. BERENDT. (R. 1876) 2	333
of meteorological tables—A. Guyot 31, 1	.53
to illustrate animal resources and fisheries of United States, exhibited	
at Philadelphia, in 1876, catalogue of—G. B. Goode3	26
to illustrate animal resources of United States, classification of—G. B. Goode	197
Collections—	
additions to. (See each annual report.)	
furnished by explorations from 1838 to 1877—S. F. BAIRD. (R. 1877) 3	323
in archæology and ethnology, circular relative to-J. HENRY 2	
made by Lt. G. K. Warren, report on—F. B. MEEK; F. V. HAYDEN	
of living reptiles, circular relating to 3	
presented to United States by foreign commissioners at Centennial Exhibition, list of. (R. 1876)	
to illustrate art of taxidermy-W. T. Hornaday. (P. 1881) 467, 4	
Collections—See Ethnographical.	
Collectors of customs to receive and transmit specimens to the Smithsonian	34
Collectors of drugs, memoranda for-J. M. FLINT. (P. 1881) 467, 4	152
College—See Girard, Hamilton, Liberia.	
Colleges in United States in correspondence with Smithsonian Inst., list of	238
Collins, Colonel. Meteorology of the Green river country. (R. 1871)	249
Colonia Tovar, Venezuela, meteorology and ethnology of—A. FENDLER. (R. 1857)	107
Colonia Toyar, Venezuela, meteorology of-A. FENDLER. (R. 1866)	214
Colorado-	
ancient remains in—E. L. BERTHOUD. (R. 1867)	215
climate of—J. Evans. (R. 1865)	209
cretaceous fossils from—C. A. WHITE. (P. 1881)	467
descriptions of new invertebrate fossils from—C. A. WHITE. (P. 1880)_	425
heights of mountains in-G. Engelmann. (R. 1862)	150
Jefferson and Clear Creek Cos., antiquities—G. L. Cannon. (R. 1877)	323
	249
	323
Color-blindness—A. Moigno. (R. 1866)	214
Color-blindness in its relations to accidents by rail and sea—F. Holmgran. (R. 1877)	399
Colored bead from mound in Florida—A. M. HARRISON. (R. 1877)	
	214
Columbia river, Chinook names of salmon in—S. B. SMITH. (P. 1881)	457
Columbia river, Salmonida of -C. Bendire. (P. 1881)	
Combustion of gun-cotton and gunpowder, products of—Lieut. von Karolyi; B. F. Craig. (R. 1864)	188

Congress, United States—Continued.	
act of, to receive residuary legacy and increase fund. (R. 1866) 27	14, 32
act of, to transfer Smithsonian library to Library of Congress. (R. 1865) 2	09, 32
acts of, (45th Congress,) relative to Smithsonian. (R. 1878)	. 34
acts of, (45th and 46th Congresses,) relative to Smithsonian, 1878-1880 (R. 1879)	
memorial of J. Henry, published by order of	35(
memorial of Regents to, asking appropriations for Museum. (R. 1866 R. 1867) 214, 2	-
memorial of Regents to, relative to new Museum building. (R. 1876).	_ 29
memorial of Regents to, relative to Smithson fund. (R. 1850)	. 2
proceedings of, relative to appropriations for Museum	_ 22
proceedings of, relative to monument of J. Henry	_ 35
proceedings of, relative to new Museum building	. 32
proceedings of, relative to Smithson bequest	. 32
proceedings of, relative to Smithsonian Institution, 1835-1877	. 32
Connecticut Academy of Arts and Sciences, planisphere of the heavens by	. 359
Connecticut, ancient implement of wood from—E. W. Ellsworth. (B. 1876)) 299
Conosaurus, memoir on—R. W. GIBBES	. 19
CONRAD, C. M., Secretary of War. Authority to officers of Quartermaster's	
Department to receive and transmit specimens to Smithsonian	
CONRAD, T. A. Chalk found in the United States. (R. 1865)	
CONBAD, T. A. Check-list of invertebrate fossils of North America. Economic and oligocene	
Consanguinity and affinity of the human family, circular—L. H. MOBGAN	. 136
Consanguinity and affinity of the human family, systems of—L. H. MORGAN -	
Considerations on electricity. (R. 1867)	
Constantinople, account of hail storm at—Com. PORTER. (R. 1870)	244
Constants of Nature—	
Part 1. Specific gravities, boiling and melting points, and chemical formulæ—F. W. CLARKE	
Part II. Table of specific heats for solids and liquids—F. W. CLARKE.	276
Part III. Tables of expansion by heat for solids and liquids—F. W. CLARKE	289
Part IV. Atomic weight determinations-G. F. BECKEB.	358
Part v. Recalculation of atomic weights-F. W. CLARKE	441
First supplement to Part 1. Specific gravities, boiling points and melting points—F. W. CLARKE	
Constants of nature—J. LE CONTE. (R. 1878)	
Constants of nature and art, tables of—C. BABBAGE. (R. 1856)	
Construction of act establishing Institution—J. McP. Berrien. (R. 1858)	
Construction of catalogues of libraries—C. C. JEWETT	
Construction of silvered glass telescope—H. DRAPER	

Contact, catalytic force or phenomena of—T. L. Phipson. (R. 1862)	
CONTAXARI, Miss E. B., present of album from Greece. (R. 1857) 107	•
Contrast of colors—A. Moigno. (R. 1866)	
Contribution to history of fresh-water alge of North America—H. C. Wood	
Contributions to Annals of Philosophy—J. Smithson. (R. 1858)	-
Contributions to history of marine algo of North America—See W. H. Harvey	
Contributions to Knowledge, vols. I-XXIII 2, 26, 38, 39, 55, 56, 76, 78, 92, 111, 112, 151, 184, 206, 211, 229, 246, 272, 284, 285, 340,	• •
Contributions to Knowledge, catalogue of, and index	478
Contributions to National Museum, and their acknowledgment. (P. 1881.) 467,	473
Contributions to natural history of— Arctic America—L. KUMLIEN	342
Fanning Islands—T. H. STREETS.	808
fresh-water fishes of North America—C. GIRARD	30
Hawaiian Islands—T. H. STREETS	808
Kerguelen Island. 1. Ornithology—E. Coues	298
Kerguelen Island. 11. Oölogy, botany, etc.—J. H. KIDDER and others	294
Lower California—T. H. STREETS	808
Contributions to North American ichthyology—	000
Part I. Review Rafinesque's memoirs—D. S. JOBDAN	305
Part 11. Notes on Cottidæ; sysopsis Siluridæ—D. S. JORDAN	
Part III. Fishes Alleghany region; synopsis Catostomidæ—D. S. Jor-	
Contributions to physical goography of United States C. Exxum In	308
Contributions to physical geography of United States—C. ELLET, Jr	
Contributions toward monograph of the pandorids—P. P. CARPENTER	202
Converging series expressing ratio between diameter and circumference of circle—W. Ferrel	233
Cook, C. Manufacture of porpoise oil. (P. 1878)	882
Cooper, J. G.—	
Distribution of forests and trees of North America, and catalogue of native trees of United States. (R. 1858) 109,	351
Forests and trees of Florida and Mexican boundary. (R. 1860)	147
Migrations and nesting habits of west coast birds. (P. 1879)	388
COPE, E. D.—	107
Genera and species of rattlesnakes	
North America batrachia and reptilia	
Zoölogical position of Texas	
Coppée, H. Report of Committee on Smithsonian Museum. (R. 1874)	
Copper and iron in salt water, preservation of—A. E. BECQUEREL. (R. 1864.) Copper region, circular, ancient mining in Lake Superior—J. HANN. (R.	188
1861)	149
Coptic language, introduction to the study of the-M. KABIS. (R. 1867)	215
Copyright books from 1846-1849, list of-C. C. JEWETT. (R. 1850)	28

Copyright system and Smithsonian library—C. C. JEWETT. (R. 1851) 51
Copyrights, circular respecting—J. HENRY. (R. 1854)75
Corbiculadæ, American, monograph of—T. PRIME
Corcoran Art Gallery, list of deposits in, by Smithsonian Institution. (R. 1874) 286
Corcoran Art Gallery, report of Committee of Regents on. (R. 1872) 271
CORCORAN, W. W. Deed of gift establishing Art Gallery. (R. 1872) 271, 329
Cordoba, Argentine Republic, account of astronomical observatory at—B. A. Gould. (R. 1878)
Corner-stone of Smithsonian building, address at laying-G. M. Dallas D, 329
Corner-stone of Smithsonian building, ceremonies at laying of. (R. 1847) H, 329
Coronado's march in search of the "seven cities of Cibola"—J. H. SIMPSON. (R. 1869)
Correspondence—
abstracts of anthropological—O. T. Mason. (R. 1880) 443
extracts from. (R. 1855, 1858-1868, 1865-1867, 1878) 77, 109, 110, 147, 149, 150, 187, 209, 214, 215, 275
meteorological—E. Foreman. (R. 1851)
relative to publication of memoir by Squier and Davis K
relative to Smithson and his bequest 228
summary of anthropological. (R. 1879) 345
Correspondents—
3 11.4 6 . (1) 1070 TO 1080\
domestic, list of. (R. 1853; R. 1872) 67, 271, 69, 238
foreign, list of64, 154, 225, 243, 309
·
foreign, list of 64, 154, 225, 243, 309 foreign, systematic index of 257 Costa Rica—
foreign, list of

CBAIG, B. F. Nitrification. (R. 1861)	140
CRAIG, B. F.; Lieut. VON KAROLYI. Products of combustion of gun-cotton	720
and gunpowder. (R. 1864)	188
Craig flounder of Europe (Glyptocephalus cynoglossus) on coast of North	
America-G. B. GOODE; T. H. BEAN. (P. 1878)	882
Crania Helvetica—F. Troyon. (R. 1864)	188
	319
Creation, visible, lecture on vastness of—S. ALEXANDER. (R. 1857)	107
Cree Indians, system of relationship of-E. A. WATKINS. (R. 1862)	150
Cremnobates, occurrence of, at San Diego, California-R. SMITH. (P. 1880)	42 5
CRESSON, J. C. Lightning discharges. (R. 1867)	215
Cretaceous fossils, check-list of-F. B. MEEK.	177
Cretaceous fossils from Arkansas and Colorado—C. A. WHITE. (P. 1881)	
Cretaceous reptiles of United States-J. LEIDY. (R. 1864)	192
Cricetodipus parvus, a rare rodent—F. W. TRUE. (P. 1881)	467
Criocardium, note on—C. A. WHITE. (P. 1879)	888
Criticisms of Dr. J. Hann, replies to the-W. FERREL. (R. 1877) 828	398
CROOK, G. Indian mode of making arrow-heads and obtaining fire. (R. 1871.)	249
Crozet flora-J. H. KIDDER	294
CRUMMELL, A. Facts respecting Liberia College. (R. 1861)	149
Crust of the earth, revolutions of the—G. PILAR. (R. 1876)	299
Crustacea-	
fresh water, circular of inquiries relative to	319
notice of, dredged off south coast of New England by Fish Commission— S. I. Smith. (P. 1880)	425
notice of new species of Willemssia group of-S. I. Smith. (P. 1879.)	888
of Hawaiian and Fanning Islands and California-T. A. STREETS	808
Crustacean destructive to submarine timber on coast of United States—S. I.	
Sмітн. (Р. 1879)	888
Crustaceans of Kerguelen Island—S. I. SMITH	294
Crustaceans, voices of-G. B. Goode. (P. 1878)	382
Cryolite of Greenland—Lewis; Quale. (R. 1866)	214
Cryptogamous plants, present state of knowledge of—W. REICHARDT. (R. 1871)	249
Crystalline rocks, metamorphism and the formation of—G. A. DAUBRÉE. (R. 1861)	149
Crystallography, explanation of principles of-A. Brezina. (R. 1872) 271,	386
Crystallophysics, explanation of principles of—A. Brezina. (R. 1872) 271,	
CULBERTSON, T. A.—	
Expedition to the Mauvaises Terres and Upper Missouri. (R. 1850)	28
Indian tribes in Upper Missouri. (R. 1850)	28
Sioux of the Upper Missouri. (R. 1850)	28
Culture, fish—See Fish.	
CUNARD, E. Free freight between United States and England. (R. 1859.) 110	829

CUNNINGHAM, K. M.; GAINES, A. S. Shell-heaps on Mobile river. (R. 1877.)
Currents, electrical, of the earth—C. MATTEUCCI. (R. 1867) 215
Currents of air-
and aerial navigation—J. HENRY. (R. 1860)
law of variation of temperature in ascending—J. HANN. (R. 1877.) 823, 89
law of variation of temperature in ascending moist—L. Sohncke. (R. 1877)
nature of—A. Colding. (R. 1877) 328, 39
relation between barometric variations and—M. PESLIN. (B. 1877)_ 323, 89
Curtiss, A. H., notes on collection of fishes made by-D. S. JORDAN. (P. 1880) 42
CUSHING, F. H. Antiquities of Orleans county, New York. (R. 1874) 28
Customs, collectors of, to receive and transmit specimens to Smithsonian 3
CUTTS, J. B. Ancient relics in northwestern Iowa. (R. 1872) 271
CUVIER, Baron G.—
history of the works of, by M. Flourens. (R. 1868) 224
memoir of, by M. Flourens. (R. 1868) 224
Memoir of René Just Haüy. (R. 1860)
Memoir of Priestley. (R. 1858)
Cybium, notes on American species of—F. Porv. (P. 1878) \$3
Cyclone in the Indian Ocean—N. PIKE. (R. 1867)
Cyclophoridæ—W. G. BINNEY
Cymatogaster rosaceus—D. S. Jordan; C. H. Gilbert. (P. 1880) 429
Cyprinidæ, notes on—D. S. Jordan
Cyprinoid fishes of San Francisco market—D. S. Jordan. (P. 1880) 425
D.
DAA, L. K. Ethnological museum of Norway. (R. 1862) 150
Dacentrus, note on—D. S. JORDAN. (P. 1880)
DA COSTA, J. M. On strain and over-action of the heart. Toner lecture No. 111. 279
DAHLBERG, R. N. and C. Ancient pottery at Pittsburg, on Des Moines river.
(R. 1879) 345
Dakota—
bibliography—S. R. Riggs

fable, A dog's revenge—S. R. Riggs. (E. 1879-80).....

grammar and dictionary—S. R. Riggs

Fort Wadsworth, Indian mounds near—A. J. Comport. (B. 1871)....

40

249

of smithsonian publications. 169
Dakota, Lincoln county, haystack mound in—A. BARRANDT. (R. 1872) 271
Dakota or Sioux Indians—A. G. BRACKETT. (R. 1876) 299
Dakota or Sioux Indians, language of—F. L. O. Rozhrig. (R. 1871) 249, 878
Dall, W. H.—
Distribution of California tertiary fossils. (P. 1878) 882
Explorations on western coast of North America. (B. 1878) 275
Fossil mollusks from later tertiaries of California. (P. 1878) 382
Index of Brachiopoda
Limpets and chitons of Alaskan and Arctic regions. (P. 1878)
Mollusks of Arctic America 342
Mollusks of Kerguelen Island 294
New mollusks from Alaska in National Museum. (P. 1878)
New species of shells from California in National Museum. (P. 1878) 382
Note on shells from Costa Rica kitchen-midden collected by Drs. Flint
and Bransford. (P. 1878) 332
On genera of chitons. (P. 1881) 467
Postpliocene fossils in coast range of California. (P. 1878) 382
Remains of later prehistoric man from caves in Alaska 318
Dallas, G. M. Address at laying corner-stone of Smithsonian building D, 829
Dalton, J. C. Origin and propagation of disease. (R. 1878) 275
Daltonism, etc.—A. Moigno. (R. 1866) 214
Danilsen, A. F. Mound in East Tennessee. (R. 1863)
Danish Royal Society of Arts and Sciences. Prize questions. (R. 1862, 1865,
1867) 150, 209, 215
DARLINGTON, W. History of Chester county, Pennsylvania. (R. 1862) 150
Darlingtonia Californica—J. Torrey
DARWIN, C. Queries about expression for anthropological inquiry. (R. 1867) 215
DAUBRÉE, G. A. Metamorphism and formation of crystalline rocks. (R.
1861) 149
DAUBRÉE, G. A. Synthetic experiments relative to meteorites. (R. 1868) 224
Davis, A. C. Antiquities of Isle Royale, Lake Superior. (R. 1874) 286
DAVIS, C. H. Law of deposit of flood-tide 33
DAVIS, C. H. Occultations in United States, 1852 29
DAVIS, E. H. On ethnological research. (R. 1866) 214
DAVIS, E. H.; SQUIER, E. G. Ancient monuments of. Mississippi valley 1
DAYTON, E. A. Explorations in Tennessee. (R. 1870) 244
Deaf mutes, sign language of-G. MALLERY. (E. 1879-80)
DEAN, C. K. Mound in Wisconsin. (R. 1872) 271
DEAN, C. W. Lightning discharges. (R. 1867) 215
Dean, J. Gray substance of medulla oblongata173
De Beaumont, E.—
Memoir of C. F. Beautemps-Beaupré. (R. 1868)

DE BEAUMONT, E.—Continued.	
Memoir of Auguste Bravais. (R. 1869)	
Memoir of Legendre. (R. 1867)	
Memoir of Oersted. (R. 1868)	224
DE BLAINVILLE, D., memoir of, by M. FLOUBENS. (B. 1865)	203
DE CANDOLLE, A.—	
Causes which limit vegetable species towards the north. (R. 1858) Probable future of the human race. (B. 1875)	
Report on transactions of Society of Physics and Natural History of Geneva for 1862. (R. 1864)	
Report on transactions of Society of Physics and Natural History of Geneva, 1878-1874. (R. 1875)	
DE CANDOLLE, A.; GRAY, J. E. On a dominant language for science. (R. 1874)	
DE CANDOLLE, P., memoir of, by M. FLOURENS. (R. 1859)	
Deduction and induction—J. VON LIEBIG. (R. 1870)	
Deed of foundation of Toner lectures—J. M. TONER. (B. 1872) 271	
Deed of foundation of Tyndall scientific fund—J. TYNDALL. (R. 1872) 271	
Deed of gift of Art Gallery-W. W. CORCOBAN. (R. 1872) 271	
DE FOREST, E. L. Methods of interpolation. Part I. (R. 1871)	
DE FOREST, E. L. Additions to memoir on methods of interpolation. Part II. (R. 1878)	
DE HART, J. N. Mounds and osteology of mound-builders of Wisconsin. (R. 1877)	823
DE IRISARRI, Guatemalan minister, recommendation of Dr. Berendt by. (R. 1865)	209
DELAFIELD, R. Report of Committee of Regents on Washington canal. (R. 1868) 224	
DELAMBRE, J. B. J., memoir of, by J. FOURIER. (R. 1864)	188
DE LA RIVE, A. A.—	
Eulogy on, by J. B. Dumas. (B. 1874)	286
Michael Faraday, his life and works. (R. 1867)	
Phenomena accompanying propagation of electricity in rarefled elastic fluids. (R. 1868)	
Report on transactions of Society of Physics and Natural History of Geneva, 1858-1859, 1872-1878. (R. 1859; R. 1874) 110,	
DE LA RIVE, A. A.; LEMSTRÖM, S. Electricity of the atmosphere and the aurora borealis. (R. 1874)	285
DE LA RUE, W. Abbreviations used in England in 1867. (R. 1867)	215
DELAUNAY, C. Essay on the velocity of light. (R. 1864) 188.	354
Delolepis, new genus of fishes, description of—T. H. Bran. (P. 1881)	
	345
DENNIS, W. C. Evaporation in Florida. (R. 1866)	
DENNIS, W. C. Fresh water in the ocean. (R. 1866)	

Deposit of—
arrow-heads near Fishkill, New York-E. M. SHEPARD. (R. 1877) 323
articles by Smithsonian in Corcoran Art Gallery. (R. 1874) 286
Beaufort Library—E. M. STANTON. (B. 1862)150
Bishop Johns' Library-E. CANBY. (R. 1862) 150
flint implements in Illinois—J. F. SNYDER. (R. 1876) 299
flint implements in Southern Illinois—C. RAU. (R. 1868) 224, 440, 370
flood-tide, law of—C. H. DAVIS33
Deposit, shell, in New Jersey—C. RAU. (R. 1864) 188, 440, 362
Deposits—See Shell deposits.
DE PRADOS, Baron. Eclipse of the sun, April 25, 1865. (R. 1864) 188
DE SAUSSURE, H.—
Electric resonance of mountains. (R. 1868) 224
Hymenoptera. (R. 1862) 150
Report on transactions of Society of Physics and Natural History of Geneva, 1870-1871. (R. 1871) 249
Synopsis of Vespidæ, (American wasps) 254
DE SCHLAGINTWEIT, H. Ethnographical collections. (R. 1862) 150
Description of—
a new fish, Apogon pandionis-G. B. Goode; T. H. Bean. (P. 1881) 467
a new fly-catcher and new petrel from Sandwich Islands—R. RIDGWAY. (P. 1881)
a new genus and species of Cottidæ—W. N. Lockington. (P. 1881) 467
a new gobioid fish (Othonops cos) from San Diego, California—R. Sмітн. (P. 1881) 467
a new owl from Porto Rico—R. RIDGWAY. (P. 1881) 467
birds—See Proceedings National Museum.
chitonidæ and acmæidæ from Panama—P. P. CARPENTER 252
country and people of Yoruba, Africa—T. J. Bowen 98
fishes—See Proceedings of National Museum.
fossil plants from China—J. S. NEWBERRY
Gobiesox rhessodon from San Diego, California—R. SMITH. (P. 1881) 467
human skull from Rock Bluff, Illinois-J. A. MEIGS, (R. 1867) 215
magnetic observatory at Smithsonian. (R. 1859) 110
meteorological instruments—L. CASELLA. (R. 1859) 110
new fishes—T. H. BEAN. (P. 1881) 467
new fishes—See D. S. Jordan; C. H. Gilbert.
new species of Centrarchide-C. L. McKAY. (P. 1881) 467
new subspecies of Loxigilla from island of St. Christopher, West Indies—G. N. LAWRENCE. (P. 1881) 467
new thrushes—R. RIDGWAY. (P. 1881) 467
observatories at Dorpat and Poulkova-C. ABBE. (R. 1867) 215, 369

Description of—Continued.
observatory at St. Martin, Isle Jesus, Canada East—C. Smallwood. (R. 1856)
plants collected by J. C. Frémont in California—J. Torrey 46
remains of extinct mammalia and chelonia from Nebraska—J. LEIDY 58
Descriptions of—
ancient works in Ohio-C. WHITTLESEY
several new races of American birds—R. RIDGWAY. (P. 1881) 467
Smithsonian anemometer—J. Henry. (R. 1860)
Squalius aliciæ from Utah Lake—P. L. Jouy. (P. 1881) 467
two new races of Myadestes obscurus—L. Stejneger. (P. 1881) 467
Desiderata, special, among North American birds, list of—R. RIDGWAY. (P. 1881)
Des Moines river, ancient pottery from-R. N. and C. Dahlberg. (R. 1879.)
Des Moines valley, mounds in—S. B. Evans. (R. 1879) 345
DESOR, E. Palafittes or lacustrian constructions, Lake Neuchâtel. (R. 1865.) 209, 360
Destruction of fish in Gulf of Mexico—J. Y. PORTER. (P. 1881) 45
Destruction of fish—See also Fish.
Destructive effect of iron rust. (R. 1861)
DEVEREUX, J. H. Ancient pottery from Arkansas. (R. 1872) 271
DEVEREUX, J. H. Catalogue of Indian relics presented to Smithsonian Institution. (R. 1872)
DEWEY, C.—
Best hours for temperature observations. (R. 1860) 14
· Best hours to find mean temperature. (R. 1857)
life of, by M. B. Anderson. (R. 1870)
The winds. (R. 1866) 214
Diagnoses des mollusques nouveaux provenant de Californie et faisant partie du Musée de l'Institution Smithsonienne—P. P. CARPENTER
Diagnoses of new forms of mollusks collected at Cape St. Lucas, Lower Cali-
fornia—P. P. CARPENTER 252
Diagnoses of new mollusks-
collected on west tropical shores of North America—P. P. CARPENTER. 252
from Reigen Mazatlan collection—P. P. CARPENTER 252
from Vancouver district—P. P. CARPENTER 252
from west coast of North America—P. P. CARPENTER 27.
from west tropical region of North America—P. P. CARPENTER 223
Diamond and other precious stones—J. BABINET. (R. 1870) 244, 377
Diary of an excursion in New Mexice—J. H. Carleton. (R. 1854) 5
Diatomacea—J. W. Bailey 23, 62
Diatomacea, instructions for collecting, preserving, and transporting—A. M. EDWARDS
Dickinson, A. B Eruption of volcano in Nicaragua. (R. 1867) 215

Berendt. (R. 1878)	275
° Oregon—G. Gibbs	
J BOWEN	
.V. W. Turner	98
. ag Smithsonian Institution-J. HENRY C	, 328
ri and Tennessee. (R. 1862)	•
earthworks of Ohio. (R. 1866)	214
por with increasing altitude—J. HANN. (R. 1877) 828	3, 898
rivers and streams—H. G. WEX. (R. 1875)	298
dreular to, relative to Morgan's research	188
1. of North America, catalogue of—R. Osten Sacken 102,	270
ons for collecting and preserving—R. OSTEN SACKEN	102
-tructions for collecting-H. LORW; R. OSTEN SACKEN. (R. 1858)	109
of Arctic America—S. H. Scudder	842
ra of North America, monographs of—	
Parts 1-111—H. Lorw	256
Part IV—R. OSTEN SACKEN	219
Directions for—	
auroral observations—J. Henry. (R. 1855)	77
collecting and preserving diptera-R. OSTEN SACKEN	102
collecting and preserving fish—T. H. BEAN. (P. 1881) 467,	464
collecting and preserving insects—A. S. PACKARD	261
collecting and preserving plants—L. F. WARD	460
collecting microscopic organisms—J. W. BAILEY	63
collecting, preserving, and transporting specimens of <i>Diatomacea</i> —A. M. EDWARDS	366
collecting, preserving, and transporting specimens of natural history-	
S. F. BAIRD. (R. 1856) 9	
constructing lightning-rods—J. HENRY	
earthquake observations—J. Henry. (R. 1855) (P. 1855)	
meteorological observations—A. Guyor; J. Henry. (R. 1855) 77, 19,	7.50
Directory of officers, collaborators, employés, etc., of the Smithsonian Institution, National Museum, Geological Survey, Bureau of Ethnology, and Fish Commission	466
Discourse on J. Henry—S. B. Dod	356
Discovery, meteorological—F. L. CAPEN. (R. 1866)	
Discovery of—	
a large meteorite in Mexico-W. M. Pierson. (R. 1878)	275
planet Neptune, report on history of the—B. A. Gould	18
stone image in Tennessee—E. M. GRANT. (R. 1870)	244

Discovery relative to magnetism, history of. (R. 1868) 18
Discussion of magnetic and meterological observations made at Girard College Observatory—A. D. BACHE:—
Part I. Eleven year period, solar diurnal variation 11
Part II. Solar diurnal variation and annual inequality 12
Part III. Influence of moon on declination13
Parts IV, V, VI. Horizontal component of magnetic force, etc 16
Parts vII, vIII, IX. Vertical force17
Parts x, xI, XII. Dip and total force 18
Parts I-XII. Complete19
Discussion of—
meteorological observations—See Schott.
Piazzi's astronomical observations—B. A. Gould. (R. 1863)18
relations of lucernarians to other acalephæ, beroids, and polypi—H. J.
Snell's barometric observations—F. H. Loud. (R. 1880) 442, 439
tables and charts of winds—A. Workor 26
Disease known as "chorea," inquiries relative to—S. W. MITCHELL. (R. 1874) 28
Disease, origin and propagation of—J. C. Dalton. (R. 1878) 276
Diseases of joints, bones, etc., bibliography of works on-W. W. KERN 80
Dispersion of cloud by an electrical discharge—D. W. NAILL. (R. 1858) 105
Distinction between tornadoes and tempests—J. B. LAMARCK. (R. 1871) 245
Distribution of—
duplicate fishes of Pacific coast, check-list of-D. S. JORDAN; P. L. JOUY. (P. 1881)
fishes of Alleghany region of South Carolina, Georgia, and Tennessee— D. S. Jordan; A. W. Brayton 308
forest trees in Montana, Idaho, and Washington—W. W. Johnson. (R. 1870)
forests and trees of North America-J. G. Cooper. (R. 1858) 109, 351
marine invertebrates from National Museum, list of—R. RATHBUN. (P. 1881)
Smithson income, majority and minority reports of Committee of Regents
on—J. A. PEARCE; J. MEACHAM. (R. 1858) 67, 329
specimens—See each annual report.
District of Columbia—
flora of-L. F. WARD. (P. 1881)
list of birds of—E. Cours; S. S. Prentiss. (R. 1861)
meteorology of—J. WISSNER. (R. 1857)107
prehistoric remains in T. R. PEALE. (R. 1872) 271
Ditrema atripes, new embiotocoid fish from coast of California—D. S. JORDAN; C. H. GILBERT. (P. 1880)
Documents, collection of historical, in Guatemala—C. H. BERENDT. (R. 1876) 299

Documents relative to origin and history of Smithsonian Institution—W. J. RHEES328
Dod, S. B. Memorial discourse on J. Henry
Dodge, G. W. Lightning discharges. (R. 1867) 215
Dodge, N. S. Memoir of C. Babbage. (R. 1878) 275
Dodge, N. S. Memoir of Sir John F. W. Herschel. (R. 1871) 249
Dog's revenge, a Dakota fable—S. R. Riggs. (E. 1879-80) 476
Dolichopodidæ, monograph of—H. Lorw
Domestic institutions in correspondence with Smithsonian Institution, list of 69, 238
Domestic institutions in correspondence with Smithsonian Institution, number of
Dominant language for science—A. DE CANDOLLE; J. E. GRAY. (R. 1874) 286 Dominica, catalogue of Ober's collections of birds from—G. N. LAWRENCE. (P. 1878) 382
Dominica, new species of Turdidæ from-G. N. LAWRENCE. (P. 1880) 425
Donati, G. B. Phenomena in telegraphic lines during aurora. (R. 1872) 271
Donations to library from foreign institutions, list of. (R. 1864) 188
Dorosoma cepedianum heterurum, (western gizzard shad,) notes on-S. Wil-
мот. (Р. 1878)
Dorpat and Poulkova, description of observatories at—C. ABBE. (R. 1867) 215, 369
Dorsey, J. O. Illustrations of method of recording Indian languages. (E. 1879-80)
Dorysomatidæ, notes on—D. S. Jordan
Douglas, S. A., eulogy on, by S. S. Cox. (R. 1861) 149
Douglass, H. M. Translation of Weismann on change of Mexican axolotl. (R. 1877) 823, 401
Downes, J. Occultations of planets and stars by the moon, 1853 54
Downes, J. Occultations visible in the United States, 1848-1852 8, 9, 10, 11, 29
DOYLE, W. E. Indian forts and dwellings, Indian Territory. (R. 1876) 298
Drainage, sanitary, of Washington city, suggestions for-G. E. WARING 849
DRAPER, H., account of telescope of, by T. W. WEBB. (R. 1864)
DRAPER, H. Construction of silvered glass telescope and its use in celestial photography180
DREUTZER, O. E. Statistics relative to Norwegian mountains, lakes, and snow line. (R. 1866)
Drift, fresh water glacial, of the Northwestern States—C. WHITTLESEY 197
Drilling in stone without metal—C. RAU. (R. 1868) 224, 440, 372
Drugs, classification of forms in which medicines and, appear and are adminis-
tered—J. M. FLINT. (P. 1881) 467, 451
Drugs, memoranda for collectors of—J. M. FLINT. (P. 1881) 467, 452
Dual character of the brain. Toner lecture No. 11—C. E. Brown-Séquard 291
Duby, P. A. Report on transactions of Society of Physics and Natural History
of Geneva. 1861. (R. 1864)

Duck, new to North American fauna—R. RIDGWAY. (P. 1881)

DUDLEY, T. Earthquake at New Madrid, Missouri. (R. 1858)	103
DUFOUR, C.; KÄMTZ. Scintillation of the stars. (R. 1861)	
Dughs, A., notes on fishes collected by, in Mexico-T. H. BEAN. (P. 1879)	
Dughs, A., notes on fishes collected by, in Mexico-D. S. Jordan. (P. 1879).	
Duke of Northumberland, account of	
Duke of Northumberland, books presented by. (R. 1859)	
Dumas, J. B. Eulogy on A. A. De La Rive. (R. 1874)	
Duncan, M. L. Translation of Holmgren on color-blindness. (R. 1877.) 323,	
Dunkirk Society for the Encouragement of Sciences. Prize questions. (R. 1865; R. 1873)	
DUNNING, E. O. Antiquities in Tennessee. (R. 1870)	
Duplicate fishes distributed by Smithsonian—T. H. BEAN. (P. 1880)	425
Duplicate shells collected by the United States Exploring Expedition under C. Wilkes	
DUPRÉ, W.; HENRY, J. Earthquakes in North Carolina, 1874. (R. 1874)	285
DUPREZ, F. Atmospheric electricity. (R. 1858)	109
Dwellings and forts, Indian, in Indian Territory—W. E. DOYLE. (R. 1876)	299
Dyeing purple, ancient and modern. (R. 1868)	187
•	
. B.	
Karth—	 .
chemistry of the—T. S. HUNT. (R. 1869) 228, 3	
electrical currents of the—C. MATTEUCCI. (R. 1867; R. 1869) 215,	
figure of the—St. M. MERINO. (R. 1868)	104
internal structure of the—J. G. BARNARD	
laws of atmospheric circulation over the—J. H. Coffin	
revolutions of crust of the—G. PILAR. (R. 1876)	
secular variation of elements of orbit of-J. N. STOCKWELL	Zó.
Earthquake—	1/16
at New Madrid, Missouri—T. Dudley. (R. 1858)	
directions—J. Henry. (R. 1855)	214
(an 1000) tables	
in Peru, August 13, 1868—J. V. CAMPBELL. (R. 1870)	
phenomena, on observations of—R. MALLET. (R. 1859)	! 1A
Earthquakes— articles on, received by Smithsonian Institution and deposited in Library	
· · · · · · · · · · · · · · · · · · ·	249
	48

Washanahan Cantinual
Karthquakes—Continued. in Guatemala—A. CANUDAS. (R. 1858) 10
in North Carolina, 1874—W. Du Pré; J. Henry. (R. 1874)
in St. Thomas—G. A. LATIMER. (R. 1867) 21
Earthwork, double-walled, in Ashtabula county, Ohio—S. D. PEET. (R. 1876) 29
Earthworks—
ancient, of Ashland county, Ohio—G. W. HILL. (R. 1877)
ancient, of Ohio, sketch of—I. DILLE. (R. 1866) 216
ancient, on upper Missouri—A. BARRANDT. (R. 1870) 24
in Tennessee—J. Jones
in Wisconsin—E. E. Breed. (R. 1872) 27
on Arkansas river—Mrs. G. KNAPP. (R. 1877) 823
Earthworks—See Mounds.
Echinoderms of Kerguelen Island—A. E. VERRILL 294
Echinoderms of northeastern coast of America—A. E. VERRILL. (P. 1879;
P. 1880)
Eclipse
annular, of May 26, 1854
of the sun, April 25, 1865—Baron DE PRADOS. (R. 1864) 186
of the sun, March 15, 1858, map of—T. HILL
of the sun, July 18, 1860—J. LAMONT. (R. 1864) 186
of the sun, September 7, 1858, in Peru—J. M. GILLISS 100
Ecliptic, obliquity of the-J. N. STOCKWELL 289
Economic geology of Trinidad-G. P. Wall; J. G. Sawkins. (R. 1856) 9
Education—
acknowledgment of books on—E. LABOULAYE. (R. 1867) 216
in the United States, project of an outline history of—F. A. PACKARD. (R. 1863) 18'
scientific, of mechanics and artisans—A. P. PEABODY. (R. 1872) 271, 380
Education, Illinois State Board of. Meteorological system for every State. (R. 1855)
Educational series of marine invertebrates distributed by United States National Museum—R. RATHBUN. (P. 1881) 467, 465, 471
Edwards, A. M. Directions for collecting, preserving, and transporting diatomacea 366
Edwards, A. M. Results of examination under microscope of Japanese infusorial earths
Edwards, V. N. Occurrence of oceanic bonito (Orcynus pelamys) in Vineyard Sound, Massachusetts. (P. 1878) 335
Edwards, W. H. Diurnal lepidoptera of Arctic America
Edwards, W. H. On lepidoptera
Eels, genitalia of male—S. T. CATTIE. (P. 1880) 428
Effect of irritation of polarized nerve—B. F. LAUTENBACH. (R. 1878) 841, 411
Effect of lightning—S. L. HILLIER. (R. 1866)

Effect of moon on the weather—J. HENRY. (R. 1871)	249
EGGERS, H. F. A. Flora of St. Croix and Virgin Islands	313
Egging expedition to Shoal Lake, Lake Winnipeg-D. Gunn. (R. 1867)	215
Eggs—	
instructions for collecting—S. F. BAIRD. (R. 1858)	139
instructions for collecting—A. NEWTON	139
instructions for collecting and preserving—T. M. BREWER	139
North American—T. M. Brewer	89
of the eight North American species of Empidonaces—T. M. Brewer. (P. 1879)	33 5
suggestions for forming collections of-A. NEWTON	139
Egleston, T.—	
Catalogue of minerals with their formulas	156
Scheme for qualitative determinations by blowpipe. (R. 1872)	271
translation by, of Brezina on crystallography	
Egypt, presentation of books on—R. LEPSIUS. (R. 1860)	
Elastic force of aqueous vapor, table for determining—J. H. COFFIN	
Electric phenomenon—W. F. GIVEN. (R. 1865)	
Electric resonance of mountains, observations on—H. DE SAUSSURE. (R. 1868.)	
Electrical—	
currents of the earth—C. MATTEUCCI. (R. 1867; R. 1869) 215,	23
discharge, dispersion of a cloud, by D. W. NAILL, (R. 1858)	109
rheometry, researches on-A. Secchi	36
Electricity—	
accounts of lightning discharges. (R. 1867)	215
and galvanism, report on recent progress in—J. Müller. (R. 1857)	
and magnetism, later views of connection of—H. Helmholtz; J. C. Maxwell. (R. 1878)	
atmospheric—F. Duprez. (R. 1858)	
considerations on. (R. 1867)	
of atmosphere and the aurora borealis—S. Lemström; A. A. De La Rive. (R. 1874)	
of induction in clouds resolving into rain, snow, and hail—F. ZANTE- DESCHI. (R. 1870)	
phenomena accompanying propagation of, in rarefied elastic fluids—A. A.	187
presence of, during fall of rain—Prof. PALMIERI. (R. 1870)	244
-	91
report or recomb brogress in the management (and recomb management)	
Electro-magnetic—	24 1
seismograph—Prof. Palmieri. (R. 1870) telegraph, Henry's contribution to—W. B. Taylor. (R. 1878) 841, 4	
telegraph—See Henry; Telegraph. Electro-physiology lectures on—C. MATTEUCCI. (R. 1865)	209

Electrotypes of engravings of shells granted by British Museum. (R. 1868) 18
Elements, table of chemical equivalents of. (R. 1864) 1864
Elephant mound in Grant county, Wisconsin-J. WARNER. (R. 1872) 27
ELLERY, R. L. J. Address of the President of the Royal Society of Victoria. (R. 1868)
ELLET, C. Physical geography of Mississippi valley13
ELLIOT, D. G. Classification and synopsis of Trochilide
Ellior, D. G. List of described species of humming birds 334
ELLIOTT, R. S. Climate of Kansas. (R. 1870) 24
ELLSWORTH, E. W. Ancient implement of wood from Connecticut. (R. 1876) 299
Embarrass, Wisconsin, pits at—E. E. Breed. (R. 1877)
Embryology of insects—L. Agassiz
Empidonaces, notes on nests and eggs of the eight North American species of— T. M. Brewer. (P. 1879)
Employés of Smithsonian Institution, National Museum, Geological Survey, Bureau of Ethnology, and Fish Commission, directory of460
Encke, J. F., memoir of, by G. HAGEN. (R. 1868) 224
Endlich, F. M.—
Analysis of water destructive to fish in Gulf of Mexico. (P. 1881) 46
Geology of Kerguelen Island 29
List of minerals in National Museum, 1873, 1879. (R. 1873; P. 1880) 275, 42
Endothyra ornata, note on—C. A. WHITE. (P. 1879) 88
ENGELHARDT, M. Formation of ice at the bottom of water. (R. 1866) 214
ENGELMANN, G. Heights of mountains in Colorado. (R. 1862) 156
Engineers, Civil, London Institution of. Prize questions. (R. 1862) 156
England—
abbreviations used in—W. DE LA RUE. (R. 1867) 216
free freights between the United States and—E. CUNARD. (R. 1859) 110
Salisbury, notice of Blackmore Museum at. (R. 1868) 224
English and French weights and measures—See Tables.
English vocabulary with comparative words in Spanish, French, and Latin 170
Engravings
of shells, electrotypes of, granted by British Museum. (R. 1863) 186
on the face of rocks in the Sierra Nevada—J. G. Bruff. (R. 1872) 273
presented to Smithsonian Institution, catalogue of—C. B. KING. (R. 1861.) 149
Entomologists, circular to 178
Entomologists, instructions to 261
Eccene and oligocene invertebrate fossils of North America, check-list of—T. A. CONBAD
EGFF, J. Habits of black bass of the Ohio. (R. 1854)70
Ephemeris of the planet Neptune, 1848-1852—S. C. WALKER 4, 5, 6, 7, 24
Epinephelus Drummond-Hayi, a new serranoid fish from the Bermudas-G. B.
GOODE: T. H. BEAN. (P. 1878) 889

Epinephelus nigritus of the southern coast, note upon—G. B. Goode; T. H. Bean. (P. 1878)
Equilibrium of a liquid mass withdrawn from the action of gravity—See J. Plateau.
Equinoxes, precession of the-J. G. BARNARD
Equinoxes, precession of the—J. N. STOCKWELL
Equivalents, chemical, of sixty-three elements, table of. (R. 1864)
ERNST, G. A. Meteorology of Caracas, South America. (R. 1867)
Erosions of the earth's surface by rivers, etc.—E. HITCHCOCK
Eruption of volcano in Nicaragua—A. B. Dickinson. (R. 1867)
Eruption of volcano of Colima—C. SARTORIUS. (R. 1869)
Eryontidæ, notice of recent—S. I. SMITH. (P. 1879)
ESPY, J. P., notice of, by A. D. BACHE. (R. 1859)
Espy, J. P. On meteorology. (R. 1847)
Essay on geographical distribution of batrachia and reptilia—E. D. Cops
Essay on velocity of light—C. DELAUNAY. (R. 1864) 188,
Establishment and officers of the Smithsonian Institution and National Museum.
Establishment of the Smithsonian Institution, Journal and By-Laws of. (R.
1858) 67,
Estes, L. C. Antiquities of Minnesota, Mississippi river, and Lake Pepin.
(R. 1866)
Estimate of population of the world—E. MAILLY. (R. 1878)
Etheostomatidæ, notes on—D. S. JORDAN
Ethmocardium, note on—C. A. WHITE. (P. 1879)
Ethnographical—
collections—T. LYMAN. (B. 1862)
collections—H. DE SCHLAGINTWEIT. (R. 1862)
collections, the Schlagintweit—H. ZISGENBALS. (R. 1867)
Ethnological—
collections of museum at Lausanne, report on—F. TROYON. (R. 1861).
department of the French Exposition, 1867. (R. 1867)
map of North America, suggestions relative to-L. H. MORGAN. (R. 1861)
map of the United States—G. GIBBS. (R. 1862)
memoir of Squier and Davis, correspondence relative to acceptance of,
for publication
museum of Norway-L. K. DAA. (R. 1862)
research—E. H. DAVIS. (R. 1866)
specimens presented by University of Christiania, Norway. (R. 1863) I
suggestions for Russian America—G. GIBBS
Ethnology and archeology, circular relative to collections in
Ethnology and philology, instructions for—G. GIBBS16
Ethnology, articles on. (R. 1867–1870–1877, 1879, 1880) 215, 244, 249, 271, 275, 28

Ethnology— Leipsic Museum of—O. T. Mason. (R. 1873)
of Arctic America—L. Kumlien
of Colonia Tovar, Venezuela, South America—A. FENDLER. (R. 1857) 107
of Indians of Red river of the North—W. H. GARDNER. (R. 1870) 244
physical, lectures on—D. Wilson. (R. 1862) 150
present state of, in relation to form of human skull-A. RETZIUS. (R.
1859) 110 Ethnology, Bureau of—
catalogue of linguistic manuscripts in library of—J. C. Pilling. (B. 1879-80)
directory of officers and employés of 466
first annual report of—J. W. Powell 476
Euchalarodus Putnami, identity of, with Pleuronectes glaber—T. H. BEAN. (P. 1878)
Eulachon or candle fish of northwest coast—J. G. Swan. (P. 1880) 425
Eulogy on—
Ampère, by F. Arago. (R. 1872)271
A. D. Bache, by J. HENRY. (R. 1870) 244, 379
A. A. De La Rive, by J. B. Dumas. (R. 1874) 286
Stephen A. Douglass, by S. S. Cox. (R. 1861) 149
Cornelius C. Felton, by T. D. Woolsey. (R. 1861) 149
Joseph Fourier, by F. Arago. (R. 1871) 249
Herschel, by F. Arago. (R. 1870) 244
Gay-Lussac, by F. Arago. (R. 1876)
La Place, by F. Arago. (R. 1874)
Quetelet, by A. MAILLY. (R. 1874) 286
James A. Pearce, by A. D. BACHE. (R. 1862) 150
Gen. J. G. Totten, by J. G. BARNARD. (R. 1865) 209
Alexander Volta, by F. Arago. (R. 1875) 298
Henry Wilson, by P. PARKER. (R. 1875)
Thomas Young, by F. ARAGO. (R. 1869)
Eulogy—See Memoir, Life, Biography.
Europe—
causes which limit vegetable species towards the north, in—A. DE CANDOLLE. (R. 1858)109
certain storms in America and, December, 1836—E. Loomis 127
man as the contemporary of the mammoth and reindeer in. (R. 1867) 215
study of high antiquity in—A. Morlot. (R. 1862; R. 1864) 150, 188
Europe, middle, fauna of, during stone age—L. RÜTIMEYER. (R. 1861) 149
EVANS, J. Climate of Colorado. (R. 1865) 209
EVANS, S. B. Notes on mounds in Des Moines valley. (R. 1879) 845
Evaporation in Florida—W. C. DENNIS. (R. 1866)

Evaporation observed at Palermo in 1865 and 1866—P. TACCHINI. (R. 1870.) 244
EVERETT, E. Report on Jewett's general stereotype catalogue of public libraries. 43
EVERETT, E. Report on organization of Smithsonian Institution. (R. 1853.) 63
EVERETT, J. D. Under-ground temperature. (R. 1874) 28
Evolution of language—J. W. Powell. (E. 1879-80)
Examination—
microscopical, of soundings—J. W. BAILEY
of J. Henry by English Scientific Commission 32
of specimens, rules for. (R. 1880) 44
of Spencer's telescope for Hamilton College. (R. 1855)
Excelsior, Minnesota, mounds near—F. H. NUTTER. (R. 1879) 34
Exchange of publications—
Agricultural Association of Milan. (R. 1868)
Chamber of Commerce of Bordeaux. (R. 1863) 18
Royal Horticultural Society, London. (R. 1861)
Exchange of specimens—
Hamburg Zoölogical Gardens. (R. 1867)
Hamilton College, Clinton, New York. (R. 1861)
W. A. LLOYD. (R. 1867)
Museum of National University of Greece. (R. 1867) 21
University of Costa Rica. (R. 1867)
American Academy of Arts and Sciences, Boston, thanks for. (R. 1855; R. 1867)
Bath and West of England Society for Encouragement of Agriculture, Arts, etc. (R. 1867)
circular relative to scientific and literary 320
Government of Bremen. (R. 1865) 20
W. Hincks. (R. 1860)
international, report on-G. H. BOEHMER477
list of. Parts 1, 11 73, 85
list of, to 1858115
Mexican Society of Geography and Statistics. (R. 1861, 1865) 149, 20
Mining Department, Melbourne. (R. 1865)
F. MÜLLER. (R. 1860)
reports on. (R. 1853–1867, 1873) 67, 75, 77, 91, 107, 109, 110, 147, 149, 150, 187, 188, 209, 214, 215, 27
J. Rosing. (R. 1865)
Royal Academy of Science, Madrid. (R. 1861)
St. Petersburg Academy of Sciences. (R. 1867) 212
St. Petersburg Academy of Sciences. (R. 1867)

Executive Committee of Regents, journal and reports of, 1846-1876	829
Exhibit by the Smithsonian Institution at Centennial Exhibition, report on	
proposed plan of—S. F. BAIRD. (R. 1875)	298
Exhibit of fisheries of United States at Berlin Fisheries Exhibition—G. B.	
Goodk	413
Exhibition, Centennial, report on—S. F. BAIRD. (R. 1876) 299,	307
Expansion, bibliography of—F. W. CLARKE	255
Expansion by heat, tables of—F. W. CLARKE	289
Expedition—	
Coast Survey, for transatlantic longitude—B. A. Gould	228
Hassler, narrative of the-L. Agassiz. (R. 1872)	271
North Pacific Surveying, contributions to natural history made in con-	
nection with the-T. H. STREETS	808
to Arctic Seas-See Hayes, Kane, McClintock.	
to Kerguelen Island—J. H. KIDDER and others 298	, 294
to Lake Winnipeg-D. GUNN. (R. 1867)	215
to Mexico, scientific. (R. 1864)	188
to the Mauvaises Terres and Upper Missouri—T. A. CULBERTSON. (R. 1850)	28
toward the North pole, scientific instructions for—J. HENRY and others.	20
(R. 1871)	249
Expeditions, account of—See each annual report.	
Expeditions contributing specimens, list of—S. F. BAIRD. (R. 1867)	215
Experimental and theoretical researches on figures of equilibrium—See J. Plates	
Experiments	
on aneroid barometers at Kew Observatory—B. STEWART. (R. 1868)	224
relative to meteorites—G. A. DAUBRÉE. (R. 1868)	
upon animal heat of fishes—J. H. KIDDER. (P. 1879)	
Explanation of principles of crystallography and crystallophysics—A. Brezina.	
(R. 1872) 271,	386
Exploration of ancient mounds in Union county, Ky.—S. Lyon. (R. 1870)	244
Explorations—	
among Indian mounds in southern Florida—S. T. WALKER. (R. 1879.)	345
and surveys, Government, report of—S. F. BAIRD. (R. 1878)	841
Arctic, lecture on—I. I. HAYES. (R. 1861)	149
articles on—See each annual report.	
botanical, in New Mexico and California, account of—A. GRAY. (R. 1849)	21
furnishing collections to National Museum, 1888-1877, list of—S. F. BAIRD. (R. 1877)	828
in Central America—C. H. BERENDT. (R. 1867)	215
	341
· · · · · · · · · · · · · · · · · · ·	442
THE ATOM PROTECT OF THE PROTECTION OF THE PROTEC	
in Greenland—L. Kumlien. (R. 1878)in New Mexico and Arizona—J. Stevenson. (R. 1880)	

Explorations—Continued.
in New York—E. G. SQUIER.
in Tennessee-E. A. DAYTON. (R. 1870) 24
in upper California in 1860—J. FEILNER. (R. 1864)
Kennicott's-Hudson's Bay Company. (R. 1863) 18
of aboriginal remains of Tennessee-J. Jones 259
of the Nile—C. HALE. (R. 1865)
of western Missouri in 1854—P. R. Hov. (R. 1864) 18
of John Xantus in Mexico-M. Romero. (R. 1862) 15
on western coast of North America—W. H. Dall. (R. 1878) 278
reports on—S. F. Baird. (R. 1851-1866, 1875-1877) 51, 57, 75, 77, 91, 107, 109, 110, 147, 149, 150, 187, 188, 209, 214, 298, 299, 223
scientific, in America, in 1852—S. F. BAIRD. (R. 1852) 57
scientific, in Mexico. (R. 1864)
Exploring expedition under Capt. C. Wilkes, duplicate shells collected by the. 193
Explosibility of coal oils—Z. Allen. (R. 1861)
Explosiveness of nitre—R. HARE
Exposition of harmonies in the solar system—S. ALEXANDER280
Exposition—See French.
Expression for anthropological inquiry, queries about—C. DARWIN. (R. 1867) 215
External appearance of the sun's disk. (R. 1866) 214
Extinct—
mammalia and chelonia of Nebraska—J. LEIDY58
reptiles—J. Leidy 192
sloth tribe of North America—J. LEIDY
species of American ox—J. Leidy41
Eye, bibliography of diseases of the—W. W. KEEN 300

F.

Facts respecting Liberia College—A. CRUMMELL. (R. 1861)	149
Fahrenheit's scale, tables for conversion of centigrade degrees to. (R. 1863)	187
Families—	
of fishes, arrangement of—T. GILL	247
of mammals, arrangement of—T. GILL	230
of mollusks, arrangement of—T. GILL	227
Family, human, systems of consanguinity and affinity of-L. H. Morgan 188,	218
Fanning Islands, natural history of—T. H. STREETS	303
Faraday, M., his life and works—A. A. DE LA RIVE. (R. 1867)	215
FARLOW, W. G.—	
Algæ of Arctic America	842
Algæ of Kerguelen Island	294
Recent progress in botany, 1879, 1880. (R. 1880) 442,	4 30
Report on water from Gulf of Mexico. (P. 1881)	467
Farming and gardening, natural history as applied to—J. G. Morris. (R. 1855.)	77
Faroe Isles, vegetable colonization of—C. MARTINS. (R. 1858)	109
FARQUHARSON, R. J. Study of skull and long bones from mound in Illinois. (R. 1874)	286
Fauna-	
and flora within living animals—J. LEIDY	44
littoral marine, of Provincetown, Mass.—R. RATHBUN. (P. 1880)	425
of middle Europe during the stone age—L. RÜTIMEYER. (R. 1861)	149
of Nebraska, ancient—J. Leidy	58
FAVRE, A. Report on transactions of Geneva Society of Physics and Natural History, July, 1876, to June, 1877. (R. 1877)	323
FAVRE, E. Biographical notice of Louis Agassiz. (R. 1878)	341
Feeling, sense of. (R. 1865)	209
FEILNER, J. Explorations in upper California in 1860. (R. 1864)	188
Feldspar, determination of, in thin sections of rocks—G. W. HAWES. (P. 1881)	467
Felton, C. C., eulogy on, by T. D. Woolsey. (R. 1861)	149
Felton, C. C.—	
Notice of Washington Irving. (R. 1859)	110
Notice of W. W. Turner. (R. 1859)	110
Report on Prof. Henry and the telegraph. (R. 1857)	107
Fendler, A., botanical explorations by, in New Mexico and California—A. GRAY. (R. 1849)	21
Fendler, A.—	
Meteorology and ethnology of Colonia Tovar, Venezuela. (R. 1857)	107
Meteorology of Colonia Tovar. (R. 1866)	214
Temperature of St. Louis, Missouri. (R. 1860)	147

Ferns, acknowledgment for-G. METTENIUS. (R. 1862)	150
FERREL, W. Converging series expressing ratio between diameter and circumference of a circle	233
FERREL, W. Reply to the criticisms of Dr. J. Hann. (R. 1877) 323,	398
Ferrel, W., theories of, as to relation between difference of pressure and velocity of wind-J. HANN. (R. 1877) 823,	398
Fever—	
a study in morbid and normal physiology—H. C. Wood	357
researches on—H. C. Wood. (R. 1878)	341
wood	282
Fevers, surgical complications and sequels of. Toner lecture No. v—W. W. Kern	300
FIALHO, A. Biographical sketch of Dom Pedro II. (R. 1876)	200
FIGANIERRE, M. Account of remarkable accumulation of bats. (R. 1868)	187
Figure of the earth—St. M. MERINO. (R. 1863)	187
Figures of equilibrium of a liquid mass withdrawn from the action of gravity— See Plateau.	
Figures, series of, for labels	164
Filices of Kerguelen Island—A. GRAY	294
Finances of the Institution, 1846-1877	329
FINCH, FLADGATE, and CLARKE. Residuary legacy of Smithson. (R. 1861.) 149,	328
FINCE, H. Antiquities in the State of Vera Cruz, Mexico. (R. 1870)	244
Fire at Smithsonian, January, 1865, report of Committee of Regents relative to—R. Wallach; J. Henry. (R. 1864)	
Fire, Indian mode of obtaining—G. CROOK. (R. 1871)	249
Fire-alarm telegraph, American, lecture on-W. F. CHANNING. (R. 1854)	75
Fire ball, meteoric, orbit and phenomena of-J. H. Coffin	22 1
Fire-proofing buildings, architecture in relation to—D. B. REID. (R. 1856)	91
First decade of United States Fish Commission—G. B. Goode. (R. 1880)	449
FISCHER, F. Scientific labors of Edward Lartet. (R. 1872)	271
Fish Commission—	
account of work of—S. F. BAIRD. (R. 1880)	
catalogue of exhibit of, at Centennial, 1876—G. B. Goode	
directory of officers and employés of	
exhibit by, at Berlin Fisheries Exhibition—G. B. Goods	413
ILIDUAD. (A. 1010)	33 }
first decade of—G. B. Goode. (R. 1880)	442
list of marine invertebrata from New England coast distributed by-	333
mollusca collected by-A. E. VERRILL. (P. 1880)	125
notice of crustacea dredged by, off south coast of New England—S. I. SMITH. (P. 1880)	

Fish—
analysis of water destructive to, in Gulf of Mexico—F. M. Endlich (P. 1881)
candle, of northwest coast—J. G. SWAN. (P. 1880)
dates of first appearance of-F. B. Hough
description of new species of, (Apogon pandionis)—G. B. Goode; T. H. Bean. (P. 1881)
destruction of, by poisonous water in Gulf of Mexico—J. Y. Porter (P. 1881)
destruction of, by polluted waters in Gulf of Mexico—W. C. W. GLAZIER. (P. 1881)
directions for collecting and preserving—T. H. BEAN
fresh, circular on shipping—S. F. BAIRD
from deep-sea fauna of western Atlantic, description of new species of (Alepocephalus Bairdii)—G. B. Goode; T. H. Bean. (P. 1879)
mortality of, in Gulf of Mexico—E. Ingersoll. (P. 1881)
mortality of, in Gulf of Mexico-M. A. Moore. (P. 1881)
new, from Alaska, with notes on genus Anarrhichas—T. H. BEAN. (P. 1879)
new, (Lopholatilus chamæleonticeps,) from south of New England—G. B. Goode; T. H. Bean. (P. 1879)
new gobioid, (Othonops eos,) from San Diego, Cal.—R. SMITH. (P. 1881)
new serranoid, (Epinephelus Drummond-Hayi,) from Bermudas and Florida—G. B. Goode; T. H. Bean. (P. 1878)
new sparoid, (Sargus Holbrookii,) from Savannah Bank, description of— T. H. BEAN. (P. 1878)
ish-culture of United States, exhibit of, at Berlin Fisheries Exhibition—G. B. Goode
ISHER, J. G. Acknowledgment of perennibranchiates. (R. 1868)
isheries—
of United States, catalogue of collection to illustrate, at Philadelphia—G. B. Goode
of United States, exhibit of, at Berlin Fisheries Exhibition—G. B. Goode.
isheries Exhibition, Berlin, additions to Museum from. (R. 1880)
iahes—
Alaskan, catalogue of-T. H. BEAN. (P. 1881)
American, in British Museum and Museum d'Histoire Naturelle, Paris, notes on typical—D. S. JORDAN. (P. 1879)
arrangement of families of-T. GILL
bibliography of-T. GILL
check-list of duplicates of, distributed by Smithsonian Institution—T. H. BEAN. (P. 1880)
collected by H. E. Nichols in British Columbia and Alaska, notes on— T. H. BEAN. (P. 1881)

Fishes—Continued.

cottoids, monograph of the—C. GIRARD 30
description of new species of North American—D. S. Jordan. (P. 1879.) 33
descriptions of and notes on—See D. S. JORDAN; C. H. GILBERT.
descriptions of deep-sea species from New England, diagnoses of two undescribed genera of flounders and genus related to Merlucius—G. B. Goode. (P. 1880)
descriptions of new species of, (Uranidea marginata, Potamocottus Bendirei,) and of Myctophum crenulare—T. H. BEAN. (P. 1881) 457
directions for collecting and preserving-T. H. BEAN. (P. 1881) 45
European, in National Museum, list of—T. H. BEAN. (P. 1879) 33
experiments upon animal heat of—J. H. KIDDER. (P. 1879)
from deep waters on south coast of New England, obtained by Fish Commission in 1880—G. B. Goode. (P. 1880)
from eastern Georgia, notes on—T. H. BEAN. (P. 1879) 3%
from eastern Mississippi, collection of—O. P. HAY. (P. 1880) 423
from Hudson's Bay, notes on—T. H. BEAN. (P. 1881) 47
from Utah Lake, notes on a collection of—D. S. JORDAN; C. H. GIL- BERT. (P. 1880)
mortality of, in Gulf of Mexico-J. P. JEFFERSON. (P. 1878) 32
mortality of, in vicinity of Tortugas—J. P. Jefferson; J. Y. Porter; T. Moore. (P. 1878)
new genus of, (Benthodesmus)-G. B. Goode; T. H. Bean. (P. 1881.)
new genus of, (Delolepis)-T. H. BEAN. (P. 1881)
North American, check-list of duplicates distributed by Smithsonian— T. H. BEAN. (P. 1880)
North American, review of Rafinesque's memoirs on—D. S. JORDAN 301
notes on—D. S. Jordan 305, 306. 206
notes on collection of, from Clackamas river, Oregon—D. S. JORDAN. (P. 1878)
of Alaska, some genera and species of—T. H. BEAN. (P. 1879) 383
of Alaska and Siberia, descriptions of new-T. H. BEAN. (P. 1881) 467
of Alleghany region of South Carolina, Georgia, and Tennessee, distribution of—D. S. JORDAN; A. W. BRAYTON
of Arctic America—T. H. BEAN
of Beaufort Harbor, North Carolina, notes on—D. S. Jordan; C. H. Gilbert. (P. 1878)
of Bermuda, catalogue of-G. B. Goode-296
of Bermuda mistakenly described as new by Günther—G. B. Goods. (P. 1878)
of California—T. H. STREETS
of California, descriptions of new genera and species of—W. N. Lock- INGTON. (P. 1879)
of east coast of North America, catalogue of—T. GILL 25

Fishes—Continued.
of Fanning Islands-T. H. STREETS 808
of Florida, description of new species of, (Seriola Stearnsii)—G. B. Goode. (P. 1879)838
of Florida, preliminary catalogue of, with new genus and three new species—G. B. Goode. (P. 1879)888
of Gulf of Mexico, collected by J. W. Velie, catalogue of and description of seven new species—G. B. Goode; T. H. Bean. (P. 1879) 338
of Hawaiian Islands—T. H. STREETS
of Kerguelen Island—T. GILL
of Mexico, descriptions of two species of, collected by Dugès—T. H. Bean. (P. 1879)
of Mexico, notes on Dugès collection of-D. S. Jordan. (P. 1879) 338
of New York—T. Gill. (R. 1856)91
of Pacific coast of United States, bibliography of-T. GILL468
of Pacific coast and Alaska, bibliography of-T. H. BEAN. (P. 1881) 467
of Pacific coast, check-list of duplicates of, distributed by Smithsonian Institution in 1881—D. S. Jordan; P. L. Jouy. (P. 1881) 467
of Pacific coast, notes on -D. S. JORDAN; C. H. GILBERT. (P. 1881) 467
of Pacific coast of United States, list of—D. S. JORDAN; C. H. GILBERT. (P. 1880)425
of Pensacola, Florida, catalogue of Stearn's collection and description of new species—G. B. Goode; T. H. Bean. (P. 1879)
of St. John's river—G. B. Goode. (P. 1879)
of Samoan Islands—T. H. STREETS
on coast of New Jersey and Long Island—S. F. BAIRD. (R. 1854) 75, 348
pediculate, of eastern coast of extratropical North America, synopsis of— T. GILL. (P. 1878) 832
Potamocottus Bendirei, description of-T. H. BEAN. (P. 1881) 467
trunk, Ostraciontidæ, a study of, with notes on American species—G. B. Goode. (P. 1879) 838
Uranidea marginata, description of-T. H. BEAN. (P. 1881) 467
Fishes, food—See G. B. Goode, T. H. Bean, D. S. Jordan, S. Stearns, F. Stein-dachner, S. Wilmot.
Fishes, food—
circular relative to—S. F. BAIRD 284
memoranda of inquiry relative to—S. F. BAIRD 231
questions relative to—S. F. BAIRD
Fishkill, New York, deposit of arrow-heads near—E. M. SHEPARD. (R. 1877.) 323
Fissirostres, oölogy of—T. M. Brewer.
FLACHENECKER, G. Indian languages. (R. 1862) 150
FLADGATE, CLARKE, and FINCH. Smithson's residuary legacy. (R. 1861.) 149, 328
Flamingo from south Florida, specimens of—G. WURDEMAN. (R. 1860) 147
Flight in the animal kingdom, phenomena of—E. J. MAREY. (R. 1869) 228

Flight, modes of, in relation to aeronautics—J. B. Petrigrew. (R. 1867) 215
FLINT, E., shells from Costa Rica kitchen-midden collected by—W. H. Dall. (P. 1878)
FLINT, J. M.—
Classification and arrangement of the materia medica collection 450
Classification of forms of drugs and medicines 451
Memoranda for collectors of drugs 452
Flint implements—
in Illinois, agricultural—C. RAU. (R. 1868)
in Illinois, deposits of—J. F. SNYDER. (R. 1876) 200
in Ohio, Holmes county—H. B. CASE. (R. 1877)
Flood-tide, law of deposit of—C. H. DAVIS33
Flora
and fauna within living animals-J. LEIDY 44
of Alaska, sketch of the-J. T. ROTHROCK. (R. 1867) 215, 367
of Crozet and Kerguelen Islands—J. H. KIDDER and others994
of North America, index of authorities for species of—S. WATSON 258
of St. Croix and the Virgin Islands—H. F. A. EGGERS
of Washington and vicinity—L. F. WARD 444
Florida—
antiquities of—J. BARTRAM. (R. 1874) 286
antiquities of—A. MITCHELL. (R. 1874)
catalogue of casts of Indian prisoners in—R. H. PRATT. (P. 1878) 332
colored bead from mound in—A. M. Habrison. (R. 1877)
Curtiss' collection of fishes from-D. S. JORDAN. (P. 1880)425
evaporation in—W. C. DENNIS. (R. 1866)
fishes of-G. B. GOODE; T. H. BEAN. (P. 1878) \$52
fishes of, preliminary catalogue and new genus and species of—G. B. Goode. (P. 1879)
forests and trees of—J. G. Cooper. (R. 1860)
gold ornament from mound in—C. RAU. (R. 1877) 323, 440, 403
Henshall's collection of fishes from—D. S. JORDAN. (P. 1880) 425
Indian mounds in—S. T. WALKER. (R. 1879)
microscopical observations in—J. W. BAILEY 28
new fish, Caulolatilus microps, from—G. B. Goode. (P. 1878) 33_
polychrome bead from—S. S. HALDEMAN. (R. 1877) 323, 404
shell-heaps at mouth of St. John's river—S. P. MAYBERRY. (R. 1877).
shell-heaps of Tampa Bay—S. T. WALKER. (R. 1879)
specimens of flamingo from—G. WURDEMAN. (R. 1860)
winds in—J. Baltzell. (R. 1866)
Flounders, diagnoses of two undescribed genera of—G. B. Goode. (P. 1880). 425
Flourens, M. J.—
Historical sketch of the Academy of Sciences, Paris. (R. 1862)150

Flourens, M. J.—Continued.
History of the works of Cuvier. (R. 1868) 22
Memoir of Ducrotay De Blainville. (R. 1865) 20
Memoir of Leopold von Buch. (R. 1862) 15
Memoir of Pyramus de Candolle. (R. 1859)
Memoir of Cuvier. (R. 1868) 22
Memoir of Magendie. (R. 1866) 21
Memoir of Geoffroy Saint-Hilaire. (R. 1861) 149
Memoir of Louis Jacques Thénard. (R. 1862) 150
The Jussieus and the natural method. (R. 1867)21
Fluctuations of level in North American lakes—C. WHITTLESEY 119
Fluids, elastic, electricity in—A. A. DE LA RIVE. (R. 1868) 18
Fly-catcher from Sandwich Islands, description of a new—R. RIDGWAY. (P. 1881)
Fog, the wind and—J. BALFOUR. (R. 1866)
Folsom, C. Report on Jewett's general stereotype catalogue of public libraries. 4
Food, relation of, to work, and its bearings on medical practice—S. HAUGHTON. (R. 1870)
Food collections, provisional classification of—G. B. Goode455
Food-fishes, inquiry relative to—S. F. BAIRD
Food-fishes, questions relative to—S. F. BAIRD
Force, thoughts on the nature and origin of-W. B. TAYLOR. (R. 1870) 244, 375
FORCE, P. Record of auroral phenomena in high northern latitudes 84
Foreign—
authors. North American conchology-W. G. BINNEY 174
Commissioners, collections presented to United States by. (R. 1876) 298
correspondents of Smithsonian Institution, list of 154, 225, 309, 469
correspondents of Smithsonian Institution, number of 290
correspondents of Smithsonian Institution, systematic index of 257
gold and silver coins, table of. (R. 1868) 224
institutions in correspondence with Smithsonian Inst., list of 64, 225 243
institutions, list of addresses of, since 1862. (R. 1865) 209
institutions making donations to Smithsonian library, list of. (R. 1864) 188
works in library of Smithsonian—See Library.
FOREMAN, E. Meteorological system and correspondence of Smithsonian Institution. (R. 1851)
FOREMAN, E. Report on meteorological system. (R. 1852) 57
Forest trees, distribution of, in Montana, Idaho, and Washington—W. W. JOHNSON. (R. 1870)
Forests and their climatic influence—A. E. BECQUEREL. (R. 1869) 228
Forests and trees of Florida and the Mexican boundary—J. G. Cooper. (R. 1860)

of a museum—L. Agarsiz. (R. 1849)	trees of the United States—J. G. Cooper. (R. 1858) 109,
of clouds over Gulf Stream—H. M. BANNISTER. (R. 1866)	· · · · · · · · · · · · · · · · · · ·
of clouds over Gulf Stream—H. M. BANNISTER. (R. 1866)	of a museum—L. Agassiz. (R. 1849)
of ice at bottom of water—M. Engelhardt. (R. 1866)	
of precipitation—J. Hann. (R. 1877)	
Forms in which drugs and medicines appear—J. M. FLINT. (P. 1881)	• • •
Formulas, chemical, tables of—F. W. CLARKE	• •
Fort, ancient, and burial ground in Tompkins Co., N. Y.—D. Trowberder. (R. 1868)	
Fort Brown, Texas, list of birds from—J. C. Merrill. (P. 1878)	Fort, ancient, and burial ground in Tompkins Co., N. Y.—D. TROWBRIDGE.
Fort Ellis, prehistoric remains near—P. W. Norris. (R. 1879)	,
Fort Marion, Florida, list of casts of heads of Indian prisoners in—R. H. PRATT. (P. 1878)	
Fort Ripley, Minn., natural history of country about—J. E. Head. (R. 1854.) Fort Wadsworth, Dakota, Indian mounds near—A. J. Comfort. (R. 1871)	Fort Marion, Florida, list of casts of heads of Indian prisoners in-R. H.
Fort Wadsworth, Dakota, Indian mounds near—A. J. Comfort. (R. 1871)	
Forts and dwellings, Indian, in Indian Territory—W. E. Doyle. (R. 1876) Fossil— corbiculadæ—T. Prime	
corbiculadæ—T. Prime	•
gasteropod, from Mexico, description of—C. A. White. (P. 1880)	
gasteropod, from Mexico, description of—C. A. White. (P. 1880)	
mollusks from later tertiaries of California—W. H. Dall. (P. 1878) 3 ox—J. Leidy	
plants from China, description of—J. S. Newberry 2 Fossils— carboniferous invertebrate, description of new species of—C. A. White. (P. 1879) 3 catalogue of rocks, minerals, ores, and—J. Locke. (R. 1854) 4 cretaceous, from Arkansas and Colorado—C. A. White. (P. 1881) 4 cretaceous invertebrate, from Kansas and Texas, description of new— C. A. White. (P. 1879) 8 from Nebraska, report on—J. Leidy. (R. 1851) 8 from Santa Barbara, California—P. P. Carpenter 9 invertebrate, from Arkansas, Wyoming, Colorado, Utah—C. A. White.	
plants from China, description of—J. S. Newberry Fossils— carboniferous invertebrate, description of new species of—C. A. White. (P. 1879)	·
carboniferous invertebrate, description of new species of—C. A. White. (P. 1879)	
carboniferous invertebrate, description of new species of—C. A. White. (P. 1879) catalogue of rocks, minerals, ores, and—J. Locke. (R. 1854) cretaceous, from Arkansas and Colorado—C. A. White. (P. 1881) cretaceous invertebrate, from Kansas and Texas, description of new—C. A. White. (P. 1879) from Nebraska, report on—J. Leidy. (R. 1851) from Santa Barbara, California—P. P. Carpenter invertebrate, from Arkansas, Wyoming, Colorado, Utah—C. A. White.	
catalogue of rocks, minerals, ores, and—J. Locke. (R. 1854)	
catalogue of rocks, minerals, ores, and—J. Locke. (R. 1854)	•
cretaceous, from Arkansas and Colorado—C. A. White. (P. 1881)	
cretaceous invertebrate, from Kansas and Texas, description of new—C. A. White. (P. 1879)	
C. A. WHITE. (P. 1879) Strom Nebraska, report on—J. LEIDY. (R. 1851) Strom Santa Barbara, California—P. P. CARPENTER Strom Strom Arkansas, Wyoming, Colorado, Utah—C. A. WHITE.	· · · · · · · · · · · · · · · · · · ·
from Nebraska, report on—J. LEIDY. (R. 1851) from Santa Barbara, California—P. P. CARPENTER ? invertebrate, from Arkansas, Wyoming, Colorado, Utah—C. A. WHITE.	
from Santa Barbara, California—P. P. CARPENTER ? invertebrate, from Arkansas, Wyoming, Colorado, Utah—C. A. WHITE.	
invertebrate, from Arkansas, Wyoming, Colorado, Utah-C. A. WHITE.	
(invertebrate, from Arkansas, Wyoming, Colorado, Utah-C. A. WHITE.
invertebrate, of North America, check-list of-T. A. Conrad 20	•
invertebrate, of North America, check-list of—F. B. MERK 177, 18	
invertebrate, of North America, cretaceous and jurassic—F. B. MEEK 17	
invertebrate, of North America, eocene and oligocene—T. A. Conrad 20	
invertebrate, of North America, miocene—F. B. MEEK18	
postpliocene, in coast range of California—W. H. Dall. (P. 1878) 33	111/01/00/14/01/01/01/01/04/04/04/01/04/04/04/04/04/04/04/04/04/04/04/04/04/
tertiary, distribution of Californian—W. H. DALL. (P. 1878) 33	
tertiary, presented by Imperial Geological Institute, Vienna. (R. 1863)	· · · · · · · · · · · · · · · · · · ·

OF SMITHSONIAN PUBLICATIONS.
FOSTER, J. W. Ancient relics in Missouri. (R. 1868)
FOSTER, J. W. Catalogue of rocks, minerals, etc. (R. 1854)
Fourier, J., eulogy on, by F. Arago. (R. 1871)
FOURIER, J. Memoir of Delambre. (R. 1864)
Fowler, J. Shell-heaps of New Brunswick. (R. 1870)
Fox, (yacht,) meteorological observations in Arctic Seas made on—F. L. McClintock
Fox, (yacht,) record of voyage of, in Arctic regions—F. L. McClintock
France, Emperor of, report to, by Minister of Public Instruction, on scientific expedition to Mexico. (R. 1864)
France—See Paris, Prize questions.
Franklin, Sir John, expedition in search of—See Kane.
Free freight between Germany and the United States by North German Lloyd— R. SCHLEIDEN. (R. 1858)
Free freight between. United States and England—E. CUNARD. (R. 1859)
Free freight between United States and Germany—Kunhardt & Co. (R. 1861)
FRÉMONT, J. C., description of plants collected by—J. TORREY
Frémontianse, plantse—J. Torrey
French —
Exposition of 1867, ethnological department of the. (R. 1867)
half-breeds of the northwest—V. HAVARD. (R. 1879)
Institute, history of—M. FLOURENS. (R. 1862)
Society of Archeology, Archeological Congress organized by. (R. 1866)
vocabulary with comparative words in English, Spanish, and Latin
weights and measures—See Tables.
Fresh fish and other animals, circular relative to shipping
Fresh-water—
alge of North America—H. C. Wood
glacial drift of northwestern States—C. WHITTLESEY
in the ocean—W. C. DENNIS. (R. 1866)
shells—See Shells.
FRIEDLÄNDER, J. Plan of a bibliography. (R. 1858)
FRIEL, J., Antiquities of Hancock county, Kentucky. (R. 1877)
Friends of the Museum, circular addressed to—S. F. BAIRD. (P. 1881) 467,
Frigate mackerel (Auxis Rochei) on New England coast—G. B. Goode. (P. 1880)
FROEBEL, J. Physical geography of North American Continent. (R. 1854.)
Frost, disintegrating effects of, on building stones—C. G. PAGE
Fruits, dates of ripening of—F. B. Hough
Fuca, Straits of, Indians of—See J. G. Swan.
Fund, Smithson, memorial of Regents to Congress relative to. (R. 1850)
Fund. Smithson, statement of, 1846-1877

Fundy, Bay of, synopsis of marine invertebrata of—W. STIMPSON	50
Funeral of J. Henry	856
Future of geology—J. Prestwich. (R. 1875)	298
Future of the human race—A. DE CANDOLLE. (R. 1875)	298

G.

Gadus cimbrius, identity of, with Rhinonemus caudacuta—G. B. Goode; T. H. Bean. (P. 1878)
GAINES, A. S.; CUNNINGHAM, K. M. Shell-heaps on Mobile river, Alabama. (R. 1877)
GALE, L. D., statement of, on telegraph. (R. 1857) 107
Galeorhinus galeus—D. S. JORDAN; C. H. GILBERT. (P. 1880)
Gales of wind and appearance of aurora, connection of—R. T. KNIGHT; J. HENRY. (R. 1871)
GALLATIN, A. Comparative vocabulary 160
GALLATIN, A. On publication of Squier and Davis' work. (R. 1847) H, E
Gallery, Art—See Corcoran.
GALT, F. L. Indians of Peru. (R. 1877)
Galvanism, recent progress in. J. Müller. (R. 1855)
Galvanometer, use of, as a measuring instrument—J. C. Poggendorf. (B. 1859)
Gangrene, bibliography of—W. W. KEEN
Gardening, natural history as applied to farming and—J. G. Morris. (R. 1855)
GABDINER, R. H.—
Barometer, rain, and snow gauges. (R. 1858) 109
Disappearance of ice. (R. 1860)
Opening and closing of Kennebec river, Maine. (R. 1858) 109
GARDNER, W. H. Indians of valley of Red river of the North. (R. 1870) 244
GARFIELD, J. A. Biographical notice of S. P. Chase and L. Agassiz. (R. 1878)
GARFIELD, J. A. Memorial address on J. Henry
GÄRKE, H. Birds of Heligoland. (P. 1879)
GARMAN, S. American Rhinobatida, synopsis and description of. (P. 1880) 425
Gas, hydrogen, as metal and—J. E. REYNOLDS. (B. 1870) 244
Gasteropod, large fossil from Puebla, Mexico—C. A. WHITE. (P. 1880) 425
Gasterosteus, description of new species of, from Schoodic Lakes, Maine— T. H. Bran. (P. 1879)

GATSCHET, A. S.—	
·	476
Klamath lake Indian conjurer's practice. (E. 1879-80)	
"The relapse" in Klamath lake dialect. (E. 1879-80)	476
Gauge, snow—W. E. Guest. (R. 1858)	109
Gauges, rain and snow—R. H. GARDINER. (R. 1858)	109
GAUTIER, Prof. Researches relative to nebulæ. (B. 1868)	187
General catalogue system for libraries—C. C. JEWETT. (R. 1850)	28
Generation, alternate and parthenogenesis in the animal kingdom—G. A. Korn-	040
HUBER. (R. 1871)	249
Generic names of recent and fossil animals, list of—S. H. SCUDDER	470
Geneva, Society of Physics and Natural History of, report on transactions of—	
July, 1858, to June, 1859—A. A. DE LA RIVE. (R. 1859)	110
" 1860, " 1861—P. Duby. (R. 1864)	188
" 1861, " 1862—A. DE CANDOLLE. (R. 1864)	188
" 1862, " 1863—Prof. MARCET. (R. 1868)	187
" 1868, " 1864—Dr. Chossat. (R. 1865)	209
" 1864, " 1865—E. PLANTAMOUR. (R. 1865)	209
1000,	214
1007	224 228
" 1868, " 1869—H. C. LOMBARD. (R. 1869)	249
" 1872, " 1878—A. A. DE LA RIVE. (R. 1874)	286
" 1878, " 1874—A. DE CANDOLLE. (R. 1875)	298
" 1874, " 1875—E. PLANTAMOUR. (R. 1877)	828
" 1875, " 1876—J. MÜLLER. (R. 1877)	828
" 1876, " 1877—A. FAVRE. (R. 1877)	828
Genitalia of male eels—S. T. CATTIE. (P. 1880)	425
Genth, F. A.; Gibbs, W. Ammonia-cobalt bases	88
Geographical distribution of batrachia and reptilia—E. D. Cope	292
Geographical manuscripts, catalogue of collection of—L. Berlandier. (R.	
1854)	75
Geography—See Physical Geography.	
	72
Geography of North American Continent—J. FROEBEL. (R. 1854)	75
Geological—	
researches in China, Mongolia, and Japan—R. Pumpelly	202
specimens, catalogue of—D. D. Owen. (R. 1854)	75
survey in Michigan, catalogue of rocks, minerals, and ores collected on-	
C. T. JACKSON. (R. 1854)	75
Geological Survey, directory of officers and employés of	466
Geological surveys—See annual reports; also Hayden, Powell, Wheeler.	
Geology—	
and history, boundary line between—E. Suess. (R. 1872)	271
economic, of Trinidad, report on survey of-G. P. WALL; J. G. SAW-	
KINS. (R. 1856)	91
of Kerguelen Island-F. M. ENDLICH	294

Geology—Continued.	
of lower Louisiana and salt deposit on Petite Anse Island—E. W. Hu	-
GARD	_ 248
past and future of—J. Prestwice. (R. 1875)	_ 296
progress in, in 1879, 1880—G. W. HAWES. (R. 1880) 44	2, 42 8
surface, illustrations of—E. HITCHCOCK	_ 90
Georgia—	
aboriginal structures in—C. C. Jones. (R. 1877) 82	
ancient mounds in—M. F. STEVENSON. (R. 1870)	_ 244
Bartow county, mound in—M. F. STEVENSON. (R. 1872)	_ 271
description of fishes of Alleghany region of—D. S. JORDAN; A. W BRAYTON	
habits of pouched rat, or salamander of-W. GESNER. (R. 1860)	_ 147
McIntosh and Early counties, mounds in-W. McKinley. (R. 1872).	_ 271
microscopical observations in-J. W. BAILEY	_ 21
notes on collection of fishes from-T. H. BEAN. (P. 1879)	_ 331
occurrence of Stricklandia salteri and Stricklandia Davidsoni in—C. A WHITE. (P. 1880)	
remarkable forms of hail stones in—S. ABICH. (R. 1869)	
shell-heaps in—D. Brown (R. 1871)	
Spalding county, antiquities of-W. B. F. BAILEY. (R. 1877)	
Gephyrsea of northeast coast of America—A. E. VERRILL. (P. 1879)	
Germany and United States, free freight between, by North German Lloyd-	
R. Schleiden. (R. 1858)	_ 109
Germany and United States, free freight between—Kunhardt & Co. (R. 1861)	
GESNER, W. Habits of pouched rat, or salamander of Georgia. (R. 1860)	
GESNER, W. Mica beds in Alabama. (R. 1879)	
GIBBONS, H. Climate of San Francisco. (R. 1854)	
GIBBES, R. W. Memoir on Mosasaurus and allied new genera, Holcodus, Conosaurus, Amphorosteus	•
Gibbs, G.—	
Archeology in the United States. (R. 1861)	149
Bibliography of Chinook jargon	161
Comparative vocabulary	170
Dictionary of Chinook jargon, or trade language of Oregon	161
Ethnological instructions	
Ethnological map of the United States. (R. 1862)	150
Ethnological suggestions	207
Indian languages. (R. 1865)	209
Indian vocabularies. (R. 1862)	150
Instructions for archeological investigations. (R. 1861)	149
Instructions for ethnology and philology	160

Gibbs, G.—Continued.	
Intermixture of races. (R. 1864)	
Language of aboriginal Indians of America. (R. 1870)	
Makah Indians	
Philological circular. (B. 1862)	
Physical atlas of North America. (R. 1866)	
Suggestions for scientific investigations in Russian America	
Gibbs, G., memoir of, by J. A. STEVENS. (R. 1878)	
GIBBS, G., and others. Recommendation of Shea's Indian linguistics. (1861)	R.
GIBBS, G., and others. Tinneh or Chepewyan Indians of British and Russis America. (R. 1866)	an
GIBBS, G. J. Stone celts in the West Indies and Africa. (R. 1877)	-
GIBBS, W.; GENTH, F. A. Researches in ammonia-cobalt bases	
Gideon's Farm, Hennepin county, Minnesota, mounds on—F. H. NUTTE (B. 1879)	R.
GILBERT, C. H.—See D. S. Jordan; C. H. Gilbert.	
GILBERT, Sir D. Notice of Smithson. (R. 1858)	67
GILL, T.—	
Arrangement of families of fishes	
Arrangement of families of mammals	
Arrangement of families of mollusks	
Bibliography of fishes	
Bibliography of fishes of Pacific coast of United States	
Bibliography of mammals	
Bibliography of mollusks	
Catalogue of fishes of east coast of North America	
Fishes of Kerguelen Island	
Fishes of New York. (R. 1856)	
Identity of the genus Leurynnis Lockington, with Lycodopsis Collet	tt.
Note on Antennariidæ. (P. 1878)	
Note on Ceratiidæ. (P. 1878)	
Note on the Latiloid genera. (P. 1881)	
Note on Maltheidæ. (P. 1878)	
Recent progress in zoölogy. (R. 1880) 44	
Status of ichthyology	-
Synopsis of pediculate fishes of the eastern coast of extratropical North	th
Synoptical tables of mammals	
GILLISS, J. M. Solar eclipse, Peru, September 7, 1858	
GILLMAN, H. Characteristics of ancient man in Michigan. (R. 1875) 29	
GILLMAN. H. Mound builders and platvenemism in Michigan. (B. 1878.) 27	•

GILMAN, W. S. Lightning discharges. (R. 1867) 21
GIRARD, C.—
Bibliography of American natural history for 1851 4
Monograph of Cottoids 3
GIRARD, C.; BAIRD, S. F. Catalogue of North American reptiles 49
Girard College observations—See A. D. Bache.
GIVEN, W. F. Remarkable electric phenomenon. (R. 1865) 20
Glacial drift (fresh water) of the northwestern States—C. WHITTLESEY 197
Glaciers, traces of, in Massachusetts and Vermont—E. HITCHCOCK 9
GLAISHER, J. Account of balloon ascensions. (R. 1863) 18
GLAZIER, W. C. W. On destruction of fish in Gulf of Mexico. (P. 1881) 46
Gliddon mummy case—C. Pickering
Globe, winds of the—J. H. Coffix 26
Glyptocephalus cynoglossus on coast of North America—G. B. Goode; T. H. BEAN. (P. 1878)
Goat, Rocky Mountain, habits of-J. C. MERRILL. (P. 1879) 33
Gobiesox rhessodon, from San Diego, California, description of—R. SMITE. (P. 1881)
Gobioid fish, description of a new, from San Diego, California—R. SMITH. (P. 1881)
Gold and silver coins, assay of, at United States Mint—J. Pollock. (R. 1868.) 22
Gold and silver coins, foreign, table of. (R. 1868) 22
Gold ornament from a mound in Florida, observations on—C. RAU. (R. 1877.) 823, 440, 400
Goode, G. B.—
Catalogue of collection to illustrate animal resources and fisheries of the United States, exhibited at Philadelphia, 1876
Catalogue of fishes of the Bermudas 290
Catalogue of United States Fish Commission's exhibit at Berlin, 1880 413
Classification of collection of animal resources 297
Clupea tyrannus of Latrobe. (P. 1878)
Description of new species of amber fish, (Seriola Stearnsii,) obtained from Florida by S. Stearns. (P. 1879)
Descriptions of new species of fishes from deep soundings on southern New England coast, with diagnoses of two undescribed genera of flounders and genus related to Merlucius. (P. 1880)
First decade of the United States Fish Commission. (R. 1880) 442
Fishes from Bermuda mistakenly described as new by Günther. (P. 1878) 332
Fishes from deep waters on south coast of New England obtained by Fish Commission in 1880. (P. 1880)
Frigate mackerel, (Auxis Rochei,) on New England coast. (P. 1880) 425
List of substances derived from animal kingdom
Methods of capture and utilization of animals

Goode, G. B.—Continued.
Notacanthus phasganorus, new species of Notacanthidæ from Grand Banks of Newfoundland. (P. 1880) 428
Occurrence of Belone latimanus in Buzzard's Bay, Mass. (P. 1878) 88
Occurrence of Canada porcupine in West Virginia. (P. 1878) 889
Occurrence of Hippocampus antiquorum on St. George's Banks. (P. 1878) 383
Organization and objects of the U.S. National Museum. (P. 1881.) 467, 459
Plan of organization and regulations of the U.S. National Museum. (P. 1881) 467, 445
Plans for installation of collections in the U.S. National Museum 472
Preliminary catalogue of fishes of St. John's river and east coast of Florida, with descriptions of new genus and three new species. (P. 1879)
Provisional classification of the food collection. (P. 1881) 467, 455
Revision of American species of genus Brevoortia, with description of new species from Gulf of Mexico. (P. 1878) 382
Scheme of classification of collections in the U.S. National Museum. (P. 1881) 467, 457
Study of trunk fishes, (Ostraciontidæ,) with notes on American species. (P. 1879) 383
Synopsis of useful and injurpus animals 29'
Taxonomic relations and geographical distribution of swordfish family, (Xiphiidæ.) (P. 1881) 467
Voices of crustaceans. (P. 1878) 882
GOODE, G. B.; BEAN, T. H
Catalogue of Stearns's fishes of Florida and descriptions of new species. (P. 1879) 888
Catalogue of Velie's fishes in Gulf of Mexico and description of new species. (P. 1879) 338
Craig flounder of Europe (Glyptocephalus cynoglossus) on coast of North America. (P. 1878)
Description of Alepocephalus Bairdii, new deep-sca fish, western Atlantic, (P. 1879) 388
Description of Argentina syrtensium, new deep-sea fish, Sable Island bank. (P. 1878) 382
Description of Caulolatilus microps from Gulf coast of Florida. (P. 1878)832
Description of gadoid fishes, Phycis Chesteri and Haloporphyrus viola, from northwestern Atlantic. (P. 1878) 832
Description of Lycodes paxillus. (P. 1879) 388
Description of new genus and species of fish, Lopholatilus chamæleonticeps, from New England. (P. 1879)
Description of new species of fish, Apogon pandionis, from mouth of Chesapeake Bay. (P. 1881)

Goode, G. B.; Bean, T. H.—Continued.	
Description of new species of fish (Lutjanus Blackfordii and Lutjanus Stearnsii) from coast of Florida. (P. 1878)	332
Identity of Brosmius brosme Americanus, Gill, with Brosmius brosme (Müller) White. (P. 1878)	832
Identity of Rhinonemus caudacuta with Gadus cimbrius. (P. 1878)	332
New genus of fishes, Benthodesmus. (P. 1881)	467
New serranoid fish, Epinephelus Drummond-Hayi, from Bermudas and Florida. (P. 1878)	332
New species of Liparus (L. ranula) from Halifax. (P. 1879)	333
Note on Platessa ferruginea and Platessa rostrata. (P. 1878)	3:32
Note upon black grouper (Epinephelus nigritus) of the Southern cost. (P. 1878)	332
Occurrence of Lycodes Vahlii on La Have and Grand Banks. (P. 1879)	8 33
Oceanic bonito on coast of United States. (P. 1878)	332
Gopher of Illinois, habits of the-J. B. PARVIN. (R. 1854)	75
Gosse, Dr. Report on transactions of the Society of Physics and Natural	
History of Geneva, 1865–1866. (R. 1866)	214
GOULD, B. A.—	
Account of astronomical observatory at Cordoba, Argentine Republic. (R. 1878)	275
Discussion of Piazzi's astronomical observations. (R. 1863)	187
List of scientific papers of A. D. Bache. (R. 1870) 244, 329,	37 9
Report on history of discovery of Neptune	18
Transatlantic longitude	223
Gould, J. Acknowledgment of birds. (R. 1867)	21.5
Government Museum—See Museum.	
Government of Bremen, exchange system. (R. 1865)	204
Graham, T., scientific work of-W. ODLING. (R. 1871)	
Grammar and dictionary of the Carib or Karif language—C. H. BERENDT.	
(R. 1878)	275
Grammar and dictionary of the Dakota language—S. R. Riggs	40
Grammar and dictionary of the Yoruba language—T. J. Bown.	98
Grammar of Yoruba language-W. W. TURNER	98
Grand Bank, new species of Notacanthidæ from-G. B. Goode. (P. 1880)	425
Grand Bank, occurrence of Lycodes Vahlii on—G. B., GOODE; T. H. BEAN. (P. 1879)	333
Grand Manan, marine invertebrata of—W. STIMPSON	50
GRANT, E. M. Account of discovery of stone image in Tennessee. (R. 1870.)	
Grasshopper, means of destroying—V. Motschulsky. (R. 1858)	
Grasshopper, wingless, of California—E. P. Vollum. (R. 1860)	147
Grasshoppers and locusts of America—A. S. TAYLOR. (R. 1858)	109
Grasshoppers, North American, circular in reference to history of-P. R.	160

Graves, ancient, and shell-heaps of California—P. Schumacher. (B. 1874) _ 2	!8 6
Gravitation, figures of equilibrium of liquid mass withdrawn from action of— See Plateau.	
Gravitation, kinetic theories of-W. B. TAYLOB. (R. 1876) 299, 39	95
GRAY, A.—	
Address at memorial of J. Henry. (R. 1878) 841, 8	56
Biographical memoir of J. Henry. (R. 1878) 841, 40)7
Biographical notice of W. H. Harvey. (R. 1867) 2	15
Memoir of John Torrey. (R. 1878) 2	
Plantse Wrightianse Texano-Neo-Mexicanse. Part 1	
Plantæ Wrightianæ Texano-Neo-Mexicanæ. Part II	
Plants of Arctic America 34	
Plants of Kerguelen Island29	
Report on Lindheimer's, Fendler's, and Wright's botanical explorations	_
	21
GRAY, A., and others. Report on organization of Smithsonian Institution. (R. 1858)	67
GRAY, A.; COPPÉE, H. Report of Committee of Regents on Smithsonian Museum. (B. 1874)	86
GRAY, A.; SARGENT, A. A.; CLYMER, H. Report of Committee on Museum.	
	99 86
Gray substance of medulla oblongata, and trapezium—J. DEAN 17	
	15 15
	15
_	07
GREEN, J. Account of a new barometer. (R. 1855) 77, 19	
Green, J.; Würdemann, W. On filling barometer tubes. (R. 1859) 1	
Green's standard barometer1	48
Greenland	
account of cryolite of—Lewis; Quale. (R. 1866) 21	14
observations in—See Kane.	
report of explorations in—L. Kumlien. (R. 1878) 84	11
Green river country, meteorology of—Col. Collins. (R. 1871) 24	49
Green river valley, rock engravings on—J. G. BRUFF. (R. 1872) 27	71
Grenada, catalogue of Ober's collection of birds of—G. N. LAWRENCE. (P. 1878) 38	B2
Grinnell expedition to Arctic seas—See Kane.	
GROSSMANN, F. E. Pima Indians of Arizona. (R. 1871) 24	19
Guadeloupe, catalogue of Ober's collection of birds of-G. N. LAWRENCE. (P.	32
	38
Quatemala—	
antiquities in—G. WILLIAMSON. (B. 1876) 29	99

Guatemala—Continued.
collection of historical documents in—C. H. BERENDT. (R. 1876) 299
earthquakes in—A. Canudas. (R. 1858)
new humming bird (Atthis Ellioti) from—B. BIDGWAY. (P. 1878) 332
sculptures of Santa Lucia Cosumalwhuapa in—S. Habel 269
Guatemalan minister, A. T. De Irisarri, letter from, recommending Dr. Berendt. (R. 1865)
Guest, W. E. Ancient Indian remains near Prescott, Canada West. (R. 1856)
Guest, W. E. Snow gauge. (R. 1858)
Guide to the flora of Washington and vicinity-L. F. WARD 444
Guild, R. A. Biographical notice of Chas. C. Jewett. (R. 1867) 215
Gulf of California, lecture on shells of-P. P. CARPENTER. (R. 1859) 110
Gulf of California, list of fishes of-D. S. JORDAN; C. H. GILBERT. (P. 1881) 467
Gulf of Mexico— analysis of water destructive to fish in—F. M. Endlich. (P. 1881) 467
destruction of fish in—W. C. W. GLAZIER. (P. 1881)
destruction of fish in—J. Y. PORTER. (P. 1881) 467.
fish mortality in—E. INGERSOLL. (P. 1881) 467
fish mortality in—M. A. MOORE. (P. 1881) 467.
notes on mortality among fishes in—S. H. Johnson. (P. 1881) 467
report on water from—W. G. FARLOW. (P. 1881)
Gulf of Mexico—See Mexico.
Gulf-stream, formation of clouds over-H. M. BANNISTER. (R. 1866) 214
Gun-cotton and gunpowder—Lieut. von Karolyi; B. F. Craig. (R. 1864.) 188
Gunn, D. Egging expedition to Shoal Lake, Lake Winnipeg. (R. 1867) 215
GUNN, D. Indian remains in Red River Settlement, Hudson's Bay Territory.
(R. 1867) 215
Guthrie, J., authority given by, to collectors of customs, etc., to receive and transmit specimens to Smithsonian
Gutta-percha and caoutchouc. (R. 1864) 186
GUYOT, A.—
Directions for meteorological observations. (R. 1855) 77, 19, 148
Meteorological tables 31
Meteorological and physical tables 153
Metric system for scientific observations. (R. 1848)I
Mountain measurements. (R. 1862)
GUYOT, A.; HENRY, J. Directions for meteorological observations. (R. 1855.) 77, 19, 148
Gyroscope, problems of rotary motion presented by—J. G. BARNARD 240

H.

Haarlem, Holland, Society of Sciences of—See Harlem.
HAAS, H. Account of lightning discharges. (R. 1867)
HABEL, S. Sculptures of Santa Lucia Cosumalwhuapa in Guatemala 269
Habits of—
beaver—F. R. Brunot. (R. 1878) 275
black bass of the Ohio—J. Eoff. (R. 1854) 75
gopher of Illinois—J. B. PARVIN. (R. 1854) 75
pouched rat or salamander of Georgia-W. GESNER. (R. 1860) 147
Rocky Mountain goat-J. C. MERRILL. (P. 1879)
species of salamander—C. MANN. (R. 1854) 75
Hæmatoma, bibliography of—W. KEEN
HAGEN, G. Memoir of Encke. (R. 1868)224
Hagen, H.—
Neuroptera. (R. 1861) 149
Pseudo-neuroptera of Kerguelen Island 294
Synopsis of North American neuroptera 134
Haidah Indians of Queen Charlotte Islands—J. G. Swan 267
HAIDINGER, W. Honorary medal to von Martius. (R. 1863)
Hail, on the electricity of induction in—F. ZANTEDESCHI. (R. 1870) 244
Hail-stones in Georgia, remarkable forms of-S. ABICH. (R. 1869) 228
Hail-storm in Texas—G. M. BACHE. (R. 1870) 244
Hail-storm on Bosphorus—Com. Porter. (R. 1870)
HAILE, J.; McHENRY, J. W. Antiquities of Jackson Co., Tenn. (R. 1874.) 286
Hainault, Belgium, Society of Science, Art, and Literature. Prize questions. (R. 1878)
Hake, description of a new-T. H. BEAN. (P. 1880) 425
HALDEMAN, S. S. Coleoptera of the United States62
HALDEMAN, S. S. On a polychrome bead from Florida. (R. 1877) 828, 404
HALE, C. Exploration of the Nile. (R. 1865) 209
HALE, E. E. Report on Jewett's general stereotype catalogue of public libraries. 47
HALE, H. Vocabulary of Chinook jargon 161
Half-breeds of the northwest—V. HARVARD. (R. 1879) 845
Haliclystus auricula, anatomy and physiology of—H. J. CLARK 242
Halifax, description of fish obtained at—See G. B. GOODE; T. H. BEAN.
Hall in Smithsonian building, report on use of-L. Agassız. (R. 1867) 215
Hall, C. F., scientific instructions to-J. HENRY; J. E. HILGARD; S. NEW-
COMB; S. F. BAIRD; F. B. MEEK; L. AGASSIZ. (R. 1871) 249
Hall, J, statement of, on telegraph. (R. 1857) 107, 115
Halliwell manuscripts, report on—C. C. JEWETT. (R. 1852) 57

Haloporphyrus viola from deep-sea fauna of northwestern Atlantic—G. B. Goode; T. H. Bean. (P. 1878)
Hamburg Zoölogieal Gardens, exchange of specimens. (R. 1867) 21
Hamilton College, Clinton, N. Y., exchange of specimens. (R. 1861) 14
Hamilton College, Clinton, N. Y., examination of Spencer's telescope. (R. 1855)
Hamilton, J. Bequest deposited in United States Treasury. (R. 1878) 275. 32
Hamilton, J. Bequest of \$1,000 to Smithsonian Institution. (R. 1872) 271, 32
HAMLIN, H. Address on death of Joseph Henry
Hamlin, H. Biographical notice of S. P. Chase. (R. 1878) 27
Hampton, Virginia, Normal and Agricultural Institute, casts of heads of Indian
boys and girls at—R. H. PRATT. (P. 1879) 33
Hann, J., reply to criticisms of, by W. FERREL. (R. 1877) 328, 39
Hann, J.—
Atmospheric pressure and rain-fall. (R. 1877) 323, 39
Diminution of aqueous vapor with increasing altitude. (R. 1877) 823, 39
Influence of rain upon the barometer. (R. 1877) 823, 39
Laws of variation of temperature in ascending currents of air. (R. 1877) 828, 39
Relation between pressure of air and velocity of wind. (R. 1877) 323, 39
Harbors, dates of opening and closing of—F. B. Hough185
Harbors, tides and tidal action in—J. E. HILGARD. (R. 1874) 286, 390
HARDISTY, W. L. The Loucheux Indians. (R. 1866) 214, 363
HARE, R.—
Explosiveness of nitre17
Letter relative to gift of apparatus. (R. 1848)
Method of forming small weights. (R. 1858) 109
On John Wise's observation of a thunder storm. (R. 1854) 75
HARGER, O. Notes on New England Isopoda. (P. 1879) 338
HARKNESS, W. Observations on terrestrial magnetism and deviation of com-
passes on iron-clads 239
Harlem, Holland, Society of Science. Prize questions. (R. 1861, 1864, 1868,
1878) 149, 188, 224, 275
Harmonies of the solar system—S. ALEXANDER 280
HARRIOTT, J. Account of the city of Washington, 1808 330
HARRIS, E. List of birds and mammalia of Missouri river. (R. 1850) 2
HARRISON, A. M. Colored bead from mound in Florida. (R. 1877) 323
HARRISON, W. Account of storm in Butler county, Kansas. (R. 1871) 249
HART, J. N. DE. Mounds and osteology of the mound builders of Wisconsin.
(R. 1877) 523
Harvey, W. H., biographical notice of, by A. GRAY. (R. 1867) 215
HARVEY, W. H.—
Lecture on marine algæ. (R. 1855)
List of arctic alge %

HABVEY, W. H.—Continued.
Marine alge. Part I. Melanospermees
Marine algae. Part 11. Rhodospermece
Marine alge. Part III. Chlorospermeæ
Marine algae. Parts 1, 11, 111. Complete
HARWOOD, A. A. Account of sarcophagus from Syria. (R. 1870)
Hassler expedition, narrative of the-L. Agassiz. (R. 1872)
HASTINGS, C.S.; HOLDEN, E.S. Synopsis of Herschel's writings. (R. 1880.) 442, HATCH, F. W. Meteorological observations at Sacramento, Cal. (R. 1854) HAUGHTON, S. Relation of food to work, and its bearing on medical practice.
(B. 1870)
Hauy, René Just, memoir of, by G. Cuvier. (R. 1860)
HAVARD, V. French half-breeds of the northwest. (R. 1879)
HAVEN, S. F. Archæology of the United States
HAVEN, S. F. Report on Jewett's general stereotype catalogue of public libraries
Hawaiian Islands, fishes of—T. H. STREETS
Hawaiian Islands, natural history of-T. H. STREETS
Hawks, G. W
Determination of feldspar in thin sections of rocks. (P. 1881) On mineralogical composition of normal mesozoic diabase upon Atlantic border. (P. 1881)
Progress in geology and mineralogy, 1879-80. (R. 1880) 442,
HAWKES, F. L., and others. On publication of Spanish works on New Mexico. (R. 1855)
Hawkins and Wale, blow-pipe apparatus of. (R. 1872)
HAY, O. P. On a collection of fishes from eastern Mississippi. (P. 1880)
HAYDEN, F. V. Notes on Indian history. (R. 1867)
HAYDEN, F. V.; MEEK, F. B. Palsontology of the upper Missouri
HAYES, I. I.—
Astronomical observations in Arctic Seas.
Atmospheric pressure, wind, temperature, tides, in Arctic Seas
Lecture on Arctic explorations. (R. 1861)
Magnetic observations in Arctic Seas
Physical observations in the Arctic Seas
Haystack mound, Lincoln county, Dakota—A. BARRANDT. (R. 1872)
HEAD, J. E. Natural history of country about Fort Ripley, Minn. (R. 1854)
Health, architecture in relation to—D. B. REID. (R. 1856)
Hearing, sense of. (R. 1866)
Heart, strain and over-action of the. Toner lecture No. 111—J. M. DA COSTA. Heat—
and light of the sun, relative intensity of-L. W. MEECH. (R. 1856.) 91
animal, of fishes, experiments upon—J. H. KIDDER. (P. 1879)

Heat—Continued.
dry, comparative action of sulphurous acid and, upon putrefactive bacteria. (P. 1881)
principles of the mechanical theory of-J. MÜLLER. (R. 1868) 22
radiant, report on state of knowledge of-B. Powell. (R. 1859) 11
recent progress in relation to the theory of—A. CAZIN. (R. 1868) 22
specific, table of—F. W. CLARKE
tables of expansion by-F. W. CLARKE 28
Heavenly bodies, results of spectrum analysis applied to—W. Huggins. (R. 1866)
Heights, circular relative to—J. HENRY
Heights of mountains in Colorado—G. Engelmann. (R. 1862) 15
Heights, quantity of rain at different—O. W. Morris; J. Henry. (R. 1855.)
HEILPRIN, A. New species of eocene mollusca from southern United States. (P. 1880)
HEINEMANN, V. Lepidoptera
Helicinidæ—W. G. BINNEY
Heligoland, birds of—H. GÄRKE. (P. 1879)
Hellwald, F. von. American migration. (R. 1866) 214
HELMHOLTZ, H. Relation of physical sciences to science in general. (B. 1871.) 249
HELMHOLTZ, H.; MAXWELL, J. C. Later views of connection of electricity
and magnetism. (R. 1878) 273
Hemipters, instructions for collecting-P. R. UHLER. (R. 1858)
Hemirhamphus rosæ from coast of California, description of—D. S. JORDAN; C. H. GILBERT, (P. 1880)
Henry and the telegraph—W. B. TAYLOB. (R. 1878) 341, 405
Henry as a discoverer—A. M. MAYER
Henry, J.—
biographical memoir of, by A. GRAY. (R. 1878) 341, 407
deposition of, in the case of Morse vs. O'Rielly, taken September, 1879 115
examination of, by English Scientific Commission 329
life and character of, by J. C. Welling 356, 338
memoir of, and sketch of his scientific work, by W. B. TAYLOR 856, 339
memorial of, by United States Congress 356
monument to, act of Congress for 356
report of Special Committee of Board of Regents on communication of— C. C. Felton. (R. 1857) 107, 115
statue of, proceedings of Congress relative to 356
statue of, report of Executive Committee on. (R. 1880) 442
HENRY, J., papers by:
Account of Priestley's lens. (R. 1859) 110
Acoustics applied to public buildings. (R. 1856) 91
Address on the Smithsonian Institution. (R. 1858) E, 67

HENRY, J., papers by:—Continued. Annual reports, 1865–1877 (reprints)	24:
Annual reports, 1847-1877—See each annual report.	
Appendix to notice of Schoenbein, the discoverer of ozone. (R. 1868)	22
Aurora directions. (R. 1855)	
Circular of instructions to observatories relative to telegraphic announce-	
ment of astronomical discoveries	
Circular relating to collections in archeology and ethnology	
Circular relative to ancient mining in Lake Superior copper region. (R. 1861).	
Circular relative to collections of birds from Middle and South America.	
Circular relative to earthquakes	
Circular relative to heights	23(
Circular respecting copyrights. (R. 1854)	
Circular respecting new report on libraries. (R. 1854)	
Circular sent with specimens presented. (R. 1872)	27
Circular to entomologists	178
Circular to officers of Hudson's Bay Company	137
Color-blindness. (R. 1877)	829
Communication relative to a publication by Prof. Morse. (R. 1857.) 107, 829,	115
Currents of atmosphere and aerial navigation. (R. 1860)	
Description of Smithsonian anemometer. (R. 1860)	
Digest of act of organization of Smithsonian Institution C	
Directions for constructing lightning-rods	
Earthquake directions. (R. 1855)	
Effect of the moon on the weather. (R. 1871)	
Eulogy on A. D. Bache. (R. 1870) 244,	
Exposition of bequest of Smithson	
History of electro-magnetic telegraph. (R. 1857) 107,	
Instructions for observations of thunder storms	
Investigation of illuminating materials. (R. 1880) 442,	
Letter to Secretary of Treasury on payment of interest in coin. (R. 1865)	
Memoir of W. W. Seaton. (R. 1866)	
Meteorological stations, cost of establishing. (R. 1858)	
Mode of testing building materials and account of the marble used in the extension of the United States Capitol. (R. 1856)	91
Notes relative to George Catlin. (R. 1872)	271
Notes to accounts of lightning discharges by G. W. Dodge and others. (R. 1867)	215
Notes to article on American migration by F. von Hellwald. (R. 1866.)	214
Notes to article on horary variations of barometer by M. Vaillant. (R.	914

HENRY, J., papers by:—Continued.
Notes to article on vitality by H. H. Higgins. (R. 1866) 21
Notes to articles on meteorology by G. Latimer and others. (R. 1871.) 24
Notice of Parker Cleaveland. (R. 1859)
On a physical observatory. (R. 1870) 24
On the "Moon hoax." (R. 1878)
Organization of local scientific societies. (R. 1875) 29
Programme of organization of Smithsonian F, 32
Queries relative to tornadoes19
Registration of periodical phenomena. (R. 1855)
Remarks appended to meteorological articles by W. C. Dennis and others. (R. 1866)
Reply to memorial on Lowe's aeronautic voyage. (R. 1860) 14
Report of Secretary, 1847-1877. (R. 1847-1877.) H, I, 21, 28, 51, 57, 67, 75, 77, 91, 107, 109, 110, 147, 149, 150, 187, 188, 209, 214, 215, 224, 228, 244, 249, 271, 275, 286, 298, 299, 328
Researches in sound. (R. 1878)
Statement in relation to the history of the electro-magnetic telegraph 11
Suggestions relative to objects of scientific investigation in Russian America
Syllabus of course of lectures on physics. (R. 1856) 9
HENRY, J., and others. Scientific instructions to Captain Hall. (R. 1871) 24
HENRY, J.; DU PRÉ, W. Earthquakes in North Carolina, 1874. (R. 1874) 28 HENRY, J.; GUYOT, A. Directions for meteorological observations. (R. 1855.)
77, 19, 146
HENRY, J.; KNIGHT, R. T. Connection of gales of wind and aurora. (R. 1871)
HENRY, J.; Morris, O. W. Quantity of rain at different heights. (R. 1855.)
HENRY, J.; RUSSELL, R. Lectures and notes on meteorology. (R. 1854) 75
HENRY, J.; WALLACH, R. Report of Committee of Regents on fire at the
Smithsonian. (R. 1864) 188, 33
Henshall, J. A., notes on collection of fishes obtained by, from east Florida— D. S. JORDAN. (P. 1880)
Herbarium captured in Tennessee—H. R. WIRTZ. (R. 1862) 150
Herpetology of Hawaiian and Fanning Islands—T. H. STREETS 303
Herschel, Sir J. F. W., memoir of, by N. S. Dodge. (R. 1871) 249
HERSCHEL, Sir J. F. W. On atoms. (R. 1862)
Herschel, Sir W., eulogy on, by F. Arago. (R. 1870) 244
Herschel, Sir W., synopsis of scientific writings of, by E. S. HOLDEN; C. S.
HASTINGS. (R. 1880) 442, 426
Hexanchus corinus, new species of notidanoid shark, description of—D. S. Jon-
DAN; C. H. GILBERT. (P. 1880)
Higgins, H. H. On vitality, with notes by J. Henry. (R. 1866)

High antiquity in Europe, study of—A. Morlor. (R. 1862; R. 1864) 156	0, 188
HILDRETH, S. P. Meteorological observations, Marietta, Ohio	120
HILGARD, E. W. Geology of lower Louisiana and salt deposit on Petite Anse Island	
HILGARD, J. E. Description of magnetic observatory at the Smithsonian Institution. (R. 1859)	
HILGARD, J. E. Tides and tidal action in harbors. (R. 1874) 286	, 39 0
HILGARD, J. E., and others. Scientific instructions to Capt. Hall. (R. 1871.)	
HILL, A. J. Map of antiquities of United States. (R. 1861)	
HILL, G. W. Ancient earthworks of Ashland county, Ohio. (R. 1877)	828
HILL, G. W. Antiquities of northern Ohio. (R. 1874)	286
HILL, T. Map of solar eclipse of March 15, 1858, and description of the occul-	
tator	101
HILLIER, S. L. Effect of lightning. (R. 1866)	214
HINCES, W. On Smithsonian exchanges. (R. 1860)	147
Hints on public architecture—R. D. Owen	P
Hippocampus antiquorum, occurrence of, on St. George's Bank—G. B. Goode. (P. 1878)	332
Hippoglossoides elassodon, new species of flounder, from Puget Sound, description of—D. S. Jordan; C. H. Gilbert. (P. 1880)	425
Hippoglossoides exilis, new flounder, from coast of California, description of—	
D. S. JORDAN; C. H. GILBERT. (P. 1880)	425
Hippoglossus vulgaris, occurrence of, at Unalaska and St. Michael's, Alaska—T. H. Bran. (P. 1879)	888
Historical documents in Guatemala—C. H. BERENDT. (R. 1876)	299
Historical manuscripts, catalogue of collection of—L. Berlandier. (R. 1854)	75
Historical sketch of Academy of Sciences of Paris—M. FLOURENS. (R. 1862.)	150
Historical sketch of Henry's contribution to electro-magnetic telegraph—W. B.	
TAYLOR. (R. 1878)	
distorical sketches of vestiges of antiquity in United States—S. F. HAVEN	71
distorical Society of Minnesota, grammar and dictionary of Dakota language.	40
distorical study of jade—S. Blondel. (R. 1876)	299
History—	040
ancient, of North America—M. Much. (R. 1871)	249
and antiquities of Santa Rosa Island, California—S. Bowers. (R. 1877.)	
and archeology, Mexican—B. MAYER	86
and climate of New Mexico—T. A. McParlin. (R. 1877) 828, and geology, boundary line between—E. Surss. (R. 1872)	
and origin of Royal Society of London—C. A. ALEXANDER. (R. 1868.)	
Indian, notes on—F. V. HAYDEN. (R. 1867)	215
of Chester county, Pennsylvania—W. Darlington. (R. 1862)	150
of discovery of planet Neptune—B. A. Gould	18
of discovery relative to magnetism. (R. 1868)	187
14	-•

History—Continued.	
of education in the United States-F. A. PACKARD. (R. 1868) 18	3
of electro-magnetic telegraph—J. HENRY. (R. 1857)10)
of fresh-water alge of North America—H. C. Wood 24	ļ
of marine alge-W. H. HARVEY 32, 48, 95, 9)
of my youth, an autobiography—F. ARAGO. (R. 1870) 24	ļ
of North American grasshoppers, circular in reference to 16	X
of petroleum or rock-oil—T. S. HUNT. (R. 1861) 14	
of Royal Institution of Great Britain—E. MAILLY. (R. 1867) 21	Į,
of Smithsonian Institution—W. J. RHEES	į
of transactions of Anthropological Society of Paris, 1865-1867-P.	
Broca. (R. 1868) 2)
of works of Cuvier-M. FLOURENS. (R. 1868)	ļ
HITCHCOCK, E.—	
Illustrations of surface geology 9	X
On erosions of earth's surface	Ì
On geology of New England	×
Traces of glaciers in Massachusetts and Vermont	
Hoax, the Moon—J. Henry. (R. 1878)	
Hodge, C. Prayer at funeral of Joseph Henry \$5	į
Hodgins, J. G. Meteorological stations of Upper Canada. (R. 1858) 10	Ķ
Hodgins, J. G. Meteorological system of Canada. (R. 1865) 20);
Hodgkinson, Eaton, memoir of, by R. Rawson. (R. 1868) 2),
Hodgson, W. B., and others. On publication of Spanish works on New Mexico. (R. 1855)	•
HOEK, M. Letter in reference to meteoric shower of November 18, 1867 21	ľ
Holcodus, memoir on—R. W. GIBBES	
Holden, E. S.—	
Index catalogue of books relating to nebulæ, clusters, etc31]
Recent progress in astronomy 1879-1880. (R. 1880) 442, 42	1
Reports of American and foreign observatories. (R. 1879) \$4	-
Studies in Central American picture-writing. (E. 1879-'80) 47	٠
Holden, E.S.; Boehmer, G.H. Report of observatories for 1880. (R. 1880) 442, 43	٤
HOLDEN, E. S.; HASTINGS, C. S. Synopsis of scientific writings of Sir Wil-	
liam Herschel. (R. 1880)	6
Holland Society of Science, Harlem. Prize questions. (R. 1861, 1864, 1868, 1878)	į
HOLMGREN, F. Color-blindness in relation to accidents by rail and sea. (R. 1877)	9
Horary variations of barometer—M. VAILLANT. (R. 1866)	1
HORNADAY, W. T. Classification of collections to illustrate taxidermy. (P. 1881)	
Hотонкіss, T. P. Indian remains in Caddo parish, Louisiana. (R. 1872) 💯	

Ноиси, Г. В.—
Dates of blossoming of plants
Dates of defoliation or fall of leaf of plants 182
Dates of first appearance of birds
Dates of first appearance of fish 182
Dates of first appearance of insects 182
Dates of first appearance of reptiles 182
Dates of foliation of plants 182
Dates of opening and closing of lakes 182
Dates of opening and closing of rivers 182
Dates of ripening of fruits 182
Periodical phenomena, 1851 to 1859185
Hough, J. Mounds in Washington county, Mississippi. (R. 1879) 846
Hours, best, to find mean temperature—C. DEWEY. (R. 1857)
Howgate Polar Expedition, natural history of Arctic America-L. Kumlien. 342
Hoy, P. R. Journal of an exploration of western Missouri. (R. 1864) 186
Hoy, P. R. On Amblystoma luridum, a salamander of Wisconsin. (R. 1854.) 76
HUBBARD, J. S. Investigations of Biela's comet. (R. 1862) 156
Hudson's Bay, notes on fishes from—T. H. BEAN. (P. 1881)
Hudson's Bay Company. Kennicott's explorations. (R. 1868) 18'
Hudson's Bay Company, circular to officers of-J. HENRY
Hudson's Bay Company, letter to officers of—G. SIMPSON 18'
Hudson's Bay Territory, observations in—B. R. Ross. (R. 1859) 110
Hudson's Bay Territory, Red river settlements, Indian remains in—D. GUNN. (R. 1867)
Huggins, W. Results of spectrum analysis applied to the heavenly bodies. (R. 1866)
Human race, probable future of the—A. DE CANDOLLE. (B. 1875) 298
Human remains from Patagonia—A. RIED. (R. 1862)
Humidity, relative, tables of-J. H. Coffin
Humming-bird, a new, (Atthis Ellioti), from Guatemala—R. RIDGWAY. (P. 1878)882
Humming-birds, described, list of-D. G. Elliot
Humming-birds, monograph of—D. G. ELLIOT
Humphreys, A. A. Method of ascertaining amount of water in rivers. (R. 1858)100
Hungarian National Museum, acknowledgment for birds. (R. 1868) 187
Hungary, prehistoric antiquities of—F. F. ROMER. (R. 1876) 299, 440, 392
HUNT, T. S. Chemistry of the earth. (R. 1869) 228, 376
Hunt, T. S. History of petroleum or rock oil. (R. 1861)
Huntington, G. C. Climate of Kelley's Island, Ohio. (R. 1866) 214
Hurricane at Tortola, St. Thomas, and Porto Rico-G. A. LATIMER. (R. 1867.) 216
Hurricane in Island of St. Thomas. (R. 1867)

HUXLEY, T. H. Principles and methods of palseontology. (R. 1869)	228
Hydrobiinæ, researches upon the—W. STIMPSON	201
Hydrogen as a gas and as a metal—J. E. REYNOLDS. (B. 1870)	244
Hygrometrical tables—A. Guyor	154
Hymenoptera— instructions for collecting—B. Clemens. (R. 1858) letter relative to catalogue of—H. DE SAUSSURE. (R. 1862) monograph of—H. DE SAUSSURE	150
of Arctic regions—S. H. Scudder and others	
Hypsometrical tables—A. GUYOT	158

I.

Ice	147
disappearance of—R. H. GARDINER. (R. 1860)	
formation of, at the bottom of water—M. Engelhardt. (R. 1866)	
observations on—D. WALKER	146
Iceland, vegetable colonization of—C. MARTINS. (R. 1858)	100
Ichthyology-	
Contributions to North American—	
Part 1. Review Rafinesque's memoirs—D. S. JORDAN	305
Part II. A. Notes on Cottidæ, (etc.) B. Synopsis Siluridæ—D. S. JORDAN	306
Part III. A. Distribution of fishes of Alleghany region, (etc.)—D. 8. JORDAN; A. W. BRAYTON. B. Synopsis of family Catostomidæ—D. S. JORDAN	
of Hawaiian and Fanning Islands and California-T. H. STREETS	30 3
status of-T. Gill.	247
Ichthyology—See Fishes.	
Icichthys Lockingtoni, new species deep-water fish from coast of California- D. S. JORDAN; C. H. GILBERT. (P. 1880)	
Icosteus senigmaticus of California, description of—W. N. Lockington. (P. 1880)	
Icterus, description of new species of, from West Indies—G. N. LAWRENCE. (P. 1880)	
Idaho, distribution of forest trees in—W. W. Johnson. (B. 1870)	
Identification of artisan and artist—N. WISEMAN. (R. 1870)	
Illinois—	
Adams county, mounds in, near Quincy—W. G. ANDERSON. (B. 1879.)	145
Albany, mounds near—R. J. FARQUHARSON. (R. 1874)	

Illinois—Continued.	
deposits of flint implements in—J. F. SNYDER. (R. 1876)	299
habits of the gopher of-J. B. PARVIN. (R. 1854)	· 7 5
Lawrence county, antiquities of-A. PATTON. (B. 1878)	275
Madison county, stone cists in, near Highland—A. OEHLER. (R. 1879.)	34 5
Mason county, antiquities of—J. Cochrane. (R. 1877)	32 3
Mercer county, ancient mounds of—T. McWhorrer. (R. 1874)	286
Pike county, mounds in—B. MITCHELL. (R. 1879)	84 5
Rock Bluff, description of human skull from—J. A. MEIGS. (R. 1867.)	215
Rock Island, shell-bed skull from—A. S. TIFFANY. (R. 1874)	286
Rock Island county, mounds in—T. THOMPSON. (R. 1879)	345
Rock Island county, antiquities of—A. Toellner. (R. 1879)	845
Rock river valley, mound builders in—J. SHAW. (R. 1877)	823
Spoon river valley, mounds in-W. H. ADAMS. (R. 1879)	845
southern, agricultural flint implements in—C. RAU. (R. 1868.) 224, 440,	370
Union county, antiquities of—T. M. PERRINE. (R. 1878)	275
Union county, mounds in, near Anna—T. M. PERRINE. (R. 1872)	271
Whitesides county, antiquities of—W. H. PRATT. (R. 1874)	286
Illinois Board of Education. Meteorological system for every State. (R. 1855.)	77
Illuminating materials, investigation of—J. HENRY. (R. 1880) 442,	389
Illustrations of method of recording Indian languages—J. O. Dorsey; A. S.	
GATSCHET; S. R. RIGGS. (E. 1879-80)	476
Illustrations of surface geology—E. HITCHCOCK	90
Image, stone, in Tennessee, account of discovery of—E. M. GRANT. (R. 1870.)	244
Imperial Academy of Sciences, Belles-lettres, and Arts, of Bordeaux. Prize	004
questions. (R. 1868)	224
Imperial Academy of Sciences, Vienna. Prize questions. (R. 1864; R. 1865.) 188,	
Imperial Geological Institute, Vienna, tertiary fossils presented by. (R. 1868.)	187
Imperial Library of Vienna, books presented by. (R. 1865)	209
Imperial Society of Natural Sciences of Cherbourg. Prize questions. (R. 1864.)	188
Imperial Society of Science, Agriculture, and Arts of Lille. Prize questions. (R. 1865)	209
Implement, wood, ancient, found in Connecticut—E. W. Ellsworth. (R. 1876)	299
Implements—	
agricultural, North American stone period—U. RAU. (R. 1868) 187,	440
flint, agricultural, in southern Illinois-C. RAU. (R. 1868) 224, 440,	37 0
flint, deposits of, in Illinois—J. F. SNYDER. (R. 1876)	299
flint, in Holmes county, Ohio-H. B. CASE. (R. 1877)	828
stone, North American—C. RAU. (R. 1872) 271, 440,	382
Improvements, recent, in chemical arts—J. C. BOOTH; C. MORFIT	27
Income, majority and minority reports on distribution of—J. A. PEARCE; J. MEACHAM. (R. 1858)	829

Incorporation, act of, of Smithsonian Institution—See Congress.
Index-catalogue of books and memoirs on nebulæ and clusters—E. S. Holden. 31
Index—
of genera of birds—S. F. BAIRD
of Smithsonian publications47
systematic and alphabetical, of publications of the Smithsonian Institution to July, 1869. (R. 1868)
systematic, to list of foreign correspondents25
to genera of recent and fossil animals—S. H. Scudder 47
to names applied to subdivisions of class Brachiopoda-W. H. DALL 30
to North American botany—S. Watson 25
to papers on anthropology published by Smithsonian Institution, 1847- 1878—G. H. Borhmer. (R. 1879) 845, 42
Indian—
boys and girls at Hampton Normal Institute, Va., catalogue of casts of
heads of—R. H. PRATT. (P. 1879)
burial in California-W. M. King. (R. 1874)
conjurer's practice, details of—A. S. GATSCHET. (E. 1879-80)47
engravings on rocks along Green river valley—J. G. Bruyy. (R. 1872.)
forts and dwellings, Indian Territory—W. E. DOYLE. (R. 1876) 29
history, notes on—F. V. HAYDEN. (R. 1867) 21
languages—G. Flachenecker. (R. 1862)
languages, method of recording—J. O. Dorsey; A. S. GATSCHET; S. R. RIGGS. (E. 1879–80)
languages—See G. Gibbs, F. L. O. Roehrig, S. R. Riggs, J. G. Shea, W. L. Hardisty.
linguistics, account of library of-J. G. SHEA. (R. 1861)
linguistics, recommendation of Shea's—G. GIBBS and others. (R. 1860.)
mode of making arrow-heads and obtaining fire—G. CROOK. (R. 1871. 24
mounds in southern Florida, explorations of—S. T. WALKER. (R. 1879) 34
mounds near Fort Wadsworth, Dakota—A. J. Comfort. (R. 1871) 24
philology-W. W. Turner. (R. 1851)
photographs, catalogue of21
portraits and sketches of scenery, catalogue of—J. M. STANLEY 55
portraits, report of Committee of Regents on Stanley's gallery of. (R. 1857)
pottery—C. RAU. (R. 1866)214, 440. 36
prisoners at St. Augustine, Florida, catalogue of casts of heads of—R. H. Pratt. (P. 1878)
relics, catalogue of cabinet of, presented by J. H. Devereux. (R. 1872.)
relics from Schoharie, New York—F. D. Andrews. (R. 1879) 34
remains in Caddo parish, Louisiana—T. P. Hotchkiss. (B. 1872) 271
remains in Red River Settlement, Hudson's Bay Territory—D. Guss.

Indian—Continued.	
remains near Prescott, Canada West-W. E. Guest. (R. 1856)	91
tribes, cessions of land by, to United States-C. C. ROYCE. (E. 1879-80)	176
tribes of upper Missouri-T. A. Culbertson. (R. 1850)	28
village (Kushkushkee) near Newcastle, Pennsylvania.—E. M. McCon- NELL. (R. 1871)	249
vocabularies—G. GIBBS. (R. 1862)	150
vocabularies received from the Wheeler survey, list of—G. Gibbs. (R. 1874)	286
Indian Ocean, cyclone in the-N. PIKE. (R. 1867)	215
Indian Territory, Indian forts and dwellings in-W. E. DOYLE. (R. 1876) 2	99
Indiana ,	
Allen county, antiquities of—R. S. Robertson. (R. 1874) 2	286
ancient burial mound in—W. PIDGEON. (R. 1867)	215
De Kalb county, antiquities of—R. S. ROBERTSON. (R. 1874) 2	286
Franklin county, mounds in—E. R. Quick. (R. 1879)	45
Knox county, antiquities of-A. PATTON. (R. 1873) 2	75
La Porte county, antiquities of—R. S. Robertson. (R. 1874) 2	86
Rush county, mounds in—F. JACKMAN. (R. 1879)	45
tornado in-J. Chappelsmith	59
Indians—	
aboriginal, of America, language of—G. Gibbs. (R. 1870) 2	44
Cree, system of relationship of—E. A. WATKINS. (R. 1862) 1	50
Dakota or Sioux, language of—F. L. O. ROEHRIG. (R. 1871)	49
Haidah, of Queen Charlotte's Islands, British Columbia—J. G. SWAN 20	57
Kutchin tribes—S. Jones. (R. 1866)	65
Loucheux—W. L. HARDISTY. (R. 1866)	65
Makah—J. G. Swan	30
mortuary customs of—H. C. YARROW. (E. 1879-80) 4	76
mythology of-J. W. Powkll. (E. 1879-80) 4	76
Navajo, sketch of the-J. Letterman. (R. 1855)	77
of British America, account of—E. Petitot. (R. 1865) 2	09
of Cape Flattery, Washington Territory-J. G. SWAN 23	20
of Peru—F. L. GALT. (R. 1877) 8	28
of valley of Red River of the North, ethnology of—W. H. GARDNER. (R. 1870)	44
of western Nevada and California, Centennial mission to—S. Powers.	99
	49
	25
	76
	99
Sioux or Dakota, language of-F. L. O. Roehrig. (R. 1871) 249, 3	78

Indians—Continued.
Tinneh or Chepewyan, of British and Russian America—G. GIBBS and others. (R. 1866)
uses of brain and marrow of animals by-T. R. PEALE. (R. 1870) 24
Indians—See California, Cape Flattery, Carib, Dakota, Haidah, Nevada, Kutchin, Loucheux, Makah, Mandan, Maya, Navajo, Peru, Portraits, Red River of the North, Shoshone, Sioux, Tinneh, Tonto.
Induction and deduction—J. von Liebig. (R. 1870) 24
Induction, electricity of, in clouds—F. ZANTEDESCHI. (B. 1870) 24
Inflammation in arteries after ligature, acupressure, and torsion. Toner lecture No. VII—E. O. SHAKESPEARE
Influence of aurora on the telegraph—W. D. SARGENT. (R. 1870) 240
Influence of rain upon the barometer—J. HANN. (R. 1877) 323. 339
Infusoria—J. W. Bailey
Infusorial earths, examination of—A. M. EDWARDS 200
INGERSOLL, E. Fish mortality in Gulf of Mexico. (P. 1881) 45
Inquiries relative to crawfish and other crustacea
Inquiries relative to disease known as chorea—S. W. MITCHELL. (R. 1874) 286
Inquiry relative to food-fishes of United States—S. F. BAIRD 231
Insect instincts and transformations, lecture on—J. G. MORRIS. (R. 1855)
Insects—
arsenic acid for protecting anatomical preparations from—J. B. S. JACK- SON. (P. 1878)
circular concerning the department of-S. F. BAIRD 448
classification of, from embryological data—L. Agassiz 16
dates of first appearance of—F. B. Hough
directions for collecting and preserving-A. S. PACKARD 261
instructions for collecting—S. F. BAIRD. (R. 1858)
of Arctic America—L. Kumlien 342
of Kerguelen Island-C. R. OSTEN SACKEN; H. A. HAGEN 294
Installation of collections in National Museum, plans for—G. B. Goode. (P. 1881) 467, 472
Instincts and transformations of insects, lecture on—J. G. Morris. (R. 1855)
Institute of Bologna, Academy of Sciences of the. Prize questions. (R. 1862.) 150
Institute of France, historical sketch of, by M. FLOURENS. (R. 1862) 150
Institute of Rupert's Land, circular of. (R. 1861)149
Institute, Royal Scientific and Library, of Lombardy. Prize questions. (R. 1865)
Institution of Civil Engineers, London. Prize questions. (R. 1862) 150
Institutions—
domestic, in correspondence with Smithsonian Institution, list of. (R. 1858) 67, 69, 238
foreign, in correspondence with Smithsonian Institution, list of. (R. 1864 · R. 1865). 188, 209, 64, 154, 225, 243, 309, 469

Institutions—Continued.
in United States and British Provinces of North America, list of—W. J. Rhees.
in which phonography is taught. (R. 1856)
scientific and literary, in United States, list of principal 33
Instructions—
for archæological investigations-G. GIBBS. (R. 1861)1
for collecting coleopters—J. L. LE CONTE. (R. 1858) 1
for collecting diatomacea—A. M. EDWARDS 30
for collecting diptera—H. Loew; R. Osten Sacken. (R. 1858) 1
for collecting hemipters—P. R. UHLEB. (R. 1858)
for collecting hymenoptera—B. Clemens. (R. 1858) 1
for collecting insects—S. F. BAIRD. (R. 1858) 1
for collecting land and fresh-water shells—J. Lxwis 36
for collecting lepidoptera—B. CLEMENS. (R. 1858)
for collecting myriapods, phalangids, etc.—H. C. Wood. (R. 1866) 2
for collecting nests and eggs of North American birds. (R. 1858) 1
for collecting neuroptera—P. R. UHLER. (R. 1858) 1
for collecting orthopters—P. R. UHLER. (R. 1858)
for observations of thunder storms—J. HENRY
for research relative to ethnology and philology of America—G. GIBBS 16
in reference to collecting nests and eggs of North American birds—T. M. Brewer
to Captain Hall for Arctic observation—J. HENRY and others. (R. 1871) 2
to observatories relative to telegraphic announcement of astronomical discoveries—J. HENRY
Instruments, meteorological,
articles on, in Smithsonian. (R. 1874)2
description of—L. CASELLA. (R. 1859) 1
description of—A. GUYOT; J. HENRY. (R. 1855) 77, 19, 14
Integrals, general, of planetary motion—S. Newcomb
Intensity of heat and light of the sun—L. W. MEECH. (R. 1856) 91, &
Interest, letter to Secretary of Treasury on payment of, in coin—J. Henry. (R. 1865)
Intermixture of races-G. GIBBS. (R. 1864)
Internal structure of the earth—J. G. BARNARD
International Archeological Congress, Antwerp, 1866. (R. 1866) 2
International code of symbols of archeology—G. DE MORTILLET; E. CHANTRE. (R. 1875)
International exchange, report on—G. H. BOEHMER 47
International Exhibition, Philadelphia—See Centennial.
International Fisheries Exhibition, Berlin, catalogue of United States exhibit at—G. B. Goode 41

Interpolation, methods of, applicable to graduation of irregular series. Parts 1, 11.—E. L. De Forest. (R. 1871; R. 1878) 249	. 275
Introduction, letters of. (R. 1865)	
Introduction to study of Coptic language—M. KABIS. (R. 1867)	
Invertebrata, marine, from New England coast distributed by United States Fish Commission, list of—A. E. VERBILL; R. RATHBUN. (P. 1879)	
Invertebrata, marine, of Grand Manan—W. STIMPSON	
Invertebrata, marine, of northeastern coast of America, notice of recent additions to—A. E. VERRILL: Part I. Annelida, etc. (P. 1879)	
Part II. Mollusca, with notes on Annelida, etc. (P. 1880)	
Part III. Catalogue of Mollusca recently added to fauna of southern New England. (P. 1880)	
Invertebrate fossils—	
check-list of-T. A. CONBAD	200
from Arkansas, Wyoming, Colorado, and Utah, descriptions of new- C. A. WHITE. (P. 1880)	
of North America, check-list of-F. B. MEEK177	183
report on-F. B. MEEK; F. V. HAYDEN	172
Invertebrates, marine, distributed by U.S. National Museum, list of—R. BATH-BUN. (P. 1881) 465, 467,	
Invertebrates—See Marine invertebrates.	
Investigation of—	
anatomy and physiology of the rattlesnake—S. W. MITCHELL	135
disturbances of horizontal component of magnetic force—A. D. BACHE	
disturbances of vertical component of magnetic force-A. D. BACHE	175
eleven year period in amplitude of solar-diurnal variation and disturbances of magnetic declination—A. D. Bache	
illuminating materials—J. HENRY. (R. 1880) 442,	
influence of moon on magnetic declination—A. D. BACHE	
orbit of Neptune and tables of its motion—S. Newcomb	
orbit of Uranus, with tables of its motion—S. NEWCOMB	
Investigations—	
anthropological, in 1879—O. T. MASON. (R. 1879) 345,	420
archæological, instructions for-G. GIBBS. (R. 1861)	149
chemical and physiological, relative to certain American vertebrata—J. JONES	
of Biela's comet—J. S. Hubbard. (R. 1862)	
scientific, in Russian America, suggestions relative to-J. HENRY	
Iowa—	
ancient relics in—J. B. Cutts. (R. 1872)	
Delaware county, mounds in—M. W. Moulton. (R. 1877)	
Muscatine county, mounds in—T. THOMPSON. (R. 1879)	345
IRRY J. R. McD. Works and character of Smithson	227

Or BEITHOUNIAN TOBELORITORS.	
Iron and copper, preservation of, in salt water—A. E. BECQUEREL. (R. 1864.)	188
Ironclad Monadnock, deviation of compasses on-W. HARKNESS	289
Iron rust, destructive effect of. (R. 1861)	149
Irradiation—Abbé Moigno. (R. 1866)	214
Irritation of polarized nerve, effect of-B. F. LAUTENBACH. (R. 1878) 841	
Irving, Washington, notice of, by C. C. FELTON. (R. 1859)	110
IRVING, W. On publication of Spanish works on New Mexico. (R. 1855)	77
IRWIN, B. J. D. Tuoson meteorite. (R. 1863)	187
Island—	
Petite Anse, salt deposit on—E. W. HILGARD	248
St. Thomas, hurricane in-G. A. LATIMER. (R. 1867)	2 15
Santa Rosa, history and antiquities of—S. Bowers. (R. 1877)	828
Islands	
Fanning, natural history of—T. H. STREETS	
Hawaiian, natural history of—T. H. STREETS	808
Kerguelen, natural history of—J. H. KIDDER and others 298	, 294
North American, meteorological stations and observers in. (R. 1868) 224	, 878
Pacific, natural history of—W. H. Prase. (R. 1862)	150
Queen Charlotte's, Haidah Indians of-J. G. Swan	267
St. Croix and Virgin, flora of-H. F. A. EGGERS	818
Shetland, Faroe, and Iceland, vegetable colonization of—C. MARTINS. (R. 1858)	109
Isle Jesus, Canada East, description of observatory on—C. SMALLWOOD. (R. 1856)	91
Isle Royale, Michigan, antiquities of—A. C. Davis. (R. 1874)	286
Isohyetal maps—See C A. Schott.	•
Isopoda of New England, notes on-O. HARGER. (P. 1879)	883
Isothermal charts—See Temperature.	
Italy—See Prize questions.	

, **J.**

JACKSON, C. T. Catalogue of rocks, minerals, and ores collected on geological
• • • • • • • • • • • • • • • • • • •
survey in Michigan. (R. 1854)
JACKSON, J. B. S. Arsenic acid for protecting anatomical preparations from insects. (P. 1878)
Jade. Study of the mineral called yu by the Chinese—S. Blondel. (R. 1876.) 299
JAMES, T. P. Mosses of Kerguelen Island
Jamin, J. Photo-chemistry. (R. 1867)
JAMIN, J. Vegetation and the atmosphere. (R. 1864)
Japan—
analysis of coals from—J. A. MacDonald
catalogue of Japanese woods from—L. F. WARD. (P. 1881) 467
examination of infusorial earth from—A. M. Edwards 202
geological researches in—R. PUMPELLY202
Government of, collection of cotton fibre presented by. (P. 1881) 467
Jargon, Chinook—
bibliography of—G. GIBBS
dictionary of—G. GIBBS; H. HALE
vocabulary of-M. Lionner; B. R. Mitchell; W. W. Turner 68
JEFFERSON, J. P. Mortality of fishes in Gulf of Mexico. (P. 1878) 352
JEFFERSON, J. P.; PORTER, J. Y.; MOORE, T. Destruction of fish in vicinity
of the Tortugas. (P. 1878) 352
JEVONS, W. S. On a national library. (R. 1878) 273
JEWETT, C. C.—
Construction of catalogues of libraries and their publication by separate
• • • • • • • • • • • • • • • • • • • •
stereotyped titles47
Stereotyped titles
Copyright books from 1846 to 1849. (R. 1850)
Copyright books from 1846 to 1849. (R. 1850)
Stereotyped titles

JOHNSON, W. W. Distribution of forest trees in Montana, Idaho, and Wash-
ington. (R. 1870)
Joints, bibliography of diseases of the—W. W. KEEN
Jones, C. C., Jr. Aboriginal structures in Georgia. (R. 1877) 828,
Jones, C. C., Jr. Primitive manufacture of spear and arrow points on Savannah river. (R. 1879)
Jones, J.—
Analysis of blood
Chemical and physiological investigations relative to vertebrata
Explorations of aboriginal remains of Tennessee
Modes of burial
Observations on the liver, pancreas, spleen, kidney, etc
On burial caves
On earthworks
On mounds
On relics
Jones, J. M. Kjækken-mædding in Nova Scotia. (R. 1868)
Jones, S. The Kutchin Indians. (R. 1866) 214
JORDAN, D. S.—
Contributions to North American ichthyology:
Part I. Review of Rafinesque's memoirs
Part II. Notes on Cottidæ, etc.; synopsis Siluridæ
Part III. Distribution of fishes of Alleghany region; synopsis of family Catostomidæ
Description of new species of Caranx (Caranx Beani) from Beaufort, North Carolina. (P. 1880)
Description of new species of North American fishes. (P. 1879)
Forgotten paper of Dr. Ayres and its bearing on nomenclature of cyprinoid fishes of San Francisco. (P. 1880)
Notes on collection of fishes from Clackamas river, Oregon. (P. 1878.)
Notes on Cottidæ, Etheostomatidæ, Percidæ, Centrarchidæ, Aphododeridæ, Dorysomatidæ, and Cyprinidæ
Notes on Curtiss' collection of fishes from Florida. (P. 1880)
Notes on Dugès' collection of fishes from Mexico. (P. 1879)
Notes on Henshall's collection of fishes from east Florida. (P. 1880)
Notes on Sema and Dacentrus. (P. 1880)
Notes on typical specimens of American fishes in British Museum and in
Museum d'Histoire Naturelle, Paris. (P. 1879)
Review of Rafinesque's memoirs on North American fishes
Synopsis of Catostomidæ
Synopsis of Siluridæ of fresh waters of North America
JORDAN, D. S.; BRAYTON, A. W. Distribution of fishes of Alleghany region of South Carolina, Georgia, and Tennessee

AN, D. B.;	GILBERT, C. H.—	
Descriptio	n of new agonoid fish (Agonus vulsus) from California. (P. 18	180)
•	n of new agonoid fish (Brachiopsis xyosternus) from Californ	nia.
_	n of new embiotocoid fish (Abeona aurora) from Monterey, Co, with notes on a related species. (P. 1880)	
-	n of new embiotocoid fish (Cymatogaster rosaceus) from co	
-	n of new embiotocoid fish (Ditrema atripes) from coast of Co. (P. 1880)	
	n of new flounder (Hippoglossoides exilis) from coast of Co. (P. 1880)	
_	n of new flounder (<i>Platysomatichthys stomias</i>) from cosst	
Description	n of new flounder (<i>Pleuronichthys verticalis</i>) from Californ notes on other species. (P. 1880)	nis,
Description	n of new flounder (Xystreurys liolepis) from Santa Catal, California. (P. 1880)	ins
Description	n of new ray (Platyrhina triseriata) from coast of Californ	nia.
Description	n of new scorpænoid fish (Sebastichthys maliger) from coast	t of
Description	n of new scorpænoid fish (Sebastichthys proriger) from Monay, California. (P. 1880)	nte-
Description	n of new species of deep-water fish (Icichthys Lockingtoni) frof California. (P. 1880)	rom
Descriptio	n of new species of <i>Hemirhampus</i> (<i>Hemirhampus Rosa</i>) frof California. (P. 1880)	rom
Description	n of new species of Nemichthys (Nemichthys avocetta) fi Sound. (P. 1880)	rom
Descriptio	n of new species of notidanoid shark (Hexanchus corinus) for coast of United States. (P. 1880)	rom
Descriptio	n of new species of Paralepis (Paralepis coruscans) from Str an de Fuca. (P. 1880)	aits
	n of new species of <i>Pomadosys</i> from Mazatlan. (P. 1881).	
Descriptio	n of new species of Ptychochilus (Ptychochilus Harfordii) finento river. (P. 1881)	rom
Descriptio	n of new species of ray (Raia rhina) from coast of Californ	n is .
Description	n of new species of ray (Raia stellulata) from Monterey, (R. 1880)	Cal-
Description	n of new species of rock cod (Sebastichthys serriceps) from california. (P. 1880)	osst
Descriptio	n of new species of rockfish (Sebastichthys carnatus) from a	

JORDAN, D. S.; GILBERT, C. H.—Continued.
Description of new species of rockfish (Schastichthys chrysomelas) from coast of California. (P. 1880) 425
Description of new species of Sebastichthys (Sebastichthys miniatus) from Monterey bay, California. (P. 1880)
Description of new species of Xenichthys (Xenichthys ocyurus) from west coast of Central America. (P. 1881) 467
Description of new species of Xiphister and Apodichthys from Monterey, California. (P. 1880)
Description of Sebastichthys mystinus. (P. 1881) 467
Description of seven new species of sebastoid fishes from coast of California. (P. 1880) 425
Description of two new species of fishes (Ascelicthys rhodorus and Scytalina cerdale) from Neah bay, Washington Territory. (P. 1880) 425
Description of two new species of flounders (Parophrys ischyurus and Hippoglossoides elassodon) from Puget Sound. (P. 1880) 425
Description of two new species of Sebastichthys (Sebastichthys entomelas and Sebastichthys rhodochloris) from Monterey bay, Cal. (P. 1880.) 425
Description of two new species of scopeloid fishes (Sudis ringens and Myctophum crenulare) from Santa Barbara channel, Cal. (P. 1880.) 425
Descriptions of five new species of fishes from Mazatlan, Mexico. (P. 1881)
Descriptions of thirty-three new species of fish from Mazatlan, Mexico. (P. 1881)467
Generic relations of Belone exilis. (P. 1880) 425
Generic relations of Platyrhina exasperata. (P. 1880)
List of fishes collected by Lt. H. E. Nichols in Gulf of California and on west coast of Lower California, with descriptions of four new species. (P. 1881)
List of fishes of Pacific coast of United States, with table showing distribution of species. (P. 1880) 425
Note on Raia inornata. (P. 1881) 467
Notes on collection of fishes from San Diego, California. (P. 1880) 425
Notes on collection of fishes from Utah Lake. (P. 1880) 425
Notes on collection of fishes made by Lt. H. E. Nichols on west coast of Mexico, with descriptions of new species. (P. 1881)
Notes on fishes of Beaufort Harbor, North Carolina. (P. 1878)
Notes on fishes of Pacific coast of United States. (P. 1881) 467
Notes on sharks from coast of California. (P. 1880) 425
Occurrence of Cephaloscyllium laticeps (Duméril), Gill, in California. (P. 1880)
Oil shark of southern California (Galeorhinus galeus). (P. 1880) 425
JORDAN, D. S.; JOUY, P. L. Check-list of duplicate fishes of Pacific, distributed by Smithsonian Institution in 1881. (P. 1881) 467
Journal of exploration of western Missouri in 1854—P. R. Hoy. (R. 1864) 188

Journey to the Yukon, Russian America—W. W. Kirby. (R. 1864)	Journals of Board of Regents—See Regents.	
Utah Lake. (P. 1881)	Journey to the Yukon, Russian America—W. W. KIRBY. (R. 1864)	188
Judiciary Committee, Senate, report of, on management of Smithsonian Institution—A. P. Butler. (R. 1855) Jupiter, orbit and tables of—J. N. Stockwell Jupiter, small planets between Mars and—Prof. Lespiault. (R. 1861) Jurassic fossils, check-list of—F. B. Meek		46 7
Jupiter, orbit and tables of—J. N. STOCKWELL Jupiter, small planets between Mars and—Prof. LESPIAULT. (R. 1861) Jurassic fossils, check-list of—F. B. MEEK		467
Jupiter, small planets between Mars and—Prof. LESPIAULT. (R. 1861)		T
Jurassic fossils, check-list of-F. B. MEEK	Jupiter, orbit and tables of-J. N. STOCKWELL	23.
	Jupiter, small planets between Mars and—Prof. LESPIAULT. (R. 1861)	144
Jussieus, the, and the natural method—M. FLOURENS. (R. 1867) 2	Jurassic fossils, check-list of-F. B. MEEK	L77
	Jussieus, the, and the natural method—M. FLOURENS. (R. 1867)	2 15

K.

KABIS, M. Introduction to study of Coptic language. (R. 1867)
Kämtz; Dufour, C. Scintillation of the stars. (R. 1861)
KANE, E. K.—
Astronomical observations in the Arctic Seas
Magnetic observations in the Arctic Seas
Meteorological observations in the Arctic Seas1
Physical observations in the Arctic Seas
Tidal observations in the Arctic Seas
Kansas—
Butler county, account of storm in—W. HARRISON. (R. 1871) 2
climate of—R. S. Elliott. (R. 1870)
coleoptera of-J. L. Le Conte
new cretaceous invertebrate fossils from—C. A. WHITE. (P. 1879) 3
Kansas City, Missouri, antiquities of—W. H. R. LYKINS. (R. 1877) \$
Karif or Carib language and people—C. H. BERENDT. (R. 1878)
KAROLYI, Lieut. von; CRAIG, B. F. Products of combustion of gun-cotton and gunpowder. (R. 1864)
KEEN, W. W. Bibliography of works on diseases3
KEEN, W. W. Surgical complications and sequels of continued fevers. Tones lecture No. v
Keller, Dr., abstract of report of, on settlements in Switzerland—A. Morlot. (R. 1868)
Kelley, O. H. Ancient town in Minnesota. (R. 1868)
Kelley's Island, Ohio, climate of—G. C. HUNTINGTON. (R. 1866)

Kennebec river, Maine, opening and closing of-R. H. GARDINER. (R. 1858.)	109
KENNEDY, J. P. Authority to naval officers to receive and transmit specimens	
to the Smithsonian Institution	84
Kennicott's explorations—Hudson's Bay Company. (R. 1863)	187
Kentucky—	
ancient mounds in—R. Peter. (R. 1872)	271
antiquities from—S. S. Lyon. (R. 1858)	109
Breckinridge county, age of stone and troglodytes of—R. S. Robertson. (R. 1874)	286
Hancock county, antiquities of-J. FRIEL. (R. 1877)	328
Lexington, ancient mound near—R. PETER. (R. 1871)	249
Union county, exploration of ancient mounds in—S. S. Lyon. (R. 1870.)	244
Kepler, his life and works, by M. BERTHRAND. (R. 1869)	228
Kerguelen Island, contributions to natural history of—	
1. Ornithology-J. H. KIDDER; E. COUES	293
11. Oölogy, botany, etc.—J. H. KIDDER and others	
Kershner collection from Kerguelen Island	294
Kew Observatory, apparatus and processes employed in verification of barome-	
ters at—J. Welsh. (R. 1859)	110
Kew Observatory, experiments on aneroid barometers made at—B. STEWART.	224
(R. 1868)	
KIDDER, J. H. Experiments on animal heat of fishes. (P. 1879)	
KIDDER, J. H. Mammals of Kerguelen Island	
KIDDER, J. H., and others. Natural history of Kerguelen Island	
KIDDER, J. H.; Coues, E. Birds of Kerguelen Island	82
Kidney, observations on—J. Jones	
Kinetic theories of gravitation—W. B. TAYLOR. (R. 1876)	
KING, C. B. Catalogue of engravings presented to Smithsonian. (R. 1861)	149
KING, W. M. Account of Indian burial in California. (R. 1874)	286
KIPP, J. Accuracy of Catlin's account of Mandan ceremonies. (R. 1872)	271
KIRBY, W. W. Journey to the Yukon, Russian America. (R. 1864)	188
KIRKWOOD, D. Asteroids between Mars and Jupiter. (R. 1876)	299
Kitchen-midden in Costa Rica, note on shells from—W. H. Dall. (P. 1878.)	882
Kitchen-midden—See Kjækken-mæddings.	105
Kjækken-mæddings in Nova Scotia—J. M. Jones. (R. 1868)	187
Kjækken-mæddings on northwest coast of America—P. SCHUMACHER. (R.	
1878; R. 1874) 275	, 286
Klamath lake dialect, details of Indian conjurer's practice in—A. S. GATSCHET. (E. 1879–80)	476
Klamath lake dialect, "The Relapse" in—A. S. GATSCHET. (E. 1879-80)	476
KLETZINSKY, Prof. Nitrogen bodies of modern chemistry. (R. 1872)	271
KNAPP, G. Earthworks on Arkansas river. (R. 1877)	

KNIGHT, E. H. Study of savage weapons at Centennial Exhibition. (B. 1879)	
845, 6	1 15
KNIGHT, G. H. New system of weights and measures. (R. 1867)	215
KNIGHT, R. T.; HENRY, J. Connection of gales of wind and appearance of aurora. (R. 1871)	249
Knowledge of cryptogamous plants, present state of-W. Reichardt. (R. 1871)	249
Kohl, J. G. On collection of charts and maps of America. (R. 1856)	
KORNHUBER, G. A. Alternate generation and parthenogenesis in the animal kingdom. (R. 1871)	249
KRON, F. J. Antiquities of Stanley and Montgomery counties, North Carolina. (R. 1874)	286
KRON, H. J. Lightning discharges. (R. 1867)	215
Kumlien, L.—	
Ethnology, mammals, and birds of Arctic America	342
Howgate Polar Expedition, natural history of Arctic America	342
Report on explorations in Greenland. (R. 1878)	341
KUNHARDT AND COMPANY. Free freight to Smithsonian Institution between United States and Germany. (R. 1861)	
Kushkushkee, Indian village near Newcastle, Penn., account of—E. M. Mc-Connell. (R. 1871)	
Kutchin Indians—S. Jones. (R. 1866) 214,	_

L.

Labels for collections, etc.	164
LABOULAYE, E. Acknowledgment for books on education. (R. 1867)	215
Lacustrian cities of Switzerland—F. TROYON. (R. 1861)	149
Lacustrian constructions or palafittes of the lake of Neuchâtel—E. DESOR. (R. 1865)	36 0
Lacustrian habitations of Switzerland—F. Troyon. (R. 1861)	149
Lacustrian settlements in Switzerland, abstract of Dr. Keller's report on—A. MORLOT. (R. 1868)	187
La Have Banks, occurrence of Lycodes vahlii on—G. B. Goode; T. H. BEAN. (P. 1879)	333
Lake Neuchâtel, lacustrian constructions of—E. Desor. (R. 1865)	506
Lake Pepin, antiquities on banks of—L. C. Estes. (R. 1866)	
Lake Superior, ancient mining on shores of-C. WHITTLESEY	
Lake Superior copper region, circular relative to ancient mining in—J. HENRY.	149
Lake Superior, Isle Royale, antiquities of—A. C. Davis. (R. 1874)	20

Latin vocabulary with comparative words in English, Spanish, and French 1	70
Latitudes and longitudes, Arctic—E. K. KANE	29
LAUGEL, A. The sun, its chemical analysis. (R. 1861)i	49
Lausanne, report on ethnological collections of Museum at—F. TROYON. (R.	**
1861)	
LAUTENBACH, B. F. Effect of irritation of a polarized nerve. (R. 1878.) 841, 45	
Law, A. E. Antiquities of Blount county, Tennessee. (R. 1874)	
Law of deposit of flood-tide—C. H. Davis	33
Law of variation of temperature in ascending currents of air—J. HANN. (R. 1877)	98
Law of variation of temperature in ascending moist currents—L. Sohncki.	•
(R. 1877)	98
LAWRENCE, G. N.—	
Birds of southwestern Mexico collected by F. E. Sumichrast 2	95
Catalogue of Ober's collection of birds of Antigua and Barbuda, with his notes. (P. 1878)	994 20-
Catalogue of Ober's collection of birds of Dominica, with his notes and observations. (P. 1878)	<u> </u>
Catalogue of Ober's collection of birds of Grenada, including others seen, but not obtained. (P. 1878)	
Catalogue of Ober's collection of birds of Guadeloupe. (P. 1878)	
Catalogue of Ober's collection of birds of Lesser Antilles, with table showing distribution. (P. 1878)	
Catalogue of Ober's collection of birds of Martinique. (P. 1878) 3	
Catalogue of Ober's collection of birds of St. Vincent, with his notes. (P. 1878)	
Description of new species of bird of family Turdides from Dominics, W. I. (P. 1880)	
Description of new species of Icterus from West Indies. (P. 1880)	
Description of new species of parrot, of genus Chrysotis, from Dominica.	
(P. 1880)	
Description of new subspecies of Loxigilla from West Indies. (P. 1881.)	
Laws of atmospheric circulation over the earth—J. H. Coffin	
LEA, I. Check-list of the shells of North America	
LE CONTE, JOHN. Constants of nature. (R. 1878)) 7 1
LE CONTE, J. L.— Classification of coleoptera of North America. Parts 1, 11	65
Coleoptera of Kansas and eastern New Mexico 1	
Instructions for collecting coleoptera. (R. 1858)	1(19
List of coleopters of North America. Part I1	40
New species of North American coleoptera. Parts 1, 11 167, 2	64
Revision of Melsheimer's catalogue of coleoptera	62
LE CONTE, JOSEPH. Lecture on coal. (R. 1857)	07
Lecture on American fire-alarm telegraph—W. F. CHANNING. (R. 1854)	75

Lectures on—	
agricultural chemistry—S. W. Johnson. (R. 1859)	110
Arctic explorations-I. I. HAYES. (R. 1861)	149
astronomy—A. Caswell. (R. 1858)	109
bridges-F. Rogers. (R. 1861)	149
bridges and roads—F. Rogers. (R. 1860)	147
the camel—G. P. Marsh. (R. 1854)	75
cancerous tumors. Toner lecture No. I-J. J. Woodward	266
coal-J. LE CONTE. (R. 1857)	107
dual character of the brain. Toner lecture No. 11-C. E. Brown-Séquard.	291
electro-physiology—C. MATTEUCCI. (R. 1865)	209
insect instincts and transformations—J. G. Morris. (R. 1855)	77
marine alge-W. H. HARVEY. (R. 1855)	77
meteoric stones—J. L. SMITH. (R. 1855)	77
meteorology-R. Russell, with notes by J. Henry. (R. 1854)	75
mollusca, or shell-fish and their allies-P. P. CARPENTER. (R. 1860.) 147	, 152
natural history as applied to farming and gardening—J. G. Morris. (R. 1855)	77
nature and cure of bites of serpents and wounds of poisoned arrows—D. Brainard. (R. 1854)	
nature of reparatory inflammation in arteries after ligature, etc. Toner	
lecture No. vii—E. O. Shakespeare	
oxygen and its combinations-G. I. CHACE. (R. 1855)	77
physical ethnology—D. Wilson. (R. 1862)	
planetary disturbances—E. S. SNELL. (R. 1855)	77
principles of linguistic science—W. D. WHITNEY. (R. 1863) 187	352
relations of time and space—S. ALEXANDER. (R. 1861)	
sanitary drainage of Washington. Toner lecture No. VIII—G. E. WARING, Jr.	
shells of Gulf of California-P. P. CARPENTER. (R. 1859)	
strain and over-action of the heart. Toner lecture No. 111-J. M. DA	
study of high antiquity in Europe-A. Morlor. (R. 1862; R. 1864.) 150), 188
study of nature and mechanism of fever. Toner lecture No. 1V—H. C. Wood	282
surgical complications and sequels of fevers. Toner lecture No. v—W. W. Keen	300
Switzerland-A. D. BACHE. (R. 1870)	
syllabus of a course of, on physics—J. HENRY. (R. 1856)	91
undulatory theory of light—F. A. P. BARNARD. (R. 1862)	150
the Union—H. Reed. (R. 1854)	75
vastness of visible creation—S. Alexander. (R. 1857)	107
zone of small planets between Mars and Juniter—E. Loomis. (R. 1854.)	

Lectures, Toner—	
No. I. Cancerous tumors—Woodward	
II. Dual character of brain—Brown-Séquard	2 91
III. Over-action of heart—Da Costa	279
IV. Study of fever-Wood.	282
v. Continued fevers—Keen	300
VI. Sub-cutaneous surgery—Adams	302
VII. Reparatory inflammation—SHAKESPEARE	321
VIII. Sanitary drainage of Washington-WARING	349
LEE, Dr. Progress of astronomical photography. (R. 1861)	149
LEE, J. C. Y. Ancient ruin in Arizona. (R. 1872)	271
Legacy—See Bequest.	
Legendre, memoir of, by E. DE BEAUMONT. (R. 1867)	215
LEIDY, J.—	
Ancient fauna of Nebraska	
Cretaceous reptiles of the United States	
Extinct sloth tribe of North America	
Extinct species of American ox	
Fauna and flora within living animals	
On parasites	44
Report on fossils from Nebraska. (R. 1851)	5.
Review of cretaceous reptiles of United States. (R. 1864)	18
LEIDY, J.; TRYON, G. W. Report on shells presented to Academy of Natural Sciences. (R. 1865)	Y-
Leipsic Museum of Ethnology—A. Schott; O. T. Mason. (R. 1873)	9
LEMSTRÖM, S.; DE LA RIVE, A. A. Electricity of the atmosphere and the aurora borealis. (R. 1874)	20
Lens, account of Priestley's—J. HENRY. (R. 1859)	
Lepidopsetta isolepis, new flat flsh, found in markets of San Francisco, note on—W. N. Lockington. (P. 1880)	
Lepidoptera—	• • • •
appendix to Morris' synopsis of—B. CLEMENS	i.j.
described, of North America, catalogue of-J. G. Morris	112
described, of North America, synopsis of. Part 1-J. G. Morris	133
diurnal, of Arctic America—W. H. Edwards	3-1- 3-2-
instructions for collecting—B. CLEMENS. (R. 1858)	Ites
introduction to Morris' synopsis of—V. HEINEMANN	13,
method of preserving-T. R. PEALE. (R. 1868)	15
nocturnal, of Arctic America—S. H. Scudder	342
notes to Morris' synopsis of-W. H. EDWARDS	13.
LEPSIUS, R. Presentation of books on Egypt. (R. 1860)	14:
Leptocardii, arrangement of families of—T. GILL	±31
LESLEY, J. P. On the classification of books. (R. 1862)	134

LESPIAULT, Prof. Small planets between Mars and Jupiter. (R. 1861) 149
Lesser Antilles, catalogue of Ober's collection of birds of-G. N. LAWBENCE.
(P. 1878) 882
Letter—
by M. Hoek in reference to meteoric shower of November 13, 1867 217
from Robert Hare relative to gift of apparatus I, 829
from Joseph Henry to Secretary Treasury on payment of interest in coin. (R. 1865)
from Richard Rush relative to Smithson. (R. 1858) 67, 828
LETTERMAN, J. Sketch of the Navajo Indians. (R. 1855) 77
Letters—
of introduction. (R. 1865) 209
on work of the Museum—B. PHILLIPS. (P. 1881) 467, 454
relative to programme of organization of Smithsonian Institution 828
relative to Smithson's bequest 328
Leurynnis, Lockington, identity of, with Lycodopsis, Collet-T. GILL. (P.
Level, fluctuations of, in North American lakes—C. Whittlesey 119
Lewis, J. Instructions for collecting land and fresh-water shells. (R. 1866.) 214, 363
Lewis, J., letter of, to G. W. Tryon, Jr., on shells 258
Lewis; Quale. Account of cryolite of Greenland. (R. 1866) 214
Lexington, Kentucky, ancient mound near—R. Peter. (R. 1871) 249
Liberia, mixed races in—E. D. BLYDEN. (R. 1870) 244
Liberia College, facts respecting—A. CRUMMELL. (R. 1861) 149
Libraries—
circular respecting new report on—J. HENRY. (R. 1854) 75
construction of catalogues of, and their publication by means of separate stereotyped titles—C. C. Jewett
••
public, in United States and British Provinces, list of—W. J. RHEES. 116, 238
public of United States, notices of—C. C. JEWETT. (R. 1849) 21, 25
report of Commission on stereotype catalogue of 47
rules for cataloguing—C. C. JEWETT 47
Library— Beaufort, deposit of—E. M. STANTON. (R. 1862)
Bishop Johns', deposit of—E. CANBY. (R. 1862) 150
Imperial, of Vienna, books presented by. (R. 1865) 209
of Bureau of Ethnology, catalogue of linguistic manuscripts in—J. C.
PILLING. (E. 1879-80) 476
of Indian linguistics, account of—J. G. SHEA. (R. 1861) 149
of National Museum, circular asking contributions to. (P. 1881) 467, 458
on a national—W. S. Jevons. (R. 1878)

Library of Congress—
act of Congress to transfer Smithsonian library to. (R. 1865) 209, 328
catalogue of publications of societies and of periodical works belonging to the Smithsonian Institution deposited in
meteorological articles received by the Smithsonian Institution and deposited in. (R. 1871; R. 1873) 249, 275
Library of Smithsonian Institution—
act of Congress to transfer, to Library of Congress. (B. 1865) 209, 328
additions to—See each annual report.
and copyright system—C. C. JEWETT. (R. 1851)51
and Halliwell manuscripts, report on—C. C. JEWETT. (R. 1852) 57
catalogue of publications of societies and other periodical works in, 117, 179
donations to, from foreign institutions. (R. 1864) 188
publications of learned societies and periodicals in. Parts 1, 11 73, 85
report on—C. C. JEWETT. (R. 1848, 1849, 1853)
report on plan of—C. C. JEWETT. (R. 1847) H
statistics of, 1846–1877 329
Lichens of Arctic America—E. Tuckerman
Lichens of Kerguelen Island—E. TUCKERMAN 294
LIEBER, F. Vocal sounds of Laura Bridgman compared with elements of phonetic language 12
Liebig, J. von. Induction and deduction. (R. 1870) 244
Life—
and character of Joseph Henry-J. C. WELLING 356, 338
and labors of Henry Gustavus Magnus. (R. 1870) 241
and scientific labors of Stefano Marianini—C. MATTEUCCI. (R. 1869) 228
and works of Michael Faraday—A. A. De LA RIVE. (R. 1867) 215
and works of Kepler-M. BERTHRAND. (R. 1869) 23
of George Catlin-J. HENRY. (R. 1872)
of Prof. Chester Dewey-M. B. Anderson. (R. 1870) 244
Life—See Biography, Eulogy, Memoirs.
Light—
and heat of sun, relative intensity of-L. W. MEECH. (R. 1856) 91,83
polar, or aurora borealis, its phenomena and laws—E. Loomis. (R. 1865.) 209
undulatory theory of, lectures on-F. A. P. BARNARD. (R. 1862) 150
velocity of, essay on—C. DELAUNAY. (R. 1864) 188, 354
Light-House Board—
investigations by, relative to illuminating materials—J. Henry. (R. 1880)42, 389
proceedings of, on death of J. Henry356
researches by, in sound—J. HENRY. (R. 1878) 341, 406
Lighting, architecture in relation to—D. B. REID. (R. 1856)91

Lightning—	
and thunder, observations on—S. MASTERMAN. (R. 1855)	77
discharges, accounts of-G. W. Dodge and others. (R. 1867)	215
effect of—S. L. HILLIER. (R. 1866)	214
Lightning-rods, directions for constructing—J. HENRY	237
Lille, Imperial Society of Science, Agriculture, and Arts of. Prize questions. (R. 1865)	209
LILLJEBORG, W. Outline of systematic review of the classification of birds. (R. 1865)	. 36 4
Limitations to use of some anthropological data—J. W. POWELL. (E. 1879-80.)	
Limpets from deep waters off eastern coast of U.SW.H. Dall. (P. 1881.)	
Limpets of Alaska and Arctic region, report on—W. H. Dall. (P. 1878)	
Lindheimer's botanical exploration in New Mexico and California, account of— A. GRAY. (R. 1849)	21
Linguistic manuscripts in library of Bureau of Ethnology, catalogue of—J. C. Pilling. (E. 1879-80)	476
Linguistic science, lectures on principles of-W. D. WHITNEY. (R. 1863.) 187	352
Linguistics, Indian, account of library of-J. G. SHEA. (R. 1861)	149
Linguistics, Indian, recommendation of Shea's—G. GIBBS and others. (R. 1861.)	149
LIONNET, M. Vocabulary of the Chinook jargon	68
Liparis ranula, new species of, obtained by U.S. Fish Commission off Halifax,	
Nova Scotia-G. B. Goode; T. H. BEAN. (P. 1879)	888
Liquids, expansion of, by heat—F. W. CLARKE	289
Liquids, specific heats of—F. W. CLARKE	276
Lisboa, M. M., books on Brazil, presented by. (R. 1865)	209
List of—	
additions to Museum—See each annual report.	
addresses of foreign institutions, 1862-1865. (R. 1865)	209
alge of Rhode Island—S. T. OLNEY	241
American correspondents	69
American libraries and public institutions—W. J. RHEES.	238
anthropological publications of Charles Rau. (P. 1881)	467
apparatus available for scientific research. (R. 1878)	34 1
articles deposited by Smithsonian Institution in Corcoran Gallery of Art. (R. 1874)	286
birds and mammalia of Missouri river—E. HARRIS. (R. 1850)	28
birds of central California, partial—L. Belding. (P. 1878)	882
birds of District of Columbia—E. Cours; S. S. Prentiss. (R. 1861)	149
casts of heads of Indian boys and girls at Hampton, Va.—R. H. PRATT. (P. 1879)	888
coleoptera of North America. Part I-J. L. LE CONTE	140
collections presented by foreign Centennial commissioners. (R. 1876)	299

List of—Continued.
described birds of Mexico, Central America, and West Indies not in Smithsonian Institution
described species of humming-birds—D. G. Elliot
desiderata among North American birds-R. Ridgway. (P. 1881) 46
domestic institutions in correspondence with Smithsonian Institution. 69, 23
duplicates of fishes of Pacific coast distributed by Smithsonian Institution in 1881—D. S. JORDAN; P. L. JOUY. (P. 1881)46
European fishes in National Museum—T. H. BEAN. (P. 1879) 33
expeditions from which specimens in Museum have been derived—S. F. BAIRD. (R. 1867)
fishes collected by Lieut. H. E. Nichols in Gulf of California—D. S. JORDAN; C. H. GILBERT. (P. 1881)
fishes of Pacific coast of United States—D. S. JORDAN; C. H. GILBERT. (P. 1880)
foreign agents of Smithsonian Institution in 1876. (See also annual reports)
foreign correspondents of Smithsonian Institution. 64, 154, 225, 243, 309, 46
foreign correspondents of Smithsonian Institution, systematic index to 25
generic names of animals—S. H. SCUDDER 47
Indian vocabularies received from the Wheeler Expedition. (R. 1874.)
institutions, libraries, colleges, and other establishments in correspondence with the Smithsonian Institution—W. J. Rhees
invertebrate fossils of North America— Cretaceous and jurassic—F. B. MEEK
Eccene and oligocene—T. A. CONBAD 20
Miocene-F. B. MEEK
marine invertebrata of New England distributed by United States Fish Commission—A. E. Verrill; R. Rathbun. (P. 1879)
marine invertebrates, mainly from New England, distributed by U.S. National Museum—R. RATHBUN. (P. 1881) 467, 465. 47
meteorological material contributed to Smithsonian Institution. (R. 1860-1865, 1867-1871) 147, 149, 150, 187, 188, 209, 215, 224, 228, 244. 24
meteorological stations and observers. (R. 1849; R. 1853-1873) 21, 67, 73
minerals in United States National Museum—F. M. ENDLICH. (R. 1873; P. 1880) 275, 42
North American batrachia and reptilia—E. D. Cope 292
North American species of myriapods of family Lysiopetalide—J. A. RYDER. (P. 1880)
official publications of the United States Government between 1868 and 1881—G. H. Boehmer
papers presented to Royal Society by James Smithson. (R. 1853) 67, 320 periodicals received by the Institution. (R. 1880)

List of—Continued.
photographic portraits of North American Indians in Smithsonian Institution 210
plants of the upper Missouri—T. C. Porter. (R. 1850) 20
plants of Washington and vicinity—L. F. WARD 460
principal literary and scientific institutions in United States, 1879 335
public libraries, institutions, and societies in United States and British
Provinces of North America—W. J. Rhees
publications of the Smithsonian Institution. (R. 1868) 74, 203, 224, 226 245, 278, 290, 801, 844, 487, 478
publications of the United States National Museum. (P. 1881) 467, 474
Regents, officers, and assistants of the Smithsonian Institution and U.S. National Museum. (See also each annual report) 466, 290
scientific papers of A. D. Bache—B. A. Gould. (R. 1870) 244, 329, 379
shells of exploring expedition193
shells of North America—I. LEA; P. P. CARPENTER; W. STIMPSON;
W. G. BINNEY; T. PRIME
species of Middle and South American birds not in United States National
Museum—R. RIDGWAY. (P. 1881) 46'
substances derived from animal kingdom—G. B. Goode 297
Literary exchanges—See Exchanges.
Literary study of jade—S. Blondel. (R. 1876)
Littoral marine fauna of Provincetown, Mass.—R. RATHBUN. (P. 1880) 428
Liver, observations on the—J. Jones82
Livermore, G. Report on Jewett's general stereotype catalogue of public libraries 47
Living animals, flora and fauna within—J. LEIDY
Lloyd, North German, free freight between Germany and the United States. (R. 1858) 109
LLOYD, W. A. Exchange of specimens. (R. 1867) 218
LLOYD, W. A. Sparrows sent to the United States. (R. 1867) 216
Locke, J. Catalogue of rocks, minerals, ores, and fossils. (R. 1854) 78
Locke, J. Observations on terrestrial magnetism 35
Lockett, S. H. Mounds in Louisiana. (R. 1872) 271
Lockington, W. N.—
Description of new chiroid fish (Myriolepis zonifer) from Monterey Bay, California. (P. 1880)
Description of new fish from Alaska (Uranidea microstoma). (P. 1880.) 425
Description of new genus and some new species of California fishes (Icosteus ænigmaticus and Osmerus attenuatus). (P. 1880) 426
Description of new genus and species of Cottidæ. (P. 1881) 467
Description of new sparoid fish (Sparus brachysomus) from Lower California. (P. 1880)

Lockington, W. N.—Continued.	
Description of new species of Agonidae (Brachyopsis verrucosus) from coast of California. (P. 1880)	425
Description of new species of <i>Prionotus (Prionotus stephanophyrs</i>) from coast of California. (P. 1880)	425
Descriptions of new genera and species of fishes from coast of California. (P. 1879)	833
Note on new flat fish (Lepidopsetta isolepis) found in markets of San Francisco. (P. 1880)	425
Remarks on species of genus Chirus found in San Francisco market. (P. 1880)	425
Review of the Pleuronectidæ of San Francisco. (P. 1879)	333
Locusts and grasshoppers of America—A. S. TAYLOR. (R. 1858)	109
Lorw, H.—	
Monographs of the diptera of North America-	
Part I. Edited by R. OSTEN SACKEN	141
Part II. Edited by R. OSTEN SACKEN	171
Part III	256
Review of North American Trypetina	256
LOEW, H.; OSTEN SACKEN, R. Instructions for collecting diptera. (R. 1858.)	109
LOGAN, T. M.—	
· Climate of California. (R. 1855)	77
· Meteorological observations at Sacramento, California. (R. 1854)	75
Meteorology of Sacramento, California. (R. 1857)	107
LOGAN, W. E. Request for duplicate shells. (R. 1859)	110
LOMBARD, H. C. Report on the transactions of Society of Physics and Natu-	
ral History of Geneva, July, 1868, to June, 1869. (R. 1869)	228
Lombardy, Royal Scientific and Literary Institute of. Prize questions. (R. 1865.)	209
London Institution of Civil Engineers. Prize questions. (R. 1862)	150
London Royal Horticultural Society. Exchange of publications. (R. 1861)	149
London, Royal Society of, origin and history of-C. A. ALEXANDER. (B. 1868.)	187
Longfellow, H. W., and others. Report of American Academy of Arts and	
Sciences on organization of Smithsonian Institution. (R. 1853)	
Long Island, fishes on coast of New Jersey and—S. F. BAIRD. (R. 1854.) 75,	348
Longitude, transatlantic—B. A. Gould	223
Longitudes, Arctic-E. K. KANE	129
Loomis, E.—	
Aurora borealis, or polar light, its phenomena and laws. (R. 1865)	209
Lecture on zone of small planets between Mars and Jupiter. (R. 1854.)	75
On certain storms in Europe and America, December, 1836	127
Report on meteorology of the United States. (R. 1847)	Ħ
Lopholatilus chamæleonticeps, new genus and species of fish from New England, description of—G. B. Goode; T. H. Bran. (P. 1879)	
· · · · · · · · · · · · · · · · · · ·	, 36 5

•	
Loud, F. H. Discussion of Prof. Snell's barometric observations. (R. 1880.) 442,	435
Louisiana, Caddo parish, Indian remains in—T. P. Hotchkiss. (R. 1872)	271
Louisiana, lower, geology of, and salt deposit of Petite Anse Island—E. W. HILGARD	2 4 8
Louisiana, mounds in—S. H. Lockett. (R. 1872)	271
LOVERING, J. Memorial of J. Henry	856
Lowe, T. S. C., aeronautic voyage of, reply to memorial on—J. HENRY. (R. 1860)	147
Lowe, T. S. C., memorial of citizens of Philadelphia relative to aeronautic voyage of, across the Atlantic. (R. 1860)	147
Loxigilla, description of new subspecies of, from West Indies-G. N. LAW- RENCE. (P. 1881)	467
LUBBOCK, J. North American archæology. (R. 1862)	150
LUBBOCK, J. Social and religious condition of lower races of man. (R. 1869.)	228
Lucernarise, and their allies, anatomy and physiology of-H. J. CLARK	242
Ludewig, H. E., and others. On publication of Spanish works on New Mexico. (R. 1855)	77
LUGGER, O. Occurrence of Canada porcupine in Maryland. (P. 1881)	467
Lunar effect on magnetic force—A. D. BACHE	195
LUPTON, N. T. Breeding habits of sea catfish (Ariopsis milberti). (P. 1878.)	882
Luray cavern, blind form of Lysiopetalides from J. A. RYDER. (P. 1880)	42 5
Luray cavern, report of visit to-O. T. MASON and others. (R. 1880) 442,	133
Lussac, Gay, eulogy on, by F. ARAGO. (R. 1876).	299
Lutjanus Blackfordii, from coast of Florida, description of—G. B. Goode; T. H. Bean. (P. 1878)	882
Lutjanus Stearnsii, from coast of Florida, description of—G. B. Goode; T. H. Bean. (P. 1878)	882
Lycodes paxillus, description of species of-G. B. Goode; T. H. BEAN. (P. 1879)	883
Lycodes Turneri, from Alaska, description of-T. H. BEAN. (P. 1878)	38
Lycodes Vahlii, occurrence of, on La Have and Grand Banks—G. B. GOODE; T. H. BEAN. (P. 1879)	888
Lycodopsis, Collett, identity of, with Leurynnis, Lockington—T. GILL. (P. 1880)	425
Lycopodiacese of Kerguelen Island—A. GRAY	294
LYKINS, W. H. R. Antiquities of Kansas City, Missouri. (R. 1877)	328
LYMAN, T. Ethnographical collections. (R. 1862)	150
Lyon, S. S. Antiquities from Kentucky. (R. 1858)	109
Lyon, S. S. Mounds in Union county, Kentucky. (R. 1870)	244
Lyon, W. B. Antiquities in New Mexico. (R. 1871)	249
Lysiopetalidse, North American species of—J. A. RYDER. (P. 1880)	425

\mathbf{M}_{-}

MCCLINTOCK, F. L. Meteorological observations in the Arctic Seas 14
McConnell, E. M. Account of old Indian village, Kushkushkee, near Newcastle, Pennsylvania. (R. 1871)
McCosh, J. Prayer at memorial service of Prof. Joseph Henry 35
MACDONALD, J. A. Analyses of Chinese and Japanese coals 20
McHenry, J. W.; Haile, J. Antiquities of Jackson Co., Tenn. (R. 1874.)
McKAY, C. L. Review of Centrarchidæ. (P. 1881) 46
McKinley, W. Mounds in McIntosh and Early counties, Georgia. (R. 1872.) 27
McMasters, S. Y. Language of Navajos said to resemble the Welsh. (R. 1865.)
McParlin, T. A. Notes on history and climate of New Mexico. (R. 1877.) 323, 39
McWhorter, T. Ancient mounds of Mercer county, Illinois. (R. 1874) 28
Mackerel, the frigate, (Auxis Rochei,) on New England coast—G. B. Goods. (P. 1880)
Madrid Royal Academy of Science. Exchanges. (R. 1861)
MAEDLER, J. H. Movement of the stars around a central point. (R. 1859) 11
Magendie, F., memoir of, by M. FLOURENS. (R. 1866)21
Magnetic-
observations at Girard College, discussion of—A. D. BACHE: Part I. Eleven year period in amplitude of solar diurnal variation 113
Part II. Solar diurnal variation in magnetic declination 12
Part III. Influence of moon on magnetic declination 13
Parts IV, V, VI. Horizontal force16
Parts vii, viii, ix. Vertical force 175
Parts x, xI, XII. Dip and total force 18
Parts 1-x11. Complete 19
observations in the Arctic Seas—E. K. KANE
observations in the Arctic Seas—I. I. HAYES
observations, on continuance of—E. SABINE. (R. 1858)100
observations on the iron-clad "Monadnock"—W. HARKNESS 235
observatory at the Smithsonian Institution, description of—J. E. HIL- GARD. (R. 1859)
storms—E. Sabine. (R. 1860)
survey of Pennsylvania, (etc.)—A. D. BACHE 166
telegraph—See Telegraph.
Magnetism— history of discovery relative to. (R. 1863)————————————————————————————————————
later views of connection of electricity and—H. Helmholtz; J. C.
MAXWELL. (R. 1878)275
terrestrial, observations on—W. HARKNESS239
terrestrial, observations on—J. Locke 35
terrestrial, observations on, in Mexico-Baron von Müller; A. Sonntag. 114

OF SMITHSONIAN PUBLICATIONS.	239
Magnus, Henry Gustavus, life and labors of. (R. 1870)	244
MAGRINI, L. Continuous vibratory movement of all matter. (R. 1868)	
MAILLY, E.—	
Estimate of population of the world. (R. 1878)	275
Eulogy on Quetelet. (R. 1874)	
History of Royal Institution of Great Britain. (R. 1867)	
Maine—	
Brunswick, results of meteorological observations at—P. CLEAVELAND.	204
Schoodic lakes, description of apparently new species of Gasterosteus from—T. H. BEAN. (P. 1879)	888
Kennebec river, opening and closing of-R. H. GARDINER. (R. 1858.)	109
Portland Society of Natural History, account of—E. C. Bolles. (R. 1867)	215
remains of walrus (?) in—C. H. Boyd. (P. 1881)	467
Makah Indians, account of the—G. GIBBS	220
Makah Indians, memoir on the-J. G. SWAN	220
Makah vocabulary—J. G. Swan	220
MALLERY, G. Sign language among North American Indians compared with	
that among other peoples and deaf-mutes. (E. 1879-80)	476
MALLET, R. On observations of earthquake phenomena. (R. 1859)	
MALLINKROOT, C. Changes of wind. (R. 1866)	
Maltheidæ, note on the—T. GILL. (P. 1878)	882
Mammals—.	
and birds of Missouri river, list of—E. HABBIS. (R. 1850)	28
arrangement of families of-T. GILL	230
bibliography of-T. GILL	280
extinct, of Nebraska—J. LEIDY	58
North American, catalogue of—S. F. BAIRD	
of Arctic America—L. Kumlien	342
of Kerguelen Island—J. H. KIDDER	
synoptical tables of—T. GILL	230
Man-	
as the contemporary of the mammoth and reindeer in middle Europe. (R. 1867)	215
ancient, in Michigan, characteristics pertaining to—H. GILLMAN. (R. 1875)	393
later prehistoric, remains of, from caves in Alaska—W. H. DALL	318
on traces of the early mental condition of—E. B. TAYLOR. (R. 1867)	215
social and religious condition of the lower races of—J. LUBBOCK. (R. 1869.)	228
Management of the Smithsonian Institution, report of Senate Judiciary Committee on—A. P. BUTLER. (R. 1855)	7 7
Mandan ceremonies, accuracy of Catlin's account of-J. KIPP. (R. 1872)	271
MANN, C. Habits of a species of salamander. (R. 1854)	75
Manners of Shoshone Indians—A. G. BRACKETT. (R. 1879)	845

Manufacture of spear and arrow points along Savannah river—C. C. Jones, Jr. (R. 1879)	345
Manuscripts, Halliwell, report on—C. C. JEWETT. (R. 1852)	5.
Manuscripts, etc., historical and geographical, catalogue of collection of—L. Berlandier. (R. 1854)	75
Manuscripts, linguistic, in library of Bureau of Ethnology, catalogue of— J. C. Pilling. (E. 1879-80)	476
Map of—	
antiquities of United States, proposed—A. J. HILL. (R. 1861)	119
solar eclipse of March 15, 1858—T. HILL	101
stars for observations on meteorites	359
stars near North Pole for observations on auroras	350
United States, ethnological, proposed—G. Gibbs. (R. 1862)	150
Maps-	
and charts of America, on a collection of-J. G. Kohl. (R. 1856)	91
catalogue of collection of-L. BERLANDIER. (R. 1854)	75
ethnological, of North America, suggestions relative to—L. H. MORGAN.	
(R. 1861)	149
of winds of the globe—S. J. COFFIN	268
Marble used in extension of U.S. Capitol, account of-J. HENRY. (R. 1856.)	91
MARCET, Prof. Report on transactions of Society of Physics and Natural History of Geneva, July, 1862, to June, 1868. (R. 1868)	
MAREY, E. J. Natural history of organized bodies. (R. 1867)	
MARRY, E. J. Phenomena of flight in the animal kingdom. (R. 1869)	
Marianini, Stefano, life and scientific labors of, by C. MATTEUCCI. (R. 1869.)	
Marietta, Ohio, meteorological observations at—S. P. HILDRETH; J. Wood	
•	
Marine alge, lecture on—W. H. HARVEY. (R. 1855)	,,
Marine algæ of North America—W. H. HARVEY: Part 1. Melanospermeæ	22
Part II. Rhodospermeæ	
Part III. Chlorospermeæ	
Parts 1, 11, 111, complete	30
Marine invertebrates— distributed by U.S. National Museum—R. RATHBUN. (P. 1881.) 465,	467
of Grand Manan—W. STIMPSON	
of New England coast distributed by United States Fish Commission— A. E. VERRILL; R. RATHBUN. (P. 1879)	333
of northeastern coast of America, notice of recent additions to, with descriptions of new genera and species—A. E. VERRILL:	
Part I. Annelida, etc. (P. 1879)	33 3
Part II. Mollusca, with notes on Annelida, etc. (P. 1880)	
Part III. Catalogue of Mollusca recently added to fauna of southern New England. (P. 1880)	
on collection and preservation of—W. STIMPSON	

OF SMITHSONIAN PUBLICATIONS.	241
Maritime disasters of the Antilles. (R. 1867)	215
Marrow of animals, uses of, among Indians—T. R. PEALE. (R. 1870)	244
Mars, secular variations of elements of orbit of—J. N. STOCKWELL	282
Mars, small planets between Jupiter and—Prof. LESPIAULT. (R. 1861)	149
Marschall, A., list of generic names supplemental to those catalogued by—S. H.	470
SCUDDER	470
MARSH, G. P. Lecture on the camel. (R. 1854)	75
MARSH, G. P., and others. On publication of Squier and Davis' ancient mon- uments. (R. 1847)	I. K
Marsipobranchii, arrangement of families of—T. GILL	-
MARTIN, S. D. Account of lightning discharges. (R. 1867)	
Martinique, catalogue of Ober's collection of birds of-G. N. LAWBENCE. (P.	
1878)	882
MARTINS, C. Vegetable colonization of the British Isles of Shetland, Faroe, and Iceland. (R. 1858)	109
Martius, C. F. P. von, honorary medal to—W. HAIDINGER. (R. 1868)	187
Martius, C. F. P. von, memoir of, by C. RAU. (R. 1869)	
Maryland, Charles county, antiquities of—O. N. BRYAN. (R. 1874)	286
Maryland, occurrence of Canada porcupine in—O. Lugger. (P. 1881)	
Mason, C. Statement relative to the telegraph. (R. 1857)	
Mason, O. T.—	
Abstracts of anthropological correspondence, 1880. (R. 1880)	442
Anthropological investigations in 1879. (R. 1879) 845,	420
Anthropological investigations in 1880. (R. 1880) 442,	432
Bibliography of anthropology. (R. 1879; R. 1880) 845,	442
Latimer collection of antiquities from Porto Rico. (R. 1876) 299,	397
Leipsic Museum of Ethnology. (R. 1878)	275
Record of progress in anthropology, 1879, 1880. (R. 1880) 442,	432
Report on Luray Cavern. (R. 1880) 442,	433
Summary of anthropological correspondence, 1879. (R. 1879) 845,	420
Massachusetts— Buzzards' Bay, occurrence of Belone latimanus in—G. B. Goode. (P.	
1878)	882
Provincetown, Cape Cod, littoral marine fauna of—R. RATHBUN. (P. 1880)	425
Vineyard Sound, occurrence of oceanic bonito in-V. N. EDWARDS.	882
MASTERMAN, S. Observations on natural phenomena, shooting stars, aurora, etc. (R. 1857)	
MASTERMAN, S. Observations on thunder and lightning. (R. 1855)	107 77
Materia medica collection, classification and arrangement of-J. M. FLINT.	-
(P. 1881) 467,	450
Materia medica section of the U.S. National Museum, memoranda for collectors of drugs for the—J. M. FLINT. (P. 1881)——————————————————————————————————	452

Matter, continuous vibratory movement of—L. MAGRINI. (R. 1868) 724
Matteucci, C.—
Electrical currents of the earth. (R. 1867; R. 1869) 215, 228
Lectures on electro-physiology. (B. 1865) 209
Life and scientific labors of Stefano Marianini. (R. 1869)
Mauvaises Terres, expedition to the upper Missouri and the—T. A. CULBERT- son. (R. 1850)
Mauvaises Terres of Nebraska, extinct mammalia and chelonia from the—J. LEIDY
MAXWELL, J. C.; HELMHOLTZ, H. Later views of connection of electricity and magnetism. (R. 1878)
Maya sculpture, remarks on ancient relic of—A. Schott. (R. 1871) 24
MAYBERRY, S. P. Gradual approach of sea upon land. (R. 1867) 21
MAYBERRY, S. P. Shell heaps at mouth of St. John's river, Florida. (R. 1877)
MAYER, A. M. Henry as a discoverer. Memorial address 356, 415
MAYER, A. M., translation by, of Delaunay's essay on velocity of light. (R. 1864)
MAYER, B. Observations on Mexican history and archeology8
Mazatlan, descriptions of new fishes from—D. S. JORDAN; C. H. GILBERT. (P. 1881)
MEACHAM, J. Minority report of Committee of Regents on distribution of Smithson income. (R. 1858)
MEADS, O. Memorial of Joseph Henry
Means available for correcting measures of sun's distance—G. B. AIRY. (R. 1859)
Means of destroying the grasshopper—V. Motschulsky. (R. 1858)
Measurements, anthropological, table of-Schenzen; Schwarz. (R. 1866) 21
Measurements, mountain—A. Guyor. (R. 1862)
Measures—
and weights, English and French, tables of. (R. 1863; R. 1864) 187, 18
and weights, metric system of, with tables—H. A. NEWTON. (R. 1865.) 209, 373
and weights, new system of, with 8 as the metrical number—G. H. KNIGHT. (R. 1867)
of sun's distance—G. B. AIRY. (R. 1859)
Measuring instrument, use of galvanometer as—J. C. Poggendorff. (R. 1859.)
Mechanics and artisans, scientific education of—A. P. PEABODY (R. 1872.) 271,380
Mechanism of fever, study of. Toner lecture No. IV-H. C. Wood 285
Medal to Von Martius—W. Haidinger. (R. 1863)
Medical practice, relation of food to work and its bearing on—S. HAUGHTON. (R. 1870)
Medicines, classification of forms in which drugs and, appear and are administered—J. M. FLINT. (P. 1881)——————————————————————————————————
Medulla oblongata and trapezium, grav substance of the-J. Dran 173

MERCH, L. W. Relative intensity of heat and light of the sun. (R. 1856.) 91,.83	
MEEK, F. B. Check-list of invertebrate fossils of North America. Oretaceous	
and jurassic	
Merk, F. B. Check-list of invertebrate fossils of North America. Miocene. 183	
MEEK, F. B, and others. Scientific instructions to Capt. Hall. (R. 1871) 249	
MERK, F. B.; HAYDEN, F. V. Palsontology of upper Missouri. Part 1 172	
MEEK, J. B. Account of tornado in Spruce Creek valley, Centre county, Penn.	
(R. 1871)	
MEIGS, J. A. Description of human skull from Rock Bluff, Illinois. (R. 1867.) 215	
Melanians of North America—G. W. TRYON, Jr. 258	
Melanospermess of North America, descriptions of—W. H. HARVEY 32	
Melbourne Mining Department. Exchange system. (R. 1865) 209	
MELSHEIMER, F. E. Catalogue of described coleoptera of the United States 62	
Melting points, bibliography of-F. W. CLARKE 255	
Melting points, tables of. Part I-F. W. CLARKE 255	
Melting points, tables of. First supplement—F. W. CLARKE 288	
Memoir—See Biography, Life, Eulogy.	
Memoir of—	
Agassiz, Louis, by R. P. STEBBINS. (R. 1878) 275	
Babbage, Charles, by N. S. Dodge. (R. 1878) 275	
Beautemps-Beaupré, C. F, by E. DE BEAUMONT. (R. 1868) 187	
Blainville, Ducrotay de, by M. FLOURENS. (R. 1865) 209	
Bravais, Auguste, by E. DE BEAUMONT. (R. 1869) 228	
Buch, Leopold von, by M. FLOURENS. (R. 1862)	
Candolle, Pyramus de, by M. FLOURENS. (R. 1859) 110	
Clark, H. J., by A. S. PACKARD242	
Cuvier, G., by M. FLOURENS. (R. 1868) 224	
Delambre, J. B., by J. Fourier. (R. 1864) 188	
Encke, J. F., by G. HAGEN. (R. 1868) 224	
Gibbs, George, by J. A. STEVENS. (R. 1873) 275	
Hauy, René Just, by G. Cuvier. (R. 1860) 147	
Henry, Joseph, by A. GRAY. (R. 1878) 841, 856, 407	
Henry, Joseph, by J. Lovering 356	
Henry, Joseph, by S. Newcomb	
Henry, Joseph, by W. B. TAYLOR	
Herschel, Sir John Frederick William, by N. S. Dodge. (R. 1871) 249	
Hodgkinson, Eaton, by R. RAWSON. (R. 1868) 224	
Legendre, A. M., by E. DE BEAUMONT. (R. 1867) 215	
Magendie, F., by M. FLOURENS. (R. 1866)214	
Martius, C. F. P. von, by C. RAU. (R. 1869) 228, 440, 251	
Oersted, H. C., by E. DE BEAUMONT. (R. 1868) 224	
Peltier, J. C. A., by F. A. Peltier. (R. 1867) 215	
,, -J & ((

Memoir of—Continued.
Priestley, J., by G. CUVIER. (R. 1858)
Saint-Hilaire, Isidore Geoffroy, by J. L. QUATREFAGES. (R. 1862) 150
Saint-Hilaire, Geoffroy, by M. FLOURENS. (R. 1861) 149
Seaton, W. W., by J. HENRY. (R. 1866)
Thénard, Louis Jacques, by M. FLOURENS. (R. 1862) 150
Torrey, John, by A. GRAY. (R. 1878)
Memoir on—
anatomy and physiology of lucernarise—H. J. CLARK 242
explosiveness of nitre—R. HARE17
extinct sloth tribe of North America—J. LEIDY
extinct species of American ox-J. LEIDY41
meteorological subjects, by J. Hann and others. Translated by C. Abbr. (R. 1877)
methods of interpolation—E. L. Dr Forest. (R. 1871, 1878) 249. 273
mosasaurus—R. W. GIBBES
Memoranda for collectors of drugs for materia medica section of the U.S. National Museum—J. M. FLINT. (P. 1881) 467, 455
Memoranda of inquiry relative to food fishes of United States 23
Memorial-
address on Joseph Henry—A. M. MAYER 356, 417
of citizens of Philadelphia relative to Lowe's aeronautic voyage across the Atlantic. (R. 1860) 147, 32
of Joseph Henry. (Published by Congress)
of Regents to Congress asking appropriations for Museum. (R. 1867.) 215, 32
of Regents to Congress relative to Smithson fund. (R. 1850; R. 1866.) 28, 214, 32
to Congress in behalf of Regents relative to new museum building—G. BANCROFT. (R. 1876)
Menhaden, gulf, (Brevoortia patronus), note on—S. STEARNS. (P. 1878) 333
Mental condition of man, early traces of—E. B. TYLOR. (R. 1867) 213
Mercury, secular variations of elements of orbit of, with tables—J. N. Stock-
MERINO, St. M. Figure of the earth. (R. 1868)
Merlucius, diagnosis of genus of fish related to—G. B. Goode. (P. 1880) 42
MERRILL, J. C. Habits of Rocky Mountain goat. (P. 1879)
MERRILL, J. C. List of birds observed near Fort Brown, Texas, 1876-1878. (P. 1878)
Mesozoic rocks of Arkansas, Wyoming, Colorado, and Utah, descriptions of new invertebrate fossils from—C. A. WHITE. (P. 1880)
Metal, hydrogen as gas and—J. E. REYNOLDS. (R. 1870)
Metallic castings of delicate natural objects. (P. 1884) 467
Metamorphism and formation of crystalline rocks—G. A. DAUBRÉE. (R. 1861.) 149
Meteoric fire-ball, orbit and phenomena of—I. H. COFFIN

OF BMIIMBONIAN I OBDIONATORO.	
Meteoric shower of November 18, 1867, letter of M. Hoek, relative to	217
Meteoric stones, lectures on—J. L. SMITH. (R. 1855)	77
Meteorite—	
in Mexico—A. Woodworth. (R. 1867)	215
in Mexico, account of—R. SIMSON. (R. 1867)	215
in Mexico, discovery of large—W. M. PIERSON. (R. 1878)	275
Tucson, account of—S. Ainsa. (R. 1868)	187
Tueson, account of—B. J. D. IRWIN. (R. 1868)	187
Meteorites—	
extract from discourse on—F. Brenndecke. (R. 1869)	228
in Mexico-J. H. CARLETON. (R. 1865)	209
in mineralogical collection of Yale College, catalogue of—G. J. Brush. (R. 1868)	224
Synthetic experiments relative to—G. A. DAUBRÉE. (R. 1868)	224
Meteorograph, universal, for detached observatories—E. H. von BAUMHAUER. (R. 1879)	845
Meteorological-	
articles in periodicals received by Smithsonian Institution and deposited in Library of Congress. (R. 1871; R. 1878) 249	, 275
correspondence of the Institution, extracts from. (R. 1866)	214
discovery-F. L. CAPEN. (R. 1866)	214
instruments, description of-L. CASELLA. (R. 1859)	110
material contributed to the Smithsonian Institution. (R. 1860-1866, 1868, 1871, 1874) 147, 149, 150, 187, 188, 209, 214, 224, 228, 244, 249	
memoirs—J. Hann; L. Sohncke; T. Reye; W. Ferrel; A. Cold- ing; M. Peslin; C. Abbe. (R. 1877) 828,	398
publications deposited in Library of Congress, classified list of. (R. 1878.)	275
reports preserved in Smithsonian Institution, classified list of. (R. 1873.)	275
stations, cost of establishment of—J. HENRY. (R. 1858)	109
stations and observers, list of. (R. 1849, 1853–1872.) 21, 67, 75, 77, 91, 107, 110, 147, 149, 150, 157, 187, 188, 209, 214, 215, 224, 228, 244, 249, 271,	•
subjects, short memoirs on, by J. Hann and others. Translated by C. ABBE. (R. 1877)	398
suggestions for Russian America—J. HENRY	207
system for every State—Illinois State Board of Education. (R. 1855.)	77
system of Canada—J. G. Hodgins. (R. 1865)	
system of Smithsonian Institution, report on—E. FOREMAN. (R. 1851;	
R. 1852) 5	1, 57
tables—A. Guyor	153
telegrams by Western Union Telegraph Company—H. SIBLEY. (R. 1862)	150
Meteorological observations—	
at Brunswick, Maine—P. CLEAVELAND.	204
at Marietta, Ohio-S. P. HILDRETH; J. WOOD	120

Meteorological observations—Continued.	
at Philadelphia, Girard College, discussion of—A. D. BACHE. 113, 121,	
132, 162, 175, 186,	195
at Providence, Rhode Island, 1881-1860—A. CASWELL	103
at Providence, Rhode Island, 1881-1876-A. CASWELL	443
at Sacramento, California—F. W. HATCH. (R. 1854)	75
at Sacramento, California—T. M. LOGAN. (R. 1854)	75
at San Francisco, California—H. GIBBONS. (R. 1854)	75
at Washington, Arkansas—N. D. SMITH	131
combined, report of American Association for Advancement of Science on system of. (R. 1851)	
directions for—A. GUYOT; J. HENRY. (R. 1855) 77, 19	
in the Arctic Seas—I. I. HAYES	
in the Arctic Seas—E. K. KANE	
in the Arctic Seas-F. L. McCLINTOCK	
results of, from 1854 to 1859. Vols. I, II	
Smithsonian, for the year 1855	
Meteorology-	
articles on. (R. 1867; R. 1870) 215	j, 244
articles on, with notes by J. Henry. (R. 1871)	
communication relative to—J. P. Espy. (R. 1847)	
lectures and notes on-R. RUSSELL; J. HENRY. (R. 1854)	75
of Caracas, South America-G. A. Ernst. (R. 1867)	
of Colonia Tovar, Venezuela—A. FENDLER. (R. 1857)	
of District of Columbia—J. WISSNER. (R. 1857)	
of Green river country, Utah—Col. Collins. (R. 1871)	
of Porto Rico-G. LATIMER. (R. 1871)	
of Russia—A. Workkoff. (R. 1872)	
of Sacramento, California—T. M. Logan. (R. 1857)	
of the United States, report on—E. Loomis. (R. 1847)	
Meteorology-See Atmospheric pressure, Temperature, Rain, Winds, etc.	
Meteors—	
articles on, preserved in Smithsonian. (R. 1874)	286
observation of—L. Andrews. (R. 1866)	214
of November 18, 1867	217
planisphere of the heavens for observations of	359
Method of—	
ascertaining amount of water in rivers—A. A. HUMPHREYS. (R. 1858.)	IU
forming small weights—R. HARE. (R. 1858)	109
preserving lepidoptera-T. R. PEALE. (R. 1863)	187
Methods of—	30 7
capture and utilization of animals—G. B. Goods	الان 194
installation in the National Museum—G. B. Goode. (P. 1881) 467,	7/0

Methods of—Continued.
interpolation applicable to graduation of irregular series—E. L. Dz
Forest. (R. 1871) 249
interpolation, additions to memoir on—E. L. DE FOREST. (R. 1878) 276
making and preserving plaster casts—A. PIRZ. (P. 1881) 46
Metric system for scientific observation—A. GUYOT. (R. 1848)
Metric system of weights and measures with tables—H. A. NEWTON. (R. 1865.) 209, 371
Metric system with 8 as base—G. H. KNIGHT. (R. 1867) 216
METTENIUS, G. Acknowledgment for ferns. (R. 1862)
Mexican axolotl, change of, to an amblystoma—A. Weismann. (R. 1877.) 823, 401
Mexican boundary, forests and trees of Florida and the-J. G. Cooper. (R. 1860.) 14
Mexican history and archeology, observations on—B. MAYER
Mexican Society of Geography and Statistics. Exchanges. (R. 1861; R. 1865.) 149, 200
Mexico-
description of large fossil gasteropod from—C. A. WHITE. (P. 1880) 428
description of new fishes from -D. S. JORDAN; U. H. GILBERT. (P. 1881.) 467
description of new species of genus Brevoortia from—G. B. Goode. (P. 1878)
eastern, earthquake in, January, 1866—C. SARTORIUS. (R. 1866) 214
explorations of John Xantus in—M. Romero. (R. 1862) 150
Guanajuato and Chapala lake, notes on Dugès' collection of fishes from— D. S. JORDAN. (P. 1879) 833
list of described birds of, not in Smithsonian collection 185
magnetic observations in—Baron von Müller; A. Sonntag 114
meteorite in—R. SIMSON. (R. 1867)
meteorite in—A. Woodworth. (R. 1867)216
meteorite in, discovery of-W. M. Pierson. (R. 1878) 276
meteorites in—J. H. CARLETON. (R. 1865) 200
notes on fishes from, collected by Lieut. H. E. Nichols—D. S. JORDAN; C. H. GILBERT. (P. 1881)467
scientific expedition to, report addressed to Emperor of France by Min- ister of Public Instruction. (R. 1864)
southwestern, birds of, collected by F. E. Sumichrast—G. N. LAWRENCE. 298
Vera Cruz, account of antiquities in-H. FINCK. (R. 1870) 244
Mexico, Gulf of, catalogue of fishes collected in, by J. W. Velie-G. B. Goode; T. H. Bean. (P. 1879)
Mexico, Gulf of, mortality of fishes in-J. P. JEFFERSON. (P. 1878) 382
Mica beds in Alabama—W. GESNER. (R. 1879)
Mica mines, ancient, in North Carolina—C. D. SMITH. (R. 1876) 299
Michigan-
ancient mining in—C. WHITTLESEY
Beaver Island, natural history of—J. J. STRANG. (R. 1854) 78

Michigan—Continued.
catalogue of rocks, minerals, and ores collected on geological survey in— C. T. Jackson. (R. 1854)
characteristics pertaining to ancient man in—H. GILLMAN. (R. 1875.) 298, 393
Isle Royale, Lake Superior, antiquities of-A. C. DAVIS. (R. 1874) 28
mound-builders and platycnemism in—H. GILLMAN. (R. 1873) 275, 393
Microscope, the. (R. 1860)14
Microscopic organisms—
directions for collecting6
directions for collecting, preserving, and transporting—A. M. Edwards. 360
notes on new species and localities of-J. W. BAILEY
Microscopical examination of Japanese infusorial earths -A. M. Edwards 20
Microscopical examination of soundings made by U.S. Coast Survey off At-
lantic coast of United States—J. W. BAILEY
Microscopical observations in South Carolina, Georgia, and Florida—J. W. BAILEY
Middle America—
birds of—S. F. BAIRD
birds of, circular for collecting 16
birds of, not in National Museum—R. RIDGWAY. (P. 1881) 46
(explanation of term)
Migration, the American—F. von Hellwald. (R. 1866)
Migrations and nesting habits of west coast birds—J. G. Cooper. (P. 1879) 33
Milan Agricultural Association. Exchange of publications. (R. 1863) 18
Milky way, bibliography of works relating to—E. S. Holden 31
MILLER, F. Mound in Trumbull county, Ohio. (R. 1877) 32
Mills, Clark, casts of heads of Indian boys and girls at Hampton, Virginia, taken by. (P. 1879)
Mills, Clark, casts of heads of Indian prisoners in Florida taken by. (P. 1878.) 33
Mineral, jade, study of—S. BLONDEL. (R. 1876)
Mineralogical collection, arrangement of—C. U. Shepard. (R. 1861) 149
Mineralogical collection of Yale College, catalogue of meteorites in the—G. J. Brush. (R. 1868)
Mineralogical composition of the normal mesozoic diabase upon Atlantic border—G. W. HAWES. (P. 1881)
Mineralogy, progress in, in 1879 and 1880—G. W. HAWES. (R. 1880) 442, 428
Minerals—
and ores collected on geological survey in Michigan, catalogue of—C. T. JACKSON. (R. 1854)
and rocks, catalogue of—J. W. FOSTER. (R. 1854)
and rocks, catalogue of—J. D. WHITNEY. (R. 1854)
catalogue of, with their formulas, etc.—T. EGLESTON156
in U.S. National Museum in 1878, list of-F. M. Experien. (R. 1878.) 275

Minerals—Continued.
in U. S. National Museum in 1879, list of-F. M. Endlich. (P. 1880.) 425
rocks, ores, and fossils, catalogue of—J. Locke. (R. 1854) 75
Mines, ancient mica, in North Carolina—C. D. SMITH. (R. 1876) 299
Mining, ancient, in Lake Superior copper region, circular relative to—J. HENRY. (R. 1861)
Mining, ancient, on shores of Lake Superior-C. WHITTLESEY 155
Mining Department, Melbourne. Exchange system. (R. 1865) 209
Minnesota—
ancient town in, account of-O. H. KELLEY. (R. 1868)
Fort Ripley, natural history of country about—J. E. HEAD. (R. 1854.) 75
Hennepin county, mounds on Gideon's farm near Excelsior—F. H. NUT- TER. (R. 1879)
Lake Pepin and Mississippi river, antiquities on banks of—L. C. Estes. (R. 1866)
Red river of the North, ethnology of Indians of—W. H. GARDNER. (R. 1870)
Minnesota Historical Society. Dakota grammar
Miocene fossils, check list of—F. B. MEEK
Miscellanea, Museum, labels, etc.—S. F. BAIRD
Miscellaneous Collections, Smithsonian. Vols. 1-xxIII. 122, 123, 124, 125, 158, 169,
191, 212, 218, 250, 278, 274, 812, 814, 315, 822, 886, 837, 416, 428, 424, 468, 475
Miscellaneous Collections, Smithsonian, catalogue and index of 478
Miscellaneous tables—A. Guyot 153
Missionary Society, Southern. Yoruba grammar and dictionary 98
Missions—See American Board of Commissioners.
Mississippi—
aboriginal lapidary in—C. RAU. (R. 1877)
eastern, collection of fishes from—O. P. HAY. (P. 1880)
mounds in—S. A. Agnew. (R. 1867) 215
Washington county, mounds in—J. Hough. (R. 1879) 345
Yazoo county, antiquities of—J. W. C. SMITH. (R. 1874) 286
Mississippi river and Lake Pepin, antiquities on banks of—L. C. Estes. (R. 1866)
Mississippi river, description of new species of fishes from—T. H. BEAN. (P. 1879)
Mississippi valley, ancient monuments of—E. G. SQUIER; E. H. DAVIS 1
Mississippi valley, physical geography of—C. ELLET, Jr 13
Missouri—
ancient relics in—J. W. FOSTER. (R. 1868) 187
antiquities in—I. DILLE. (R. 1862)
Kansas City, antiquities of—W. H. R. LYKINS. (R. 1877) 828
New Madrid, earthquake at—T. DUDLEY. (R. 1858)

ALPHABETICAL INDEX

Missouri-Continued.
prehistoric evidences in-G. C. BROADHEAD. (R. 1879) 345
St. Louis, ancient mound at—T. R. PEALE. (R. 1861)
St. Louis, temperature of—A. FENDLER. (R. 1860)
upper, expedition to Mauvaises Terres and—A. Culbratson. (R. 1850.)
upper, list of plants of—T. C. Porter. (R. 1850)
upper, palseontology of—F. B. MEEK; F. V. HAYDEN
western, exploration of-P. R. Hov. (R. 1864)
Missouri river, ancient earthworks on—A. BARRANDT. (R. 1870)
Missouri river, list of birds and mammalia of—E. HARRIS. (R. 1850) 2
MITCHELL, A. Antiquities of Florida. (R. 1874) 28
MITCHELL, B. Mounds in Pike county, Illinois. (R. 1879)
MITCHELL, B. R.; TURNER, W. W. Vocabulary of the jargon of Oregon 6
MITCHELL, S. S. Funeral sermon of Joseph Henry
MITCHELL, S. W. Inquiries relative to disease known as chorea. (R. 1874) 28
MITCHELL, S. W. Researches upon the venom of the rattlesnake135
MITCHELL, S. W.; MOREHOUSE, G. R. Researches upon anatomy and physi-
ology of respiration in chelonia159
Mitla, drawings of—J. G. SAWKINS
Mixed races in Liberia—E. D. BLYDEN. (R. 1870)
Mobile river, shell-heaps on—A. S. GAINES; K. M. CUNNINGHAM. (R. 1877.)
Modern theory of chemical types—C. M. WETHERILL. (R. 1863)
Modes of flight in relation to aeronautics—J. B. Pettigrew. (R. 1867) 215
Moigno, F. N. Accidental or subjective colors. (R. 1866)214
Molina, L., Costa Rican minister, recommending Dr. Berendt. (R. 1865) 20
Mollusca of Northeastern coast of America. Parts I, II, III—A. E. VERBILL. (P. 1879; P. 1880)
Mollusca, or shellfish, and their allies, lectures on—P. P. CARPENTEE. (R. 1860)
Molluscoids of Arctic America—A. E. VERBILL
Mollusks-
arrangement of families of—T. GILL 227
bibliography of—T. GILL
eocene, some new species of, from southern United States—A. HEILPRIS. (P. 1880)
new forms of, from Alaska—W. H. Dall. (P. 1878)
of Arctic America—W. H. Dall
of Kerguelen Island-W. H. DALL
of western North America—P. P. CARPENTER252
Monadrock, ironclad, deviation of compasses on-W. HARKNESS
Mongolia, geological researches in—R. PUMPELLY
Monograph of—
American corbiculadæ, recent and fossil—T. PRIME145

•	
Monograph of—Continued.	
bats of North America—H. Allen	.65
cottoids of North America—C. GIRARD	30
dolichopodidæ—H. Loew	171
ortalidæ—H. Loew	256
strepomatidæ, (American melanians)—G. W. TRYON 2	53
tipulidæ—C. R. Osten Sacken	219
wasps-H. Dr Saussure 2	54
Monographs of diptera of North America—	
Part I-H. LOEW; R. OSTEN SACKEN 1	.41
Part II-H. LOEW; R. OSTEN SACKEN	.71
Part III—H. Loew	!56
Part IV-R. OSTEN SACKEN2	19
Montana, distribution of forest trees in-W. W. Johnson. (R. 1870)	244
Montana, prehistoric remains in—P. W. Norris. (R. 1879)	845
Monterey Bay, California, new fishes from—D. S. JORDAN; C. H. GILBERT. (P. 1880)	42 5
	856
Monuments—	
aboriginal, of State of New York-E. G. SQUIER.	15
ancient, of the Mississippi valley—E. G. SQUIER; E. H. DAVIS	1
and antiquities of Denmark, preservation of J. A. Worsaae. (R. 1879.)	34 5
Moon-	
effect of, on the magnetic force—A. D. BACHE	175
effect of, on the weather—J. HENRY. (R. 1871)	249
influence of, on magnetic declination—A. D. BACHE	182
occultations of planets by—See Downes, J.	
Moon-hoax, on the—J. HENRY. (R. 1873)	275
MOORE, M. A. Fish mortality in the Gulf of Mexico. (P. 1881)	467
MOORE, T., and others. Destruction of fish near the Tortugas. (P. 1878)	38 2
Morehouse, G. W.; Mitchell, S. W. Anatomy and physiology of respira-	
	L59
Мовгіт, С; Воотн, J. C. Recent improvements in chemical arts	27
Morgan, L. H.—	
Circular in reference to degrees of relationship among different nations 1	138
Proposed ethnological map of North America. (R. 1861)	149
Systems of consanguinity and affinity of the human family2	218
Morgan, L. H., circular to diplomatic agents by State Department relative to research of	188
MORIN, A. Warming and ventilating buildings. Parts 1, 11. (R. 1878; R. 1874) 275, 286, 4	13 9
Morlot, A.—	_ _
Abstract of Dr. Keller's report on lacustrian settlements in Switzerland.	
(R. 1868)	187

MORLOT, A.—Continued. Archæology. (R. 1861)	140
General views on archæology. (R. 1860)	
Morris, J. G.—	U , 100
Catalogue of described lepidoptera of North America	118
Lecture on insect instincts and transformations. (R. 1855)	
Lecture on natural history as applied to farming and gardening. (R. 1855.)	
Synopsis of described lepidoptera	
MORRIS, O. W.; HENRY, J. Quantity of rain at different heights. (R. 1855.)	
Morse, S. F. B., account of origin and development of invention of—W. B.	
TAYLOR. (R. 1878)	
Morse, S. F. B., communication relative to a publication of—J. HENRY. (R. 1857)	
Mortality—	
of fishes in Gulf of Mexico—E. INGERSOLL. (P. 1881)	467
of fishes in Gulf of Mexico—J. P. JEFFERSON. (P. 1878)	
of fishes in Gulf of Mexico—S. H. Johnson. (P. 1881)	
of fishes in Gulf of Mexico-M. A. Moore. (P. 1881)	
of fishes in vicinity of Tortugas—J. P. JEFFERSON; T. MOORE; J. Y. PORTER. (P. 1878)	
tables of, methods of interpolation applied to graduation of—E. L. Dr	
FOREST. (R. 1871; R. 1878): 249	
MORTILLET, G. DE; CHANTRE, E. International code of symbols for charts of prehistoric archæology. (R. 1875)	
MORTON, S. G., and others. On publication of Squier and Davis' ancient mon-	
uments. (R. 1847)	
Mortuary customs of North American Indians—H. C. YARROW. (E. 1879-80.)	
Mosasaurus and the three allied genera, Holcodus, Conosaurus, and Amphoros-	
teus, memoir on—R. W. GIBBES	14
Motion, planetary, general integrals of—S. NEWCOMB	281
Motion, rotary, problems of, presented by gyroscope, precession of equinoxes,	242
and pendulum—J. G. BARNARD	
Motschulsky, V. On means of destroying the grasshopper. (R. 1858)	
Moulton, M. W. Mounds in Delaware county, Iowa. (R. 1877)	323
Mound in—	971
Dakota, Lincoln county—A. BARRANDT. (R. 1872)	293
Florida, colored bead from—A. M. HARRISON. (R. 1877)	
Florida, gold ornament from—C. RAU. (R. 1877) 823, 440,	
Indiana—W. Pidgeon. (R. 1867)	
Kentucky, near Lexington—R. Peter. (R. 1871)	149
Missoull, No. 10015—1. 16. 1 BRDB. (16. 1001)	323
Ohio, Trumbull county—F. MILLER. (R. 1877)	323
Pennsylvania, western—W. M. TAYLOR. (R. 1877)	الهي

Mound in-Continued.
Tennessee, near Chattanooga-M. C. READ. (B. 1867)
Tennessee, east—A. F. Danilsen. (R. 1868)
Wisconsin-C. K. DEAN. (R. 1872)
Wisconsin, Grant county-J. WARNER. (R. 1872)
Mound-builders—
and platycnemism in Michigan—H. GILLMAN. (R. 1878) 275,
in Rock river valley, Illinois—J. SHAW. (R. 1877)
of Wisconsin, mounds and osteology of-J. N. DE HART. (R. 1877)
Mounds and osteology of the mound-builders of Wisconsin—J. N. DE HART. (R. 1877)
Mounds in-
Dakota, Fort Wadsworth-A. J. COMFORT. (R. 1871)
Florida, (southern,) explorations among—S. T. WALKER. (R. 1879)
Georgia-M. F. Stephenson. (R. 1870)
Georgia, Bartow county-M. F. STEPHENSON. (R. 1872)
Georgia, McIntosh and Early counties—W. McKinley. (R. 1872)
Illinois, Mercer county—T. McWhorter. (R. 1874)
Illinois, near Albany, skull and bones from—R. J. FARQUHARSON. (R. 1874)
Illinois, Pike county—B. MITCHELL. (B. 1879)
Illinois, Rock Island county—T. Thompson. (R. 1879)
Illinois, Spoon river valley—W. H. Adams. (R. 1879)
Illinois, Union county, near Anna-T. M. PERRINE. (R. 1872)
Indiana, Franklin county—E. R. Quick. (R. 1879)
Indiana, Rush county—F. JACKMAN. (R. 1879)
Iowa, Delaware county-M. W. Moulton. (B. 1877)
Iowa, Des Moines valley, notes on—S. B. Evans. (R. 1879)
Iowa, Muscatine county-T. Thompson. (R. 1879)
Kentucky—R. PETER. (R. 1872)
Kentucky, exploration of—S. S. Lyon. (R. 1870)
Kentucky, Union county, exploration of—S. S. Lyon. (R. 1870)
Louisiana—S. H. Lockett. (R. 1872)
Louisiana, Moorehouse parish—B. H. Brodnax. (R. 1879)
Minnesota, Hennepin county, on Gideon's farm—F. H. NUTTER. (R. 1879)
Mississippi—S. A. AGNEW. (R. 1867)
Mississippi, Washington county—J. Hough. (R. 1879)
New Brunswick and New England, notes on—S. F. BAIRD. (P. 1881.)
Ohio, Trumbull county—F. MILLER. (R. 1877)
Pennsylvania, (western)—W. M. TAYLOR. (R. 1877)
Tennessee-J. Jones
Wisconsin, Grant county—M. STRONG. (R. 1879)
Wisconsin, Winnebago county—T. Armstrong. (R. 1879)
TI LOUVILLE IT ILLIUUNEU UVULLE "TALALALALALALALA "TALAUULLI IT LULUULLE IT LULUULLE IT LULUULLE IT LULUULLE I

Mountain, Black, North Carolina, topography of—T. L. OLINGMAN. (R. 1866.)	77
Mountain measurements—A. Guyor. (R. 1862)	150
Mountains, electric resonance of—H. DE SAUSSURE. (R. 1868)	24
Mountains in Colorado, heights of-G. Engelmann. (R. 1862)	150
Mountains, lakes, and the snow-line, Norwegian—O. E. DREUTZER. (R. 1866.)	214
Movement of the stars around a central point—J. H. MAEDLER. (R. 1859)	110
MUCH, M. Ancient history of North America. (R. 1871)	249
MUDGE, B. F. Account of lightning discharges. (R. 1867)	215
MÜLLER, Baron von. Observations on terrestrial magnetism in Mexico	114
MÜLLER, F. Smithsonian exchanges. (R. 1860)	147
MÜLLER, J.—	
Principles of the mechanical theory of heat. (R. 1868)	224
Report on recent progress in physics—electricity. (R. 1856; R. 1857.) 91,	, 107
Report on recent progress in physics—galvanism. (R. 1855; R. 1857; R. 1858) 77, 107,	, 109
Report on transactions of Geneva Society of Physics and Natural History, June, 1875, to July, 1876. (R. 1877)	
Mummy and skulls from Patagonia—A. REID. (R. 1862)	
	208
MURPHEY, H. C., and others. On publication of Spanish works on New Mexico.	
(R. 1855)	11
Musci of Kerguelen Island—T. P. JAMES.	294
Museum—	• 40
at Lausanne, ethnological collections of—F. TROYON. (R. 1861)	
Blackmore, Salisbury, England, notice of. (R. 1868)	
British, statistics of—S. F. BAIRD. (R. 1850)	25
d'Histoire Naturelle, Paris, notes on typical American fishes in—D. S. JORDAN. (P. 1879)	
Hungarian National. Acknowledgment for birds. (R. 1868)	187
miscellanea—S. F. BAIRD	164
of ethnology, Leipsic-A. Schott; O. T. Mason. (R. 1878)	275
of National University of Greece. Exchange of specimens. (B. 1867.)	215
of Norway, Ethnological-L. K. DAA. (R. 1862)	
of Toronto University, objects of. (R. 1865)	
on formation of a-L. Agassiz. (R. 1849)	21
Smithsonian—See Museum, U.S. National.	
Museum, U. S. National—	
acts and resolutions of Congress relative to—See Congress.	
additions to, list of—See each annual report.	
birds of Middle and South America not in—R. RIDGWAY. (P. 1881)	457
Bulletins of—See Bulletins.	
catalogue and index of the Proceedings and Bulletins of	
catalogue of collection of Japanese cotton fibre presented to. (P. 1881.)	467

Museum, U. S. National—Continued.
catalogue of collection of Japanese woods presented to—L. F. WARD. (P. 1881) 467
catalogue of Old World birds in-R. RIDGWAY. (P. 1881) 467, 462
circular asking contributions to library of. (P. 1881) 467, 458
circulars of—See Circulars.
classification of collections of—G. B. Goode. (P. 1881) 467, 457
classification of collections of, to illustrate taxidermy—W. T. Hornaday. (P. 1881) 467, 456
collections furnished to, by explorations, 1888 to 1877—S. F. BAIRD. (R. 1877) 823
contributions to, and their acknowledgment. (P. 1881) 467, 473
directory of officers, collaborators, and employés of 466
distribution by, of marine invertebrates—R. RATHBUN. (P. 1881.) 467, 465, 471
establishment and officers of Smithsanian Institution and. (P. 1881.) 467, 449
letters on work of—B. PHILLIPS. (P. 1881)
list of publications of. (P. 1881) 467, 478, 474
memorial of Regents of Smithsonian Institution to Congress asking appropriations for. (B. 1867)215, 329
memorial to Congress in behalf of Regents of Smithsonian Institution
for new building for-G. BANCROFT. (R. 1876) 299, 829
organization and objects of—G. B. Goode. (P. 1881) 467, 459
outline of scheme of classification for collections in—G. B. Goods. (P., 1881)
plan of organization and regulations of—G. B. Goode. (P. 1881) 467, 445
plans for installation of collections in—G. B. Goode. (P. 1881) 467, 472 proceedings in Congress relative to 328
Proceedings of. Vols. 1-1v, 1878-1881 332, 333, 425, 467
report of Committee of Regents on-A. GRAY; H. COPPÉE. (R. 1874.) 286
report of Committee of Regents on—A. GRAY; A. A. SARGENT; H. CLYMER. (R. 1876)
report on, and statistics of British Museum—S. F. BAIRD. (R. 1850) 28
reports of Building Commission of, for 1879, 1880. (R. 1879; R. 1880.) 845, 442, 409, 434
reports on, 1858-1867, 1878-1877-S. F. BAIRD. (R. 1858-1867, 1878-1877) 67, 75, 77, 91, 107, 109, 110, 147, 149, 150, 187, 188, 209, 214, 215, 275, 286, 298, 299, 328
Myadestes obscurus, description of two new races of—L. STEJNEGER. (P. 1881.) 467
Myctophum crenulare, description of—T. H. BEAN. (P. 1881) 467
Myctophum crenulare from Santa Barbara channel, California, description of—
D. S. JORDAN; C. H. GILBERT. (P. 1880)
Myological specimens, rapid preparation of—F. PLATEAU. (P. 1881) 467

Myriapods, list of North American species of, belonging to family of Lysiope- talidæ—J. A. RYDER. (P. 1880)	425
Myriapods, phalangidse, etc., instructions for collecting—H. C. Wood. (R. 1866)	214
Myriolepis zonifer, a new chiroid fish from Monterey Bay, Calfornia, description of—W. N. Lockington. (P. 1880)	425
Mythology of North American Indians-J. W. Powell. (E. 1879-80)	476
Myths, Indian. (E. 1879-80)	476

M.

NAILL, D. W. Dispersion of a cloud by electrical discharge. (R. 1858) !	(F)
Naples, Italy, Royal Institute for Encouragement of Natural, Economical, and	75
Technological Sciences. Prize questions. (R. 1878)	
Narrative of the Hassler expedition—L. Agassiz. (R. 1872) 271, 8	
National Academy of Sciences, bequest to—A. D. BACHE. (R. 1872) 271, \$	3
National library, on a-W. S. JEVONS. (B. 1878) 2	.75
National Museum—See Museum, U. S. National.	
Natural history—	
applied to farming and gardening, lecture on—J. G. Morris. (R. 1855.)	77
bibliography of American—C. GIBARD	18
directions for collecting, preserving, and transporting specimens of-	
S. F. BAIRD. (R. 1856)	34
explorations in, in the United States, 1851—S. F. BAIRD. (R. 1851)	51
of Arctic America—L. Kumlien	12
of Beaver Island, Michigan-J. J. STRANG. (R. 1854)	75
of country about Fort Ripley, Minnesota-J. E. HEAD. (R. 1854)	
of Fanning Islands—T. H. STREETS	
of fresh-water fishes of North America-C. GIRARD	30
of Hawaiian Islands, etc.—T. H. STREETS	
of Kerguelen Island-J. H. KIDDER and others 29	
of lower California—T. H. STREETS	03
of organized bodies-E. J. MAREY. (R. 1867) 2	
of Pacific Islands-W. H. PEASE. (R. 1862)	
suggestions relative to investigations in, in Russian America—S. F. BAIRD	
Natural History Society—See Geneva.	
Natural method, the Jussieus and the-M. FLOURENS. (R. 1867)	15
Naturalists, generic names of animals employed by-S. H. Scupper 47	

and art, on tables of the constants of—C. Babbage. (R. 1856)	32
and origin of force—W. B. TAYLOB. (R. 1870) 244, 87 of reparatory inflammation in arteries after ligature. Toner lecture No. VII—E. O. SHAKESPKARE 32 Nature—See Constants of Nature. Navajo Indians, language of, said to resemble Welsh—S. Y. McMasters. (R.	
of reparatory inflammation in arteries after ligature. Toner lecture No. VII—E. O. SHAKESPKARE	75
VII—E. O. SHAKESPKARE	
Navajo Indians, language of, said to resemble Welsh-S. Y. McMasters. (R.	:1
)9
	77
Naval officers, authority to, to receive and transmit specimens to Smithsonian 8	34
Navigation of Ohio and other rivers, improvement of-C. ELLET 13	3
Navy Department, authority given by, to naval officers to receive and transmit specimens to the Smithsonian Institution8	3 4
Neah Bay, Washington Territory, description of two new species of fishes from—D. S. JORDAN; C. H. GILBERT. (P. 1880) 42	25
Nebraska-	
ancient fauna of-J. LEIDY5	8
palsontology of-F. B. MEEK; F. V. HAYDEN	2
report on fossils from—J. LEIDY. (R. 1851)	51
Nebulæ, index-catalogue of books relative to—E. S. Holden 31	.1
Nebulæ, researches relative to the—Prof. GAUTIER. (R. 1863) 18	37
Nebular hypothesis, bibliography of the—E. S. HOLDEN	11
Nematoda of northeast coast of North America-A. E. VERRILL. (P. 1879) 88	38
Nemertina of northeast coast of America—A. E. VERRILL. (P. 1879) 88	38
Nemichthys avocetta, new species of Nemichthys, description of—D. S. Jon- DAN; C. H. GILBERT. (P. 1880)	25
Neptune—	
ephemerides of, 1848, 1846-49, 1850, 1851, 1852-S. C. WALKER. 4, 5, 6, 7, 2	4
history of discovery of—B. A. Gould1	8.
investigation of orbit of, with tables of its motion—S. NEWCOMB 19	9
researches relative to—S. C. WALKER	3
secular variations of elements of orbit of-J. N. STOCKWELL 23:	2
Nereis-Boreali-Americana—W. H. HARVEY: Part I. Melanospermeæ	2
	:3
Part III. Chlorospermece 9	5
•	6
Neritidæ, fresh-water—W. G. BINNEY	
Nerve, polarized, effect of irritation of—B. F. LAUTENBACH. (B. 1878) 841, 41.	
Nervous system of Rana pipiens, anatomy of—J. WYMAN 4	
Nests—	
and eggs of American birds, instructions for collecting—S. F. BAIRD. (R. 1858) 109, 139	9

Nests-Continued.	
and eggs of the eight North American species of empidonaces—T. M. Brewer. (P. 1879)	333
instructions for collecting and preserving-T. M. BREWEB	
of west coast birds-J. G. Cooper. (P. 1879)	33 3
Netherlands, the, Royal Academy of. Prize questions. (R. 1861)	149
Neuchâtel, lake of, palafittes or lacustrian constructions of—E. Desor. (R. 1865)	360
Neuroptera—	
instructions for collecting—P. R. UHLER. (R. 1858)	109
North American, synopsis of—H. HAGEN. (R. 1861) 149,	134
of Arctic America—S. H. SCUDDER	342
of North America, synopsis of—H. HAGEN	134
of North America, synopsis of—P. R. UHLER	134
pseudo-, of Kerguelen Island-H. A. HAGEN	294
Nevada, western, Centennial mission to Indians of—S. Powers. (R. 1876) New—	299
classification of clouds—A. Poëv. (R. 1870)	244
species and localities of microscopical organisms, notes on—J. W. BAILEY.	
species of North American coleoptera. Parts 1, 11-J. L. LE CONTE. 167,	
system of weights and measures, with 8 as metrical number—G. H. Knight. (R. 1867)	
tables for determining values of coefficients in perturbative function of planetary motion—J. D. RUNKLE	
New Brunswick— Bay of Fundy, marine invertebrata of region about mouth of—W. STIMP-	
shell heaps in-J. Fowler. (R. 1870)	
shell mounds of—S. F. BAIRD. (P. 1881)	
New England— crustacea dredged off south coast of—S. I. Smith. (P. 1880)	
fishes from south coast of, descriptions of new genera and species of—G. B. Goods. (P. 1880)	
frigate mackerel (Auxis Rochei) on coast of-G. B. Goode. (P. 1880)	
geology of—E. Hitchcock	
marine invertebrata from, distributed by U. S. Fish Commission—A. E. VERRILL; R. RATHBUN. (P. 1879)	
marine invertebrates from, distributed by U.S. National Museum-R. RATHBUN. (P. 1881) 467, 465,	•
mollusca added to fauna of-A. E. VERRILL. (P. 1880)	
shell mounds on coast of—S. F. BAIRD. (P. 1881)	1 5;
New Hurmony, Indiana, account of tornado near, April 80, 1852-J. CHAPPEL-	59
NEW HAVEN JOURNAL. Account of lightning discharges. (R. 1867)	

New Jersey-	
artificial shell deposits in—C. RAU. (R. 1864) 188, 440,	362
fishes on the coast of Long Island and—S. F. BAIRD. (R. 1854) 75,	348
stone age in—C. C. Abbott. (R. 1875)	394
New Jersey Historical Society, address of Prof. Henry before the	B
New Madrid, Missouri, earthquake at—T. DUDLEY. (R. 1858)	109
New Mexico-	
account of Lindheimer's, Fendler's, and Wright's botanical explorations	•
in—A. Gray. (R. 1849)	21
antiquities in—W. B. Lyon. (R. 1871)	249
communications relative to publication of Spanish works on—E. Robinson; H. E. Ludewig; E. G. Squier; H. C. Murphry; W. B. Hodgson; W. Irving; W. H. Prescott; J. Sparks; G. Bancroft; F. L. Hawks. (R. 1855)	7 7
description of new cretaceous Pinna from-C. A. WHITE. (P 1880)	425
diary of excursion in-J. H. CARLETON. (R. 1854)	75
eastern, coleoptera of Kansas and-J. L. LE CONTE	126
notes on the history and climate of-T. A. McParlin. (R. 1877) - 828,	396
plants of, collected by C. Wright. Parts I, II-A. GRAY 2	2, 42
report of explorations in Arizona and—J. Stevenson. (R. 1880)	442
New York-	
aboriginal monuments of—E. G. SQUIER	15
Clinton, Humilton College. Exchange of specimens. (R. 1861)	149
fishes of—T. GILL. (R. 1856)	91
Fishkill, deposit of arrow-heads near—E. M. Shepard. (R. 1877)	828
New York, explosion of nitre in, July, 1845-R. HARE	17
Orleans county, antiquities of—F. H. CUSHING. (R. 1874)	286
Schoharie, Indian relics from-F. D. Andrews. (R. 1879)	845
Tompkins county, ancient fort and burial ground in—D. TROWBRIDGE.	
(R. 1863)	187
University of State of, Regents of. Acknowledgment of specimens. (R.	000
Name of St. David die of family along from Objects and beginning	209
NEWBERRY, J. S. Description of fossil plants from Chinese coal-bearing rocks.	202
Newcastle, Pennsylvania, account of old Indian village near—E. M. McCon- NELL. (R. 1871)	249
Newcomb, S.—	
General integrals of planetary motion	
Investigation of orbit of Neptune, with tables of its motion	
Investigation of orbit of Uranus, with tables of its motion	262
Memorial address on Joseph Henry	856
NEWCOMB, S., and others. Scientific instructions to Capt. Hall. (R. 1871)	249
Newfoundland, new species of fishes from Grand Banks of—G. B. Goode. (P. 1880)	425
NEWTON, A. Suggestions for forming collections of birds' eggs	189

ALPHABETICAL INDEX

NEWTON, A. Suggestions for saving parts of skeletons of birds. (R. 1860). NEWTON, H. A. Metric system of weights and measures, with tables. (1	R.
1865)	19, 37
Nicaragua—	•
antiquities of—E. G. SQUIER. (R. 1850)	
archæological researches in—J. F. Bransford	
eruption of volcano in—A. B. DICKINSON. (R. 1867)	21
list of fishes collected by, in Gulf of California and on west coast of low California—D. S. JORDAN; C. H. GILBERT. (P. 1881)	
notes on fishes collected by, in British Columbia and southern Alaska- T. H. BEAN. (P. 1881)	
notes on fishes collected by, on west coast of Mexico—D. S. JORDAN; (H. GILBERT. (P. 1881)	
NICKLÉS, F. J. Scientific Congress of Carlsruhe, 1858. (R. 1860)	_ 14
Nile, explorations of the—C. HALE. (R. 1865)	
Nitre, memoir on explosiveness of—R. HARE	
Nitrification—B. F. CRAIG. (R. 1861)	
Nitrogen bodies of modern chemistry-Prof. KLETZINSKY. (R. 1872)	
Nomenclator zoölogicus. Alphabetical list of generic names employed by na uralists. Part I—S. H. SCUDDER	} -
Nomenclature of— certain North American birds, revisions of—R. RIDGWAY. (P. 1880) - clouds	_ 347
North American birds—R. Ridgway	
Norris, P. W. Prehistoric remains in Montana. (R. 1879)	. 3 4i
North America—	364
abbreviations of names of States and Territories of	
algæ of, fresh-water, history of—H. C. Wood, Jr	. 341
algæ of, marine—See W. H. Harvey.	
ancient aboriginal trade in—C. RAU. (R. 1872) 271, 440	
ancient history of—M. Much. (R. 1871)	, 24 9
atlas of, physical, proposed—G. GIBBS. (R. 1866)	, 211
batrachia and reptiles of, check-list of—E. D. COPE	, 23 4
bats of, monograph of—H. ALLEN	100
birds added by L. Belding to fauna of-R. RIDGWAY. (P. 1881)	9 01
birds of, catalogue of—S. F. BAIRD	TOO
birds of, instructions for collecting nests and eggs of	400
birds of, nomenclature and catalogue of-R. RIDGWAY	422
birds of, review of. Part I-S. F. BAIRD	Ter
botany of, bibliographical index to—S. WATSON	255
coleoptera of, classification of. Parts I, II—J. L. LE CONTE 136	263
coleoptera of, list of-J. L. LE CONTE	140

North America—Continued.	
coleoptera of, new species of. Parts I, II-J. L. LE CONTE 167,	264
combined meteorological system for, report of Committee of American	
Association for Advancement of Science on. (R. 1851)	51
conchology of, bibliography of. Parts I, II-W. G. BINNEY 142,	174
contributions to natural history of fresh-water fishes of. Part I. Monograph of cottoids—U. GIRARD	
diptera of, catalogue of-C. R. Osten Sacken 102,	270
diptera of, monographs of-H. LOEW; R. OSTEN SACKEN. 141, 171, 219	
duck new to fauna of-R. RIDGWAY. (P. 1881)	467
explorations on western coast of-W. H. Dall. (R. 1873)	275
extinct sloth tribe of—J. LEIDY	72
fishes of east coast of, catalogue of-T. GILL	283
fishes of Pacific coast of, duplicates of, distributed in 1881—D. S. Jor-DAN; P. L. JOUY. (P. 1881)	467
forests and trees of, distribution of-J. G. Cooper. (R. 1858) 109,	351
grasshoppers of, circular relative to history of	163
Indians of, uses of brain and marrow of animals among—T. R. PEALE. (R. 1870)	244
lakes of, fluctuations of level of-C. WHITTLESEY	119
lepidoptera of, catalogue of-J. G. Morris	118
lepidoptera of, synopsis of-J. G. Morris	
libraries, public, in, list of-W. J. RHEES	
mammals of, catalogue of—S. F. BAIRD	105
map of, ethnological, suggestions relative to-L. H. MORGAN. (R. 1861.)	149
meteorological stations and observers in. (R. 1868) 224,	373
mollusks of western-P. P. CARPENTER.	
myriapods of, belonging to family Lysiopetalide—J. A. RYDER. (P. 1880)	
neuroptera of, synopsis of-H. HAGEN	134
orthoptera of, catalogue of-S. H SCUDDER	
rain and snow fall of, tables of-C. A. SCHOTT 222,	
reptiles of, catalogue of—S. F. BAIRD; C. GIRARD	
shells of, catalogue of—I. Lea; P. P. Carpenter; W. Stimpson; W. G. Binney; T. Prime	
shells of, circular in reference to collecting	176
shells of, land and fresh-water. Parts I-IV-W. G. BINNEY; T. BLAND. 143, 144, 194,	
siluride of, synopsis of—D. S. Jordan	
stone implements of—C. RAU. (R. 1872) 271,	
North American—	
archeology-J. Lubbock. (R. 1862)	150
hirds catalogue of S F BAIRD	108

North American—Continued.	
birds, catalogue of-R. RIDGWAY. (P. 1880) 425, 4	2:
birds, desiderata among-R. RIDGWAY. (P. 1881)	461
Continent, physical geography of-J. FROEBEL. (R. 1854)	71
fishes, duplicates of, distributed by Smithsonian Institution—T. H. Bran. (P. 1880)	
Indians, catalogue of portraits of—J. M. STANLEY	
Indians, list of photographic portraits of 2	
stone implements—C. RAU. (R. 1872)	
stone period, agricultural implements of—C. RAU. (R. 1868) 187, 3	
tortoises, land, of genus Xerobates-F. W. TRUE. (P. 1881)	
North American Ichthyology, contributions to— Part 1. Review Rafinesque's memoirs—D. S. Jordan	
Part II. A. Notes on Cottidæ, (etc.) B. Synopsis Siluridæ—D. S. JORDAN	
Part III. A. Distribution of fishes of Alleghany region, (etc.)—D.S. JORDAN; A. W. BRAYTON. B. Synopsis of family Cutostomide—D. S. Jordan 3	IOE
North American Oölogy. Part I. Raptores and fissirostres-T. M. BREWER.	89
North Carolina-	
ancient mica mines in—C. D. SMITH. (R. 1876)	20%
Beaufort, description of new species of fish (Caranx Beani) from—D. S. JORDAN. (P. 1880)	42 5
Beaufort, notes on fishes of-D. S. JORDAN; C. H. GILBERT. (P. 1878.)	83:
Black Mountain, topography of-T. L. CLINGMAN. (R 1855)	T
earthquakes in, in 1874—W. Du Pré; J. Henry. (R. 1874)	280
Lenoir county, antiquities in J. M. SPAINHOUR. (R. 1871)	249
Montgomery county, antiquities of-F. J. KRON. (R. 1874)	286
occurrence of Physis regius in-T. H. BEAN. (P. 1880)	12
Stanley county, antiquities of-F. J. KRON. (R. 1874)	
North German Lloyd, free freight between Germany and United States by- R. SCHLEIDEN. (R. 1858)	
North Pacific Surveying Expedition, contributions to natural history made in connection with—T. H. STREETS	
North Pole, expedition toward the, scientific instructions to—J. HENRY; J. B. HILGARD; S. NEWCOMB; S. F. BAIRD; F. B. MEEK; L. AGASSIZ. (R. 1871)	149
North Pole, map of stars near the, for observations on the aurora	50
Northern Hemisphere, winds of the-J. H. Coffin	52
Northern latitudes, record of auroral phenomena observed in-P. Force	B4
	23
Northumberland, Duke of, account of the	30
Northumberland, Duke of, presentation of books by. (R. 1859) 1	10

Northwest coast of America, kjökken-möddings on the—P. SCHUMACHER. (R. 1878)	275
Norton, E. On wasps	254
Norway, Ethnological Museum of-L. K. DAA. (R. 1862)	150
Norway, University of Christiania, ethnological specimens from. (R. 1868)	187
Norwegian mountains, lakes, and the snow-line—O. E. DREUTZER. (R. 1866.)	214
Notacanthus phasganorus from Newfoundland—G. B. Goode. (P. 1880)	425
Note on occurrence of Phycis regius in South Carolina—T. H. BEAN. (P. 1880.)	425
Note on the latiloid genera—T. GILL. (P. 1881)	467
Notes—	
of egging expedition to Shoal Lake, Lake Winnipeg-D. Gunn. (R. 1867)	215
on aboriginal shell mounds of New Brunswick and New England—S. F. BAIRD. (P. 1881)	467
on American migration—F. von Hellwald. (R. 1866)	214
on fishes—See D. S. Jordan; C. H. Gilbert; also, Proceedings National Museum.	
on fishes of Pacific coast of United States-D. S. JORDAN; C. H. GIL-	
винт. (Р. 1881)	467
on history and climate of New Mexico-T. A. McParlin. (R. 1877.) 828,	396
on horary variations of barometer—M. VAILLANT; J. HENRY. (R. 1866.)	214
on Indian history-F. V. HAYDEN. (R. 1867)	215
on life and character of Joseph Henry—J. C. WELLING 856,	338
on mortality among fishes of Gulf of Mexico—S. H. Johnson. (P. 1881.)	467
on new species and localities of microscopical organisms—J. W. BAILEY.	63
on physical geography of North America—J. G. Cooper. (R. 1858.) 109,	351
on Salmonidæ of upper Columbia—C. BENDIRE. (P. 1881)	467
on some Costa Rican birds—R. RIDGWAY. (P. 1881)	467
on Tinneh or Chepewyan Indians of British and Russian America—G.	
GIBBS and others. (R. 1866) 214,	365
on Tonto Apaches—C. Smart. (R. 1867)	215
relative to George Catlin-J. HENRY. (R. 1872)	271
to accounts of lightning discharges, by G. W. Dodge and others—J. HENRY. (R. 1867)	215
to article on vitality, by H. H. Higgins-J. HENRY. (R. 1866)	214
to articles on meteorology, by G. Latimer and others—J. HENRY. (R. 1871)	249
to lectures on meteorology, by R. Russell—J. HENRY. (R. 1854)	75
Notice, biographical, of-	
Agassiz, L., by J. A. GARFIELD. (R. 1878)	275
Agassiz, L., by P. PARKER. (R. 1878)	275
Blackmore Museum, Salisbury, England. (R. 1868)	224
Chase, S. P., by J. A. GARFIELD. (R. 1873)	275

Notice, biographical, of—Continued.
Chase, S. P., by H. Hamlin. (R. 1878) 275
Cleaveland, Parker, by J. HENRY. (R. 1859) 110
Espy, James P., by A. D. BACHE. (R. 1859)
Harvey, W. H., by A. Gray. (R. 1867) 215
Irving, Washington, by C. C. FELTON. (R. 1859)
Jewett, C. C., by R. A. Guild. (R. 1867)215
Rush, Richard, by J. A. PEARCE. (R. 1859) 110
Schoenbein, Christian Frederic, the discoverer of ozone. (R. 1868) 224
Smithson, James, by D. GILBERT, President of the Royal Society. (R. 1853)
Smithson, James, by W. R. Johnson; J. R. McD. Irby 32
Smithson, James, by W. J. RHEES. (R. 1879) 349, 330
Turner, W. W., by C. C. FELTON. (R. 1859) 110
Würdemann, G., by A. D. BACHE. (R. 1859)110
Notice of recent additions to marine invertebrates of northeast coast of America— A. E. Verrill. (P. 1880) 425
Notices of public libraries in the United States—C. C. JEWETT 25
Nova Scotia
birds of—Blackiston; T. Bland; J. R. Willis. (R. 1858) 109
fishes obtained at—See G. B. Goode; T. H. Bean.
kjökken-möddings in—J. M. Jones. (R. 1863) 187
Numeration, report on improved system of—W. B. TAYLOR. (B. 1867) 215
Nuovi Lincei, Pontifical Academy of the. Prize questions. (R. 1865; R. 1867) 209, 215
Nutation, phenomena of—J. G. BARNARD
NUTTER, F. H. Mounds on Gideon's farm, near Excelsior, Hennepin county, Minnesota. (R. 1879)

O.

Ober, F. A.—
birds of Antigua and Barbuda collected by, catalogue of—G. N. Law- RENCE. (P. 1878)8
birds of Dominica collected by, catalogue of-G. N. LAWRENCE. (P. 1878.). 8
birds of Grenada collected by, catalogue of-G. N. LAWRENCE. (P. 1878.) 33
birds of Guadeloupe collected by, catalogue of—G. N. LAWRENCE. (P. 1878)
birds of Lesser Antilles collected by, catalogue of—G. N. LAWRENCE. (P. 1878)
birds of Martinique collected by, catalogue of—G. N. LAWRENCE. (P. 1878)8
birds of St. Vincent collected by, catalogue of—G. N. LAWRENCE. (P. 1878) 8
OBER, F. A. Ornithology of the Caribbee Islands. (R. 1878) 84
Obsequies of Joseph Henry
Observations-
astronomical—W. HARKNESS 2
astronomical, discussion of Piazzi's—B. A. Gould. (R. 1868) 18
astronomical, in Arctic Seas-E. K. KANE
in Hudson's Bay Territory—B. R. Ross. (R. 1859)
magnetic, in the Arctic Seas—E. K. KANE 9
physical, in Arctic Seas—I. I. HAYES
physical, in Arctic Seas. Parts I-IV-E. K. KANE 19
scientific, metric system for—A. GUYOT. (R. 1848)
Observations-See Magnetic, Meteorological, Physical, Tidal.
Observations, meteorological-
at Brunswick, Maine-P. CLEAVELAND 20
at Providence, Rhode Island—A. CASWELL 103, 44
at Sacramento, California—F. W. HATCH. (R. 1854)
at Sacramento, California—T. M. Logan. (R. 1854)
at Washington, Arkansas—N. D. Smith 13
combined, report of committee of American Association for Advance- ment of Science on system of. (R. 1851)
directions for - A. Guyot1
for year 1855 9
in the Arctic Seas—E. K. KANE10
Observations of—
earthquake phenomena—R. Mallet. (R. 1859)
Siredon lichenoides—W. E. CARLIN. (P. 1881)
temperature, best hours for—C. DEWEY. (R. 1860) 14
thunder storm, John Wise's-R. HARE. (R. 1854)
thunder storms, instructions for—J. HENRY 23

Observations on—
auroras, map of stars near North Pole for 350
Batis maritima of Linneus-J. Torrey
electric resonance of mountains—H. DE SAUSSURE. (R. 1868) 22
gold ornament from mound in Florida—C. RAU. (R. 1877) 323, 440, 403
ice—D. Walker
Mexican history and archeology—B. MAYER
natural phenomena, shooting stars, aurora, etc.—S. MASTERMAN. (R. 1857)
prehistoric mounds of Grant county, Wisconsin-M. Strong. (R. 1876.) 29
terrestrial magnetism—J. Locke 2
terrestrial magnetism and deviation of compasses of iron-clad "Monad-nock"—W. HARKNESS
terrestrial magnetism in Mexico—Baron von Müllen; A. Sonntag 11
thunder and lightning—S. MASTERMAN. (R. 1855)
Observatories—
at Dorpat and Poulkova, description of—C. ABBE. (R. 1867) 215, 365
detached, universal meteorograph for—E. H. von BAUMHAUER. (R. 1879.) 84
instructions to, relative to telegraphic announcements of astronomical discoveries—J. Henry
reports of, 1879—E. S. HOLDEN. (R. 1879)
reports of, 1880—E. S. HOLDEN; G. H. BOEHMER. (R. 1880) 442, 436
Observatory—
at Cordova, Argentine Republic, account of—B. A. Gould. (R. 1873.) 27
at St. Martin, Isle Jesus, Canada East, description of—C. SMALLWOOD. (R. 1856)
Girard College—See Bache, A. D.
Kew, verification of barometers at-J. WELSH. (R. 1859) 110
magnetic, at Smithsonian Institution, description of—J. E HILGARD
(R. 1859) 110
physical, on a—J. Henry. (R. 1870) 244
Toronto, instructions by, for aurora observations
Observers, meteorological, of Smithsonian Institution, list of. (R. 1868) 224, 373
Observers—See Meteorological stations and observers.
Occultations—
in 1852—C. H. DAVIS
of planets and stars by the moon, 1858—J. Downes
visible in the United States during the years 1848-1852-J. Downer.
8, 9, 10, 11, 29 Occultator, account of—T. HILL
Ocean, fresh water in the—W. C. DENNIS. (R. 1866) 214
Ocean, Indian, cyclone in the—N. PIKE (R. 1867)
Ocean, Indian, cyclone in the N. I Inc. (10. 1001) 11.11.11.11.11.11.11.11.11.11.11.11.11.

One to	Edwards. (P 1878)
	ER, A. Stone cists near Highland, Madison county, Illinois. (R. 1879.)
	ed, H. C., memoir of, by E. DE BEAUMONT. (R. 1868)
Office	rs— and establishment of the Smithsonian Institution and National Museum.
	collaborators, employés, etc., of Smithsonian Institution, National Museum, Geological Survey, Bureau of Ethnology, and Fish Commission, directory of
	and Regents of Smithsonian Institution, 1846, list of
Ohio-	
	ancient earthworks in, sketch of—I. DILLE. (R. 1866)
	ancient works in, descriptions of—C. WHITTLESEY
	Ashland county, earthworks in—G. W. HILL. (R. 1877)
	Ashtabula county, double-walled earthwork in—S. D. PEET. (R. 1876.)
	Holmes county, flint instruments in—H. B. CASE. (R. 1877)
	Kelley's Island, climate of-G. C. HUNTINGTON. (R. 1866)
	Marietta, meteorological observations at—S. P. HILDRETH; J. WOOD
	northern, antiquities of—G. W. HILL. (R. 1874)
	Perry county, antiquities of—W. Anderson. (R. 1874)
	Trumbull county, mound in—F. MILLER. (R. 1877)
Ohio 1	river, habits of black bass of the-J. Eoff. (R. 1854)
Ohio 1	river, suggestions for improvement of navigation of—C. Ellet, Jr
Oil—	
	coal, explosibility of—Z. ALLEN. (R. 1861)
	porpoise, manufacture of—C. Cook. (P. 1878)
	rock, or petroleum, history of—T. S. HUNT. (R. 1861)
Oil-sh	ark of southern California—D. S. JORDAN; C. H. GILBERT. (P. 1880.)
Old-V	Vorld birds in National Museum, catalogue of—R. RIDGWAY. (P. 1881.)
Oligo	cene fossils, check-list of—T. A. CONRAD
Olmos	s, Peru, account of solar eclipse at—J. M. GILLISS
OLMS:	red, D. Aurora borealis, recent secular period of
OLNE	Y, S. T. List of Rhode Island algæ
Omah	a myth—J. O. Dorsey. (E. 1879-80)
0ōlog	y, North American. Part 1. Raptores and fissirostres-T. M. Brewer.
0ŏlog	y of Kerguelen Island—J. H. KIDDER; E. COUES
Openi	ng and closing of Kennebec river, Maine-R. H. GARDINER. (R. 1858.)
_	esaurus ventralis, remarks on osteology of—R. W. Shuffldt. (P. 1881.)
Orbit-	
	and phenomena of meteoric fire-ball-J. H. COFFIN
	of Neptune, investigation of, with tables of its motion—S. NEWCOMB
	of Uranus, investigation of, with tables of its motion—S. NEWCOMB

Orbits, planetary, secular variations of—J. N. STOCKWELL. (R. 1871) 249, 2	32
Orcynus pelamys (oceanic bonito), occurrence of, in Vineyard Sound, Massa-	
chusetts—V. N. Edwards. (P. 1878)	132
Oregon—	
fishes of, notes on collection of—D. S. JORDAN. (P. 1878)	132
jargon of—B. R. MITCHELL; W. W. TURNER	68
trade-language of-G. GIBBS	161
Ores collected on geological survey in Michigan, catalogue of—C. T. JACKSON. (R. 1854)	75
Ores, rocks, minerals, and fossils, catalogue of—J. Locke. (R. 1854)	īó
Organic bases—A. BAUER. (R. 1872)	7.1
Organisms—See Microscopic.	
Organization and objects of the United States National Museum—G. B. Goods.	
(P. 1881) 467, 4	59
Organization, plan of, and regulations of the U.S. National Museum-G.B.	
Goode. (P. 1881)467. 4	45
Organization of—	
local scientific societies—J. Henry. (R. 1875)	29 6
Smithsonian Institution, programme of P, J,	K)
Smithsonian Institution, report of American Academy of Arts and Sciences on—E. EVERETT; J. SPARKS; B. PEIRCE; H. W. LONG-FELLOW; A. GRAY. (R. 1858)	67
Smithsonian Institution, report of committee of Regents on B, L.	
Organized bodies, natural history of—E. J. MAREY. (R. 1867)	
O'Rielly, Henry, deposition of J. Henry in case of S. F. B. Morse versus	
Origin—	
and history of the Smithsonian Institution-W. J. RHEES 3	28
and propagation of disease—J. C. Dalton. (R. 1878)	
of force—W. B. TAYLOR. (R. 1870)	
Ornament, gold, from a mound in Florida, observations on—C. RAU. (R. 1877.) 323, 440. 46	
Ornithology—	
of Caribbee Islands—F. A. OBER. (R. 1878)	
of Hawaiian and Fanning Islands and California-T. H. STREETS 3	
of Kerguelen Islands—J. H. KIDDER; E. COUES	
Ortalide, monograph of—H. Loew 2	56
O' moptera—	•
bibliography of-S. H. SCUDDER	
mediactions for conecuing—1. As. On Las. (15. 1806)	()) ~~
of North America, catalogue of—S. H. SCUDDER))
Obboana, v. m. moodile of lightning discussing co. (16, 1001)	15
Osmerus attenuatus, new species of California fish, description of—W. N. Lock-	•3."
ington. (Р. 1880)	25

OSTEN SACKEN, R.—
Bibliography of diptera 270
Catalogue of described diptera of North America 102, 270
Diptera of Kerguelen Island 294
Directions for collecting and preserving diptera 102
Monograph of Tipulida 219
Monographs of diptera of North America. Part IV 219
OSTEN SACKEN, R.; LOEW, H. Instructions for collecting diptera. (R. 1858.) 109
OSTEN SACKEN, R.; LOEW, H. Monographs of diptera of North America.
Parts I, II 141, 171
Osteology and mounds of the mound-builders of Wisconsin-J. N. DE HART. (R. 1877)
. ,
Ostraciontidæ (trunk fishes), a study of, with notes on American species—G. B. GOODE. (P. 1879) 888
Othonops eos, a new gobioid fish from San Diego, Cal.—R. SMITH. (P. 1881.) 467
Outline of scheme of classification of collections in U. S. National Museum—
G. B. Goode. (P. 1881) 467, 457
Outline of systematic review of the class of birds—W. LILLJEBORG 364
OWEN, D. D. Catalogue of geological specimens. (R. 1854) 75
OWEN, D. D. Report on building stones. (R. 1847) H, 829
Owen, R. D. Hints on public architectureP
Owl, description of a new, from Porto Rico—R. RIDGWAY. (P. 1881) 467
Ox, American, extinct species of—J. LEIDY 41
Oxygen and its combinations, lecture on—G. I. CHACE. (R. 1855) 77
Ozone and antozone-C. M. WETHERILL. (R. 1864) 188, 355
Ozone, notice of C. F. Schoenhein, the discoverer of (R. 1868)

P.

Pacific const—	
fishes from, descriptions of new-D. S. JORDAN; C. H. GILBERT. (P. 1881)	467
fishes from, distributed in 1881—D. S. JORDAN; P. L. JOUY. (P. 1881.)	467
fishes of—See D. S. Jordan; C. H. Gilbert.	
fishes of, bibliography of-T. H. BEAN. (P. 1881)	467
fishes of, bibliography of-T. GILL	463
fishes of, notes on-D. S. JORDAN; C. H. GILBERT. (P. 1881)	467
Pacific Islands, natural history of-W. H. PEASE. (R. 1862)	150
PACKARD, A. S. Directions for collecting and preserving insects	261
PACKARD, A. S. Memoir of H. J. Clark.	24:
PACKARD, F. A. Project of outline history of education in the United States. (R. 1868)	187
PAGE, C. G. Report on effect of frost on building stones. (R. 1847) H.	329
Palsontology-	
list of generic names employed in—S. H. SCUDDER	470
of the upper Missouri—F. B. MEEK; F. V. HAYDEN	L72
principles and methods of—T. H. HUKLEY. (R. 1869)	228
Palafittes or lacustrian constructions of the lake of Neuchâtel—R. Desor.	
(R. 1865)	
Palenque tablet in the U.S. National Museum—C. RAU	
Palermo, evaporation observed at, in 1865, 1866—P. TACCHINI. (R. 1870)	
PALMIERI, Prof. Electro-magnetic seismograph. (R. 1870)	
PALMIERI, Prof. Presence of electricity during fall of rain. (R. 1870)	
Panama, shells of, review of C. B. Adams' catalogue of—P. P. CARPENTER	
Pancreas, observations on the—J. Jones	
Pandoridse, contributions towards a monograph of the—P. P. CARPENTER	202
Paralepis coruscans, new species of Paralepis, description of—D. S. Jordan; C. H. Gilbert. (P. 1880)	
Parasites-J. Leidy	#
Paris—	
Academy of Sciences, historical sketch of the—M. Flourens. (R. 1862.)	150
Anthropological Society of, history of transactions of, 1865 to 1867—P. Broca. (R. 1868)	224
Exposition—See French Exposition.	
Muséum d'Histoire Naturelle, typical specimens of American fishes in	-
the—D. S. JORDAN. (P. 1879)	
PARKER, P. Biographical notice of L. Agassiz. (R. 1878) 275.	
PARKER, P. Eulogy on Henry Wilson. (R. 1875) 298,	£.A
Parophrys ischyurus, new species of flounders from Puget Sound, description of—D. S. JORDAN; C. H. GILBERT. (P. 1880)	4 25

Parrot, new species of, of genus Chirus, from Dominica-G. N. LAWRENCE. (P. 1880)
Parthenogenesis in the animal kingdom-G. A. KORNHUBER. (R. 1871)
PARVIN, J. B. Habits of the gopher of Illinois. (R. 1854)
Past and future of geology-J. PRESTWICH. (R. 1875)
Patagonia, human remains from—A. RIED. (R. 1862)
Patent Office, U.S., meteorological observations under the direction of. Vols.
I, II 157, 1
PATTERSON, R.; SHARPLESS, T. Phonography. (R. 1856)
PATTON, A. Antiquities of Knox county, Indiana. (R. 1878)
PATTON, A. Antiquities of Lawrence county, Illinois. (R. 1873)
PRABODY, A. P. Scientific education of mechanics and artisans. (R. 1872.) 271, 3
PEALE, T. R.—
Ancient mound in St. Louis, Missouri. (R. 1861)
Method of preserving lepidoptera. (R. 1868)
Prehistoric remains in vicinity of city of Washington, D. C. (R. 1872.)
Uses of brain and marrow of animals among Indians of North America. (R. 1870)
Pearce, J. A., eulogy on, by A. D. BACHE. (R. 1862)
PEARCE, J. A. Notice of Richard Rush. (R. 1859)
PEARCE, J. A. Report of Committee of Regents on distribution of Smithso-
nian income. (R 1858)
PRASE, W. H. Acknowledgment for shells, etc. (R. 1861)
PEASE, W. H. Natural history of Pacific Islands. (R. 1862)
Pedro II, Dom, biographical sketch of, by A. FIALHO. (R. 1876)
Pret, S. D. Double-walled earthwork in Ashtabula county, Ohio. (R. 1876.)
PRIRCE, B., and others. Report of American Academy of Arts and Sciences
on organization of Smithsonian Institution. (R. 1858)
PELTIER, F. A. Memoir of J. C. A. Peltier. (R. 1867)
Peltier, J. C. A., memoir of, by F. A. PELTIER. (R. 1867)
Peltier, J. C. A., scientific researches of. (R. 1867)
Pendulum, problems of rotary motion presented by the—J. G. BARNARD 2
Pennsylvania—
Centre county, Spruce creek valley, account of a tornado in—J. B. Merk. (R. 1871)
Chester county, birds of—V. BARNARD. (R. 1860)
Chester county, history of-W. Darlington. (R. 1862) 1
magnetic survey of-A. D. BACHE
Newcastle, old Indian village, Kushkushkee, near—E. M. McConnell. (R. 1871)
western, ancient mound in-W. M. TAYLOR. (R. 1877)
Pensacola, Florida, description of new species of umber fish (Seriola Stearnsii)
from—G. B. Goode. (P. 1879)

Pepin, Lake, antiquities on the banks of-L. C. Estes. (R. 1866) 21
Perca flavescens, notes on—F. STEINDACHNER. (P. 1878) 38
Percide, notes on-D. S. JORDAN 20
Perennibranchiates, acknowledgment of—J. G. FISHER. (R. 1868) 18
Periodical phenomena—
directions for registry of14
observations on, 1851 to 1859—F. B. Hough18
registration of—J. HENRY. (R. 1855)
registry of 65, 14
Periodicals—
deposited in Library of Congress, meteorological articles in. (R. 1878) 27
in Library of Smithsonian Institution. Parts 1, 11; foreign works; 1866. 73, 85, 117, 17
received by the Institution, list of. (R. 1880) 442, 43
received in reading-room of Smithsonian Institution in 1853, check-list of.
PERRINE, T. M. Antiquities of Union county, Illinois. (R. 1878) 27
PERRINE, T. M. Mounds near Anna, Union county, Illinois. (R. 1872) 27
Persistence of images—F. N. Moigno. (R. 1866)
Perspiration and respiration, apparatus for testing results of—M. PETTENKOPER. (R. 1864)
Peru-
earthquake in, August 18, 1868—J. V. CAMPBELL. (R. 1870) 24
Indians of-F. L. GALT. (R. 1877) 32
Olmos, total eclipse of sun, September 7, 1858, observed near—J. M. GILLISS
PESLIN, M. Relation between barometric variations and general atmospheric currents. (R. 1877)
PETER, R. Ancient mounds in Kentucky. (R. 1872) 27
PETER, R. Ancient mound near Lexington, Kentucky. (R. 1871) 24
Petite Anse Island, salt deposit on—E. W. HILGARD
PETITOT, E. Account of the Indians of British America. (R. 1865) 200
Petrel, supposed new, from Sandwich Islands, description of—R. RIDGWAY. (P. 1881)
Petroleum collections, circular in reference to—S. F. BAIRD. (P. 1881) 467, 447
Petroleum or rock oil, history of—T. S. HUNT. (B. 1861)
PETTENKOFER, M. Apparatus for testing the results of perspiration and res-
piration. (R. 1864) 188
PETTIGREW, J. B. Modes of flight in relation to aeronautics. (R. 1867) 215
Phænogamia of Kerguelen Island—A. Gray
Phalangidæ, instructions for collecting—H. C. Wood. (R. 1866) 214
Phenomens—
accompanying the propagation of electricity in highly rarefled elastic fluids—A. De La Rive. (R. 1868)
in telegraphic lines during the aurora borealis—G. B. Donatt. (R. 1872) 271

Phenomena—Continued.
natural, observations on—S. MASTERMAN. (R. 1857) 10
of contact, studies on—T. L. Phipson. (R. 1862) 15
of flight in animal kingdom—E. J. MAREY. (R. 1869) 22
of meteoric fire-ball—J. H. Coffin 22:
of precession and nutation—J. G. BARNARD
periodical registration of—J. Henry. (R. 1855)7
Philadelphia—
exhibit at—See Centennial.
magnetic and meteorological observations at—See A. D. Bache.
memorial of citizens of, relative to Lowe's aeronautic voyages across
Atlantic. (R. 1860)
Philadelphia Academy of Natural Sciences, report on shells presented to—J. LEIDY; G. W. TRYON. (R. 1865)
PHILLIPS, B. Two letters on work of the Museum. (P. 1881) 467, 45
Philological circular—G. GIBBS. (R. 1862)
Philology, Indian—W. W. TURNER. (R. 1851)
Philology, instructions for research relative to—G. GIBBS 16
Philosophical Society of Washington— Bulletins of. Vols. 1-111. March, 1871, to June, 188042
memoir of Joseph Henry read before the—W. B. TAYLOB 856, 33
notes on life and character of Joseph Henry read before the—J. C.
Welling 856, 33
proceedings of, on death of Joseph Henry 85
Philosophy, annals of, contributions to, by J. Smithson. (R. 1858) 67, 88
Philosophy, Batavian Society of Experimental, of Rotterdam. Prize questions.
(R. 1861) 14
Phipson, T. L. The catalytic force or studies on the phenomena of contact.
(R. 1862) 15
Phlegmasia, disease of, bibliography of works on—W. W. KEEN 80
Phonetic language, vocal sounds of L. Bridgman compared with elements of— F. LIEBER
Phonography—T. Sharpless; R. Patterson. (R. 1856)
Phonography, institutions in which, is taught. (R. 1856)
Photo-chemistry—J. Jamin. (B. 1867) 21
Photographs of Indians, list of21
Photography, astronomical, progress in—Dr. LEE. (R. 1861) 14
Photography, celestial, use of silvered-glass telescope in—H. Draper 18
Phycis Chesteri from deep-sea fauna of northwestern Atlantic, description of—
G. B. GOODE; T. H. BEAN. (P. 1878) 88
Phycis Earlii, new hake, from South Carolina, description of—T. H. BEAN. (P. 1880)
Phycis regius from North Carolina, note on occurrence of—T. H. BEAN. (P. 1880)

Physical—
atlas of North America, proposed—G. GIBBS. (R. 1866)
ethnology, lectures on—D. WILSON. (R. 1862)
geography of Mississippi valley—C. Ellet, Jr.
geography of North America—J. G. Cooper. (R. 1858) 109, 353
geography of North American Continent-J. Frozbel. (R. 1854) 7
observations in the Arctic Seas—I. I. HAYES
observations in the Arctic Seas—E. K. KANE 190
observations in the Arctic Seas—F. L. McCLINTOCK 14
observatory, on a-J. HENRY. (R. 1870) 24
sciences, on the relation of, to science in general—H. Helmholtz. (R. 1871) 24
tables-A. Guyot
Physics—
present fundamental principles of—F. J. Pisko. (R. 1879)
recent progress in, 1879 and 1880—G. F. BARKER. (R. 1880) 442, 42
report on recent progress in, electricity, galvanism—J. MÜLLER. (R. 1855–1858)
syllabus of a course of lectures on—J. HENRY. (R. 1856)
terrestrial, articles on—J. V. CAMPBELL; Prof. PALMIERI; W. W. Johnson; W. D. SARGENT. (R. 1870)
Physics and Natural History, Geneva Society of-See Geneva.
Physiological investigations of vertebrata—J. Jones
Physiology-
electro-, lectures on—C. MATTEUCCI. (R. 1865)
of fever—H. C. Wood
of lucernarians—H. J. CLABK
of rattlesnake—S. W. MITCHELL
of respiration in chelonia—S. W. MITCHELL; G. B. MOREHOUSE 15
Piazzi's astronomical observations, discussion of—B. A. Gould. (B. 1863) 18
Pickering, C. Gliddon mummy case in Museum of Smithsonian Institution. 20
Picture writing, Central American, studies in—E. S. Holden. (E. 1879-80.)
PIDGEON, W. Ancient burial mound in Indiana. (R. 1867) 21
PIERSON, W. M. Discovery of large meteorite in Mexico. (R. 1978) ?
PIKE, N. Cyclone in Indian Ocean. (R. 1867)
PILAR, G. Revolutions of crust of the earth. (R. 1876)
Pile work antiquities of Olmutz. (R. 1866)
PILLING, J. C. Catalogue of linguistic manuscripts in library of Bureau of Ethnology. (E. 1879–80)47
Pima Indians of Arizona—F. E. GROSSMAN. (R. 1871) 24
Pinna, cretaceous, from New Mexico, description of new—C. A. WHITE. (P. 1880)
Pirz, A. Methods of making and preserving plaster casts. (P. 1881)
Pisces, class, arrangement of—T. GILL

OF SMITHSONIAN PUBLICATIONS.	275
Pisko, F. J. Present fundamental principles of physics. (R. 1879)	845
Pitcher plant, new, (Darlingtonia Californica,) from California-J. Torrey	61
Pits at Embarrass, Wisconsin—E. E. BREED. (R. 1877)	828
Pittsburg, on Des Moines river, composition of ancient pottery found at—R. N. Dahlberg; C. Dahlberg. (R. 1879)	845
Plan of—	
a bibliography—J. Friedländer. (R. 1858)	109
organization and regulations of U.S. National Museum—G.B. Goode. (P. 1881) 467,	445
research upon the atmosphere—C. M. WETHERILL. (R. 1866)	214
Smithsonian exhibit at Centennial—S. F. BAIRD. (R. 1875)	298
Planet—See Neptune, Uranus.	
Planetary—	
disturbances, lecture on—E. S. SNELL. (R. 1855)	77
motion, general integrals of—S. NEWCOMB	281
motion, perturbative function of, tables for determining values of coefficients in—J. D. RUNKLE), 94
orbits, secular variations of elements of—J. N. STOCKWELL. (R. 1871.)	232
Planetoids—See Asteroids.	
Planets-	
between Mars and Jupiter—Prof. LESPIAULT. (R. 1861)	149
between Mars and Jupiter, lecture on—E. Loomis. (R. 1854)	75
occultations of—See Downes, J.	
Planisphere of the visible heavens for observations of meteoric displays	359
Plans for the installation of collections in the U.S. National Museum—G.B. Goode. (P. 1881) 467,	472
Plans of the Smithsonian building—R. D. Owen	P
Plantse Frémontianse—J. Torrey	46
Plantæ Wrightianæ Texano-Neo-Mexicanæ. Parts 1, 11-A. GRAY 22	ł, 4 2
PLANTAMOUR, E. Report on transactions of the Society of Physics and Natural History of Geneva, July, 1864, to June, 1865; July, 1874, to June, 1875. (R. 1865; R. 1877)	202
Plants—	, 020
collected by J. C. Frémont in California—J. Torrey	46
cryptogamous, present state of knowledge of-W. REICHARDT. (R. 1871.)	
dates of blossoming of—F. B. Hough	182
dates of defoliation or fall of leaf of—F. B. Hough	182
dates of foliation of—F. B. Hough	182
directions for collecting and preserving—L. F. WARD	
of Arctic America, descriptions of—A. GRAY	842
of Texas, New Mexico, and Mexico, collected by C. Wright—A. GRAY.	22
of upper Missouri, list of—T. C. Porter. (R. 1850)	28
of Washington, check-list of-L. F. WARD	_

Plants—See Torrey, J., Gray, A.
Plaster casts—
methods of making and preserving—A. PIRZ. (P. 1881) 46
of antique and modern statues—W. J. STONE. (R. 1855) 7
of heads of Indian boys and girls at Hampton, Virginia—R. H. PRATT. (P. 1879)
of heads of Indian prisoners at St. Augustine, Florida—R. H. PRATT. (P. 1878)
PLATEAU, F. Rapid preparation of myological specimens. (P. 1881) 467
PLATEAU, J. Experimental and theoretical researches on the figures of equilibrium of a liquid mass withdrawn from the action of gravity. Parts I-VI. (R. 1868-1866)
Platessa ferruginea, note on-G. B. Goode; T. H. Bean. (P. 1878) 335
Platessa rostrata, note on-G. B. Goode; T. H. BEAN. (P. 1878) 33
Platycnemism in Michigan—H. GILLMAN. (R. 1878) 275, 393
Platyrhina exasperata, generic relations of—D. S. Jordan; C. H. Gilbert. (P. 1880)
Platyrhina triseriata, new ray from coast of California, description of—D. S. JORDAN; C. H. GILBERT. (P. 1880) 425
Platysomatichthys stomias, new flounder from coast of California, description of—D. S. JOBDAN; C. H. GILBERT. (P. 1880)425
Pleuronectes glaber, identity of, with Euchalarodus Putnami—T. H. BEAN. (P. 1878)
Pleuronectides of San Francisco, review of-W. N. LOCKINGTON. (P. 1879) 833
Pleuronichthys verticalis, new flounder from coast of California, description of—D. S. JORDAN; C. H. GILBERT. (P. 1880) 425
POE, O. M. Account of lightning discharges. (R. 1867) 215
Poër, A. New classification of clouds. (R. 1870) 244
POEY, F. Notes on American species of genus Cybium. (P. 1878) 822
POGGENDORFF, J. C. Use of galvanometer as measuring instrument. (R. 1859.) 110
Points, boiling and melting—F. W. CLARKE
Points, spear and arrow, primitive manufacture of, along line of Savannah
river—C. C. Jones, Jr. (R. 1879)
Poison—See Venom.
Poisoned arrows, lecture on nature and cure of wounds of—D. BRAINARD. (R. 1854)
Poisonous water in Gulf of Mexico, destruction of fish by—J. Y. PORTER. (P. 1881) —————————————————————————————————
Polar expedition—See Hall, Capt.
Polar expedition, the Howgate, contributions to natural history of Arctic America made in connection with—L. Kumlien 342
Polar light, or aurora borealis, its phenomena and laws—E. Loomis. (R. 1865.) 209
Polaris expedition, scientific instructions for—J. HENRY; J. E. HILGARD; S. NEWCOMB: S. F. BAIRD: F. B. MERK: T. AGASSIZ. (R. 1871)

·
Polarized nerve, effect of irritation of-B. F. LAUTENBACH. (R. 1878) 841, 411
Pollock, J. Assay of coins at the Mint of the United States. (R. 1868) 224
Polychrome bead from Florida—S. S. HALDEMAN. (R. 1877) 828, 404
Polypetalæ, bibliographical index of—S. WATSON 258
Polypi, relations of lucernarians to—H. J. CLARK 242
Polyzoa of northeastern coast of America, notice of recent additions to—A. E. VERRILL. (P. 1879)888
Pomadasys, description of new species of, from Mazatlan, Mexico—D. S. Jon-DAN; C. H. GILBERT. (P. 1881) 467
Pontifical Academy of the Nuovi Lincei, Rome. Prize questions. (R. 1865; R. 1867)
Poole, H. Cone-in-cone. (R. 1868) 187
Popocatepetl, examination of—Baron von Müller; A. Sonntag 114
Population of the world, estimate of the—E. MAILLY. (R. 1873) 275
Porcupine, Canada, occurrence of, in Maryland-O. Lugger. (P. 1881) 467
Porcupine, Canada, occurrence of, in West Virginia-G. B. Goode. (P. 1878.) 832
Porifers of northeast coast of America—A. E. VERRILL. (P. 1879) 838
Porpoise oil, manufacture of—C. Cook. (P. 1878) 882
PORTER, Commodore. Account of hail storm on the Bosphorus. (R. 1870) 244
PORTER, J. Y. Destruction of fish by poisonous water in Gulf of Mexico. (P. 1881) 467
PORTER, J. Y.; JEFFERSON, J. P.; MOORE, T. Destruction of fish near the Tortugas. (P. 1878)
PORTER, T. C. List of plants of upper Missouri. (R. 1850) 28
Portland Society of Natural History, account of—E. C. Bolles. (R. 1867) 215
Porto Rico—
description of new owl from-R. RIDGWAY. (P. 1881) 467
great hurricane at—G. A. LATIMER. (R. 1867) 215
Latimer collection of antiquities from—O. T. MASON. (R. 1876) 397
meteorology of-G. LATIMER. (R. 1871) 249
Portraits-
of North American Indians, catalogue of—J. M. STANLEY 53
of North American Indians, Stanley, report of Committee of Regents on. (R. 1857) 107
photographic, of North American Indians, in gallery of Smithsonian Institution, list of
Potamocottus Bendirei, description of—T. H. BEAN. (P. 1881) 467.
Pottery—
ancient, from Phillips county, Arkansas—J. H. Devereux. (R. 1872.) 271
ancient, from Pittsburg, on Des Moines river—R. N. Dahlberg; C. Dahlberg. (R. 1879) 345
Indian—C. RAU. (R. 1866)
Pouched rat, or salamander of Georgia, habits of—W. GESNER. (R. 1860) 147
Poulkova and Dorpat, description of observatories at—C. ABBE. (R. 1867.) 215, 369

Powell, B. Reports on state of knowledge of radiant heat. (R. 1859) 110
Powell, J. W.—
Evolution of language. (E. 1879-80)
Limitations to use of some anthropological data. (E. 1879-80) 476
Mythology of North American Indians. (E. 1879-80) 476
Report of Director of Bureau of Ethnology. (E. 1879-80) 476
Wyandot government. (E. 1879-80) 47
Powers, S. Centennial mission to Indians of western Nevada and California. (R. 1876)
PRATT, R. H. Catalogue of casts of heads of Indian boys and girls at Hampton Institute, Virginia. (P. 1879)
PRATT, R. H. Catalogue of casts of heads of Indian prisoners at St. Augustine, Florida. (P. 1878)
PRATT, W. H. Antiquities of Whiteside county, Illinois. (R. 1874) 28
Prayer at funeral of Joseph Henry—C. Hodge 35
Prayer at memorial of Joseph Henry-J. McCosh
Prayer at memorial of Joseph Henry—B. SUNDERLAND
Precession of the equinoxes—J. N. STOCKWELL 22
Precession of the equinoxes, problems presented by-J. G. BARNARD 240, 310
Precious stones—J. BABINET. (R. 1870)
Precipitation—
influence of rain upon the formation of—J. HANN. (B. 1877) 823, 395
of rain and snow in the United States, charts of-C. A. Schott 374
of rain and snow in the United States, tables of-C. A. Schott 222, 355
Prehistoric—
antiquities of Hungary—F. F. Romer. (R. 1876) 299, 440, 392
archeology, international code of symbols for charts of—G. DE MORTIL- LET; E. CHANTRE. (R. 1875)
evidences in Missouri—G. C. BROADHEAD. (R. 1879) 345
man, remains of, from caves in Alaska—W. H. DALL 318
mounds of Grant county, Wisconsin-M. STRONG. (R. 1876) 29
remains in vicinity of city of Washington, D. C.—T. R. PEALE. (R. 1872)
PRENTISS, S. S.; COUES, E. List of birds of the District of Columbia. (R. 1861)
Preparation of large myological specimens—F. J. PLATEAU. (P. 1881) 467
Preparations, dry, on Semper's method of making—J. A. RYDER. (P. 1881.) 467
Prescott, Canada West, ancient Indian remains near—W. E. Guest. (R. 1856)
PRESCOTT, W. H., and others. On publication of Spanish works on New Mexico. (R. 1855)
Present state of ethnology in relation to form of human skull—A. RETZIUS. (R. 1859)

OF SMITHSONIAN PUBLICATIONS.	279
Presentation of—	
books—Duke of Northumberland. (R. 1859)	110
books-Imperial Library of Vienna. (R. 1865)	209
books on Brazil—M. M. LISBOA. (R. 1865)	209
books on Egypt—R. LEPSIUS. (R. 1860)	147
collections by foreign governments. (R. 1876)	
engravings—C. B. KING. (R. 1861)	149
specimens—University of Christiania, Norway. (R. 1868)	187
Preservation—	
and collection of marine invertebrates—W_STIMPSON	84
of antiquities and national monuments in Denmark—J. J. A. Worsaar. (R. 1879)	8 4 5
of copper and iron in salt water—A. E. BECQUEREL. (R. 1864)	188
of wood. (R. 1864)	188
Preserving—	
diptera, directions for-R. OSTEN SACKEN	102
fish, directions for-T. H. BEAN. (P. 1881) 467,	464
insects, directions for—A. S. PACKARD	261
lepidoptera, method of-T. R. PEALE. (R. 1868)	187
nests and eggs, instructions for-T. M. Brewer	189
plants, directions for-L. F. WARD	460
specimens of diatomacea—A. M. EDWARDS	366
specimens of natural history—S. F. BAIRD. (R. 1856) 9	1, 34
Pressure and velocity of the wind—J. HANN. (R. 1877) 828	3, 398
Pressure of the air—J. HANN and others. (R. 1877) 828	3, 898
PRESTWICH, J. Past and future of geology. (R. 1875)	.298
Priestley, J., memoir of, by G. CUVIER. (R. 1858)	109
Priestley's lens, account of—J. HENRY. (R. 1859)	110
PRIME, T. Monograph of American corbiculade	145
PRIME, T., and others. Check-list of shells of North America	128
Primitive manufacture of spear and arrow points along line of Savannah river— C. C. Jones, Jr. (R. 1879)	
Prince Regent's Inlet, meteorological observations in-F. L. McCLINTOCK	
Princeton, New Jersey, address at, by J. HENRY	
Principles—	
and methods of palsontology—T. H. HUXLEY. (R. 1869)	228
of crystallography and crystallophysics, explanation of—A. Brezina. (R. 1872)	
of mechanical theory of heat—J. MÜLLER. (R. 1868)	-
of physics, present fundamental—F. J. Pisko. (R. 1879)	
of subcutaneous surgery—W. ADAMS	
Prionotus stephanophrys, new species of Prionotus from coast of California,	
description of—W. N. Lockington. (P. 1880)	425

Prize questions—	
Academy of Sciences of the Institute of Bologna. (R. 1862)	150
Batavian Society of Experimental Philosophy of Rotterdam. (R, 1861.)	149
Dunkirk Society for Encouragement of Sciences, Letters, and Arts. (R. 1865)	209
Holland Society of Science, Harlem. (R. 1861, 1864, 1867, 1878) 149, 188, 224,	275
Imperial Academy of Sciences of Vienna. (R. 1864; R. 1865) 188,	209
Imperial Society of Natural Sciences of Cherbourg. (R. 1864)	188
Imperial Society of Science, Agriculture, and Arts of Lille. (R. 1865.)	209
London Institution of Civil Engineers. (R. 1862)	150
Pontifical Academy of Nuovi Lincei, Rome. (R. 1865; R. 1867) 209,	
Royal Academy of Netherlands. (R. 1861)	149
Royal Academy of Science, Literature, and Fine Arts, Brussels. (R. 1878.)	_
Royal Danish Society of Sciences. (R. 1862, 1865, 1867) 150, 209,	215
Royal Institute for Encouragement of Natural, Economical, and Technological Science, Naples. (R. 1878)	275
Royal Prussian Academy of Sciences. (R. 1864)	
Royal Scientific and Literary Institute of Lombardy. (R. 1865)	
Society for Encouragement of Science, Literature, and Art, Dunkirk, France. (R. 1878)	
Society of Arts and Sciences of Utrecht. (R. 1861; R. 1862) 149,	
Society of Science, Art, and Literature of Hainaut, Mons, Belgium. (R. 1873)	
Problems of rotary motion, presented by gyroscope, precession of equinoxes, and pendulum—J. G. BARNARD	
Proceedings of—	
Board of Regents—See Regents.	
Establishment, 1853-1854. (R. 1853) 67,	329
U. S. National Museum. Vols. I-IV, 1878-1881 332, 333, 425,	
U. S. National Museum, catalogue and index of	478
Products of combustion of gun-cotton and gunpowder—Lieut. von Karolyi; B. F. Chaig. (R. 1864)	188
Productus giganteus, note on occurrence of, in California—C. A. WHITE. (P. 1880)	425
Programme of organization of Smithsonian Institution P, J,	328
Progress—	
in physics—See J. Müller.	
in science, record of—See Record.	
of astronomical photography—Dr. Lee. (R. 1861)	149
Project of outline history of education in United States—F. A. PACKARD. (R. 1868)	187
Promotion of science in the United States, Tyndall trust for the. (R. 1872)	271
Propagation of disease—J. C. Dalton. (R. 1873)	
Prospectus of a Bibliographia Americana—H. STEVENS. (B. 1848)	_

Providence, Rhode Island, meteorological observations at—A. CASWELL: 1881 to 1860
Provincetown, Massachusetts, experiments made at, upon animal heat of fishes— J. H. Kidder. (P. 1879)
Provincetown, Massachusetts, littoral marine fauna of-R. RATHBUN. (P. 1880.) 49
Provincial Society of Arts and Sciences, Utrecht. Prize questions. (R. 1862.) 1
Provisional classification of the food collections—G. B. Goode. (P. 1881.) 467, 45
Prussian Royal Academy of Sciences. Prize questions. (R. 1864)
Psychrometric observations 1
Psychrometrical tables—J. H. Coffin
Pterophryne, on proper specific name of—T. Gill. (P. 1878) 83
Ptychochilus Harfordi, new species of Ptychochilus, description of—D. S. Jor- DAN; C. H. GILBERT. (P. 1881)
Public—
architecture, hints on—R. D. OWEN
buildings, acoustics applied to—J. HENRY. (R. 1856)
instruction, minister of, report to Emperor of France by, on scientific expedition to Mexico. (R. 1864)
libraries of the United States, notices of-C. C. JEWETT. (R. 1849) 21, 2
libraries—See Libraries.
Publications
exchange of—Agricultural Association of Milan. (R. 1863) 1
exchange of—Chamber of Commerce of Bordeaux. (R. 1868) 1
of learned societies and periodicals in Smithsonian library. 73, 85, 117, 17 of Smithsonian Institution, list of. (R. 1866.) 74, 203, 226, 245, 278, 290, 801, 844, 487, 4
of U.S. Government, 1868-1881-G. H. Boehmer 4
of U.S. National Museum, list of. (P. 1881) 467, 47
periodical received in reading-room of Smithsonian Institution
report on, 1858-1866—S. F. BAIRD. (R. 1858-1866.) 67, 75, 77, 91, 107, 109, 110, 147, 149, 150, 187, 188, 209, 2
Puebla, Mexico, description of large fossil gasteropod from—C. A. WHITE. (P. 1880)
Puget Sound, description of two new species of flounders (Parophrys ischyurus and Hippoglossoides elassodon) from—D. S. JORDAN; C. H. GILBERT. (P. 1880)
Pulmonata geophila, monograph of—W. G. BINNEY; T. BLAND
Pulmonata limnophila, monograph of-W. G. BINNEY 1
PUMPELLY, R. Geological researches in China, Japan, and Mongolia 20
Purple dyeing, ancient and modern. (R. 1868) 1

Q.

QUALE; LEWIS. Account of the cryolite of Greenland. (R. 1866)	214
Qualitative determinations by the blow-pipe—T. EGLESTON. (R. 1872)	271
Quartermasters, authority to, to receive and transmit specimens to Smithsonian.	34
QUATREFAGES, J. L. Memoir of Isidore Geoffroy St. Hilaire. (B. 1862)	150
Queen Charlotte's Islands, Haidah Indians of-J. G. SWAN	267
Queries about expression for anthropological inquiry—C. DARWIN. (R. 1867.)	215
Queries relative to tornadoes—J. HENRY	190
Questions relative to the food fishes of the United States—S. F. BAIRD	234
Questions—See Prize questions.	
Quetelet, L. A. J., eulogy on, by E. MAILLY. (R. 1874)	286
QUICK, E. R. Mounds in Franklin county, Indiana. (R. 1879)	845
Quillehute Indians of west coast of Washington Territory, method of taking	
surf smelt by-J. G. SWAN. (P. 1880)	425
Quincy, Illinois, mounds near—W. G. Armstrong. (R. 1879)	845

R.

Rabbit catching the sun in a trap (Omaha myth)—J. O. Dorsey. (E. 1879-80.) 476
Race, the human, probable future of-A. DE CANDOLLE. (R. 1875) 298
Races-
intermixture of—G. GIBBS. (R. 1864)188
lower, social and religious condition of the—J. LUBBOCK. (R. 1869) 228
mixed, in Liberia—E. D. BLYDEN. (R. 1870)
Radiant heat, reports on state of knowledge of—B. Powell. (R. 1859) 110
Radiates of Arctic America—A. E. VERRILL 342
Radiation, lecture on—J. TYNDALL. (R. 1868)
Rafinesque's memoirs on North American fishes, review of-D. S. Jordan 305
Raia inornata, note on—D. S. JORDAN; C. H. GILBERT. (P. 1881) 457
Raia rhina, new species of ray from coast of California, description of—D. S. JORDAN; C. H. GILBERT. (P. 1880)
Raia stellulata, new species of ray from Monterey, California, description of— D. S. JORDAN; C. H. GILBERT. (P. 1880)
Rain-
and atmospheric pressure—J. HANN. (R. 1877) 328, 398
and barometric minima—T. REYE. (R. 1877) 823, \$98
and snow, tables of precipitation in—C. A. Schott 222, 353

Rain—Continued.
articles on, preserved in Smithsonian. (R. 1874) 286, 353
gauges—R. H. GARDINER. (R. 1858) 109
influence of, upon the barometer—J. HANN. (R. 1877) 828, 898
observations 157
presence of electricity during fall of-Prof. PALMIERI. (R. 1870) 244
quantity of, at different heights—O. W. Morris; J. Henry. (R. 1855.) 77
snow, or hail, electricity of induction in strata of atmosphere surrounding cloud resolving into—F. ZANTEDESCHI. (R. 1870) 244
Rain-charts of United States, 1, 2, 8: summer, winter, and year—C. A. Schott. 374
Rana pipiens, anatomy of nervous system of—J. WYMAN
Raptores, oŏlogy of—T. M. Brewer.
Rat, pouched, of Georgia, habits of-W. GESNER. (R. 1860) 147
RATHBUN, R. Littoral marine fauna of Provincetown, Cape Cod, Massachusetts. (P. 1880) 425
RATHBUN, R. Marine invertebrates from New England, distributed by U. S. National Museum, list of. (P. 1881) 467, 465, 471
RATHBUN, R.; VERRILL, A. E. Marine invertebrata of New England, distributed by Fish Commission. (P. 1879)838
Ratio between diameter and circumference of a circle—W. FERREL 233
Rattlesnake, researches upon venom of—S. W. MITCHELL
Rattlesnakes, bibliography of—S. W. MITCHELL 185
Rattlesnakes, genera and species of—E. D. Cope
RAU, C.—
Agricultural flint implements in southern Illinois. (R. 1868.) 224, 440, 370
Agricultural implements of North American stone period. (R. 1868.) 187, 440
Ancient aboriginal trade in North America. (R. 1872) 271, 440, 385
Anthropological publications, list of. (P. 1881) 467
Archæological collection of the U.S. National Museum 287
Artificial shell-deposits in New Jersey. (R. 1864) 188, 440, 362
Drilling in stone without metal. (R. 1868) 224, 440, 372
Gold ornament from mound in Florida, observations on. (R. 1877.)
828, 440, 403
Indian pottery. (R. 1866)
Memoir of C. F. P. Von Martius. (R. 1869) 228, 440, 251
North American stone implements. (R. 1872) 271, 440, 382
Palenque tablet in the U.S. National Museum
Stock-in-trade of an aboriginal lapidary. (R. 1877) 828, 440, 402
Translation of Baegert's account of the Indians of southern California. (R. 1868; R. 1864) 187, 188, 440, 361
Translation of Romer's antiquities of Hungary. (R. 1876) 299, 440, 392
RAWSON, R. Memoir of Eaton Hodgkinson. (R. 1868) 224
READ, M. C. Ancient mound near Chattanooga, Tennessee. (R. 1867) 215

Reading-room of Smithsonian Institution, periodicals received in, 1858 Q
Recalculation of atomic weights. Constants of Nature Part v-F. W. CLARKE. 441
Receipts of specimens—See annual reports.
Recent—
improvements in the chemical arts—J. C. BOOTH; C. MORFIT 27
progress in physics—J. MÜLLER. (R. 1855–1858) 77, 91, 107, 109
progress in relation to the theory of heat—A. CAZIN. (R. 1868) 224
progress in science, reports on—See Record of recent progress.
researches on secular variations of planetary orbits—J. N. STOCKWELL. (R. 1871)
Recommendation of Shea's Indian linguistics—G. GIBBS and others. (R. 1861.) 149
Record of-
auroras in higher northern latitudes—P. Force 84
meteorological material preserved in Smithsonian. (R. 1874) 286
monthly meteorological reports preserved in Smithsonian Institution. (R. 1878) 275
Record of recent progress in science, 1879 and 1880-
Anthropology—O. T. MASON. (R. 1880)
Astronomy—E. S. Holden. (R. 1880) 442, 427
Botany—W. G. FARLOW. (R. 1880)
Chemistry—G. F. BARKER. (R. 1880)
Geology—G. W. HAWES. (R. 1880) 442, 428
Mineralogy—G. W. HAWES. (R. 1880) 442, 428
Physics—G. F. BARKER. (R. 1880) 442, 429
Zoölogy—T. GILL. (R. 1880) 442, 431
Recording Indian languages, method of—J. O. Dorsey; A. S. GATSCHET; S. R. RIGGS. (E. 1879-80)
Records and results of magnetic survey of Pennsylvania—A. D. BACHE 166
Red river of the North, ethnology of Indians of valley of—W. H. GARDNER. (R. 1870)
Red river settlement, Hudson's Bay Territory, Indian remains in—D. Gunn. (R. 1867)
REED, H. Lecture on "the Union." (R. 1854) 75
Refraction of sound—W. B. TAYLOB. (R. 1875) 298
Regents of Smithsonian Institution— list of, 1846
list of, 1846-1879. (See also each annual report) 329
memorial of Board of, relative to Smithson fund. (R. 1850; R. 1866.) 28, 214, 329
memorial of Board of, to Congress asking appropriations for museum. (R. 1867)
memorial to Congress in behalf of, relative to new museum building— G. BANGROFT. (R. 1876)
proceedings of Board of. (See also each annual report) A, G, H, I, M, 329
proceedings of Board of, on death of Joseph Henry
historiane or moure or our common or a confin mount

Regents of Smithsonian Institution—Continued.
proceedings of Board of, relative to the electro-magnetic telegraph 115, 82
report of Committee of, on Corcoran Art Gallery. (R. 1872) 271, 82
report of Committee of, on fire at Smithsonian—R. WALLACH; J. HENRY. (R. 1864) 188, 829
report of Committee of, on Museum — A. GRAY; H. COPPÉE. (R. 1874) 286, 829
report of Committee of, on Museum—A. GRAY; A. A. SARGENT; H. CLYMER. (R. 1876) 299, 829
report of Committee of, on Stanley's gallery of Indian portraits. (R. 1857) 107, 829
report of Committee of, on communication of J. Henry relative to electromagnetic telegraph—C. C. Felton. (R. 1857) 107, 829
reports of Committee of, on distribution of Smithsonian income. J. A. Pearce; J. Meacham. (R. 1858) 67, 329
Regents of University of State of New York. Acknowledgment for specimens. (R. 1865)
Registration of periodical phenomena—J. HENRY. (R. 1855) 77
Registry of periodical phenomena 148, 65
Regulations of the Smithsonian Institution, 1872 260
Regulations of the U.S. National Museum—G. B. Goode. (P. 1881) 467, 445
REICHARDT, W. Present state of knowledge of cryptogamous plants. (R. 1871.) 249
Reid, D. B. Architecture in relation to ventilation, warming, lighting, fire- proofing, acoustics, and health. (R. 1856)
Reigen Mazatlan collection of mollusks—P. P. CARPENTER 252
Reindeer, man as the contemporary of, in middle Europe. (R. 1867) 218
Relation—
between barometric variations and atmospheric currents—M. PESLIN. (R. 1877) 828, 898
of food to work, and its bearing on medical practice—S. HAUGHTON. (R. 1870)
of physical sciences to science in general—H. Helmholtz. (R. 1871) 249
of time and space, lecture on—S. ALEXANDER. (R. 1861)
Relationship
circular respecting degrees of-L. H. MOBGAN
system of, of Cree Indians—E. A. WATKINS. (R. 1862) 150
systems of—L. H. Morgan 218
Relative intensity of heat and light of sun-L. W. MEECH. (R. 1856) 91, 83
Relic, ancient, of Maya sculpture, remarks on—A. Schott. (R. 1871) 249
Relies
ancient, in Missouri-J. W. Foster. (R. 1868) 187
ancient Indian-J. Jones 259
ancient, of northwestern Iowa-J. B. Currs. (R. 1872) 271
presented by J. H. Devereux. (R. 1872) 271

Religion of Shoshone Indians—A. G. BRACKETT. (R. 1879)	145
Religious condition of lower races of man—J. Lubbock. (R. 1869)	228
Remains—	
aboriginal, of Tennessee-J. Jones	259
ancient, in Colorado-E. L. BERTHOUD. (R. 1867)	215
human, from Patagonia-A. RIED. (R. 1862)	150
Indian, in Caddo parish, Louisiana-T. P. HOTCHKISS. (R. 1872)	
of later prehistoric man from caves in Alaska—W. H. Dall 3	
of walrus (?) in Maine—C. H. Boyd. (P. 1881)	
prehistoric, in Montana—P. W. Norris. (R. 1879)	
prehistoric, in vicinity of city of Washington, D. C.—T. R. PEALE. (R. 1872)	
Zapotec, account of—J. G. SAWKINS; B. MAYER	
Remains—See Ancient remains.	
Remarkable forms of hail-stones in Georgia—S. ABICH. (R. 1869)	998
Remarks—	
	908
concerning nature of currents of air—A. Colding. (R. 1877) 328,	
on ancient relic of Maya sculpture—A. SCHOTT. (R. 1871)	
on Cara gigantesca of Yzamal in Yucatan—A. Schott. (R. 1869) on kjökken-möddings on northwest coast of America—P. Schumacher.	
	275
on osteology of Opheosaurus ventralis—R. W. SHUFELDT. (P. 1881)	
on physical geography of North America—J. FROEBEL. (R. 1854)	
on species of genus Chiurus—W. N. Lockington. (P. 1880)	
Reminiscences of Joseph Henry by S. B. Dop	856
Reparatory inflammation in arteries after ligature, etc., nature of. Toner lecture No. VII—E. O. SHAKESPEARE	32 1
Reply to criticisms of J. Hann by W. FERREL. (R. 1877) 823,	39 8
Report—See Henry, J., Baird, S. F., Regents.	
Report of—	
American Academy of Arts and Sciences on organization of Smithso-	
nian—E. Everett; J. Sparks; B. Pierce; H. W. Longfellow;	47
A. GRAY. (R. 1858)	
architect—B. S. ALEXANDER. (R. 1854)	10
architects, 1879, 1880—Cluss and Schulze. (R. 1879; R. 1880) 345, 442, 409,	434
Board of Regents of Smithsonian Institution, 1847-1887 G, H, I, 21, 28, 51, 57, 67, 75, 77, 91, 107, 109, 110, 147, 149, 150, 187, 188, 209, 214, 215, 224, 228, 244, 249, 271, 275, 286, 298, 299, 828, 841, 345,	442
Bureau of Ethnology, first-J. W. Powell. (E. 1879-80)	176
chemist—F. W. TAYLOR. (R. 1880)	449
Commission on general stereotype catalogue of public libraries. (R. 1850.) 28	
Committee of American Association on system of combined meteoro-	
logical observations. (R. 1851)	51

Report of—Continued.
Executive Committee (See also each annual report) 829
Executive Committee on Henry statue. (R. 1880) 445
experiments on animal heat of fishes—J. H. KIDDER. (P. 1879) 888
explorations in New Mexico and Arizona-J. STEVENSON. (R. 1880) 442
Government explorations and surveys. (R. 1878) 84
National Museum Building Commission for 1879. (R. 1879) 845, 405
National Museum Building Commission for 1880. (R. 1880) 442, 434
Secretary of Smithsonian Institution, 1847–1877—J. HENRY. (R. 1847–1877) F, H, I, 21, 28, 51, 57, 67, 75, 77, 91, 107, 109, 110, 147, 149, 150, 187, 188, 209, 214, 215, 224, 228, 244, 249, 271, 275, 286, 298, 299, 828
Secretary of Smithsonian Institution, 1878-1880—S. F. BAIRD. (R. 1878-1880)
Senate Judiciary Committee on management of Smithsonian—A. P. But- Ler. (R. 1855)
Smithsonian Institution—See Report of Board of Regents.
visit to Luray Cavern-O. T. Mason. (R. 1880) 442, 433
Report of Committee of Regents of Smithsonian Institution on-
best use of new hall of the Institution—L. Agassiz. (R. 1867) 215, 829
communication from S. F. B. Morse—C. C. Felton. (R. 1857) 107, 829
Corcoran Art Gallery. (R. 1872) 271, 829
electro-magnetic telegraph—C. C. FELTON. (R. 1857) 107, 829
fire of January, 1865—R. WALLACH; J. HENRY. (R. 1864) 188, 829
Greek album
Museum—A. Gray; H. Coppér. (R. 1874)
Museum—A. Gray; A. A. SARGENT; H. CLYMER. (B. 1876) 299
organization of Institution B, L, 82
Stanley's gallery of Indian portraits. (R. 1857) 107, 829
system of accounts. (R. 1866) 214
Washington canal—R. DELAFIELD. (R. 1868) 224, 829
Report on—
building stones—D. D. OWEN 829
catalogue system—C. C. JEWETT. (R. 1849)2
Centennial Exhibition—S. F. BAIRD. (R. 1876) 299, 307
contents of bottles of water from Gulf of Mexico—W. G. FARLOW. (P. 1881) 467
copyright system—C. C. JEWETT. (R. 1851) 51
effect of frost on stones—C. G. PAGE 829
ethnological collections of Museum of Lausanne—F. Troyon. (R. 1861) 149
exploration of ancient mounds in Union county, Kentucky—S. S. Lyon. (R. 1870)
fishes of New Jersey coast, as observed in 1854—S. F. BAIRD. (R.
1854)
fossils from Nebrasku—J. LEIDY. (R. 1851)

Report on—Continued.
Halliwell manuscripts—C. C. JEWETT. (R. 1852)
history of discovery of Neptune—B. A. Gould 18
improved system of numeration—W. B. TAYLOR. (R. 1867) 215
international exchanges—G. H. BOEHMER 477
lacustrian settlements—A. Morlot. (R. 1868) 187
library-C. C. JEWETT. (R. 1848-1858) I, 21, 28, 51, 57, 67
limpets and chitons of Alaskan and Arctic regions—W. H. Dall. (P. 1878)
magnetic telegraph 115
meteorological system—E. FOREMAN. (R. 1852)
meteorology of the United States—E. Loomis. (R. 1847) H
mollusca of west coast of North America-P. P. CAPENTER 252
Museum and statistics of British Museum—S. F. BAIRD. (R. 1850) 28
paleontological collections made by Lieut. G. K. Warren—F. B. MEEK; F. V. HAYDEN
plan of library-C. C. JEWETT. (R. 1847)
proposed exhibit by Smithsonian Institution at the Centennial Exhibition—S. F. BAIRD. (R. 1875) 298
public libraries of United States—C. C. JEWETT. (R. 1849) 21
publications, exchanges, explorations, and Museum, 1851-1877—S. F. BAIRD. (R. 1851-1877) 51, 57, 67, 75, 77, 91, 107, 109, 110, 147, 149, 150, 187, 188, 209, 214, 215, 224, 228, 244, 249, 271, 275, 286, 298, 299, 328 recent progress in physics: electricity, galvanism—J. Müller. (R. 1855-
1857) 77, 91, 107
shell heaps of Tampa Bay, Florida—S. T. WALKER. (R. 1879) 345
shells presented to Academy of Natural Sciences—J. LEIDY; G. W.
TRYON. (R. 1865)
state of knowledge of radiant heat—B. Powell. (R. 1859) 110 survey of economic geology of Trinidad—G. P. Wall; J. G. Sawkins.
(R. 1856) 91 system of accounts. (R. 1866) 214
use of new hall in Smithsonian building—L. Agassiz. (R. 1867) 215
Report to Emperor of France by Minister of Public Instruction on scientific
expedition to Mexico. (R. 1864)188
Reports, meteorological, preserved in Smithsonian Institution, list of. (R. 1878.) 275
Reports of—
astronomical observatories, 1879—E. S. Holden. (R. 1879) 845, 410
astronomical observatories, 1880—E. S. Holden; G. H. Boehmer. (R. 1880)
Committees of Regents, 1846-1877
Secretary of Smithsonian Institution, J. Henry, 1865-1877 343
Smithsonian Institution, catalogue and index of478

Reports on transactions of Geneva Society of Physics—See Geneva.	
Reptiles—	
circular relating to collections of living	320
cretaceous—J. Leidy	192
cretaceous, review of-J. LEIDY. (R. 1864)	188
dates of first appearance of-F. B. Hough	182
extinct—J. Leidy	192
North American, catalogue of. Part I. Serpents—S. F. BAIRD; C. GIRARD	49
Reptilia, check-list of North American—E. D. Cope	292
Request for bison by Bern Museum. (R. 1865)	
Request for shells—W. R. Logan. (R. 1859)	
Research—	
ethnological—E. H. DAVIS. (R. 1866)	214
relative to ethnology and philology of America, instructions for—G.	•
Gibbs	160
upon the atmosphere, plan of a-C. M. WETHERILL. (R. 1866)	
Researches—	
archæological, at Concise—F. TROYON. (R. 1861)	149
archæological, in Nicaragua—J. F. Bransford	383
by officers of Hudson's Bay Company, letter of Sir G. Simpson	
geological, in China, Mongolia, and Japan—R. PUMPELLY	202
in sound—J. HENRY. (R. 1878)	
on ammonia-cobalt bases-W. GIBBS; F. A. GENTH.	88
on electrical rheometry—A. SECCHI	36
on figure of equilibrium of a liquid mass withdrawn from action of gravity—See J. Plateau.	
on secular variations of planetary orbits—J. N. STOCKWELL. (R. 1871.)	249
relative to nebulæ—Prof. GAUTIER. (R. 1868)	187
relative to planet Neptune—S. C. WALKER	3
scientific, list of apparatus available for. (R. 1878)	841
scientific, of Peltier-F. A. Peltier. (R. 1867)	215
upon anatomy and physiology of Chelonia—S. W. MITCHELL; G. R.	159
upon fever—H. C. Wood	857
	201
upon venom of the rattlesnake—S. W. MITCHELL	
Residuary legacy—See Smithson.	
Resolutions of Congress—See Congress.	
Resolutions of Regents—See each annual report.	
Resonance, electric, of mountains, observations on the—H. DE SAUSSURE. (R. 1868)	224
Respiration, apparatus for testing results of—M. Pettenkofer. (R. 1864)	188

Respiration in the Chelonia, researches on—S. W. MITCHELL; G. R. MORE-HOUSE
Results of—
magnetic survey of Pennsylvania—A. D. BACHE 166
meteorological observations at Brunswick, Maine-P. CLEAVELAND 204
meteorological observations at Marietta, Ohio—S. P. HILDRETH; J. WOOD
meteorological observations at Providence, Rhode Island, 1831-1860,
1881-1876—A. CASWELL 103, 443
meteorological observations in the United States, 1854-1859 157, 182
precipitation in rain and snow in United States—C. A. Schott 222, 353
spectrum analysis applied to heavenly bodies—W. Huggins. (R. 1866.) 214
RETZIUS, A. Present state of ethnology in relation to form of the skull. (R. 1859)
Review of—
American birds. Part I. North and Middle America—S. F. BAIRD 181
American species of genus Scops—R. RIDGWAY. (P. 1878) 332
classification of birds—W. LILLJEBORG. (R. 1865) 209, 364
cretaceous reptiles of the United States—J. LEIDY. (R. 1864) 188
genera and species of family Centrarchide—C. L. McKAY. (P. 1881) 467
genus Centurus—R. RIDGWAY. (P. 1881) 467
pleuronectids of San Francisco—W. N. Lockington. (P. 1879) 833
Rafinesque's memoirs on North American fishes—D. S. JORDAN 305
Revisions of nomenclature of North American birds—R. RIDGWAY. (P. 1880.) 425
Revolutions of crust of the earth—G. PILAR. (R. 1876)
REYE, T. Rain-fall and barometric minima. (R. 1877) 823, 898
REYNOLDS, J. E. Hydrogen as gas and metal. (R. 1870) 24
Rhees, W. J.—
Catalogue and index of publications of the Smithsonian Institution, U.S. National Museum, etc., 1846–1881 478
James Smithson and his bequest. (R. 1879) 345, 330
Journals of Board of Regents, reports, statistics, etc., of Smithsonian Institution
List of public libraries, institutions, and societies in United States and British Provinces116
List of public libraries, institutions, and societies in United States in cor-
respondence with Smithsonian Institution 238
Origin and history of Smithsonian Institution
Scientific writings of James Smithson, (edited by)327
Rheometry, electrical, researches on—A. SECCHI
Rhinobatidæ, American, synopsis and descriptions of—S. GARMAN. (P. 1880.) 425
Rhinonemus caudacuta, identity of with Gadus cimbrius—G. B. Goode; T. H. BEAN. (P. 1878)

method of ascertaining amount of water in—A. A. Humphreys. (R. 1858)

Roads and bridges, lectures on—F. Rogers. (R. 1860; R. 1861)_____ 147, 149

109

Robertson, R. S.—
Age of stone and troglodytes of Breckenridge county, Kentucky. (R. 1874)
Antiquities of Allen and De Kalb counties, Indiana. (R. 1874) 28
Antiquities of La Porte county, Indiana. (R. 1874) 28
Antiquities of Nashville, Tennessee. (R. 1877) 82
Robinson, E., and others. On publication of Spanish works on New Mexico. (R. 1855)
Robinson, E., and others. On publication of Squier and Davis' ancient monuments
Rock Bluff, Illinois, description of human skull from-J. A. MEIGS. (R. 1867.) 21
Rock-oil, or petroleum, history of-T. S. HUNT. (R. 1861)14
Rock river valley, Illinois, mound builders in-J. SHAW. (R. 1877) 82
Rocks—
catalogue of—J. W. Foster. (R. 1854)
catalogue of—C. T. Jackson. (R. 1854)
catalogue of—J. Locke. (R. 1854)
catalogue of—J. D. WHITNEY. (R. 1854)
crystalline, metamorphism, and formation of—G. A. DAUBBÉE. (R. 1861.)
in Green river valley, Indian engravings on face of—J. G. Bruff. (R. 1872)
Rocky Mountain goat, habits of the-J. C. MEBRILL. (P. 1879) 33
Rodent, rare, (Cricetodipus parvus)-F. W. TRUE. (P. 1881) 46
ROEHRIG, F. L. O. Language of Dakota or Sioux Indians. (R. 1871) 249, 376
ROGERS, F. Lectures on roads and bridges. (R. 1860; R. 1861) 147, 149
Rogers, W. B. Memorial address on Joseph Henry
Rome, Pontifical Academy—See Prize questions.
Romer, F. F. Prehistoric antiquities of Hungary. (B. 1876) 299, 440, 392
Romero, M. Explorations of John Xantus in Mexico. (B. 1862)
Rosing, J. Exchange system. (R. 1865)
Ross, B. R. Eastern Tinneh Indians. (R. 1866) 214, 365
Ross, B. R. Observations in Hudson's Bay Territory. (B. 1859)
Rotary motion, problems of—J. G. BARNARD 240
ROTHBOCK, J. T. Sketch of flora of Alaska. (R. 1867) 215, 367
Rotterdam, Batavian Society of Experimental Philosophy of. Prize questions. (R. 1861)
Royal—
Academy of Netherlands. Prize questions. (R. 1861)
Academy of Science, Madrid. On exchanges. (R. 1861) 149
Danish Society of Sciences. Prize questions. (R. 1862, 1865, 1867.) 150, 209, 215
Horticultural Society, London. Exchange of publications. (R. 1861) 149
Institution of Great Britain, history of—E. MAILLY. (R. 1867) 216
Prussian Academy of Sciences. Prize questions. (R. 1864) 1864

,
Royal—Continued.
Scientific and Literary Institute of Lombardy. Prize questions. (R. 1865.) 209
Society of London, list of Smithson's papers presented to. (R. 1858.) 67, 880
Society of London, notice of Smithson, by D. Gilbert, president of. (R. 1858) 67, 880
Society of London, origin and history of-C. A. ALEXANDER. (R. 1868.) 187
Society of Victoria, address of president of—R. L. J. Ellery. (R. 1868.) 224
ROYCE, C. C. Cessions of land by Indian tribes to United States. (E. 1879-80.) 476
Ruin, ancient, in Arizons-J. C. Y. LEE. (R. 1872) 271
Ruins-
at Savannah, Tennessee-J. P. Stelle. (R. 1870) 244
in Hardin county, Tennessee—J. P. Stelle. (R. 1870) 244
in White river cañon, Pima county, Arizona-R. T. Burr. (R. 1879.) 845
Rules-
for cataloguing libraries—C. C. JEWETT 47
for examination of specimens. (R. 1880) 442
of distribution of Smithsonian publications290
RUNKLE, J. D. New tables for determining perturbation of planets 79
RUNKLE, J. D. Supplement for asteroid perturbations 94
Rupert's Land, Institute of, circular of the. (R. 1861) 149
Rush, R.—
correspondence of, relative to Smithson828
letter from, relative to James Smithson. (R. 1853) 67, 328
notice of, by J. A. PEARCE. (R. 1859)
RUSSELL, R. Lectures on meteorology, with notes by J. HENRY. (R. 1854) 75
Russia, meteorology in—A. Woeikoff. (R. 1872) 271
Russian America—
ethnological investigations in, suggestions for-G. GIBBS 207
suggestions relative to objects of scientific investigation in—J. HENRY 207
Tinneh Indians of—G. GIBBS; B. Ross; W. L. HARDISTY; S. JONES. (R. 1866)
Yukon, journey to the—W. W. KIBBY. (R. 1864) 188
RÜTIMEYER, L. Fauna of middle Europe during the stone age. (R. 1861). 149 RYDER, J. A.—
Camaraphysema, a new type of sponge, (P. 1880)
List of North American species of myriapods belonging to family of Lysiopetalidæ, with description of blind form from Luray Cave, Virginia. (P. 1880)
On Semper's method of making dry preparations. (P. 1881) 467

S.

Sabine, E. Magnetic storms. (R. 1860) 1	47
SABINE, E. On continuance of magnetic observations. (B. 1858) 1	09
Sable Island Bank, description of Argentina syrtensium, new deep-sea fish from-	
G. B. GOODE; T. H. BEAN. (P. 1878)	82
Sacken, Osten—See Osten Sacken.	
Sacramento, California—	
meteorological observations at—F. W. HATCH. (R. 1854)	
meteorological observations at—T. M. Logan. (R. 1854)	75
meteorology of-T. M. Logan. (R. 1857)	107
Sacramento river, description of new species of Ptychochilus (Ptychochilus Harfordi) from—D. S. JORDAN; C. H. GILBERT. (P. 1881)4	16 7
Saint Augustine, Florida, catalogue of casts of heads of Indian prisoners at— R. H. Pratt. (P. 1878)	:32
Saint Christopher, West Indies, description of new subspecies of saxigilla from—G. N. LAWRENCE. (P. 1881)	
Saint Croix Island, flora of-H. F. A. Eggers	313
Saint George's Banks, occurrence of Hippocampus antiquorum on—G. B. Goods. (P. 1878))32
Saint Hilaire, Étienne Geoffroy, memoir of, by M. FLOUBENS. (R. 1861) 1	
Saint Hilaire, Isidore Geoffroy, memoir of-J. L. QUATREFAGES. (R. 1862) 1	50
Saint John's river, Florida, catalogue of fishes of-G. B. Goode. (P. 1879) 8	33
Saint John's river, Florida, shell-heaps at mouth of—S. P. MAYBERRY. (R. 1877)	<u>523</u>
Saint Louis, Missouri, ancient mound in—T. R. PEALE. (R. 1861)1	
Saint Louis, Missouri, temperature of—A. FENDLER. (R. 1860)1	47
Saint Martin, Isle Jesus, Canada East, description of observatory at—C. SMALL-WOOD. (R. 1856)	91
Saint Michael's, Alaska, occurrence of Hippoglossus vulgaris at—T. H. Bran. (P. 1879)	
Saint Michael's, Alaska, occurrence of Stichous punctatus at—T. H. Bran. (P. 1878)	
Saint Petersburg Academy of Sciences. Exchange system. (R. 1867) 2	
Saint Thomas—	
earthquakes in—G. A. LATIMER. (R. 1867)	
hurricane in the island of—G. A. LATIMER. (R. 1867) 2	15
whirlwind at, August 21, 1871—A. Colding. (B. 1877) 328, 8	98
Saint Vincent, catalogue of Ober's collection of birds of—G. N. LAWRENCE. (P. 1878)	32
Salamander—	
manife of a species of to mann. (The took)	75
inhabiting Wisconsin—P. R. Hoy. (R. 1854)	
or pouched rat, of Georgia, habits of—W. GESNER. (R. 1860)	47

Bran. (P. 1878)
Savannah river, primitive manufacture of spear and arrow points along line of— C. C. Jones, Jr. (R. 1879)
Sawkins, J. G., drawings of Zapotec remains by
SAWKINS, J. G.; WALL, G. P. Economic geology of Trinidad. (R. 1856)
Scenery, catalogue of sketches of-J. M. STANLEY
Scheme for qualitative determinations by the blow-pipe—T. EGLESTON. (R. 1872)
Scheme of classification for collections of U.S. National Museum—G.B. Goode. (P. 1881)467,
SCHERZER; SCHWARZ. Table of anthropological measurements. (R. 1866)
SCHLAGINTWEIT, H. DE. Ethnographical collections. (R. 1862)
Schlagintweit ethnographic collection, account of the—H. ZISGENBALS. (R. 1867)
Schleiden, R. Free freight between Germany and United States by North German Lloyd. (R. 1858)
Schoenbein, C. F., notice of, by J. HENRY. (R. 1868)
Schoharie, New York, Indian relics from-F. D Andrews. (R. 1879)
Schoodic Lakes, Maine, description of new species of Gasterosteus from—T. H. BEAN. (P. 1879)
SCHOTT, A. Remarks on ancient relic of Maya sculpture. (R. 1871)
Schott, A. Remarks on the Cara gigantesca in Yucatan. (R. 1869)
SCHOTT, A.; MASON, O. T. Leipsic Museum of Ethnology. (R. 1878)
SCHOTT, C. A.—
Base-chart of the United States
Discussion of Caswell's meteorological observations at Providence, Rhode Island
Discussion of Cleaveland's meteorological observations at Brunswick, Maine
Discussion of Hayes' physical observations in the Arctic Seas
Discussion of Hildreth's and Wood's meteorological observations at Marietta, Ohio
Discussion of Kane's astronomical observations in the Arctic Seas
Discussion of Kane's magnetic observations in the Arctic Seas
Discussion of Kane's meteorological observations in the Arctic Seas
Discussion of Kane's physical observations in the Arctic Seas
Discussion of Kane's tidal observations in the Arctic Seas
Discussion of McClintock's meteorological observations in the Arctic
Discussion of Smith's meteorological observations made near Washing- ton, Arkansas
Tables, distribution, and variation of atmospheric temperature
Tables of rain and snow in the United States 222.

SCHOTT, C. A.; EVERETT, J. D. Underground temperature. (R. 1874)	280
Schulze—See Cluss & Schulze.	
SCHUMACHER, P. Ancient graves and shell-heaps of California. (R. 1874)	286
SCHUMACHER, P. Kjökken-möddings on the northwest coast of North America. (R. 1878)	276
	214
Science—	,
	286
in general, relation of the physical sciences to—H. HELMHOLTZ. (R. 1871.)	248
recent progress in—See Record.	
trust fund for promotion of, in the United States-J. TYNDALL. (R. 1872)	271
Science, Agriculture, and Arts, Society of, Lille—See Prize questions.	
Science, Art, and Literature, Society of, Hainaut—See Prize questions.	
Sciences, Harlem Society of—See Prize questions.	
Sciences, Letters, and Arts, Society of, Dunkirk—See Prize questions.	
Sciences, Natural, Society of, Cherbourg—See Prize questions.	
Sciences, Royal Danish Society of—See Prize questions.	
Scientific—	
character and researches of Smithson-W. R. Johnson	B27
Congress of Carlsruhe, 1858-F. J. Nicklés. (R. 1860)	147
cooperation—University of Toronto. (R. 1861)	149
education of mechanics and artisans—A. P. PEABODY. (R. 1872) 271, 3	80
exchanges—See Exchanges.	
expedition to Mexico by French Government. (R. 1864) 1	188
instructions to Capt. Hall—J. HENBY; J. E. HILGARD; S. NEWCOMB; S. F. BAIRD; F. B. MEEK; L. AGASSIZ. (R. 1871)	249
investigation in Russian America, suggestions relative to-J. HENRY 20	
labors of Edward Lartet-F. FISCHER. (R. 1872.) 2	271
observations, on metric system for-A. Guyor. (R. 1848)	1
progress, reports of. (R. 1880) 442, 427, 428, 429, 430, 431, 48	
researches of J. C. A. Peltier—F. A. Peltier. (R. 1867) 2	215
societies, local, organization of—J. HENRY. (R. 1875) 2	298
writings of Sir William Herschel—E. S. Holden; C. S. Hastings. (R. 1880)	26
	327
Scientific and Literary Institute of Lombardy—See Prize questions.	
Scintillation of the stars—C. Dufour; Kämtz. (R. 1861)	149
Scops, review of American species of genus—R. RIDGWAY. (P. 1878) 3	
SCUDDER, S. H.—	
	189
Catalogue of orthoptera of North America	89
Nomenclator zoölogicus. An alphabetical list of generic names of recent and fossil animals	

SCUDDER, S. H., and others, Insects of Arctic America	342
Sculpture, Maya, remarks on ancient relic of—A. Schott. (R. 1871)	249
Sculptures of Santa Lucia Cosumalwhuapa in Guatemala—S. HABEL	269
Scytalina cerdale, new species of fish from Neah Bay, Washington Territory,	
description of—D. S. JORDAN; C. H. GILBERT. (P. 1880).	
Sea, gradual approach of, upon land—S. P. MAYBERRY. (R. 1867)	
Seas, the northern—J. BABINET. (R. 1869)	
Seaton, W. W., memoir of, by J. HENRY. (R. 1866)	214
Sebastichthys carnatus, new species of rockfish from coast of California, description of—D. S. JORDAN; C. H. GILBERT. (P. 1880)	425
Sebastichthys chrysomelas, new species of rockfish from coast of California, description of—D. S. JORDAN; C. H. GILBERT. (P. 1880)	425
Sebastichthys entomelas, new species of Sebastichthys from Monterey Bay, California, description of—D. S. JORDAN; C. H. GILBERT. (P. 1880)	425
Sebastichthys maliger, new scorpsenoid fish from coast of California, description of—D. S. JORDAN; C. H. GILBERT. (P. 1880)	
Sebastichthys miniatus, new species of Sebastichthys from Monterey Bay, Cali-	
fornia, description of—D. S. JORDAN; C. H. GILBERT. (P. 1880)	425
•	
Sebastichthys proriger, new scorpsenoid fish from Monterey Bay, California, description of—D. S. JORDAN; C. H. GILBERT. (P. 1880)	425
Sebastichthys rhodochloris, new species of Sebastichthys from Monterey Bay, California, description of—D. S. JOBDAN; C. H. GILBERT. (P. 1880)	
Sebastichthys serriceps, new species of rock cod from coast of California, description of—D. S. JOBDAN; C. H. GILBERT. (P. 1880)	
Sebastoid fishes, description of seven new species of—D. S. JORDAN; C. H. GIL- BERT. (P. 1880)	
SECCHI, A. Researches on electrical rheometry	
Secretary of Smithsonian Institution, report of—See Henry, J., Baird, S. F.	
Secretary of Treasury, letter to, on payment of interest in coin-J. HENRY. (R.	
	209
Secular period of aurora borealis—D. Olmsted	81
Secular variations of elements of planetary orbits—J. N. STOCKWELL. (R. 1871)	232
Seismograph, electro-magnetic-Prof. PALMIERI. (R. 1870)	244
Sema, note on—D. S. Jordan. (P. 1880)	425
Semper's method of making dry preparations—J. A. RYDER. (P. 1881)	467
Senate Judiciary Committee, report of, on management of Smithsonian—A. P. BUTLER. (R. 1855)	77
Senses, the. Sense of feeling, sense of smell. (R. 1865)	209
Senses, the. Sense of taste, sense of hearing, sense of sight. (R. 1866)	214
Séquard—See Brown-Séquard.	
Series, converging, expressing ratio of diameter and circumference of a circle— W. FERREL	233

Seriola Stearnsii, new species of amber fish from Pensacola, Florida, descrip-
tion of—G. B. Goode. (P. 1879)
Sermon at funeral of Joseph Henry—S. S. MITCHELL 356
Serpent venom, bibliography of—S. W. MITCHELL 185
Serpents, catalogue of—S. F. BAIRD; C. GIRARD
Serpents, lecture on nature and cure of bites of—D. Brainard. (R. 1854) 75
Seven cities of Cibola, Coronado's march in search of the—J. H. SIMPSON. (R. 1869)
Sexual characters of eels—S. T. CATTIE. (P. 1880)425
Shad, western gizzard, Dorosoma cepedianum heterurum, notes on—S. WILMOT. (P. 1878)
SHAKESPEARE, E. O. The nature of reparatory inflammation in arteries after ligature, etc. Toner lecture No. VII
Shark, the oil, of southern California, (Galeorhinus galeus)—D. S. JORDAN; C. H. GILBERT. (P. 1880)
SHARPLESS, T.; PATTERSON, R. Phonography. (R. 1856) 91
SHAW, J. Mound-builders in Rock river valley, Illinois. (R. 1877) 828
SHEA, J. G. Account of library of Indian linguistics. (R. 1861) 149
Shea's Indian linguistics, recommendation of-G. GIBBS and others. (R. 1861.) 149
Shell-bed skull—A. S. TIFFANY. (R. 1874)
Shell-deposits, artificial, of the United States-D. G. BRINTON. (R. 1866.) 214, 440
Shell-deposits, artificial, in New Jersey-C. RAU. (R. 1864) 188, 862
Shell engravings granted by British Museum. (R. 1863) 187
Shellfish and their allies, lectures on-P. P. CARPENTER. (R. 1860) 147, 152
Shell-heap in Georgia-D. Brown. (R. 1871) 249
Shell-heaps-
and ancient graves of California—P. SCHUMACHER. (R. 1874) 286
at mouth of St. John's river, Florida—S. P. MAYBERRY. (R. 1877) 828
in New Brunswick-J. Fowler. (R. 1870) 244
of Tampa Bay, Florida—S. T. WALKER. (R. 1879) 845
on Mobile river, Alabama—A. S. GAINES; K. M. CUNNINGHAM. (R. 1877)828
Shell-mounds—See Mounds.
Shells-
acknowledgment for-W. H. Prase. (R. 1861) 149
bibliography of. Parts I, II-W. G. BINNEY 142, 174
duplicate, collected by United States exploring expedition—C. WILKES. 193
from California, description of new species of, in U.S. National Museum— W. H. Dall. (P. 1878)832
from Costa Rica kitchen-midden-W. H. Dall. (P. 1878)
land and fresh-water, instructions for collecting—J. LEWIS. (R. 1866) 214, 363
land and fresh-water, of North America:
Part I. Pulmonata geophila—W. G. BINNEY; T. BLAND 194

Shells—Continued.
land and fresh-water, of North America:
Part II. Pulmonata limnophila and Thallasophila—W. G. BINNEY. 14
Part III. Ampullariidæ, Valvatidæ, (etc.)—W. G. BINNEY 19
Part IV. Strepomatidæ (American melanians)-G. W. TRYON, Jr 25
letter on—J. Lewis 2
North American, check-list of-I. Lea; P. P. CARPENTER; W. STIMP-
son; W. G. Binney; T. Prime
North American, circular in reference to collecting 17
of Gulf of California, lectures on the-P. P. CARPENTER. (B. 1859) 11
of Panama, collected by C. B. Adams, descriptions of new species and
varieties of Chitonidæ and Acmæidæ in-P. P. CARPENTER 2
presented to Academy of Natural Sciences, report on—J. LEIDY; G. W. TRYON. (R. 1865)
request for duplicate—W. E. Logan. (R. 1859)
Shells—See Mollusca.
SHEPARD, C. U. Arrangement of mineralogical collection. (R. 1861)
SHEPARD, E. M. Deposit of arrow-heads near Fishkill, New York. (R. 1877.) 33
SHERMAN, W. T. Memorial address on J. Henry
SHERMAN, W. T. Report of National Museum Building Commission, 1879, 1880. (R. 1879; R. 1880) 345, 442, 409, 43
Shetland, vegetable colonization of—C. MARTINS. (R. 1858)
Shipping fresh fish and other animals, circular relative to—S. F. BAIRD 36
Shoal Lake, Lake Winnipeg, egging expedition to-D. Gunn. (R. 1867) 2
Shooting stars, observations on—S. MASTERMAN. (R. 1857)
Short memoirs on meteorological subjects, translated by C. ABBE. (R. 1877.) 328, 3
Shoshone Indians, religion, superstitions, and manners of—A. G. BRACKETT. (R. 1879)
SHUFELDT, R. W. Remarks upon osteology of Opheosaurus ventralis. (P. 1881.)
Siberia, descriptions of new fishes from—T. H. BEAN. (P. 1881)
SIBLEY, H. Meteorological telegrams by Western Union Telegraph Company. (R. 1862)
Sight, sense of. (R. 1866)
Sign language among North American Indians—G. MALLERY. (E. 1879-80.)
Sign language of deaf mutes-G. MALLERY. (E. 1879-80)
Siluridæ of fresh waters of North America, synopsis of-D. S. JOBDAN 30
Silver coins, assay of, at United States Mint-J. Pollock. (R. 1868) 22
Silver coins, foreign, table of. (R. 1868)
Silvered-glass telescope, on construction of—H. DRAPER18
SIMPSON, G. Letter to officers of Hudson's Bay Company relative to scientific researches
SIMPSON, J. H. Coronado's march in search of the seven cities of Cibola. (R. 1869)

Description of new species of gobiesox (Gobiesox rhessodon) from San

Occurrence of species of cremnobates at San Diego, California. (P. 1880.)

467

425

Diego, California. (P. 1881)

Sioux—See Dakota.

Sketch of—

Skull—

SMITH, R.—

SMITH, S. B. On the Chinook names of salmon in Columbia river. (P. 1881.)
Smith, S. I.—
Crustaceans of Kerguelen Island
Notice of crustacea dredged off south coast of New England in 1880. (P. 1880)
Notice of new species of Willemæsia group of crustacea (recent Eryontidæ.) (P. 1879)
Occurrence of Chelura terebrans (crustacean destructive to timber) on coast of United States. (P. 1879)
Smithson fund, memorial of Regents to Congress relative to the. (R. 1850)
Smithson, James—
act of Congress to receive residuary legacy of. (R. 1866) 214,
catalogue of library of
character and researches of-W. R. Johnson
contributions to Annals of Philosophy by. (R. 1853) 67,
exposition of bequest of-J. HENRY
letter from Richard Rush relative to. (R. 1858) 67,
letters from Fladgate, Clark, and Finch, relative to. (R. 1861) 149,
life of, by W. J. RHEES. (R. 1879) 845, 3
list of papers presented to Royal Society by. (B. 1858) 67,
notice of, by D. Gilbert, president of the Royal Society. (R. 1858) 67,
personal effects of 828,
scientific writings of
will of. (R. 1858) 67, 328, 1
will of, fac-simile of
works and character of—J. R. McD. IRBY
Smithson, James, and his bequest—W. J. RHEES. (R. 1879) 845, 3
Smithsonian—
building, report on use of new hall in—L. Agassiz. (R. 1867) 2
Bulletin No. 1. Vocabulary of the jargon
Contributions to Knowledge—See Contributions.
Miscellaneous Collections—See Miscellaneous Collections.
publications—See Publications.
publications, catalogue and index of—W. J. RHEES 47
•
Reports—See Reports.
Smithsonian Institution— act to establish—See Congress.
address on the—J. Henry. (R. 1858)
business arrangements of the 32 directory of officers, collaborators, and employés of the 46
establishment and officers of the
Journals of Regents, reports of committees, statistics, etc.—W. J. RHEES. 32
magnetic observatory at, description of—J. E. HILGARD. (R. 1859) 11

Smithsonian Institution—Continued.	
origin and history of-W. J. RHEES.	328
report of American Academy of Arts and Sciences, on the. (R. 1858)	67
report of Senate Judiciary Committee on management of-A. P. BUTLER.	
(R. 1855)	77
views and plans of the-R. D. Owen	P
Snake Indians, religion, superstition, and manners of—A. G. BRACKETT. (R. 1879)	845
SNELL, E. S. Lecture on planetary disturbances. (R. 1855)	77
Snell's barometric observations, discussion of—F. H. Loud. (R. 1880) 442,	435
Snow-	
charts—See Rain.	
electricity during fall of-F. Zantedeschi. (R. 1870)	244
gauge-W. E. Guest. (R. 1858)	109
gauges-R. H. GARDINER. (R. 1858)	109
line, Norwegian-O. E. DREUTZER. (R. 1866)	214
observations of	157
tables of precipitation in—C. A. SCHOTT	353
SNYDER, J. F. Deposit of stone implements in Illinois. (R. 1876)	299
Social and religious condition of the lower races of man-J. LUBBOCK. (R. 1869.)	228
Societies—	
list of, in United States and British Provinces—W. J. RHEES	116
local scientific, organization of-J. HENRY. (R. 1875)	2 98
proceedings of, relative to Joseph Henry	856
publications of, in Smithsonian library, 1854, 1856, 1858, 1866. 73, 85, 117,	179
Societies—See Holland, Belgium, Publications.	
Society-	
Batavian, of Experimental Philosophy, Rotterdam. Prize questions. (R. 1861)	149
Bath and West of England, for Encouragement of Agricultural Arts, etc. Exchange system. (R. 1867)	215
for Encouragement of Science, Dunkirk—See Prize questions.	
Society of—	
Arts and Sciences, Utrecht. Prize questions. (R. 1861)	149
Natural History, Portland, Maine, account of—E. C. Bolles. (R. 1867.)	215
Natural Sciences, Cherbourg—See Prize questions.	
Physics and Natural History of Geneva—See Geneva.	
Science, Agriculture, and Arts of Lille—See Prize questions.	
Science, Art, and Literature, Hainaut—See Prize questions.	
Sciences, Harlem—See Prize questions.	
Sciences, Letters, and Arts, Dunkirk—See Prize questions.	
Sciences, Royal Danish—See Prize questions.	
Victoria, address of president—R. L. J. ELLERY. (R. 1868)	224

Source, L. Law of variation of temperature in ascending moist currents of air. (R. 1877)	
Solar—	,
diurnal variation in magnetic declination—A. D. BACHE	i, 121
diurnal variation in vertical component of magnetic force—A. D. BACHE.	175
eclipse of April 25, 1865—Baron DE PRADOS. (R. 1864)	
eclipse of July 18, 1860—J. LAMONT. (R. 1864)	
system, harmonies of the—S. ALEXANDER	
Solids, specific heat of—F. W. CLARKE	
Solids, tables of expansion of, by heat—F. W. CLARKE	
Solitary wasps, synopsis of—H. DE SAUSSURE	
Sonntag, A. Observations on terrestrial magnetism in Mexico	
Sonora, plants collected in, by C. Wright—A. GRAY	
Sound, refraction of—W. B. TAYLOR. (R. 1875)	
Sound, researches in—J. HENRY. (R. 1878)841	
Soundings, microscopical examination of—J. W. BAILEY 2	
Sounds, vocal, of Laura Bridgman—F. LIEBEB	
South America—	
birds of, not in U. S. National Museum—R. RIDGWAY. (P. 1881)	457
Caracas, meteorology of—G. A. Ernst. (R. 1867)	
circular relative to collections of birds from	
rain-full in—C. A. Schott22	
travels in—S. Habel	
South Carolina—	
description of new hake from—T. H. BEAN. (P. 1880)	425
fishes of—D. S. Jordan; A. W. Brayton	
microscopical observations in—J. W. BAILEY	
Southern Baptist Missionary Society. Yoruba grammar and dictionary	98
Southern States, antiquities in the—H. C. WILLIAMS. (R. 1880)	244
Southern States, and quities in the—II. O. WILLIAMS. (II. 1000) Southern States, microscopical observations in the—J. W. BAILEY	
Space and time, lecture on the relations of—S. ALEXANDER. (R. 1861)	
SPAINHOUR, J. Antiquities in Lenoir county, North Carolina. (R. 1871)	
Spanish vocabulary, with comparative words in English, French, and Latin	
Spanish works on New Mexico, by Buckingham Smith, communications relative to publication of. (R. 1855)	
SPARKS, J., and others. On publication of Spanish works on New Mexico. (R. 1855)	
SPARKS, J., and others. Report of American Academy of Art and Sciences on organization of Smithsonian Institution. (R. 1858)	
Aparoid fish, new, (Sparus brachysomus,) from lower California—W. N. Lock- INGTON. (P. 1880)	
Sparrows sent to United States—W. A. LLOYD. (R. 1867)	215

Sparus brachysomus, new sparoid fish, from lower California, description of—	
W. N. Lockington. (P. 1880)	425
Specific gravity—	
bibliography of-F. W. CLARKE	255
of sea water—D. Walker	146
tables. Constants of Nature, Part I-F. W. CLARKE	255
tables. First supplement to Constants of Nature, Part I-F. W. CLARKE.	288
Specific heat, bibliography of-F. W. CLARKE	276
Specific heat tables. Constants of Nature, Part II-F. W. CLARKE	276
Specimens—	
acknowledged by British Museum. (R. 1865)	209
acknowledged by Chicago Academy of Sciences. (R. 1867)	215
authority to collectors of customs to receive and transmit to Smithsonian Institution	84
authority to naval officers to receive and transmit to Smithsonian Institution	84
authority to quartermasters of War Department to receive and transmit	
to Smithsonian Institution	84
derived from expeditions—S. F. BAIRD. (R. 1867)	215
exchange of—Hamburg Zoölogical Gardens. (R. 1867)	215
exchange of—W. A. Lloyd. (R. 1867)	215
exchange of—University of Costa Rica. (R. 1867)	215
exchange of—University of Greece. (R. 1867)	215
from Brazil, presented—M. M. Lisboa. (R. 1865)	209
large myological, rapid preparation of—F. PLATEAU. (P. 1881)	467
of flamingo from south Florida, presented—G. Wurdeman. (R. 1860.)	147
of natural history, directions for collecting, preserving, and transport-	721
ing—S. F. Baird. (R. 1856) 91	. 34
presented, circular sent with—J. HENRY. (R. 1872)	271
receipts and distribution of. (R. 1880)	442
rules for examination of. (R. 1880)	442
Spectrum analysis applied to the heavenly bodies—W. Huggins. (R. 1866)	214
Spencer's telescope, examination of, for Hamilton College. (R. 1855)	77
	80
Spheres, tangencies of—B. ALVORD	
Spleen, observations on—J. JONES	82
Sponge, a new type of, (Camaraphyeema)—J. A. RYDER. (P. 1880)	425
Spruce Creek valley, Pennsylvania, account of tornado in—J. B. MEEK. (R. 1871)	249
Squalius aliciæ from Utah Lake—P. L. Jour. (P. 1881)	467
SQUIER, E. G. Aboriginal monuments of State of New York	15
SQUIER, E. G. Antiquities of Nicaragua. (R. 1850)	28
SQUIER, E. G., and others. On publication of Spanish works on New Mexico. (R. 1855)	77

SQUIER, E. G.; DAVIS, C. H. Ancient monuments of the Mississippi valley.
Squier, E. G.; Davis, C. H., correspondence relative to memoir by. (R. 1847.) H, E
Stanley's gallery of Indian portraits, report of Committee of Regents on the. (R. 1857)
STANLEY, J. M. Catalogue of portraits of North American Indians 53
STANTON, E. M. Deposit of Beaufort Library. (R. 1862) 150
Star clusters, bibliography of—E. S. HOLDEN
Stars-
index-catalogue of—E. S. HOLDEN 311
movement of, around a central point—J. H. MAEDLER. (R. 1859) 110
near the North Pole, map of, for observations on auroras 350
scintillation of the—C. DUFOUB; KÄMTZ. (R. 1861)
State Department. Circular to diplomatic agents relative to Morgan's research. 13
Statement and exposition of harmonies of the solar system—S. ALEXANDER 280
Stations, meteorological, of Smithsonian Institution (See, also, List) 373
Statistics of—
British Museum—S. F. BAIRD. (R. 1850) 28
Norwegian mountains, lakes, and snow line-O. E. DREUTZER. (R. 1866.) 214
Smithsonian Institution, finances, exchanges, etc., 1846-1877 329
Statue of Joseph Henry, proceedings of Congress relative to 356
Statues, on plaster casts of—W. J. STONE. (R. 1855)
STEARNS, J. Translation of Babinet on diamonds and precious stones. (R. 1870)
Stearns, S., catalogue of fishes from Pensacola, collected by—G. B. Goode; T. H. Bean. (P. 1879)
Stearns, S., description of new species of amber fish (Seriola Stearnsii) obtained near Pensacola, by—G. B. Goode; T. H. Bran. (P. 1879) 353
STEARNS, S. Note on Gulf menhaden, Brevoortia patronus. (P. 1878) \$32
STEBBINS, R. P. Memoir of Louis Agassiz. (R. 1873) 275
STEINDACHNER, F. Note on Perca flavescens. (P. 1878) 332
STEJNEGER, L. Description of two new races of Myadestes obscurus. (P. 1881.) 467
STELLE, J. P. Account of aboriginal ruins in Hardin Co., Tennessee. (R. 1870.) 244
STELLE, J. P. Account of aboriginal ruins at Savannah, Tennessee. (R. 1870.) 244
STEPHENSON, M. F. Account of ancient mounds in Georgia. (R. 1870) 244
STEPHENSON, M. F. Mounds in Bartow county, Georgia. (R. 1872) 271
Stereotype catalogue of public libraries, report of Commissioners on. (R. 1850.) 28,47
Stereotyped titles for library catalogues—C. C. JEWETT
STEVENS, H. Prospectus of a bibliographia Americana. (R. 1848)
STEVENS, J. A. Memoir of George Gibbs. (R. 1873) 275
STEVENSON, J. Report of explorations in New Mexico and Arizona. (R. 1880.) 442
STEWART, B. Experiments on aneroid barometers at Kew Observatory. (R. 1868.) 224
Stichseus punctatus, occurrence of, at St. Michael's, Alaska—T. H. BEAN. (P.

STIMPSON, W.—	
Check-list of the shells of North America	28
On collection and preservation of marine invertebrates	84
Researches upon the Hydrobiinæ and allied forms 20	
Synopsis of the marine invertebrata of Grand Manan	
Stock-in-trade of an aboriginal lapidary—C. RAU. (R. 1877) 823, 440, 40	02
Stockholm Academy of Sciences. Acknowledgment of birds. (R. 1867) 2	
STOCKWELL, J. N. Secular variations of elements of orbits of the eight principal planets. (R. 1871)	32
STONE, W. J. On plaster casts of statues. (R. 1855)	77
Stone—	
celts in the West Indies and Africa—G. J. GIBBS. (R. 1877)	128
cists near Highland, Madison county, Illinois—A. OEHLER. (R. 1879.) 8	
drilling in, without metal—C. RAU. (R. 1868) 224, 440, 3	
image in Tennessee, account of discovery of-E. M. GRANT. (R. 1870.) 2	
implements, North American—C. RAU. (R. 1872) 271, 440, 36	
Stone age—	
and the troglodytes of Breckinridge county, Kentucky-R. S. ROBERT-	286
fauna of middle Europe during the-L. RÜTIMEYER. (R. 1861) 1	149
in New Jersey—C. C. Abbott. (R. 1875) 298, 39	94
Stone period, North American, agricultural implements of the—C. RAU. (R. 1868) 187, 4	
Stones, precious—J. BABINET. (R. 1870) 244, 8	377
Storm in Butler county, Kansas, June 28, 1871—W. HARRISON. (R. 1871) 2	
Storms—	
in Europe and America, December, 1836—E. Loomis	27
magnetic—E. Sabine. (R. 1860) 1	147
of 1859—J. H. Coffin1	182
thunder, instructions for observations of—J. HENRY 2	285
Strain and over-action of the heart. Toner lecture No. 111-J. M. Da Costa. 2	279
Strait of Juan de Fuca, Indians of-J. G. SWAN	220
Strait of Juan de Fuca, description of new species of Paralepis (Paralepis corus- cans) from—D. S. Jordan; C. H. Gilbert. (P. 1880)	125
STRANG, J. J. Natural History of Beaver Island, Michigan. (R. 1854)	75
STREETS, T. N. Contributions to natural history of Hawaiian Islands, Fanning	
Islands, and lower California	808
Strepomatidæ (American melanians)—G. W. TRYON 2	53
Stricklandia Davidsoni, note on the occurrence of, in Georgia—C. A. WHITE. (P. 1880)	12 5
Stricklandia Salteri, note on the occurrence of, in Georgia-C. A. WHITE.	12 5
STRONG, M. Antiquities in Wisconsin. (R. 1877)	328

STRONG, M. Observations on the prehistoric mounds of Grant county, Wisconsin. (R. 1876)
Structure, internal, of the earth—J. G. BARNARD
Structure of cancerous tumors. Toner lecture No. 1-J. J. Woodward
Structures, aboriginal, in Georgia-C. C. Jones, Jr. (R. 1877) 328
Study of—
chionis minor—J. H. KIDDER; E. COUES
high antiquity in Europe, lecture on—A. Monlot. (R. 1862; R. 1864.) 150
jade—S. Blondel. (R. 1876)
mortuary customs of North American Indians—H. C. YABROW. (E. 1879-80)
nature and mechanism of fever. Toner lecture No. IV—H. C. WOOD
phenomena of contact—T. L. Phipson. (R. 1862)
physiology of fever—H. C. Wood.
savage weapons at the Centennial—E. H. KNIGHT. (R. 1879) 845,
skulls and long bones from mounds in Illinois—R. J. FARQUHARSON. (R. 1874)
trunk fishes, Ostraciontidæ—G. B. Goode. (P. 1879)
Studies in Central American picture writing—E. S. Holden. (E. 1879-80)
Subcutaneous surgery. Toner lecture No. VI-W. ADAMS
Sudis ringens, new species of scopeloid fish from Santa Barbara channel, California, description of—D. S. JORDAN; C. H. GILBERT. (P. 1880)
Suess, E. Boundary line between geology and history. (R. 1872)
Suggestions—
for ethnological researches in Russian America—G. GIBBS:
for forming collections of birds' eggs—A. NEWTON
for meteorological observations in Russian America—J. HENRY
for sanitary drainage of Washington city—G. E. WARING
for saving parts of the skeletons of birds—A. NEWTON. (R. 1860)
relative to ethnological map of North America—L. H. Morgan. (R. 1861.
relative to objects of scientific investigations in Russian America—J. HENRY; S. F. BAIRD
to beginners in botany—L. F. WARD
Sulphurous acid, action of, upon putrefactive bacteria. (P. 1881)
Sumichrast, F., birds of Mexico, collected by—G. N. LAWRENCE
Summary of anthropological correspondence previous to 1880—O. T. MASON.
(R. 1879)
Sun— chemical analysis of the—A. LAUGEL. (R. 1861)
disk of the, external appearance of. (R. 1866)
eclipse of the, September 7, 1858—J. M. GILLISS
eclipse of the, March 15, 1868—T. HILL
eclipse of the, April 25, 1865—M. DE PRADOS. (R. 1864)
relative intensity of heat and light of the-L. W. MEECH. (R. 1856.) 91

OF SMITHSUNIAN PUBLICATIONS.	อบช
SUNDEBLAND, B. Prayer at memorial of Joseph Henry	356
Sun's distance, means available for correcting measures of the—G. B. AIRY. (R. 1859)	
Superstitions of Shoshone Indians—A. G. BRACKETT (R. 1879)	84 5
Supplement, asteroid, to new tables of planetary motion—J. D. RUNKLE	94
Surgery, subcutaneous. Toner lecture No. VI-W. ADAMS	302
Surgical complications and sequels of continued fevers. Toner lecture No. v—W. W. KEEN.	300
Survey—	
in Michigan, catalogue of rocks, minerals, and ores collected in the—O. T. Jackson. (R. 1854)	75
magnetic, of Pennsylvania and parts of adjacent States—A. D. BACHE.	166
of economic geology of Trinidad—G. P. Wall; J. G. Sawkins. (R. 1856)	91
of State of New York—E. G. SQUIER	15
Surveying expedition, North Pacific, natural history of—T. H. STREETS	303
Surveys, Government—See Explorations.	
SWAN, J. G.—	
Eulachon, or candlefish of northwest coast. (P. 1880)	425
Haidah Indians of Queen Charlotte's Islands, British Columbia	
Makah Indians of Cape Flattery, Strait of Fuca	220
Makah vocabulary	220
Surf smelt of the northwest coast, and mode of taking them by the Quillehute Indians of west coast of Washington Territory. (P. 1880.)	425
Sweden, Stockholm, Academy of Sciences. Acknowledgment of birds. (R. 1867.)	215
Switzerland—	
Concise, archæological researches made at—F. Troyon. (R. 1861)	149
Crania helvetica—F. Troyon. (R. 1864)	188
lacustrian cities of—F. Troyon. (R. 1861)	149
lacustrian settlements in, abstract of Dr. Keller's report on—A. MORLOT. (R. 1868)	187
Lake of Neuchâtel, palafittes or lacustrian constructions of—E. DESOR. (R. 1865)	360
Lausanne, Cantonal Museum at, antiquarian and ethnological collections of the—F. Troyon. (R. 1861)	149
lecture on-A. D. BACHE. (R. 1870)	244
Syllabus of a course of lectures on physics—J. Henry. (R. 1856)	91
Symbols for charts of prehistoric archeology, international code of—G. DE MORTILLET; E. CHANTRE. (R. 1875)	298
Symbols used by James Smithson	
Synopsis of—	
American Rhinobatidæ—S. GARMAN. (P. 1880)	425
American wasps, solitary wasps—H. DE SAUSSURE	254
catostomide—D. S. Jordan	808

Synopsis of—Continued.
marine invertebrata of Grand Manan—W. STIMPSON 50
North American lepidoptera—J. G. Morris———————————————————————————————————
North American neuroptera-H. HAGEN; P. R. UHLER 134
pediculate fishes of east coast of extra tropical North America—T. GILL. (P. 1878)———————————————————————————————————
scientific writings of Sir William Herschel—E. S. Holden; C. S. Has- TINGS. (R. 1880) 442, 426
siluride of fresh waters of North America—D. S. JORDAN 306
trochilidæ—D. G. Klliott
useful and injurious animals—G. B. GOODE
Synoptical tables of characters of mammals—T. GILL
Synthetic experiments relative to meteorites—G. A. DAUBRÉE. (R. 1868) 224
Syria, Beirut, sarcophagus from—A. A. HARWOOD. (R. 1870) 244
System—
meteorological, of Smithsonian Institution—E. FOREMAN. (R. 1851; R. 1852)
of accounts adopted by Board of Regents. (R. 1866) 214
of combined meteorological observations, report on, by committee of American Association for Advancement of Science. (R. 1851) 51
of numeration, improved, report on—W. B. TAYLOR. (R. 1867) 215
of weights and measures, new-G. H. KNIGHT. (R. 1867) 215
solar, harmonies of the—S. ALEXANDER 200
Systematic—
index of Smithsonian publications. (B. 1868) 224, 478
index to list of foreign correspondents257
list of batrachia and reptilia—E. D. Cope 292
review of classification of birds—W. LILLJEBORG. (B. 1865) 209, 364
Systems of consanguinity and affinity of the human family—L. H. Morgan. 218
Systems of relationship of Cros Indians_R A WATERTER (R 1969)

T.

Table of—
anthropological measurements-Scherzer; Schwarz. (R. 1866)
chemical equivalents of sixty-three elements. (R. 1864)
distribution of birds-G. N. LAWRENCE. (P. 1878)
foreign gold and silver coins. (R. 1868)
Tables-
asteroid—J. D. Runkle
barometrical—A. GUYOT
for conversion of centigrade degrees to Fahrenheit's scale. (R. 1863)
for determining values of coefficients in perturbative function of planetary motion which depend on ratio of mean distances—J. D. RUNKLE
hygrometrical—A. Guyor
hypsometrical—A. Guyor
meteorological—A. Guyor31,
metric—H. A. NEWTON. (B. 1865) 209,
miscellaneous—A. Guyot
mortality, methods of interpolation applied to. Parts I, II—E. L. DE FOREST. (R. 1871; R. 1878) 249,
of atmospheric temperature—C. A. Schott
of constants of nature and art-C. BABBAGE. (R. 1856)
of expansion by heat. Constants of Nature, Part III-F. W. CLARKE.
of meteorological corrections—A. GUYOT
of motion of Neptune—S. Newcomb
of motion of Uranus—S. Newcomb
of precipitation in rain and snow—C. A. Schott 222,
of variations of elements of orbits of the eight principal planets—J. N. STOCKWELL
of weights and measures, English and French. (R. 1863-1865.) 187, 188,
of winds of the globe-J. H. COFFIN; S. J. COFFIN
physical—A. Guyot
psychrometrical—J. H. Coffin
specific gravity. Constants of Nature, Part 1, and supplement—F. W. CLARKE
specific heat. Constants of Nature, Part II-F. W. CLARKE
thermometrical—A. Guyor
Tablet, Palenque—C. RAU 3
TACCHINI, P. Evaporation observed at Palermo in 1865 and 1866. (R. 1870.)
Tampa Bay, Florida, shell-heaps of—S. T. WALKER. (R. 1879)
Tangencies of circles and of spheres—B. ALVORD
Taste, sense of. (R. 1866)

Taxidermy, classification of collections of U.S. National Museum to illustrate	
art of-W. T. Hornaday. (P. 1881) 467, 45	6
TAYLOR, A. S. Grasshoppers and locusts of America. (R. 1858) 10)9
TAYLOR, F. W. Report of chemist of Smithsonian Institution. (R. 1880) 4	12
TAYLOR, W. B.—	
Henry and the telegraph. (R. 1878) 841, 40)5
Kinetic theories of gravitation. (R. 1876) 299, 39) 5
Memoir on the scientific work of Joseph Henry	19
Refraction of sound. (R. 1875)	98
Report on improved system of numeration. (R. 1867) 2	15
Thoughts on the nature and origin of force. (R. 1870) 244, 37	15
TAYLOR, W. M. Ancient mound in western Pennsylvania. (R. 1877) 3	23
Telegrams, astronomical, circular relative to—J. HENRY	3
Telegrams, meteorological, by Western Union Telegraph Company—H. Sibler.	
(R. 1862)	
Telegrams read at memorial of Joseph Henry	56
Telegraph—	
American fire-alarm, lecture on the—W. F. CHANNING. (R. 1854)	
electro-magnetic, deposition of J. Henry in relation to. (R. 1857) 107, 1	
electro-magnetic, Henry's contribution to—W. B. TAYLOR. (R. 1878.) 841, 40	
electro-magnetic, Morse's invention—W. B. TAYLOR. (R. 1878) 841, 40)5
electro-magnetic, proceedings of Board of Regents in relation to. (R. 1857)	15
Henry and the-W. B. TAYLOR. (R. 1878)	
history of the-J. HENRY. (R. 1857)	
influence of aurora on the-W. D. SARGENT. (R. 1870)	
lines, phenomena in, during auroras-G. B. Donati. (R. 1872) 2	
report of Committee of Regents on the. (R. 1857) 107, 115, 3	29
statement of S. P. Chase relative to the. (R. 1857) 107, 115, 3	29
statement of L. D. Gale relative to the. (R. 1857) 107, 115, 3	29
statement of James Hall relative to the. (R. 1857) 107, 115, 3	29
statement of J. Henry relative to history of. (R. 1857) 107, 115, 3	29
statement of Charles Mason relative to the. (R. 1857) 107, 115, 3	29
Telegraphic announcements of astronomical discoveries—J. HENRY 2	63
Telescope—	
Henry Draper's, account of-T. W. WEBB. (R. 1864)	88
silvered-glass, construction of, and its use in celestial photography—H.	
Draper	90
Spencer's, examination of, for Hamilton College, Clinton, N.Y. (R. 1855.)	77
Temperature—	
chart of United States for the year-O. A. Schott 381,3	88
charts of United States, summer, winter, and year-C. A. SCHOTT 3	87

Temperature—Continued.
in ascending currents of air, laws of variation of-J. HANN. (R. 1877.) 828, 898
in ascending moist currents of air, laws of variation of—L. Sohncke. (R. 1877) 828, 898
mean, best hours to find—C. DEWEY. (R. 1857; R. 1860) 107, 147
observations in the Arctic Seas—I. I. HAYES 196
observations in the Arctic Seas—E. K. KANE
observations in the Arctic Seas-F. L. McClintock 146
of St. Louis, Missouri-A. Fendler. (R. 1860)
tables for United States and adjacent parts of America—C. A. Schott 277
underground—C. A. Schott; J. D. Everett. (R. 1874) 286
Temperature—See Kane, Hayes, Meteorology, etc.
Tempests and tornadoes, distinction between-J. B. Lamarck. (R. 1871) 249
Tennessee
aboriginal remains of -J. Jones 259
Alleghany region of, fishes of—D. S. Jordan; A. W. Bratton 808
antiquities in-I. DILLE. (R. 1862)
antiquities in—E. O. DUNNING. (R. 1870)
antiquities of—W. M. CLARK. (R. 1877)
antiquities of-J. Jones 259
antiquities of—D. F. WRIGHT. (R. 1874) 286
Blount county, antiquities of-A. E. LAW. (R. 1874) 286
Chattanooga, ancient mound near-M. C. READ. (R. 1867) 218
East, mound in—A. F. DANILSEN. (R. 1868)
explorations in—E. A. DAYTON. (R. 1870)
Hardin county, aboriginal ruins in—J. P. STELLE. (R. 1870) 244
herbarium captured in—H. R. WIRTZ. (R. 1862)
Jackson county, antiquities of-J. HAILE; J. W. McHENRY. (R. 1874.) 286
Nashville, antiquities of—R. S. Robertson. (R. 1877) 328
Savannah, aboriginal ruins at-J. P. STELLE. (R. 1870) 244
stone image in, discovery of—E. M. GRANT. (R. 1870) 244
Terrestrial magnetism— observations on—W. HARKNESS
observations on—J. Locke 35
observations on—Baron von Müller; A. Sonntag 114
Terrestrial magnetism—See Bache, A. D.
Terrestrial physics, articles on. (R. 1870)
Tertiary fossils, presented—Imperial Geological Institute, Vienna. (R.
1863) 187
Testing building materials, on mode of—J. Henry. (R. 1856) 91
Testing results of perspiration and respiration, apparatus for—M. PETTENKOFER. (R. 1864)

Texas—	
hail storm in—G. M. BACHE. (R. 1870)	244
new cretaceous invertebrate fossils from, descriptions of—C. A. WHITE. (P. 1879)	833
ornithology of, notes on—J. C. MERRILL. (P. 1878)	832
plants of, collected by C. Wright. Parts I, II-A. GRAY 2	2, 42
zoölogical position of—E. D. Cope	412
Thalassophila—W. G. BINNEY	143
Thénard, L. J., memoir of, by M. FLOURENS. (R. 1862)	150
Theoretical researches—See Plateau, J.	
Theories, kinetic, of gravitation—W. B. TAYLOR. (B. 1876)	39 5
Theory—	
mechanical, of heat, principles of—J. MÜLLER. (R. 1868)	224
modern, of chemical types—C. M. WETHERILL. (R. 1868)	187
of heat, recent progress in relation to—A. CAZIN. (R. 1868)	224
Thermometer scales, table for conversion of centigrade to Fahrenheit. (R. 1863.)	187
Thermometer, wet and dry bulb, tables for—J. H. COFFIN	87
Thermometric observations	157
Thermometrical tables—A. GUYOT	153
THOMPSON, T. Mounds in Iowa and Illinois. (R. 1879)	345
Thoughts on the nature and origin of force—W. B. TAYLOR. (R. 1870) 244,	375
Thrushes, description of two new, from United States—R. RIDGWAY. (P. 1881.)	467
Thunder—	
and lightning, observations on—S. MASTERMAN. (R. 1855)	77
storm, on John Wise's observation of a-R. HARE. (R. 1854)	75
storms, instructions for observations of-J. HENRY	23 5
Tidal observations—	
at Wolstenholm Sound-Com. SAUNDERS	130
in the Arctic Seas—I. I. HAYES	196
in the Arctic Seas—E. K. KANE	130
Tide, flood, law of deposit of the—C. H. DAVIS	33
Tides and tidal action in harbors—J. E. HILGARD. (R. 1874) 286,	390
TIFFANY, A. S. Shell-bed skull from Illinois. (R. 1874)	236
Timber of submarine structures, occurrence of Chelura terebrans, a crustacesa destructive to—S. I. SMITH. (P. 1879)	333
Time and space, lecture on relations of—S. ALEXANDER. (R. 1861)	
Tinneh or Chepewyan Indians of British America—G. GIBBS; W. L. HARD- ISTY; S. JONES; B. R. Ross. (R. 1866)	36 5
Tipulide, monograph of-R. OSTEN SACKEN	
TOELLNER, A. Antiquities of Rock Island county, Illinois. (R. 1879)	
Tokio, University of, catalogue of Japanese woods presented by—L. F. WARD. (P. 1881)	

Tompkins county, New York, ancient fort and burial ground in—D. Trow- BRIDGE. (R. 1868)
TONER, J. M. Deed of foundation of Toner lectures. (R. 1872) 271, 82
Toner lectures—
No. 1. On the structure of cancerous tumors and the mode in which adjacent parts are invaded—J. J. Woodward
II. Dual character of the brain—C. K. Brown-Séquard 293
III. On strain and over-action of the heart—J. M. Da Costa 279
. IV. A study of the nature and mechanism of fever-H. C. Wood 282
v. On the surgical complications and sequels of the continued fevers, with a bibliography of works on diseases of the joints, bones, larynx, the eye, gangrene, hæmatoma, phlegmasia, etc.—W. W. Kekn
VI. Subcutaneous surgery, its principles, and its recent extension in practice—W. ADAMS
VII. The nature of reparatory inflammation in arteries after ligature, acupressure, and torsion—E. O. Shakespeare
VIII. Suggestions for the sanitary drainage of Washington city—G. E. Waring, Jr
Tonto Apaches, notes on the—C. SMART. (R. 1867) 216
Topography of Black Mountain, North Carolina-T. L. CLINGMAN. (R. 1855.) 7
Tornado in Spruce creek valley, Centre county, Pennsylvania—J. B. MEEK. (B. 1871) 249
Tornado near New Harmony, Indiana, April 30, 1852—J. CHAPPELSMITH 59
Tornadoes and tempests, distinction between—J. B. LAMARCK. (R. 1871) 249
Tornadoes, queries relative to—J. HENRY 190
Toronto—
Observatory, instructions by, for observations of aurora
Observatory, map of stars near North Pole used by 850
University. Objects of the museum. (R. 1865) 209
University. Scientific cooperation. (R. 1861) 149
Torrey, John, memoir of, by A. GRAY. (R. 1878) 278
Tobrey, J.—
Observations on the Batis maritima60
On the Darlingtonia Californica, a new pitcher plant from northern California
Plantæ Frémontianæ, or descriptions of plants collected by J. C. Frémont in California
Tortoises, North American land, of genus Xerobates—F. W. TRUE. (P. 1881.) 467
Tortola, great hurricane at—G. A. LATIMER. (R. 1867) 215
Tortugas, destruction of fish in vicinity of the—J. P. JEFFERSON; J. Y. Por- TER; T. Moore. (P. 1878)
Totten, Gen. J. G., eulogy on, by J. G. BARNARD. (R. 1865) 209
Traces of early mental condition of man—E. B. TYLOB. (R. 1867) 215

Trade—
ancient aboriginal—C. BAU. (R. 1872) 271, 440, 385
language of Oregon, dictionary of the—G. GIBBS
language of Oregon, vocabulary of the—B. R. MITCHELL; W. W. TURNER.
Transactions—See Geneva Society, Paris Anthropological Society.
Transatlantic longitude—B. A. Gould
Transfer of Smithsonian library to Library of Congress, act of Congress to authorize. (R. 1865) 209, 328, 328
Transformations, lecture on insect instincts and—J. G. Morris. (R. 1855) 7
Transit of Venus expedition to Kerguelen Island.—J. H. KIDDER and others. 293, 294
Transporting specimens of Diatomaceæ, directions for-A. M. EDWARDS 366
Transporting specimens of natural history, directions for—S. F. BAIRD. (R. 1856)
Trapezium, gray substance of the—J. DEAN
Travels in Central and South America—S. HABEL 269
Treasury Department, authority given by, to collectors of customs and others to receive and transmit specimens to the Smithsonian
Treasury, Secretary of, letter to, on payment of interest in coin—J. Herry. (R. 1865)
Treasury of United States, account of, with Smithson fund 328, 329
Trees—
forest, distribution of, in Montana, Idaho, and Washington—W. W. Johnson. (R. 1870)
of Florida and the Mexican boundary—J. G. Cooper. (R. 1860) 147
of North America, distribution of-J. G. Cooper. (R. 1858) 109, 351
Trinidad, economic geology of—G. P. Wall; J. G. Sawkins. (R. 1856) 91 Trochilida—
classification and synopsis of—D. G. ELLIOT
in collection of U. S. National Museum, catalogue of-R. RIDGWAY.
(P. 1880)425
list of—D. G. Elliot
Troglodytes of Breckinridge county, Kentucky—R. S. Robertson. (R. 1874.) 286
Troglodytes or cave dwellers of the valley of the Vézère—P. Broca. (R. 1872.) 271
TROWBRIDGE, D. Ancient fort and burial ground in Tompkins county, New York. (R. 1868)
TROYON, F.—
Archæological researches made at Concise. (R. 1861)
Lacustrian cities of Switzerland. (R. 1861)
On the Crania helvetica. (R. 1864)
Report on ethnological collection of Museum at Lausanne, Switzerland. (R. 1861)
TRUE, F. W. On North American land tortoises of genus Xerobates. (P. 1881.) 467
TRUE, F. W. On the rare rodent, Cricetodipus parvus. (P. 1881) 467

Tyrannus, synopsis of genus—R. RIDGWAY. (P. 1878)

U.

UHLER, P. R.—
. Circular in reference to the history of North American grasshoppers 16
Instructions for collecting hemipters. (R. 1858)
Instructions for collecting neuroptera. (R. 1858) 16
Instructions for collecting orthopters. (R. 1858) 10
Synopsis of neuroptera of North America
Underground temperature—C. A. SCHOTT; J. D. EVERETT. (R. 1874) 2
Union, lectures on the-H. REED. (R. 1854)
United States—
animal resources and fisheries of the, catalogue of collection to illustrate the—G. B. Goode
animal resources of the, catalogue of collection to illustrate the—G. B. Goode
antiquities of the, proposed map of—A. J. HILL. (R. 1861)
archæology of the—S. F. HAVEN
base chart of the—C. A. SCHOTT 41
chalk found in the-T. A. CONRAD. (R. 1865)
coleoptera of the, catalogue of the—F. E. MELSHEIMER
collections presented to the, by foreign commissioners, list of. (R. 1876.)
education in the, project of outline history of-F. A. PACKARD. (R. 1863.)
ethnological map of the, proposed—G. GIBBS. (B. 1862)1
exhibit of fisheries and fish culture of the, at Berlin, 1880-G. B. Goods. 41
explorations in the—See Baird, S. F.
fishes, food, of the—See Baird, S. F.
fishes of Pacific coast of the-D. S. JORDAN; C. H. GILBERT. (P. 1880.)
fishes of Pacific coast of the, bibliography of-T. H. Bran. (P. 1881.)
fishes of Pacific coast of the, bibliography of-T. GILL
fishes of Pacific coast of the, notes on-D. S. JORDAN; C. H. GILBERT.
(P. 1881) 4
free freight between England and the—E. CUNARD. (R. 1859) 11
free freight between Germany and the—Kunhardt & Co. (R. 1861) 19
free freight between Germany and the, by North German Lloyd-R. Schleiden. (R. 1858)
fund for promotion of science in the-J. TYNDALL. (R. 1872) 2
institutions, scientific and literary, in the, list of the principal 33
libraries, institutions, and societies in the, list of-W. J. RHEES 116, 23
libraries, public, in the, notices of—C. C. JEWETT 2
libraries, public, of the, report on—C. C. JEWETT. (R. 1849)
limpets and chitons from deep waters off eastern coast of the—W. H. DALL. (P. 1881)

United States—Continued.
meteorology of the, report on—E. Loomis. (R. 1847)
microscopic examinations of soundings off Atlantic coast of the—J. W. BAILEY
mollusca of the southern, some new species of eocene—A. Heilprin. (P. 1880)425
occultations visible in the—See Downes, J.
physical geography of the—C. ELLET, Jr18
rain and snow fall in the, tables of—C. A. SCHOTT 222, 353
rain charts of the, 1, 2, 8: summer, winter, year—C. A. Schott 374
reptiles, cretaceous, of the-J. LEIDY
reptiles, cretaceous, of the, review of-J. LEIDY. (R. 1864) 188
shell-deposits, artificial, of the-D. G. BRINTON. (R. 1866) 214
sparrows sent to the—W. A. LLOYD. (R. 1867)
temperature chart of the, for the year—C. A. SCHOTT 381, 388
temperature charts of the, 1, 2, 8: summer, winter, year—C. A. Schott. 387
temperature tables of the—C. A. SCHOTT
thrushes from the, description of two new-R. RIDGWAY. (P. 1881) 467
trees of the, catalogue of—J. G. Cooper. (R. 1858) 109, 851
United States Exploring Expedition, duplicate shells collected by the 193
United States Fish Commission—See Fish Commission.
United States Government, list of publications of, 1868-1881-G. H. BOEHMER. 477
United States National Museum—See Museum, U. S. National.
United States Patent Office, results of meteorological observations under the
direction of the. Vols. I, II 157, 182
United States Treasury—See Treasury.
Universal meteorograph for detached observatories—E. H. Von BAUMHAUER. (R. 1879)
University—
National, of Greece, Museum of. Exchange of specimens. (R. 1867.) 215
of Christiania, Norway, ethnological specimens presented by. (R. 1868.) 187
of Costa Rica. Exchange of specimens. (R. 1867) 215
of State of New York, Regents of the. Acknowledgment for specimens. (R. 1865)
of Tokio, catalogue of Japanese woods presented by—L. F. WARD. (P. 1881)467
of Toronto, objects of museum of. (R. 1865) 209
of Toronto, scientific cooperation. (R. 1861) 149
Uranidea marginata, new species of fish, description of-T. H. BEAN. (P. 1881.) 467
Uranidea microstoma, new fish from Alaska, description of—W. N. Locking- Ton. (P. 1880)425
Uranus, investigation of orbit of, with tables of its motion—S. NEWCOMB 262
Uranus, secular variation of elements of orbit of-J. N. STOCKWELL 282

Urine, observations on the—J. Jones	82
Use of galvanometer as a measuring instrument—J. C. Poggendorff. (R. 1859.)	110
Uses of brain and marrow of animals among Indians—T. R. PEALE. (R. 1870.)	244
Utah, descriptions of new invertebrate fossils from—C. A. WHITE. (P. 1880.)	425
Utah lake, description of new species of Squalius (Squalius aliciæ) from—P. L. Joux. (P. 1881)	467
Utah lake, notes on collection of fishes from—D. S. JORDAN; C. H. GILBERT. (P. 1880)	425
Utrecht, Provincial Society of Arts and Sciences. Prize questions. (R. 1861; R. 1862)	, 150

V.

VAILLANT, M.; HENRY, J. Horary variations of the barometer. (R. 1866) 214
Values of $b^{(i)}$ and its derivatives—J. D. Runkle
Values of coefficients in perturbative function of planetary motion depending on ratio of mean distances—J. D. Runkle
Valvatidæ—W. G. BINNEY 144
Van Rensselaer harbor, Arctic Seas, observations at—See Kane.
Vapor, aqueous, on the diminution of, with increasing altitude in the atmosphere—J. Hann. (R. 1877)
Vapor, aqueous, table for determining elastic force of—J. H. Corrin 87
Vastness of the visible creation, lectures on—S. ALEXANDER. (R. 1857) 107
Vegetable colonization of Shetland Islands, Faroe Islands, and Iceland—C. MARTINS. (R. 1858)
Vegetable species, causes which limit, towards the north—A. DE CANDOLLE. (R. 1858)
Vegetation and the atmosphere—J. JAMIN. (R. 1864)188
Velie, J.W., catalogue and description of fishes collected by, in Gulf of Mexico-G. B. Goode; T. H. Bean. (P. 1879) 333
Velocity of light, essay on the—C. DELAUNAY. (R. 1864) 188, 354
Velocity of the wind—J. Hann and others. (R. 1877) 823, 398
Venezuela, Colonia Tovar, meteorology and ethnology of—A. FENDLEE. (R. 1857; R. 1866)
Venom of rattlesnake, researches on the—S. W. MITCHELL135
Venom of serpents, bibliography of—S. W. MITCHELL
Ventilating and warming buildings. Parts I, II—A. MORIN. (R. 1873; R. 1874) 275, 286, 439
Ventilation, architecture in relation to—D. B. REID. (R. 1856)91
Venus, secular variations of elements of orbit of—J. N. STOCKWELL 222

Venus—See Transit of Venus.	
Vera Cruz, Mexico, antiquities in-H. Finck. (R. 1870)	244
Verification of barometers at the Kew Observatory—J. WELSH. (R. 1859)	110
VERRILL, A. E.—	
Annelids, echinoderms, and anthozoa of Kerguelen Island	294
Annelids, molluscoids, and radiates of Arctic America	842
Marine invertebrata of New England. (P. 1879)	838
Mollusca recently added to fauna of southern New England, catalogue of. (P. 1880)	425
Mollusca, with notes on annelida, echinodermata, etc. (P. 1880)	425
Recent additions to marine invertebrata of northeast coast of America, with new genera and species. Parts 1, 11, 111. (P. 1879; P. 1880.) 888	, 425
VERRILL, A. E.; RATHBUN, R. Marine invertebrata of northeast coast of America distributed by Fish Commission. (P. 1879)	833
Vertebrata, chemical and physiological investigations relative to-J. Jones	82
Vespidæ, synopsis of-H. Dr Saussure	254
Vestiges of antiquity in the United States—S. F. HAVEN	71
Vézère, valley of the, cave-dwellers of the-P. Broca. (R. 1872)	271
Vibratory movement of matter-L. MAGRINI. (R. 1868)	224
Victoria Royal Society, address of the president—R. L. J. ELLERY. (R. 1868.)	224
Vienna— Imperial Academy of Sciences of. Prize questions. (R. 1864; R. 1865.) 188 Imperial Geological Institute, tertiary fossils presented by. (R. 1863) Imperial Library of, books presented. (R. 1865)	, 209 187 209
Views and plans of Smithsonian Institution—R. D. Owen	P
Vineyard Sound, Massachusetts, occurrence of oceanic bonito in—V. N. EDWARDS. (P. 1878)	832
Virgin Islands, flora of the-H. F. A. EGGERS	818
Virginia—	
Hampton, catalogue of casts of heads of Indian boys and girls at—R. H. PRATT. (P. 1879)	333
Luray Cave, blind myriapod from—J. A. RYDER. (P. 1880)	425
Luray Cave, report of visit to-O. T. Mason and others. (R. 1880.) 442,	433
Vitality-H. H. HIGGINS; J. HENRY. (R. 1866)	214
Viviparide-W. G. BINNEY	144
Vocabularies, Indian—G. GIBBS. (R. 1862)	150
Vocabularies, Indian, received from the Wheeler expedition, list of. (R. 1874.)	286
Vocabulary—	
comparative—A. GALLATIN	160
comparative, English, Spanish, French, Latin	170
Makah-J. G. SWAN	220
of Chinook jargon, or trade language of Oregon-G. GIBBS	161
of Chinook jargon, or trade language of Oregon—B. R. MITCHELL; W. W. TURNER; G. GIBBS	68

Vocal sounds of Laura Bridgman—F. LIEBER	12
Voices of crustaceans—G. B. Goode. (P. 1878)	832
Volcano-	
in Nicaragua, eruption of—A. B. DICKINSON. (R. 1867)	215
of Colima, eruption of—C. SARTORIUS. (R. 1869)	228
of Popocatepetl, examination of—Baron von Müller; A. Sonntag	114
Vollum, E. P. On the wingless grasshopper of California. (R. 1860)	147
Volta, Alexander, eulogy on, by F. ARAGO. (R. 1875)	298
Von Baumhauer, E. H. Universal meteorograph for detached observatories. (R. 1879)	
Von Buch, Leopold, memoir of, by M. FLOURENS. (R. 1862)	
Von Hellwald, F. The American migration. (R. 1866)	214
VON KAROLYI, Lieut.; CRAIG, B. F. Products of combustion of gun-cotton	i
and gunpowder. (R. 1864)	168
VON LIEBIG, J. Induction and deduction. (R. 1870)	. 244
Von Martius, C. F. P., honorary medal to-W. HAIDINGER. (R. 1868)	. 187
Von Martius, C. F. P., memoir of, by C. RAU. (R. 1869) 228, 440	, 251
Von Müller, Baron. Observations on terrestrial magnetism in Mexico	. 114

W.

WAITE, M. R. Memorial of Joseph Henry	256
WALE; HAWKINS. Blow-pipe apparatus. (R. 1872)	
WALKER, D. Observations on Arctic auroras and specific gravity of sea water.	146
WALKER, D. Observations on ice	146
WALKER, S. C. Ephemerides of planet Neptune, 1848, 1846-1849, 1850, 1851, 1852	24
WALKER, S. C. Researches relative to planet Neptune	3
WALKER, S. T. Explorations among Indian mounds in southern Florids. (R. 1879)	345
WALKER, S. T. Shell-heaps of Tampa Bay, Florida. (R. 1879)	345
WALL, G. P.; SAWKINS, J. G. Report on survey of economic geology of Trinidad. (R. 1856)	91
WALLACH, R.; HENRY, J. Report of Committee of Regents on the fire at Smithsonian. (R. 1864)	188
Walrus, remains of, in Maine—C. H. Boyd. (P. 1881)	467
	345
War Department, authority given by, to officers of Quartermaster's Department to receive and transmit specimens to Smithsonian	34

WARD, L. F.—	
Barometer tube breaking suddenly. (R. 1866)	214
Catalogue of collection of Japanese woods presented by University of Tokio, Japan. (P. 1881)	467
Check-list of plants of Washington and vicinity	
Directions for collecting and preserving plants	
Guide to flora of Washington and vicinity	
WARING, G. E., Jr. Suggestions for sanitary drainage of Washington city. Toner lecture No. VIII	849
Warming and ventilating buildings. Parts I, II—A. MORIN. (R. 1878; R. 1874) 275, 286,	439
Warming, architecture in relation to—D. B. REID. (R. 1856)	91
WARNER, J. Big elephant mound in Grant county, Wisconsin. (R. 1872)	•
Warren, G. K., report on collections made by-F. B. MEEK; F. V. HAYDEN.	
WARTMANN, E. Report on transactions of the Society of Physics and Natural	224
Washington, Arkansas, meteorological observations near—N. D. SMITH	
Washington, D. C.—	
canal of, report of Committee of Regents on—R. Delafield. (R. 1868.)	224
flora of, and vicinity, guide to the—L. F. WARD	•
notice of, in Harriot's travels	880
notice of, in Isaac Weld's travels	880
prehistoric remains in vicinity of—T. R. PEALE. (R. 1872)	271
washington Philosophical Society, bulletin of. Vols. I-III; March, 1871, to June, 1880	349 428
Washington Territory—	120
fishes of—T. H. BEAN. (P. 1881)	467
forest trees in, distribution of—W. W. Johnson. (R. 1870)	244
Indians of—J. G. Swan	220
method of taking surf smelt by Indians of—J. G. SWAN. (P. 1880) Neah Bay, description of two new species of fishes (Ascelicthys rhodorus and Scytalina cerdale) from—D. S. Jobdan; C. H. Gilbert. (P. 1880)	42 5
Wasps, American solitary, synopsis of—H. DE SAUSSURE	254
Wasps, translation of De Saussure's monograph of-E. Norton-	
Water—	
formation of ice at bottom of-M. Engelhardt. (R. 1866)	214
from Gulf of Mexico, report on-W. G. FARLOW. (P. 1881)	467
in rivers, diminution of—H. G. WEX. (R. 1875)	298
in rivers, method of ascertaining the amount of—A. A. HUMPHREYS. (R. 1858)	109
polluted, in Gulf of Mexico, destruction of fish by—W. C. W. GLAZIER. (P. 1881)	467

Water—Continued.
salt, preservation of copper and iron in—A. E. BECQUEREL. (R. 1864.)
sea, specific gravity of—D. WALKER14
WATKINS, E. A. System of relationship of Cree Indians. (R. 1862) 15
WATSON, S. Bibliographical index to North American botany. Part 1. Polypetalæ
Weapons, savage, at Centennial Exhibition, study of—E. H. KNIGHT. (R. 1879) 345, 41
Weather, effect of moon on the-J. HENRY. (R. 1871) 24
WEBB, T. W. Account of Henry Draper's telescope. (R. 1864)
Weights, atomic, recalculation of. Constants of Nature, Part v-F. W. CLARKE. 4
Weights, small, method of forming—R. HARE. (R. 1858)
Weights and measures—
English and French, tables of. (R. 1868; R. 1864) 187, 18
metric system of, with tables—H. A. NEWTON. (R. 1865) 209, 37
new system of, with 8 as the metrical number—G. H. KNIGHT. (B. 1867.) 21
WEISMANN, A. Change of Mexican axolotl to amblystoma. (R. 1877) 323, 403
WELD, I. Account of city of Washington, 1807 33
Welling, J. C. Notes on the life and character of Joseph Henry 856, 33
Welsh, language of Navajos said to resemble—S. Y. McMasters. (R. 1865.) 20
Welsh, J. Construction and verification of standard barometers. (R. 1859.) 110
West Indies—
birds of the—See Lawrence, G. N., Ober, F. A.
list of described birds of the, not in Smithsonian collection 185
loxigilla from, description of new subspecies of—G. N. LAWRENCE. (P. 1881)
stone celts in the-G. J. GIBBS. (R. 1877)
West Virginia, occurrence of Canada porcupine in-G. B. Goode. (P. 1878.) 332
Wet and dry bulb thermometer, tables for-J. H. Coffin 87
WETHERILL, C. M.—
Modern theory of chemical types. (R. 1863) 187
Ozone and antozone. (R. 1864) 188, 355
Plan of a research upon the atmosphere. (R. 1866)214
WEX, H. G. Diminution of the water of rivers and streams. (R. 1875) 238
Wheeler expedition, list of Indian vocabularies received from the. (R. 1874.) 286
Wheeler expedition—See Explorations.
Whirlwind at St. Thomas, August 21, 1871—A. Colding. (R. 1877) 823, 398
Wніте, С. А.—
Description of new cretaceous pinna from New Mexico. (P. 1880) 425
Description of very large fossil gasteropod from State of Puebla, Mexico.
(P. 1880) 425
Descriptions of new cretaceous invertebrate fossils from Kansas and Texas. (P. 1879)

WHITE, C. A.—Continued.
Descriptions of new invertebrate fossils from the mesozoic and cenozoic
rocks of Arkansas, Wyoming, Colorado, and Utah. (P. 1880) 49
Descriptions of new species of carboniferous invertebrate fossils. (P. 1879.)
Note on acrothele. (P. 1880) 42
Note on Criocardium and Ethmocardium. (P. 1879)
Note on Endothyra ornata. (P. 1879) 38
Note on occurrence of Productus giganteus in California. (P. 1880) 42
Note on occurrence of Stricklandia salteri and Stricklandia Davidsoni in Georgia. (P. 1880) 42
On certain cretaceous fossils from Arkansas and Colorado. (P. 1881) 46
White river canon, Pima county, Arizona, ruins in-R. T. Burr. (R. 1879.) 84
WHITNEY, J. D. Catalogue of rocks, minerals, etc. (R. 1854)
WHITNEY, J. D. Cave in Calaveras county, California. (R. 1867) 21
WHITNEY, W. D. Lectures on principles of linguistic science. (R. 1863.) 187, 35
WHITTLESEY, C
Ancient mining on shores of Lake Superior 15
Descriptions of ancient works in Ohio
On fluctuations of level in North American lakes 11
On fresh-water glacial drift of the northwestern States 19
Wilkes, C., duplicate shells of exploring expedition under 19
Will of Smithson. (R. 1853)
Willemœsia group of crustacea, (recent Eryontidæ,) notice of new species of the—S. I. Smith. (P. 1879)
WILLIAMS, H. C. On antiquities in some of the southern States. (R. 1870.) 24
WILLIAMSON, G. Antiquities in Guatemala. (R. 1876) 29
WILLIS, J. R.; BLACKISTON; BLAND. Birds of Nova Scotis. (R. 1858) 10
WILLIS, J. R.; BLAND. Birds of Bermuda. (R. 1858) 10
WILMOT, S. Notes on western gizzard shad, Dorosoma cepedianum heterurum. (P. 1878)
Wilson, D. Lectures on physical ethnology. (R. 1862) 15
Wilson, Henry, eulogy on, by P. PARKER. (R. 1875) 29
Wind-
and fog-J. Balfour. (R. 1866) 21
articles on, preserved in Smithsonian. (R. 1874) 28
changes of—C. Mallinikroot. (R. 1866) 21
connection of gales of, and appearance of aurora—R. T. KNIGHT; J. HENRY. (R. 1871)24
observations of, in the Arctic Seas-I. I. HAYES19
observations of, in the Arctic Seas—E. K. KANE
observations of, in the Arctic Seas-F. L. McCLINTOCK 14
observations of, in the United States 15
relation between difference of pressure and velocity of -J. HANN. (R.
1877) 823, 89

Winds—	
in Florida-J. BALTZELL. (R. 1866)	214
of the globe—J. H. Coffin	268
of the northern hemisphere—J. H. Coffin	52
upper, importance of attention to the—C. DEWEY. (R. 1866)	214
Winnipeg Lake, egging expedition to—D. Gunn. (R. 1867)	
Wirtz, H. R. Herbarium captured in Tennessee. (R. 1862)	
Wisconsin—	
antiquities in—M. Strong. (R. 1877)	323
antiquities of—I. LAPHAM	70
earthworks in—E. E. BREED. (R. 1872)	271
Grant county, elephant mound in-J. WARNER. (R. 1872)	271
Grant county, observations on prehistoric mounds of—M. STRONG. (R. 1876)	299
mound in—C. K. DEAN. (R. 1872)	271
mounds and osteology of mound-builders of-J. N. DE HART. (R. 1877.)	328
mounds in-W. G. Anderson. (R. 1879)	34 5
salamander of-P. R. Hov. (R. 1854)	75
Winnebago county, mounds in—T. Armstrong. (R. 1879)	345
Wise, J., on observation by, of a thunder storm—R. HARE. (R. 1854)	.75
WISEMAN, N. Identification of the artisan and artist. (R. 1870)	244
WISSNER, J. Meteorology of the District of Columbia. (R. 1857)	107
WITHERS, R. E. Memorial address on Joseph Henry	856
Workoff, A. Discussion of winds of the globe	268
Workoff, A. Meteorology in Russia. (R. 1872)	271
Wolstenholm Sound, tidal record at—Com. SAUNDERS	180
Wood, ancient implement of, from Connecticut—E. W. Elisworth. (R. 1876)	299
Wood, preservation of. (R. 1864)	188
Wood, H. C.—	
Bibliography of alge	24 1
Contribution to the history of the fresh-water alge of North America	241
Fever. A study in morbid and normal physiology. (R. 1878) 341,	357
Instructions for collecting myriapods, phalangids, etc. (R. 1866)	214
Study of nature and mechanism of fever. Toner lecture No. IV	282
Wood, J. Meteorological observations at Marietta, Ohio	120
Woods of Japan, collection of, presented to U.S. National Museum, catalogue of—L. F. WARD. (P. 1881)	467
WOODWARD, J. J. Structure of cancerous tumors and mode in which adjacent	
parts are invaded	
WOODWORTH, A. Meteorite in Mexico. (R. 1867)	
(400 2000) and a contract (400 2000) and a contract (400 2000)	149
Work, relation of food to, and its bearing on medical practice—S. HAUGHTON. (R. 1870)	244

OF SMITHSONIAN PUBLICATIONS.	327
Works published by Smithsonian Institution to January, 1866	208
World, estimate of population of the—E. MAILLY. (R. 1873)	275
Worsaar, J. J. A. Preservation of antiquities and national monuments in Denmark. (R. 1879)	845
Wounds of poisoned arrows, lecture on nature and cure of—D. Brainard. (R. 1854)	75
Wright, C., account of botanical explorations of, in New Mexico and California—A. Gray. (R. 1849)	21
Wright, C., account of plants collected by, in Texas and New Mexico. Parts 1, 11—A. GRAY	2, 42
WRIGHT, D. F. Antiquities of Tennessee. (R. 1874)	286
WRIGHT, G. Account of lightning discharges. (R. 1867)	215
Writings, scientific, of Sir William Herschel, synopsis of—E. S. Holden; C. S. Hastings. (R. 1880) 442,	426
Writings, scientific, of James Smithson	827
Würdemann, G., notice of, by A. D. BACHE. (R. 1859)	110
WÜRDEMANN, G. Specimens of flamingo, etc., from south Florida. (R. 1860.)	147
WÜRDEMANN, W.; GREEN, J. On filling barometer tubes. (R. 1859)	110
Wyandot government—J. W. Powell. (E. 1879-80)	476
WYMAN, J. Anatomy of nervous system of Rana pipiens	45
Wyoming, new invertebrate fossils from—C. A. WHITE. (P. 1880)	425
• • • • • • • • • • • • • • • • • • •	
X.	
Xantus, John, explorations of, in Mexico—M. Romero. (R. 1862) Xenichthys ocyurus, new species of Xenichthys from west coast of Central	
America, description of—D. S. JORDAN; C. H. GILBERT. (P. 1881)	
Xerobates, land tortoises of genus—F. W. TRUE. (P. 1881)	
Xiphister, description of new species of, from Monterey, California—D. S. Jor- DAN; C. H. GILBERT. (P. 1880)	
Xystreurys liolepis, new flounder from Santa Catalina Island, California, description of—D. S. Jordan; C. H. Gilbert. (P. 1880)	

Y.

Yacht "Fox," meteorological observations made on-F. L. McCLINTOCK 14
Yale College, catalogue of meteorites in—G. J. BRUSH. (R. 1868) 224
YARROW, H. C. Study of mortuary customs of the North American Indians. (E. 1879-80)
Yellowstone river, prehistoric remains near the—P. W. Norris. (R. 1879) 345
Yoruba, Africa—
country and people of, description of—T. J. Bowen
language of, bibliography of98
language of, grammar and dictionary of-T. J. Bowkn; W. W. TURNER. 98
language of, grammar and dictionary of—Southern Baptist Mission-
YOUNG, C. B. Translation of Morin on warming and ventilating buildings. (R. 1878; R. 1874) 275, 286, 439
Young, C. B. Translation of Von Baumhauer on a universal meteorograph. (R. 1879)
Young, Thomas, eulogy on, by F. ARAGO. (R. 1869) 228
Young, W. J. Cloud bursts. (R. 1867) 215
Yu, (Chinese name of jade,) study of—S. BLONDEL. (R. 1876) 299
Yucatan, remarks on Cara gigantesca of Yzamal in-A. Schott. (R. 1869) 228
Yukon, Russian America, journey to the-W. W. KIRBY. (R. 1864) 188
Yzamal, Yucatan, remarks on Cara gigantesca of—A. Schott. (R. 1869) 228

Z.

ZANTEDESCHI, F. R. Electricity of induction in aerial strata of atmosphere surrounding cloud resolving into rain, etc. (R. 1870)	244
Zapotec remains, notice of—B. MAYER; J. G. SAWKINS	86
ZISGENBALS, H. Schlagintweit ethnographic collection. (R. 1867)	215
Zone of small planets between Mars and Jupiter, lecture on the—E. Loomis. (R. 1854)	75
Zoölogical—	
Gardens, Hamburg. Exchange of specimens. (R. 1867)	215
position of Texas—E. D. Cope	412
Record, list of generic names supplemental to those indexed in—S. H. SCUDDER	470
Zoölogicus, Nomenclator. A list of generic names of recent and fossil ani- mals—S. H. Scudder	470
Zoölogy, list of generic names employed in-S. H. Scudder	470
Zoölogy, record of recent progress in, 1879, 1880—T. GILL. (R. 1880) 442,	431



•				
		•	•	
	•			
			•	
•				



		-	
	• •	•	
	•		
		_	
		•	
		_	•
		•	
		•	•
		•	
		•	
		•	
			•
	•	•	
_	•		
•			
		•	
			•
		•	
		-	



